

# The contribution of blowing snow to cloud properties and the atmospheric radiative budget over Antarctica

Stefan Hofer,<sup>1\*</sup> Charles Amory,<sup>2</sup> Christoph Kittel,<sup>2</sup> and Trude Storelvmo<sup>1</sup>

<sup>1</sup>Department of Geosciences, University of Oslo, Oslo, Norway

<sup>2</sup>Laboratory of Climatology, Department of Geography, University of Liège, Belgium

\*Corresponding author: Stefan Hofer, stefan.hofer@geo.uio.no

Antarctica and its surroundings are one of the main areas for biases in climate models, mostly due to uncertainties in the representation of clouds. Over the Antarctic Ice Sheet, temperature inversions and strong temperature gradients between the cold interior and the edges lead to strong katabatic downslope winds, transporting snow and moisture from the interior towards the peripheral regions at the southern edge of the Southern Ocean's storm track. These blowing snow layers are usually 100-200 m thick, but can reach a thickness of more than 500 m and can be advected offshore from Antarctica over open ocean waters (Scarchili et al., 2010; Palm et al., 2017). However, the impacts of moisture and wind-induced snow mass transport (i.e. ice nucleating particles) on cloud structure and development over Antarctica has not been thoroughly investigated and most state-of-the-art climate models do not account for its presence. Here, we use a regional climate model with a newly developed fully active blowing snow scheme and satellite data, to show that accounting for drifting snow notably alters the spatial distribution, vertical structure and radiative contribution of clouds over Antarctica and its periphery. Additionally, our results indicate that the advection of blowing snow and air with a higher humidity content over the Southern Ocean also impacts clouds and their microphysics in areas outside of Antarctica. While our study area is limited to 60S, our results highlight the need to study the impact of missing blowing snow processes on the future evolution of clouds not just over Antarctica, but potentially also over the Southern Ocean, an area with significant uncertainties in future climate projections.

## Introduction

**First paragraph:** Main facts about Antarctica and clouds there Radiative effects, influence on surface temperature (SEB). a bit about melt (even in winter; see Kuipers-Munneke 2014).

Antarctica is the largest body of ice on the planet and it contains the coldest places on Earth. It contains ice mass equivalent to roughly **XX m** of global sea level rise, roughly 10 times more than the Greenland Ice Sheet (**cite XX**). The interior plateau of the Antarctica reaches up to elevations of **XX m**, and is marked by very cold and dry conditions (**cite XX**). However, there are marked differences in the present climatic conditions between East and West Antarctica, and especially between the edges and the inner plateau. While West Antarctica, and especially the northwards reaching mountainous Antarctic peninsula receive ample precipitation and are rather cloudy, weather systems never reach the interior of Antarctica, where most of the precipitation falls as diamond dust under extremely cold "clear" sky conditions (**cite XX**).

Due to strong radiative cooling in the interior plateau, strong and perpetual katabatic winds emerge, redistributing snow mass from the interior towards the edges and ice shelves, where the up to **XX m** high plateau slopes steeply towards sea level (**CITE XX**).

**Second paragraph:** describing BS as near-surface clouds + source of moisture and condensation nuclei for additional cloud formation in the lower atmosphere

**Third paragraph:** What are the open questions? i) Does BS lead to improved representation of cloud properties in the model and ii) how does it affect the representation of clouds?

Clouds are known to notably affect the present and future climates of polar ice sheets [**Izeboud2020**, **Hahn2019**, 1, 2, 3]. Clouds have the ability to amend incoming shortwave and longwave fluxes, depending on the cloud phase, height and particle size distribution, impacting the rate of surface melt and snowpack warming. Blowing snow, while not accounted for in most global and regional climate models, can change the vertical structure and radiative impact of clouds, most notably because blowing snow sublimation changes the atmospheric humidity and temperature distribution (**cite Louis 2020**). Blowing snow particles can also act as ice nucleating particles

for cloud formation, which also impacts the longevity, structure and cloud-phase distribution within pre-existing clouds. Additionally, optically thick blowing snow layers can act as a cloud themselves, increasing the atmospheric longwave emissivity and shortwave transparency of the atmosphere. However, so far very little is known about how clouds are influenced by blowing snow processes in climate models, and how accounting for blowing snow over the current climate influences key polar cloud-, and therefore climate processes.

**Fourth paragraph:** How are we planning to address the questions? Two sets of simulations + satellite products

Here, we use two regional climate model simulations spanning the period of 1979-2019, one with a dynamic representation of blowing snow and one without, to assess the impact of accounting for blowing snow on Antarctic clouds and radiative fluxes. We compare our two simulations to satellite products of cloud cover and the ERA5 reanalysis product, to show whether accounting for drifting snow only amends or also improves the representation of polar clouds. However, due to the remote location and complications of detecting cloud structure and micro-physics from satellites over highly reflective surfaces, we don't expect to comprehensively address whether blowing snow improves cloud representation over Antarctica. Nevertheless, our results deliver a clear indication that accounting for blowing snow over polar ice sheets changes the 3D-structure of clouds, their phase and ultimately their contribution to the surface energy budget. In conclusion, not accounting for drifting snow in future projections of the Antarctica climate and sea level rise contribution might significantly bias the drawn conclusions.

## Results

**Potential subtitle here**




Figure 1: **Difference in temperature and cloud properties between MAR with and without blowing snow.** A) Cross-section of temperature differences between MAR with blowing snow turned on, and MAR without blowing snow (positive means MARb is warmer), along the path shown in **MISSING FIGURE XX**. B) Same as panel A), but showing the difference in cloud cover (in %) between the two simulations. C) Same as panel A) and B), but for the difference in the cloud radiative effect ( $Wm^{-2}$ ).




Figure 2: **Difference in radiative components at the surface between MAR with and without blowing snow.** A) Difference in incoming shortwave radiation (SWD) at the surface in  $Wm^{-2}$ . Red color indicates a greater downwelling shortwave flux in MAR with active blowing snow parameterisation. B) Same A) but for the downwelling longwave flux at the surface. C) Same as A) and B), but for the difference in the net radiation at the surface ( $R = SWD * (1 - \alpha) + LWD - LWU$ ).



## Main References

- [1] Stefan Hofer et al. “Decreasing cloud cover drives the recent mass loss on the Greenland Ice Sheet”. In: *Science Advances* 3.6 (2017). DOI: 10.1126/sciadv.1700584.
- [2] Stefan Hofer et al. “Cloud microphysics and circulation anomalies control differences in future Greenland melt”. In: *Nature Climate Change* 9.7 (2019), pp. 523–528. DOI: 10.1038/s41558-019-0507-8.
- [3] K. Van Tricht et al. “Clouds enhance Greenland ice sheet meltwater runoff”. In: *Nature Communications* (2016). DOI: 10.1038/ncomms10266.
- [4] X. Fettweis. “Reconstruction of the 1979-2006 Greenland ice sheet surface mass balance using the regional climate model MAR”. In: *The Cryosphere* 1.1 (Oct. 2007), pp. 21–40. DOI: 10.5194/tc-1-21-2007.
- [5] Xavier Fettweis et al. “Estimating the Greenland ice sheet surface mass balance contribution to future sea level rise using the regional atmospheric climate model MAR”. In: *The Cryosphere* 7.2 (2013), pp. 469–489. DOI: 10.5194/tc-7-469-2013.
- [6] Xavier Fettweis et al. “Reconstructions of the 1900–2015 Greenland ice sheet surface mass balance using the regional climate MAR model”. In: *The Cryosphere* (2017). DOI: 10.5194/tc-11-1015-2017.
- [7] Hubert Gallée and Guy Schayes. “Development of a Three-Dimensional Meso- $\gamma$  Primitive Equation Model: Katabatic Winds Simulation in the Area of Terra Nova Bay, Antarctica”. In: *Monthly Weather Review* 122.4 (1994), pp. 671–685. DOI: 10.1175/1520-0493(1994)122<0671:DOATDM>2.0.CO;2.
- [8] Hubert Gallée and Hubert Gallée. “Simulation of the Mesocyclonic Activity in the Ross Sea, Antarctica”. In: *Monthly Weather Review* (1995).
- [9] Christoph Kittel et al. “Sensitivity of the current Antarctic surface mass balance to sea surface conditions using MAR”. In: *Cryosphere* 12.12 (2018), pp. 3827–3839. DOI: 10.5194/tc-12-3827-2018.
- [10] Alison Delhasse et al. “Brief communication: Impact of the recent atmospheric circulation change in summer on the future surface mass balance of the Greenland Ice Sheet”. In: *The Cryosphere* 12.11 (Oct. 2018), pp. 3409–3418. DOI: 10.5194/tc-12-3409-2018.
- [11] C. Lang, X. Fettweis, and M. Erpicum. “Stable climate and surface mass balance in Svalbard over 1979-2013 despite the Arctic warming”. In: *The Cryosphere* 9.1 (Jan. 2015), pp. 83–101. DOI: 10.5194/tc-9-83-2015.
- [12] Cécile Agosta et al. “Estimation of the Antarctic surface mass balance using the regional climate model MAR (1979-2015) and identification of dominant processes”. In: *Cryosphere* 13.1 (2019), pp. 281–296. DOI: 10.5194/tc-13-281-2019.



- [13] A. Delhasse et al. “Brief communication: Evaluation of the near-surface climate in ERA5 over the Greenland Ice Sheet”. In: *The Cryosphere* 14.3 (2020), pp. 957–965. DOI: 10.5194/tc-14-957-2020.
- [14] Hubert Gallée, Gilbert Guyomarc’h, and Eric Brun. “Impact Of Snow Drift On The Antarctic Ice Sheet Surface Mass Balance: Possible Sensitivity To Snow-Surface Properties”. In: *Boundary-Layer Meteorology* 99.1 (Apr. 2001), pp. 1–19. DOI: 10.1023/A:1018776422809.
- [15] V. Vionnet et al. “The detailed snowpack scheme Crocus and its implementation in SURFEX v7.2”. In: *Geoscientific Model Development* 5.3 (May 2012), pp. 773–791. DOI: 10.5194/gmd-5-773-2012.
- [16] IPCC. “Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]” In: *IPCC* (2014). DOI: 10.1017/CBO9781107415324. arXiv: arXiv:1011.1669v3.
- [17] M. Tedesco and X. Fettweis. “Unprecedented atmospheric conditions (1948–2019) drive the 2019 exceptional melting season over the Greenland ice sheet”. In: *The Cryosphere* 14.4 (2020), pp. 1209–1223. DOI: 10.5194/tc-14-1209-2020.

## Materials and Methods

### MAR

### MODIS

### AVHRR

### ERA5

### Modèle Atmosphérique Régional (MAR)

For the downscaling of coarse-resolution CMIP5 and CMIP6 data we used the Modèle Atmosphérique Régional (MAR), an open-source and widely used polar regional climate model [1,

2, 4, 5, 6, 7, 8, 9, 10, 11, 12]. MAR consists of a hydrostatic dynamical core which solves the primitive equation set [7, 8]. A full description of the model setup, the underlying physical parameterizations and evaluation of MAR for polar climates are described in [1, 2, 4, 6, 7, 8, 9, 12]. In this study we used the MARv3.9.6 version, evaluated in A. Delhasse et al. [13], and the source code of MAR for the reproduction of this study is available via the MAR homepage at <http://mar.cnrs.fr>.

Within MAR, the snow and ice properties at the ice sheet-atmosphere interface are calculated in the Soil Ice Vegetation Atmosphere Transfer module (SISVAT) [7]. This module calculates the main snowpack based on the snow module CROCUS [14, 15], but also handles the mass and energy exchange between the atmosphere (e.g. radiation, precipitation, temperature) and the bare-ice surfaces, the snowpack and the Arctic tundra that surrounds the GrIS [4, 7].

For the 6 CMIP5 and 5 CMIP6 future projections we downscaled we prescribed the boundary conditions in exactly the same manner and also used the MAR version and setup throughout. Overall, MAR was forced at its lateral boundaries (pressure, wind speed, temperature, specific humidity), at the top of the stratosphere (temperature, wind speed) and at the ocean surface (sea ice concentration, sea surface temperature) every 6 hours using GCM and ERA-Interim reanalysis fields [4, 5, 9, 12]. We ran MAR at a spatial resolution of 15 km x 15 km on a polar stereographic projection, which represents a significant increase in resolution compared to previous GrIS regional climate projections with MAR in Xavier Fettweis et al. [5], and which was used in the IPCC AR5 [16]. The MAR setup used in this study has been thoroughly compared to observations from weather stations, observed radiative fluxes, satellite cloud cover, satellite albedo and melt extent, ablation and SMB in-situ measurements [1, 4, 6, 13, 17].

## **Acknowledgments**

This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant agreement No. 758005). Computational resources have been provided by the Consortium des Equipements de Calcul Intensif (CECI), funded by the Fonds de la Recherche Scientifique de Belgique (F.R.S.-FNRS) under grant no. 2.5020.11 and the Tier-1 supercomputer (Zenobe) of the Fédération Wallonie Bruxelles infrastructure funded by the Walloon Region under the grant agreement no. 1117545. This work was also supported by the Fonds de la Recherche Scientifique (FNRS) and the Fonds Wetenschappelijk Onderzoek-Vlaanderen (FWO) under the EOS Project n° O0100718F. We thank Katherine Thayer-Calder and William Lipscomb for providing 6-hourly output data from CESM2.

## **Author contributions**

S.H., C.K., C.A., X.F., C.L. and A.T. designed the study. S.H. analyzed the data and wrote the manuscript. C.L. provided the analysis for the supplementary material. X.F. did the MAR simulations. All authors discussed the final version of manuscript.

## **Competing interests**

The authors declare that they have no competing interests.

## **Code and data availability**

All the code used for the analysis in this study is available upon request from the corresponding author (stefan.hofer@geo.uio.no). All the MAR model results are available for download on <ftp://ftp.climato.be/fettweis/MARv3.9/ISMIP6/GrIS/> in the framework of the ISMIP6 exercise (<https://tc.copernicus.org/articles/14/2331/2020/>).