

Introduction to Bayesian linear regression with brms — Part II: Bayesian Inference

Stefano Coretta

18/01/2020

Road map

1. Statistical inference.

Road map

1. Statistical inference.
2. Frequentist inference.

Road map

1. Statistical inference.
2. Frequentist inference.
3. Bayesian inference:

Road map

1. Statistical inference.
2. Frequentist inference.
3. Bayesian inference:
 - 3.1 Posterior distributions.


Road map

1. Statistical inference.
2. Frequentist inference.
3. Bayesian inference:
 - 3.1 Posterior distributions.
 - 3.2 ROPE.

Road map

1. Statistical inference.
2. Frequentist inference.
3. Bayesian inference:
 - 3.1 Posterior distributions.
 - 3.2 ROPE.
 - 3.3 Bayes factors.

STATISTICAL INFERENCE



Hypothesis testing

We want to know two (interrelated) things:

- Is there **evidence** for the hypothesis H ?

Hypothesis testing

We want to know two (interrelated) things:

- Is there **evidence** for the hypothesis H ?
- What is the **strength** of the evidence?

Hypothesis testing

- We test H against empirical data (**hypothesis testing**).

Hypothesis testing

- We test H against empirical data (**hypothesis testing**).
 - It is important to decide in advance the details of the analysis.


Hypothesis testing

- We test H against empirical data (**hypothesis testing**).
 - It is important to decide in advance the details of the analysis.
 - Even when you think you are not making decisions, the model is.

Hypothesis testing

- We test H against empirical data (**hypothesis testing**).
 - It is important to decide in advance the details of the analysis.
 - Even when you think you are not making decisions, the model is.
- Inference is ultimately a **long-term endeavour** (via accumulation of knowledge).

Hypothesis testing



FREQUENTIST INFERENCE

Frequentist statistics

- **Frequentist statistics** is based on the frequentist interpretation of probability.

Frequentist statistics

- **Frequentist statistics** is based on the frequentist interpretation of probability.
- **Frequentist probability of an event** is the *relative frequency* of occurrence of an event within an infinite set of imagined repetitions of that event.

Frequentist statistics

- **Frequentist statistics** is based on the frequentist interpretation of probability.
- **Frequentist probability of an event** is the *relative frequency* of occurrence of an event within an infinite set of imagined repetitions of that event.
- Most of modern science is based on *frequentism*.

Frequentist statistics

- **Frequentist statistics** is based on the frequentist interpretation of probability.
- **Frequentist probability of an event** is the *relative frequency* of occurrence of an event within an infinite set of imagined repetitions of that event.
- Most of modern science is based on *frequentism*.
 - lme4 package.

Frequentist statistics

- **Frequentist statistics** is based on the frequentist interpretation of probability.
- **Frequentist probability of an event** is the *relative frequency* of occurrence of an event within an infinite set of imagined repetitions of that event.
- Most of modern science is based on *frequentism*.
 - lme4 package.
 - **Null Hypothesis Significance Testing.**

Null Hypothesis Significance Testing

- NHST is based on falsification.

Null Hypothesis Significance Testing

- NHST is based on falsification.
 - Rejection of the Null Hypothesis (H_0).

Null Hypothesis Significance Testing

- NHST is based on falsification.
 - Rejection of the Null Hypothesis (H_0).
 - No direct “support/evidence” for hypotheses.

Null Hypothesis Significance Testing

- NHST is based on falsification.
 - Rejection of the Null Hypothesis (H_0).
 - No direct “support/evidence” for hypotheses.
- P -value (between 0 and 1).

Null Hypothesis Significance Testing

- NHST is based on falsification.
 - Rejection of the Null Hypothesis (H_0).
 - No direct “support/evidence” for hypotheses.
- P -value (between 0 and 1).
 - Probability of obtaining an estimate as extreme or more extreme, *assuming H_0 is true*.

Null Hypothesis Significance Testing

- NHST is based on falsification.
 - Rejection of the Null Hypothesis (H_0).
 - No direct “support/evidence” for hypotheses.
- P -value (between 0 and 1).
 - Probability of obtaining an estimate as extreme or more extreme, *assuming H_0 is true*.
 - Should be as low as possible.

Null Hypothesis Significance Testing

- Significance is dichotomous.

Null Hypothesis Significance Testing

- Significance is dichotomous.
 - $p < \alpha$ = “significant”.

Null Hypothesis Significance Testing

- Significance is dichotomous.
 - $p < \alpha$ = “significant”.
 - $p \geq \alpha$ = “non-significant”.

Null Hypothesis Significance Testing

- Significance is dichotomous.
 - $p < \alpha$ = “significant”.
 - $p \geq \alpha$ = “non-significant”.
- No degrees of significance.

Null Hypothesis Significance Testing

- Significance is dichotomous.
 - $p < \alpha$ = “significant”.
 - $p \geq \alpha$ = “non-significant”.
- No degrees of significance.
- “Significance” is a concept that makes sense only within frequentist statistics (NHST).

(Frequentist) confidence intervals

<https://rpsychologist.com/d3/ci/>

BAYESIAN INFERENCE

Bayesian statistics is based on the Bayesian interpretation of the **Bayes theorem**.

$$P(\theta \mid d) = \frac{P(d \mid \theta) \times P(\theta)}{P(d)}$$

- Evidence for any H (even the null).

- Evidence for any H (even the null).
- Strength of evidence.

Bayesian inference

- Evidence for any H (even the null).
- Strength of evidence.
- Capitalise on previous knowledge.

Three ways of doing hypothesis testing with Bayesian statistics:

- Inference from the **posterior**.

Three ways of doing hypothesis testing with Bayesian statistics:

- Inference from the **posterior**.
- Inference using a **Region Of Practical Equivalence** (ROPE).

Three ways of doing hypothesis testing with Bayesian statistics:

- Inference from the **posterior**.
- Inference using a **Region Of Practical Equivalence** (ROPE).
- Inference using the **Bayes factor**.

Inference from the posterior

1. Formulate a **hypothesis**:

Inference from the posterior

1. Formulate a **hypothesis**:

- H: Condition B decreases reaction times relative to Condition A.

Inference from the posterior

1. Formulate a **hypothesis**:
 - H: Condition B decreases reaction times relative to Condition A.
2. Choose **model specification** (including priors).

Inference from the posterior

1. Formulate a **hypothesis**:
 - H: Condition B decreases reaction times relative to Condition A.
2. Choose **model specification** (including priors).
3. Collect data.

Inference from the posterior

1. Formulate a **hypothesis**:
 - H: Condition B decreases reaction times relative to Condition A.
2. Choose **model specification** (including priors).
3. Collect data.
4. Calculate the **posterior** (fit the model):

Inference from the posterior

1. Formulate a **hypothesis**:
 - H: Condition B decreases reaction times relative to Condition A.
2. Choose **model specification** (including priors).
3. Collect data.
4. Calculate the **posterior** (fit the model):
 - Condition B 95% CI = [-80, -15] ms.

Inference from the posterior

1. Formulate a **hypothesis**:
 - H: Condition B decreases reaction times relative to Condition A.
2. Choose **model specification** (including priors).
3. Collect data.
4. Calculate the **posterior** (fit the model):
 - Condition B 95% CI = [-80, -15] ms.
5. **Inference**:

Inference from the posterior

1. Formulate a **hypothesis**:
 - H: Condition B decreases reaction times relative to Condition A.
2. Choose **model specification** (including priors).
3. Collect data.
4. Calculate the **posterior** (fit the model):
 - Condition B 95% CI = [-80, -15] ms.
5. **Inference**:
 - The posterior suggests that Condition B decreases reaction times by 15 to 80 ms at 95% confidence.

Inference from the posterior

1. Formulate a **hypothesis**:
 - H: Condition B decreases reaction times relative to Condition A.
2. Choose **model specification** (including priors).
3. Collect data.
4. Calculate the **posterior** (fit the model):
 - Condition B 95% CI = [-80, -15] ms.
5. **Inference**:
 - The posterior suggests that Condition B decreases reaction times by 15 to 80 ms at 95% confidence.
 - The posterior is **compatible with H**.

Inference from the posterior

1. Formulate a **hypothesis**:

Inference from the posterior

1. Formulate a **hypothesis**:

- H: Condition B decreases reaction times relative to Condition A
by 100 ms.

Inference from the posterior

1. Formulate a **hypothesis**:
 - H: Condition B decreases reaction times relative to Condition A *by 100 ms.*
2. Choose **model specification** (including priors).

Inference from the posterior

1. Formulate a **hypothesis**:
 - H: Condition B decreases reaction times relative to Condition A *by 100 ms.*
2. Choose **model specification** (including priors).
3. Collect data.

Inference from the posterior

1. Formulate a **hypothesis**:
 - H: Condition B decreases reaction times relative to Condition A *by 100 ms.*
2. Choose **model specification** (including priors).
3. Collect data.
4. Calculate the **posterior** (fit the model):

Inference from the posterior

1. Formulate a **hypothesis**:
 - H: Condition B decreases reaction times relative to Condition A *by 100 ms.*
2. Choose **model specification** (including priors).
3. Collect data.
4. Calculate the **posterior** (fit the model):
 - Condition B 95% CI = [-80, -15] ms.

Inference from the posterior

1. Formulate a **hypothesis**:
 - H: Condition B decreases reaction times relative to Condition A *by 100 ms.*
2. Choose **model specification** (including priors).
3. Collect data.
4. Calculate the **posterior** (fit the model):
 - Condition B 95% CI = [-80, -15] ms.
5. **Inference**:

Inference from the posterior

1. Formulate a **hypothesis**:
 - H: Condition B decreases reaction times relative to Condition A *by 100 ms.*
2. Choose **model specification** (including priors).
3. Collect data.
4. Calculate the **posterior** (fit the model):
 - Condition B 95% CI = [-80, -15] ms.
5. **Inference**:
 - The posterior suggests that Condition B decreases reaction times by 15 to 80 ms at 95% confidence.

Inference from the posterior

1. Formulate a **hypothesis**:
 - H: Condition B decreases reaction times relative to Condition A *by 100 ms.*
2. Choose **model specification** (including priors).
3. Collect data.
4. Calculate the **posterior** (fit the model):
 - Condition B 95% CI = [-80, -15] ms.
5. **Inference**:
 - The posterior suggests that Condition B decreases reaction times by 15 to 80 ms at 95% confidence.
 - The posterior is **not compatible with H**.

H0 vs H1

- H1 states that Condition B increases segment duration (alternative hypothesis), while H0 states that Condition B does not increase segment duration (null hypothesis, null effect).

H0 vs H1

- H1 states that Condition B increases segment duration (alternative hypothesis), while H0 states that Condition B does not increase segment duration (null hypothesis, null effect).
 - $H_1 : \beta > 0$

H0 vs H1

- H1 states that Condition B increases segment duration (alternative hypothesis), while H0 states that Condition B does not increase segment duration (null hypothesis, null effect).
 - $H_1 : \beta > 0$
 - $H_0 : \beta = 0$

Inference with a ROPE

Region Of Practical Equivalence (ROPE):

- Define a region around $\beta = 0$ that practically corresponds to a null effect.

Inference with a ROPE

Region Of Practical Equivalence (ROPE):

- Define a region around $\beta = 0$ that practically corresponds to a null effect.
 - For example: $[-5, +5]$ ms ($-5 \geq \beta \leq +5 =$ null effect).

Inference with a ROPE

Region Of Practical Equivalence (ROPE):

- Define a region around $\beta = 0$ that practically corresponds to a null effect.
 - For example: $[-5, +5]$ ms ($-5 \geq \beta \leq +5$ = null effect).
 - This ROPE has a width of 10 ms.

Inference with a ROPE

Region Of Practical Equivalence (ROPE):

- Define a region around $\beta = 0$ that practically corresponds to a null effect.
 - For example: $[-5, +5]$ ms ($-5 \geq \beta \leq +5$ = null effect).
 - This ROPE has a width of 10 ms.
- Collect data until the 95% CI of β has a width equal to or smaller than the width of the ROPE.

Inference with a ROPE

Region Of Practical Equivalence (ROPE):


- Define a region around $\beta = 0$ that practically corresponds to a null effect.
 - For example: $[-5, +5]$ ms ($-5 \geq \beta \leq +5$ = null effect).
 - This ROPE has a width of 10 ms.
- Collect data until the 95% CI of β has a width equal to or smaller than the width of the ROPE.
 - Choose a minimal sample size (ideally based on a prospective power analysis).

Inference with a ROPE

Region Of Practical Equivalence (ROPE):

- Define a region around $\beta = 0$ that practically corresponds to a null effect.
 - For example: $[-5, +5]$ ms ($-5 \geq \beta \leq +5$ = null effect).
 - This ROPE has a width of 10 ms.
- Collect data until the 95% CI of β has a width equal to or smaller than the width of the ROPE.
 - Choose a minimal sample size (ideally based on a prospective power analysis).
 - Collect data and check 95% CI. If the width is greater than the ROPE, collect more data and repeat (*sequential testing*).

Inference with a ROPE



- We focus on the estimate **precision** of β .

- We focus on the estimate **precision** of β .
- Independent from the value of β .

- We focus on the estimate **precision** of β .
- Independent from the value of β .
- Higher precision means greater confidence in the estimated value of β .

The Bayes factor is the ratio of the likelihood of H_1 to the likelihood of H_2 .

$$BF_{12} = \mathcal{L}(H_1) / \mathcal{L}(H_2)$$

Bayes Factor

BF	$p(M1 D)$	evidence
1–3	0.5–0.75	weak
3–20	0.75–0.95	positive
20–150	0.95–0.99	strong
> 150	> 0.99	very strong

```
priors <- c(  
  prior(normal(0, 500), class = Intercept),  
  prior(cauchy(0, 15), class = sigma),  
  prior(normal(0, 750), class = b, coef = "vowel_o"),  
  prior(normal(0, 750), class = b, coef = "vowel_u"),  
  prior(cauchy(0, 15), class = sd),  
  prior(lkj(2), class = cor)  
)
```

Bayes Factor

```
f1_3_bf <- brm(  
  f1 ~ 1 + vowel + (1 + vowel | speaker),  
  family = gaussian(),  
  prior = priors,  
  data = f_end,  
  chains = 4,  
  iter = 2000,  
  file = "./cache/f1_3_bf",  
  save_all_pars = TRUE  
)
```

Bayes Factor

```
priors <- c(  
  prior(normal(0, 500), class = Intercept),  
  prior(cauchy(0, 15), class = sigma),  
  prior(normal(0, 750), class = b, coef = "vowel_o"),  
  prior(normal(0, 750), class = b, coef = "vowel_u"),  
  prior(normal(0, 750), class = b, coef = "c2_placevelar"),  
  prior(cauchy(0, 15), class = sd),  
  prior(lkj(2), class = cor)  
)
```

Bayes Factor

```
f1_3_place <- brm(  
  f1 ~ 1 + vowel + c2_place + (1 + vowel + c2_place | sp  
  family = gaussian(),  
  prior = priors,  
  data = f_end,  
  chains = 4,  
  iter = 2000,  
  file = "./cache/f1_3_place",  
  save_all_pars = TRUE  
)
```

Bayes Factor

```
bf <- bayes_factor(f1_3_bf, f1_3_place)
```

```
## Iteration: 1
```

```
## Iteration: 2
```

```
## Iteration: 3
```

```
## Iteration: 4
```

```
## Iteration: 5
```

```
## Iteration: 6
```

```
## Iteration: 7
```

```
## Iteration: 8
```

```
## Iteration: 9
```

```
## Iteration: 1
```

```
## Iteration: 2
```

```
## Iteration: 3
```

```
## Iteration: 4
```


Bayes Factor

```
bf
```

```
## Estimated Bayes factor in favor of f1_3_bf over f1_3_place: 0.58852
```

- BFs are *VERY* sensitive to priors and data.

Bayes Factor

- BFs are **VERY** sensitive to priors and data.
- Always calculate and report BFs by comparing models with increasingly narrower priors (at least 3-4).

- BFs are **VERY** sensitive to priors and data.
- Always calculate and report BFs by comparing models with increasingly narrower priors (at least 3-4).
- It's important to run *sensitivity analyses* that assess the influence of the priors on the posterior (not only if you use BFs, but always).

THE END

