

An introduction to GAM(M)s

Stefano Coretta

12/07/2018

Linear models


Time travel...

Linear models

$$y = 3 + 2x$$


where $x = (2, 4, 5, 8, 10, 23, 36)$

Linear models



Linear models


$$y = 3 + 2x$$



Linear models

- ▶ In science, we have x and y ...
- ▶ for example, vowel duration and VOT

Linear models




Linear models


- ▶ The formula: $y = \beta_0 + \beta_1 x$
 - ▶ β_0 is the **intercept**
 - ▶ β_1 is the **slope**
- ▶ We know x and y
 - ▶ we need to estimate β_0, β_1
- ▶ We can add more predictors
 - ▶ $y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \dots + \beta_n x_n$
- ▶ code in R

```
lm(y ~ x, data)
```



Linear models




Linear models



Linear models with non-linear data



Linear models with non-linear data



Linear models with non-linear data

How to account for non-linearity in a linear model?


- ▶ Use **higher-degree polynomials**

- ▶ quadratic: $y = \beta_0 + \beta_1x + \beta_2x^2$


- ▶ cubic: $y = \beta_0 + \beta_1x + \beta_2x^2 + \beta_3x^3$

- ▶ n th: $y = \beta_0 + \beta_1x + \beta_2x^2 + \beta_3x^3 + \dots + \beta_nx^n$

Linear models with non-linear data



Linear models with non-linear data



Linear models with non-linear data

