

Quantifying vocal fold activity: two new methods for analysing electroglottographic data

Stefano Coretta


The University of Manchester

New developments in speech sensing and imaging, Lisbon, 23rd June 2018


1


Background: Electroglottography

- EGG (Fabre, 1957; Scherer & Titze, 1987; Rothenberg & Mahshie, 1988)
- Purpose: estimation of vocal folds contact area (VFCA)
- How: based on modulations of a current that travels the neck generated by the opening and closing of the vocal folds




3






- · Pros:
 - non-invasive
 - relatively simple signal
- · Cons:
 - · Herbst et al. (2014), Hampala et al. (2016)


Background: EGG signal



Background: EGG signal




Background: Contact quotient



Background: Wavegrams

- · CQ reduces dimentionality of EGG signal
- Herbst et al. (2010) propose the wavegram as a multidimentional account of the EGG signal

Background: Wavegrams




Background: Tracegram and Wavegram GAMs

- limitations
 - CQ is not precise (Baken, 1992; Herbst et al., 2017)
 - wavegram cannot be accessed statistically
- two new techniques
 - · tracegram
 - wavegram GAMs

Methods

- 4 phonetically trained BE speakers (1 F, 3 M)
- [α] in modal and breathy voice
 - 10 × 2 = 20 tokens per speaker
 - · 80 tokens
- · equipment
 - · Glottal Enterprises EG2-PCX2 unit
 - Movo LV4-O2 Lavalier microphone (sample rate 44100 Hz, 16-bit)
- · analysis window
 - · 500 ms portion centred around mid point of each token


Results: CQ




Results: CQ

```
## Linear mixed model fit by REML. t-tests use Satterthwaite's method [
## lmerModImerTest1
## Formula: contact quotient ~ phonation + (1 + phonation | speaker)
     Data: tracegram
##
##
## REML criterion at convergence: -24474.4
##
## Scaled residuals:
      Min 10 Median 30 Max
##
## -7.1831 -0.5597 0.0237 0.6202 5.3121
##
## Random effects:
                    Variance Std.Dev. Corr
## Groups Name
## speaker (Intercept) 0.003305 0.05749
          phonationbreathy 0.005009 0.07077 -0.19
##
## Residual
                           0.000976 0.03124
## Number of obs: 5999, groups: speaker, 5
##
## Fixed effects:
           Estimate Std. Error df t value Pr(>|t|)
##
## (Intercept) 0.50512 0.02572 4.00001 19.643 3.96e-05 ***
## phonationbreathy -0.06246 0.03166 3.99996 -1.973 0.12
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Correlation of Fixed Effects:
##
            (Intr)
## phontnbrthy -0.190
```

Results: Tracegram




Results: Tracegram




Results: Wavegram GAM

- generalised additive mixed models (Wood, 2006; Sóskuthy, 2017; van Rij et al., 2017)
 - · non-linear multidimensional data
- statistical testing of wavegram data
 - heat-map plots: time, period, amplitude

Results: Wavegram GAM (modal)



Results: Wavegram GAM (breathy)



Results: Wavegram GAM

```
## phonation gam null: amplitude ~ s(sequence, k = 8) + s(sample) + ti(sequence, sample,
##
      k = 8) + s(sequence, speaker phon, bs = "fs", m = 1, k = 8)
##
## phonation gam: amplitude ~ phonation + s(sequence, k = 8) + s(sample) + s(sequence,
      by = phonation, k = 8) + s(sample, by = phonation) + ti(sequence,
##
##
      sample, k = 8) + ti(sequence, sample, by = phonation, k = 8) +
      s(sequence, speaker phon, bs = "fs", m = 1, k = 8)
##
##
## Chi-square test of ML scores
## ----
                 Model
                           Score Edf Difference Df p.value Sig.
##
## 1 phonation_gam_null -53190.37 10
## 2
         phonation gam -66983.42 18 13793.050 8.000 < 2e-16 ***
##
## AIC difference: 27741.14, model phonation gam has lower AIC.
```

Discussion

- · CQ performed badly for speaker 03
- Tracegrams
 - · non-resource-intensive method for visualising fold activity
- · Wavegram GAMs
 - assessing fold activity data statistically

The end

Thanks!

References

- Baken, Ronald J. 1992. Electroglottography. *Journal of Voice* 6(2). 98–110.
- Fabre, P. 1957. Un procede electrique percutane d'inscrition de l'accolement glottique au cours de la phonation: glottographie de haute frequence. Premiers resultats. Bulletin de l'Académie nationale de médecine 141. 66.
- Hampala, Vít, Maxime Garcia, Jan G. Švec, Ronald C. Scherer & Christian T. Herbst. 2016. Relationship between the electroglottographic signal and vocal fold contact area. *Journal of Voice* 30(2). 161–171. doi:10.1016/j.jvoice.2015.03.018.

- Herbst, Christian T., W. Tecumseh S. Fitch & Jan G. Švec. 2010. Electroglottographic wavegrams: A technique for visualizing vocal fold dynamics noninvasively. *The Journal of the Acoustical Society of America* 128(5). 3070–3078. doi:10.1121/1.3493423.
- Herbst, Christian T., Jörg Lohscheller, Jan G. Švec, Nathalie Henrich, Gerald Weissengruber & W. Tecumseh S. Fitch. 2014. Glottal opening and closing events investigated by electroglottography and super-high-speed video recordings. *Journal of Experimental Biology* 217(6). 955–963. doi:10.1242/jeb.093203.
- Herbst, Christian T., Harm K. Schutte, Daniel L. Bowling & Jan G. Svec. 2017. Comparing chalk with cheese—the EGG contact quotient is only a limited surrogate of the closed quotient. *Journal of Voice* 31(4). 401–409.

- Rothenberg, Martin & James J. Mahshie. 1988. Monitoring vocal fold abduction through vocal fold contact area. *Journal of Speech, Language, and Hearing Research* 31(3). 338–351.
- Scherer, Ronald C. & Ingo R. Titze. 1987. The abduction quotient related to vocal quality. *Journal of Voice* 1(3). 246–251.
- Sóskuthy, Márton. 2017. Generalised additive mixed models for dynamic analysis in linguistics: a practical introduction. arXiv preprint arXiv:1703.05339.
- van Rij, Jacolien, Martijn Wieling, R. Harald Baayen & Hedderik van Rijn. 2017. itsadug: Interpreting time series and autocorrelated data using GAMMs. R package version 2.3.
- Wood, Simon. 2006. Generalized additive models: An introduction with R. CRC Press.