Vowel duration and tongue root advancement: Results from an exploratory study of the relation between voicing and vowel duration

Stefano Coretta¹

The University of Manchester^{a)}

Put your abstract here. Abstracts are limited to 200 words for regular articles and

100 words for Letters to the Editor. Please no personal pronouns, also please do not

use the words new'' and/ornovel" in the abstract. An article usually includes an

abstract, a concise summary of the work covered at length in the main body of the

5 article.

a)stefano.coretta@manchester.ac.uk

6 I. INTRODUCTION

This paper reports a previously undocumented correlation between vowel duration and degree of tongue root advancement. In an exploratory study of the articulatory correlates of stop voicing, it has been found that tongue root advancement—a known mechanism that facilitates voicing during stop closure—can be implemented not only during the closure of a stop, but even during the production of the vowel preceding a stop. Moreover, vowel acoustic duration turned out to be linearly correlated with degree of tongue root advancement, such that longer vowels show greater tongue root advancement.

One of the known differences in supra-glottal articulation between voiced and voiceless 14 stops concerns the position of the tongue root relative to the front-back axis of the oral tract. It has been observed that the tongue root is more advanced in voiced than in voiceless stops (Kent and Moll, 1969; Perkell, 1969; Westbury, 1983). This gesture has been interpreted 17 as a mechanism to ensure voicing can be maintained during the closure of the stop. The 18 realisation of vocal fold vibration (i.e. voicing) requires the air pressure in the supra-glottal cavity to be lower than the air pressure below the glottis. During the production of voiced obstruents, the supra-glottal pressure increases due to the immittance of air in the supra-21 glottal cavity. Such pressure increase can hinder the ability to sustain voicing during closure, to the point that voicing ceases if the supra-glottal pressure is higher than the sub-glottal pressure (Ohala, 2011). One of the possible articulatory solution to counterbalance the increase in pressure during the closure of a voiced stop is to expand the supra-glottal cavity by advancing the root of the tongue.

Tongue root advancement has also been reported as a mechanism for ensuring a short voice onset time (Ahn and Davidson, 2016).

An extensive number of studies show that, cross-linguistically, vowels tend to be longer 29 when followed by voiced obstruents than when they are followed by voiceless obstruents (Chen, 1970; Durvasula and Luo, 2012; Esposito, 2002; Farnetani and Kori, 1986; Fowler, 31 1992; House and Fairbanks, 1953; Hussein, 1994; Klatt, 1973; Lampp and Reklis, 2004; Lisker, 1974; Peterson and Lehiste, 1960). This phenomenon, know as the voicing effect, has been reported in a variety of languages, including (but not limited to) English, German, Hindi, Russian, Italian, Arabic, and Korean (see Maddieson and Gandour 1976 for a more comprehensive list). A common stance in the literature is that the magnitude of the voicing effect differs depending on the language (although see Laeufer 1992), and that this 37 phenomenon is not a universal tendency, since the duration of vowels is not affected by the voicing of the following obstruents in some languages, like Polish and Czech (Keating, 1984). Although several attempts have been made to explain the voicing effect, an account that survives all empirical data is still lacking (Durvasula and Luo, 2012; Sóskuthy, 2013).

To summarise, tongue root advancement, shorter VOT duration, and longer vowel durations are all correlates of voicing. Moreover, tongue root advancement and shorter VOT show a link. In this paper, I will report the results from an exploratory study which fill the gap in this picture of correlation, by showing that tongue root advancement is also linked to longer vowel durations.

47 II. METHODOLOGY

48 A. Participants

Eleven native speakers of Italian (5 females, 6 males) and 6 native speakers of Polish (3 females, 3 males) were recorded in the Phonetics Laboratory at the University of Manchester and in a private location in Italy (see ??). The Italian speakers of this study are from Northern Italy (three from the north-west and one from north-east). The Polish group was more heterogeneous, with two speakers from the West (Poznań), and two from the East (Warsaw and Przasnysz). Ethical clearance was obtained for this study from the University of Manchester (REF 2016-0099-76). The participants received a monetary compensation.

56 B. Materials

Disyllabic words of the form $C_1V_1C_2V_2$ were used as targets, where $C_1 = /p/$, $V_1 = /p/$, $V_1 = /p/$, $V_1 = /p/$, $V_1 = /p/$, $V_2 = /p/$, $V_3 = /p/$, $V_4 = /p/$, $V_5 = /p/$, $V_6 = /p/$, $V_7 = /p/$, $V_8 = /p/$, $V_8 = /p/$, $V_8 = /p/$, $V_9 = /p/$

only central/back vowels (low /a/, mid /o/, and high /u/) were included in the target words.

The use of back and central vowels (with the exclusion of mid/high front vowels) had the advantage of facilitating the identification of the consonantal gesture of C2. Since the original motive for the exploratory study was to study possible difference in the articulation of closure in voiceless vs. voiced stops, only coronal and velar stops were chosen as target consonants since, of course, the closure of labial consonants cannot be imaged with ultrasonography.

The target words were embedded in a frame sentence, Dico X lentamente 'I say X slowly' for Italian, and Mówię X teraz 'I say X now' for Polish. The similarity of prosodic structure of these sentences ensured better comparability between the two languages.

C. Equipment and procedure

76

An Articulate Instruments LtdTM system was used for this study. The system is made
of a TELEMED Echo Blaster 128 unit, an Articulate Instruments LtdTM P-Stretch synchronisation unit, and a FocusRight Scarlett Solo pre-amplifier (see ??). A TELEMED
C3.5/20/128Z-3 ultrasonic transducer (20mm radius, 2-4 MHz) and a Movo LV4-O2 Lavalier microphone were used respectively the acquisition of ultrasonic and audio data. Stabilisation of the ultrasonic transducer was ensured by means of a metallic headset designed
by Articulate Instruments LtdTM (2008). The transducer was placed in contact with the
sub-mental triangle, aligned with the mid-sagittal plane. The headset holds the transducer
and it keeps it in a constant position and inclination relative to the sub-mental triangle, thus
allowing head movements without the need for post-processing correction. The acquisition
of the mid-sagittal ultrasonic and audio signals was achieved with the software Articulate

with Microsoft Windows 7. The synchronisation of the ultrasonic and audio signals was performed by AAA after recording by means of a synchronisation signal produced by the P-Stretch unit. The ranges of the ultrasonic settings were: frames per second = 43-68, number 91 of scan lines = 88–114, pixel per scan line = 980–988, field of view = 71–93°, pixel offset = 109-263, depth (mm) = 75-180. The audio signal was recorded at 22050 Hz (16-bit). The head set with the prob was fitted at the beginning of the experimental session. Then 94 the hard palate was imaged by recording the participant swallowing water (Epstein and Stone, 2005). A between-speaker reference coordinate system was derived from imaging the occlusal plane, which corresponds to the trace of a metallic bite plate inserted in the mouth of the participant (Scobbie et al., 2011). The participant gently bites on the bite plate while pressing the tongue against it. The participant then started reading six repetitions of the sentences with the target words. These were presented on the screen in a randomised 100 order across participants, although the order was kept the same for each repetition within 101 participant due to design constraints of the AAA software.

Assistant Advanced (AAA, v2.17.2) running on a Hawlett-Packard ProBook 6750b laptop

D. Data processing and analysis

1. Acoustic data

103

104

The audio data was subject to force alignment using the SPeech Phonetisation Alignment and Syllabification software (SPPAS) (Bigi, 2015). The outcome of the automatic alignment was then manually corrected, according to the criteria in Table I, which were mainly based

TABLE I. List of measurements extracted from the acoustic data.

landmarks		criteria
vowel onset	(V1 onset)	appearance of higher formants in the spectrogram
		following the release of $/p/$ (C1)
vowel offset	(V1 offset)	disappearance of the higher formants in the
		spectrogram preceding the target consonant (C2)
consonant onset	(C2 onset)	corresponds to V1 offset
closure onset	(C2 closure onset)	corresponds to V1 offset
consonant offset	(C2 offset)	appearance of higher formants of the vowel
		following C2 (V2); corresponds to V2 onset
consonant release	(C2 release)	automatic detection (Ananthapadmanabha et al.,
		2014)

on properties of the the spectrogram. The release C2 was detected automatically in Praat

(Boersma and Weenink, 2016) by means of the burst-detection algorithm described in Anan
thapadmanabha et al. (2014). The durations of the following intervals were extracted from

the annotated acoustic landmarks using a scripted procedure in Praat: vowel duration (V1

onset to V1 offset), consonant duration (V1 offset to V2 onset), and closure duration (V1

offset to C2 release).

2. Ultrasonic data

114

Mid-sagittal tongue contours were obtained from the ultrasonic data according to the fol-115 lowing method. Smoothing splines were automatically fitted to the visible tongue contours 116 in AAA. Manual correction was then applied in cases of clear tracking errors. The time of 117 maximum tongue displacement within consonant closure was then calculated in AAA follow-118 ing the method described in Strycharczuk and Scobbie (2015), which is based on the velocity 119 of tongue displacement along a vector. To obtain the time of maximum displacement, a fan-120 like frame consisting of 42 equidistant radial lines is used as the coordinate system. The origin of the 42 fan-lines coincides with the centre of the ultrasonic transducer, such that 122 each fan-line is parallel to the direction of the ultrasonic scan lines. Tongue displacement 123 was thus calculated as the displacement of the fitted splines along the fan-line vectors. The time of maximum tongue displacement was the time of greater displacement along the vector 125 that showed the greatest standard deviation of displacement. The fan-lines from which the 126 relevant vector was chosen were restricted to the fan-lines corresponding to the tongue tip for coronal consonants, and to the fan-lines corresponding to the tongue dorsum for velar 128 consonants.

The polar coordinates of the tongue contours were exported from two time points: the onset of C2 closure, and the time of maximum tongue displacement (which is always within C2
closure). The contours were normalised within speaker by applying offsetting and rotation
relative to the participant's occlusal plane (Scobbie et al., 2011).

- ¹³⁴ Although there is a tendency in the articulatory literature to suggest that labial consonants do not affect
- lingual articulations, ? report tongue body lowering in the context of labial stops.

- Ahn, S., and Davidson, L. (2016). "Tongue root positioning in English voiced obstruents:
- Effects of manner and vowel context," The Journal of the Acoustical Society of America
- 140(4), 3221–3221.

136

- Ananthapadmanabha, T. V., Prathosh, A. P., and Ramakrishnan, A. G. (2014). "Detection
- of the closure-burst transitions of stops and affricates in continuous speech using the plosion
- index," The Journal of the Acoustical Society of America 135(1), 460–471.
- ¹⁴³ Articulate Instruments LtdTM (2008). "Ultrasound stabilisation headset users manual: Re-
- vision 1.4" Edinburgh, UK: Articulate Instruments Ltd.
- Bigi, B. (2015). "SPPAS Multi-lingual approaches to the automatic annotation of speech,"
- The Phonetician **111–112**, 54–69.
- Boersma, P., and Weenink, D. (2016). "Praat: doing phonetics by computer [Computer
- program]. Version 6.0.23" http://www.praat.org/.
- chen, M. (1970). "Vowel length variation as a function of the voicing of the consonant
- environment," Phonetica **22**(3), 129–159.
- Durvasula, K., and Luo, Q. (2012). "Voicing, aspiration, and vowel duration in Hindi,"
- Proceedings of Meetings on Acoustics 18, 1–10.
- Epstein, M. A., and Stone, M. (2005). "The tongue stops here: Ultrasound imaging of the
- palate," The Journal of the Acoustical Society of America 118(4), 2128–2131.

- Esposito, A. (2002). "On vowel height and consonantal voicing effects: Data from Italian,"
- Phonetica 59(4), 197–231.
- Farnetani, E., and Kori, S. (1986). "Effects of syllable and word structure on segmental
- durations in spoken Italian," Speech communication 5(1), 17–34.
- Fowler, C. A. (1992). "Vowel duration and closure duration in voiced and unvoiced stops:
- There are no contrast effects here," Journal of Phonetics **20**(1), 143–165.
- House, A. S., and Fairbanks, G. (1953). "The influence of consonant environment upon the
- secondary acoustical characteristics of vowels," The Journal of the Acoustical Society of
- America 25(1), 105-113.
- Hussein, L. (1994). "Voicing-dependent vowel duration in Standard Arabic and its acquisi-
- tion by adult american students," Ph.D. thesis, The Ohio State University.
- Keating, P. A. (1984). "Universal phonetics and the organization of grammars," UCLA
- Working Papers in Phonetics **59**.
- Kent, R. D., and Moll, K. L. (1969). "Vocal-tract characteristics of the stop cognates,"
- Journal of the Acoustical Society of America 46(6B), 1549–1555.
- Klatt, D. H. (1973). "Interaction between two factors that influence vowel duration," The
- Journal of the Acoustical Society of America 54(4), 1102–1104.
- Laeufer, C. (1992). "Patterns of voicing-conditioned vowel duration in French and English,"
- Journal of Phonetics 20(4), 411–440.
- Lampp, C., and Reklis, H. (2004). "Effects of coda voicing and aspiration on Hindi vowels,"
- The Journal of the Acoustical Society of America 115(5), 2540–2540.

- Lisker, L. (1974). "On "explaining" vowel duration variation," in Proceedings of the Lin-
- guistic Society of America, pp. 225–232.
- Maddieson, I., and Gandour, J. (1976). "Vowel length before aspirated consonants," in
- UCLA Working papers in Phonetics, Vol. 31, pp. 46–52.
- Ohala, J. J. (2011). "Accommodation to the aerodynamic voicing constraint and its phono-
- logical relevance," in *Proceedings of the 17th International Congress of Phonetic Sciences*,
- pp. 64–67.
- Perkell, J. S. (1969). Physiology of Speech production: Results and implication of quantitative
- cineradiographic study (Cambridge, MA: MIT Press).
- Peterson, G. E., and Lehiste, I. (1960). "Duration of syllable nuclei in english," The Journal
- of the Acoustical Society of America **32**(6), 693–703.
- Scobbie, J. M., Lawson, E., Cowen, S., Cleland, J., and Wrench, A. A. (2011). "A common
- co-ordinate system for mid-sagittal articulatory measurement," in QMU CASL Working
- 189 *Papers*, pp. 1–4.
- Sóskuthy, M. (2013). "Phonetic biases and systemic effects in the actuation of sound
- change," Ph.D. thesis, University of Edinburgh.
- 192 Strycharczuk, P., and Scobbie, J. M. (2015). "Velocity measures in ultrasound data. Gestu-
- ral timing of post-vocalic /l/ in English," in Proceedings of the 18th International Congress
- of Phonetic Sciences, pp. 1–5.
- Westbury, J. R. (1983). "Enlargement of the supraglottal cavity and its relation to stop
- consonant voicing," The Journal of the Acoustical Society of America **73**(4), 1322–1336.