

# Cáncer de endometrio localizado

Maximiliano Ink  
Residencia de Oncología Clínica  
Instituto de Oncología *Á. H. Roffo* - UBA





# Introducción

El cáncer de endometrio es el principal tumor maligno del tracto genital femenino en países desarrollados, y el segundo en frecuencia en nuestro país, por detrás del cáncer de cérvix.





# Epidemiología

- Incidencia 3,4%
- Mortalidad 2,2%
- Sobrevida a 5 años 80%

Cancer of the Endometrium - Cancer Stat Facts [Internet]. SEER. Available from: <https://seer.cancer.gov/statfacts/html/corp.html>

How JA, Jazaeri AA, Westin SN, Lawson BC, Klopp AH, Soliman PT, et al. Translating biological insights into improved management of endometrial cancer. Nat Rev Clin Oncol [Internet]. 2024;21(November):781–800. Available from: <http://dx.doi.org/10.1038/s41571-024-00934-7>

# Factores de riesgo



Obesidad



Hiperestrogenemia



Menarca temprana



Menopausia tardía



Sme. de Lynch



# Manifestaciones clínicas

## Workup

Ecografía TV

Histeroscopia + biopsia


Laboratorio con hemograma,  
función renal y hepática

RMN de abdomen y pelvis

Tomografía de tórax (y abdomen y  
pelvis)

# Clasificación histológica

- Tipo 1: endometroide
- Tipo 2: no endometroide → células claras, seroso, carcinosarcoma



**POLEmut**

**5-15%**  
**Endometroide**  
**Alto grado**  
**Estadíos tempranos**  
**Excelente pronóstico**

**NSMP**

**30-40%**  
**Bajo grado**  
**Pronóstico intermedio**

**dMMR**

**25-30%**  
**Endometroide**  
**Alto grado**  
**Sme. Lynch**  
**Pronóstico intermedio**

**p53mut**

**5-15%**  
**Cualquier histología**  
**Alto grado**  
**Pésimo pronóstico**

# Cirugía



## CRITERIA FOR CONSIDERING FERTILITY-SPARING OPTIONS FOR MANAGEMENT OF ENDOMETRIAL CARCINOMA (All criteria must be met)

- Well-differentiated (grade 1) endometrioid adenocarcinoma on dilation and curettage (D&C) confirmed by expert pathology review
- Disease limited to the endometrium on MRI (preferred) or transvaginal ultrasound<sup>i</sup>
- Absence of suspicious or metastatic disease on imaging
- No contraindications to medical therapy or pregnancy
- Patients should undergo counseling that fertility-sparing option is NOT standard of care for the treatment of endometrial carcinoma

- Consultation with a fertility expert prior to therapy
- Recommend molecular evaluation of tumor and evaluation for inherited cancer risk [\(UN-1\)](#)
- Ensure negative pregnancy test

## PRIMARY TREATMENT

- Continuous progestin-based therapy:
  - Megestrol
  - Medroxyprogesterone
  - Levonorgestrel IUD (preferred for fertility preservation)
- Weight management/lifestyle modification counseling<sup>t</sup>

## SURVEILLANCE

Endometrial evaluation every 3–6 mo (either D&C or endometrial biopsy)

Complete response by 6 mo

Encourage conception (with continued surveillance/ endometrial sampling every 6–12 mo and consider maintenance progestin-based therapy if patient is not actively trying to conceive)

TH/BSO with staging<sup>d,e</sup> after childbearing complete or progression of disease on endometrial sampling [\(ENDO-1\)](#)

- Ovarian preservation may be considered in select patients who are premenopausal

Endometrial cancer present at 6–12 mo<sup>i,u</sup>

TH/BSO with staging<sup>d,e</sup> [\(ENDO-1\)](#)

- Ovarian preservation may be considered in select patients



# Estadificación

## Stage I: Tumour confined to the corpus uteri and/or ovary

- **IA:** Disease limited to the endometrium OR non-aggressive histological subtype with <50% DMI and no or focal LVSI
  - **IA1:** Non-aggressive histological subtype limited to the endometrium or endometrial polyp
  - **IA2:** Non-aggressive histological subtype with <50% DMI
  - **IA3:** Low-grade endometrioid carcinoma limited to the uterus and ovary
- **IB:** Non-aggressive histological subtype with ≥50% DMI here and no or focal LVSI

## Stage IV: Spread to bladder mucosa and/or intestinal mucosa and/or distant metastases

- **IVA:** Invasion of the bladder mucosa and/or the intestinal bowel mucosa
- **IVB:** Abdominal peritoneal metastasis beyond the pelvis
- **IVC:** Distal metastases, including metastasis to any extra- or intra-abdominal lymph nodes above the renal vessels, or to the lungs, liver, brain or bone



# En el consultorio...

- ❖ ¿Qué riesgo de recaída presenta el paciente?
- ❖ ¿Tiene indicado adyuvancia?
- ❖ ¿Cuál es el esquema adyuvante más apropiado?
- ❖ ¿Cómo lo controlarían?



# Estratificación de riesgo

**Table 2. EC risk groups**

| Risk group             | Description <sup>a</sup>                                                                                                                                                                                                                                                                                                                                                                                        |
|------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Low risk               | Stage IA (G1-G2) with endometrioid type (dMMR <sup>b</sup> and NSMP) and no or focal LVSI<br>Stage I/II <i>POLE</i> mut cancer; for stage III <i>POLE</i> mut cancers <sup>c</sup>                                                                                                                                                                                                                              |
| Intermediate risk      | Stage IA G3 with endometrioid type (dMMR and NSMP) and no or focal LVSI<br>Stage IA non-endometrioid type (serous, clear-cell, undifferentiated carcinoma, carcinosarcoma, mixed) and/or p53-abn cancers without myometrial invasion and no or focal LVSI<br>Stage IB (G1-G2) with endometrioid type (dMMR and NSMP) and no or focal LVSI<br>Stage II G1 endometrioid type (dMMR and NSMP) and no or focal LVSI |
| High-intermediate risk | Stage I endometrioid type (dMMR and NSMP) any grade and any depth of invasion with substantial LVSI<br>Stage IB G3 with endometrioid type (dMMR and NSMP) regardless of LVSI<br>Stage II G1 endometrioid type (dMMR and NSMP) with substantial LVSI<br>Stage II G2-G3 endometrioid type (dMMR and NSMP)                                                                                                         |
| High risk              | All stages and all histologies with p53-abn and myometrial invasion<br>All stages with serous or undifferentiated carcinoma including carcinosarcoma with myometrial invasion<br>All stage III and IVA with no residual tumour, regardless of histology and regardless of molecular subtype <sup>b</sup>                                                                                                        |

Estadío – Grado histológico – Invasión miometrial – LVSI – Perfil molecular

# FIFTEEN-YEAR RADIOTHERAPY OUTCOMES OF THE RANDOMIZED PORTEC-1 TRIAL FOR ENDOMETRIAL CARCINOMA

CARIEN L. CREUTZBERG, M.D., Ph.D.,<sup>\*</sup> REMI A. NOUT, M.D.,<sup>\*</sup> MARNIX L. M. LYBEERT, M.D.,<sup>†</sup>  
CARLA C. WÁRLÁM-RODENHUIS, M.D.,<sup>‡</sup> JAN J. JOBSEN, M.D., Ph.D.,<sup>§</sup> JAN-WILLEM M. MENS, M.D.,<sup>¶</sup>  
LUDY C. H. W. LUTGENS, M.D., Ph.D.,<sup>\*\*</sup> ELISABETH PRAS, M.D., Ph.D.,<sup>††</sup>  
LONNEKE V. VAN DE POLL-FRANSE, Ph.D.,<sup>‡‡</sup> AND WIM L. J. VAN PUTTEN, M.Sc.<sup>||</sup>  
FOR THE PORTEC STUDY GROUP



- Grade 1 ; myometrial invasion  $\geq 1/2$
- Grade 2
- Grade III; myometrial invasion  $< 1/2$

TAH-BSO without lymphadenectomy







Table 1. Patient characteristics after central pathology review

| Characteristic           | No. of patients (%) |               |
|--------------------------|---------------------|---------------|
|                          | RT (n = 354)        | NAT (n = 360) |
| Age (y)                  |                     |               |
| <60                      | 93 (26)             | 108 (30)      |
| 60–70                    | 136 (38)            | 134 (37)      |
| >70                      | 125 (35)            | 118 (33)      |
| Mean (SD)                | 66.3 (SD 9)         | 65.7 (SD 9)   |
| Range                    | 41–85               | 43–90         |
| Myometrial invasion      |                     |               |
| <50%                     | 138 (39)            | 156 (43)      |
| ≥50%                     | 216 (61)            | 204 (57)      |
| Revised histologic grade |                     |               |
| 1                        | 198 (56)            | 197 (55)      |
| 2                        | 49 (14)             | 39 (11)       |
| 3                        | 32 (09)             | 54 (15)       |
| ND*                      | 75 (21)             | 70 (19)       |
| Revised FIGO 1988 stage  |                     |               |
| IB Grade 1 <sup>†</sup>  | 60 (17)             | 74 (21)       |
| IB Grade 2 <sup>‡</sup>  | 56 (16)             | 47 (13)       |
| IB Grade 3               | 22 (06)             | 35 (10)       |
| IC Grade 1               | 138 (39)            | 123 (34)      |
| IC Grade 2 <sup>‡</sup>  | 68 (19)             | 62 (17)       |
| IC Grade 3               | 10 (03)             | 19 (05)       |
| Vascular space invasion  |                     |               |
| Present                  | 22 (06)             | 19 (05)       |
| HIR                      |                     |               |
| No                       | 170 (48)            | 178 (49)      |
| Yes                      | 184 (52)            | 182 (51)      |



# **PORTEC-1**

Overall survival





**ARTICLE**

Clinical Study

# Ten-year results of the PORTEC-2 trial for high-intermediate risk endometrial carcinoma: improving patient selection for adjuvant therapy

B. G. Wortman<sup>1</sup>, C. L. Creutzberg<sup>1</sup>, H. Putter<sup>2</sup>, I. M. Jürgenliemk-Schulz<sup>3</sup>, J. J. Jobsen<sup>4</sup>, L. C. H. W. Lutgens<sup>5</sup>, E. M. van der Steen-Banasik<sup>6</sup>, J. W. M. Mens<sup>7</sup>, A. Slot<sup>8</sup>, M. C. Stenfort Kroese<sup>9</sup>, B. van Triest<sup>10</sup>, H. W. Nijman<sup>11</sup>, E. Stelloo<sup>12</sup>, T. Bosse<sup>12</sup>, S. M. de Boer<sup>1</sup>, W. L. J. van Putten<sup>13</sup>, V. T. H. B. M Smit<sup>12</sup> and R. A. Nout<sup>1</sup> for the PORTEC Study Group

## PORTEC – 2 Design

### TAH / BSO **no LND**

Eligibility:

High intermediate risk group

-age>60+ IC G1-2 or IB G3

-stage Ila

**N=427 pts**

### EBRT

**N=214**

### Brachytherapy

**N=213**

Primary endpoint: rate of vaginal relapse

Secondary endpoints: OS, QOL

30

## PORTEC-2

Table 1. Patient and tumour characteristics and pathology review



Intention to treat population

| <i>n</i> = 427             | EBRT ( <i>n</i> = 214) |       | VBT ( <i>n</i> = 213) |       |
|----------------------------|------------------------|-------|-----------------------|-------|
| <b>Age</b>                 |                        |       |                       |       |
| </=60 years                | 8                      | 3.7%  | 8                     | 3.8%  |
| 60-70 years                | 109                    | 50.9% | 99                    | 46.5% |
| >70 years                  | 97                     | 45.3% | 106                   | 49.8% |
| <b>FIGO 1988 stage</b>     |                        |       |                       |       |
| IB                         | 19                     | 8.9%  | 16                    | 7.5%  |
| IC                         | 172                    | 80.4% | 171                   | 80.3% |
| IIA                        | 23                     | 10.7% | 26                    | 12.2% |
| <b>Myometrial invasion</b> |                        |       |                       |       |
| < 50%                      | 31                     | 14.5% | 28                    | 13.1% |
| > 50%                      | 183                    | 85.5% | 185                   | 86.9% |
| <b>Grade</b>               |                        |       |                       |       |
| 1                          | 99                     | 46.3% | 103                   | 48.4% |
| 2                          | 97                     | 45.3% | 94                    | 44.1% |
| 3                          | 18                     | 8.4%  | 16                    | 7.5%  |
| <b>LVSI</b>                |                        |       |                       |       |
| Present                    | 25                     | 11.7% | 21                    | 9.9%  |
| Absent                     | 189                    | 88.3% | 192                   | 90.1% |

Confirmed-HIR at pathology review


| <i>n</i> = 344                      | EBRT ( <i>n</i> = 175) |       | VBT ( <i>n</i> = 169) |       |
|-------------------------------------|------------------------|-------|-----------------------|-------|
| <b>Grade</b>                        |                        |       |                       |       |
| 1                                   | 148                    | 84.6% | 134                   | 79.3% |
| 2                                   | 16                     | 9.1%  | 23                    | 13.6% |
| 3                                   | 11                     | 6.3%  | 12                    | 7.1%  |
| <b>Substantial LVSI</b>             |                        |       |                       |       |
| Yes                                 | 9                      | 5.1%  | 7                     | 4.1%  |
| No                                  | 160                    | 91.4% | 156                   | 92.3% |
| Missing <sup>a</sup>                | 6                      | 3.4%  | 6                     | 3.6%  |
| <b>Molecular subgroup</b>           |                        |       |                       |       |
| POLE                                | 10                     | 5.7%  | 6                     | 3.6%  |
| MSI                                 | 41                     | 23.4% | 36                    | 21.3% |
| NSMP                                | 103                    | 58.9% | 96                    | 56.8% |
| TP53 <sup>b</sup>                   | 10                     | 5.7%  | 15                    | 8.9%  |
| Double classifiers                  | 4                      | 2.3%  | 6                     | 3.6%  |
| Missing <sup>a</sup>                | 7                      | 4.0%  | 10                    | 5.9%  |
| <b>L1CAM expression<sup>c</sup></b> |                        |       |                       |       |
| > 10%                               | 4                      | 2.3%  | 14                    | 8.3%  |
| < 10%                               | 168                    | 96.0% | 151                   | 89.3% |
| Missing <sup>a</sup>                | 3                      | 1.7%  | 4                     | 2.4%  |

## PORTEC-2




**VR: 3,4 vs 2,4%.  $p = 0,55$**   
**PR: 6,3 vs 0,9%.  $p = 0,004$**

## PORTEC-2



**67 vs 69%.  $P = 0.72$**



# Adjuvant chemoradiotherapy versus radiotherapy alone for women with high-risk endometrial cancer (PORTEC-3): final results of an international, open-label, multicentre, randomised, phase 3 trial




*Stephanie M de Boer, Melanie E Powell, Linda Mileskin, Dionyssios Katsaros, Paul Bessette, Christine Haie-Meder, Petronella B Ottevanger, Jonathan A Ledermann, Pearly Khaw, Alessandro Colombo, Anthony Fyles, Marie-Helene Baron, Ina M Jürgenliemk-Schulz, Henry C Kitchener, Hans W Nijman, Godfrey Wilson, Susan Brooks, Silvestro Carinelli, Diane Provencher, Chantal Hanzen, Ludy C H W Lutgens, Vincent T H B M Smit, Naveena Singh, Viet Do, Romerai D'Amico, Remi A Nout, Amanda Feeney, Karen W Verhoeven-Adema, Hein Putter, Carien L Creutzberg, on behalf of the PORTEC study group\**



## PORTEC-3 trial design



### ➤ High risk Endometrial Cancer (HREC)




- uniform treatment schedule
- upfront pathology review
- quality of life analysis

# PORTEC-3


|                                       | Chemoradiotherapy<br>(n=330) | Radiotherapy<br>alone (n=330) |
|---------------------------------------|------------------------------|-------------------------------|
| <b>Age at randomisation (years)</b>   |                              |                               |
| Median                                | 62.4 (56.5-67.9)             | 62.0 (55.8-68.2)              |
| <60                                   | 128 (39%)                    | 140 (42%)                     |
| 60-69                                 | 144 (44%)                    | 128 (39%)                     |
| ≥70                                   | 58 (18%)                     | 62 (19%)                      |
| <b>Participating groups</b>           |                              |                               |
| NCRI (UK)                             | 82 (25%)                     | 95 (29%)                      |
| DGOG (Netherlands)                    | 72 (22%)                     | 66 (20%)                      |
| ANZGOG (Australia and<br>New Zealand) | 60 (18%)                     | 58 (18%)                      |
| MaNGO (Italy)                         | 52 (16%)                     | 46 (14%)                      |
| CCTG (Canada)                         | 36 (11%)                     | 29 (9%)                       |
| Fedegyn (France)                      | 28 (9%)                      | 36 (11%)                      |
| <b>FIGO stage</b>                     |                              |                               |
| Stage IA                              | 39 (12%)                     | 38 (12%)                      |
| Stage IB                              | 59 (18%)                     | 59 (18%)                      |
| Stage II                              | 80 (24%)                     | 90 (27%)                      |
| Stage III                             | 152 (46%)                    | 143 (43%)                     |
| <b>Histological grade and type</b>    |                              |                               |
| EEC grade 1                           | 68 (21%)                     | 56 (17%)                      |
| EEC grade 2                           | 59 (18%)                     | 73 (22%)                      |
| EEC grade 3                           | 90 (27%)                     | 95 (29%)                      |
| Serous                                | 53 (16%)                     | 52 (16%)                      |
| Clear cell                            | 29 (9%)                      | 33 (10%)                      |
| Mixed                                 | 17 (5%)                      | 13 (4%)                       |
| Other                                 | 14 (4%)                      | 8 (2%)                        |

|                              |           |           |
|------------------------------|-----------|-----------|
| <b>Myometrial invasion</b>   |           |           |
| <50%                         | 116 (35%) | 123 (37%) |
| ≥50%                         | 212 (65%) | 206 (63%) |
| Missing                      | 2 (<1%)   | 1 (<1%)   |
| <b>LVI</b>                   |           |           |
| Yes                          | 197 (60%) | 192 (58%) |
| No                           | 133 (40%) | 138 (42%) |
| <b>WHO performance score</b> |           |           |
| 0-1                          | 323 (99%) | 324 (99%) |
| ≥2                           | 5 (2%)    | 5 (2%)    |
| Missing                      | 2 (<1%)   | 1 (<1%)   |
| <b>Comorbidity</b>           |           |           |
| Diabetes                     | 45 (14%)  | 36 (11%)  |
| Hypertension                 | 116 (35%) | 104 (32%) |
| Cardiovascular               | 29 (9%)   | 20 (6%)   |

(Table 1 continues on next column)




**OS: 82 (QTRT) vs 77%. HR 0.76, 95% CI 0.54–1.06;  $p=0.109$ .**  
**FFS: 75,5 vs 68,6%. HR 0.71, 0.53–0.95;  $p=0.022$**



**E III 69,3 vs 58%. HR 0.66, 95% CI 0.45-0.97; p=0.014.**


A




# Molecular Classification of the PORTEC-3 Trial for High-Risk Endometrial Cancer: Impact on Prognosis and Benefit From Adjuvant Therapy

Alicia León-Castillo, MD<sup>1</sup>; Stephanie M. de Boer, MD<sup>2</sup>; Melanie E. Powell, MD<sup>3</sup>; Linda R. Mileschkin, MBBS<sup>4</sup>; Helen J. Mackay, MD<sup>5</sup>; Alexandra Leary, MD, PhD<sup>6</sup>; Hans W. Nijman, MD, PhD<sup>6,7</sup>; Naveena Singh, MD, MBBS<sup>8</sup>; Pamela M. Pollock, PhD<sup>9</sup>; Paul Bessette, MD<sup>10</sup>; Anthony Fyles, MD<sup>11</sup>; Christine Haie-Meder, MD<sup>12</sup>; Vincent T. H. B. M. Smit, MD, PhD<sup>1</sup>; Richard J. Edmondson, MD<sup>13</sup>; Hein Putter, MD<sup>14</sup>; Henry C. Kitchener, MD<sup>13</sup>; Emma J. Crosbie, MD, PhD<sup>13</sup>; Marco de Bruyn, PhD<sup>7</sup>; Remi A. Nout, MD<sup>2</sup>; Nanda Horeweg, MD, PhD<sup>2</sup>; Carien L. Creutzberg, MD, PhD<sup>2</sup>; and Tjalling Bosse, MD, PhD<sup>1</sup> on behalf of the TransPORTEC consortium

1. 93 (22.7%) p53abn
2. 51 (12.4%) POLEmut
3. 137 (33.4%) MMRd
4. 129 (31.5%) NSMP

**A**

|                 |     |     |     |     |    |    |
|-----------------|-----|-----|-----|-----|----|----|
| No. at risk:    |     |     |     |     |    |    |
| p53abn          | 93  | 72  | 57  | 49  | 44 | 32 |
| <i>POLE</i> mut | 51  | 50  | 50  | 49  | 48 | 37 |
| MMRd            | 137 | 124 | 112 | 102 | 96 | 74 |
| NSMP            | 129 | 122 | 113 | 105 | 94 | 69 |

**B**



|                 |     |     |     |     |     |    |
|-----------------|-----|-----|-----|-----|-----|----|
| No. at risk:    |     |     |     |     |     |    |
| p53abn          | 93  | 87  | 71  | 61  | 52  | 37 |
| <i>POLE</i> mut | 51  | 51  | 50  | 49  | 48  | 37 |
| MMRd            | 137 | 136 | 128 | 115 | 108 | 85 |
| NSMP            | 129 | 125 | 122 | 118 | 110 | 85 |



**TP53 → RFS 48% / OS 54.0%**



***POLE*mut → RFS y OS 98%**



**dMMR → RFS 72% / OS 81%**

**NSMP → RFS 74% / OS 88,5%**



**C****POLEmut EC****D****POLEmut EC**

**E****MMRd EC****F****MMRd EC**

**G****NSMP EC****H****NSMP EC**

ORIGINAL ARTICLE

# Adjuvant Chemotherapy plus Radiation for Locally Advanced Endometrial Cancer

Daniela Matei, M.D., Virginia Filiaci, Ph.D., Marcus E. Randall, M.D.,  
David Mutch, M.D., Margaret M. Steinhoff, M.D., Paul A. DiSilvestro, M.D.,  
Katherine M. Moxley, M.D., Yong M. Kim, M.D., Ph.D., Matthew A. Powell, M.D.,  
David M. O'Malley, M.D., Nick M. Spirtos, M.D., William Small, Jr., M.D.,  
Krishnansu S. Tewari, M.D., William E. Richards, M.D., John Nakayama, M.D.,  
Ursula A. Matulonis, M.D., Helen Q. Huang, M.S., and David S. Miller, M.D.

**GOG 258**

**Table 1. Characteristics of the Patients.\***

| Characteristic                          | Chemoradiotherapy<br>(N=370) | Chemotherapy<br>Only<br>(N=366) |
|-----------------------------------------|------------------------------|---------------------------------|
| Mean age (range) — yr                   | 60.5 (31–88)                 | 60 (31–85)                      |
| Race — no. (%)†                         |                              |                                 |
| White                                   | 291 (78.6)                   | 279 (76.2)                      |
| Black                                   | 37 (10.0)                    | 42 (11.5)                       |
| Asian, other, or not specified          | 42 (11.4)                    | 45 (12.3)                       |
| GOG performance status score — no. (%)‡ |                              |                                 |
| 0                                       | 278 (75.1)                   | 268 (73.2)                      |
| 1                                       | 88 (23.8)                    | 96 (26.2)                       |
| 2                                       | 4 (1.1)                      | 2 (0.5)                         |
| FIGO stage — no. (%)§                   |                              |                                 |
| I or II                                 | 6 (1.6)                      | 10 (2.7)                        |
| IIIA                                    | 70 (18.9)                    | 81 (22.1)                       |
| IIIB                                    | 12 (3.2)                     | 13 (3.6)                        |
| IIIC1                                   | 189 (51.1)                   | 166 (45.4)                      |
| IIIC2                                   | 90 (24.3)                    | 93 (25.4)                       |
| IVA                                     | 3 (0.8)                      | 3 (0.8)                         |

**Histology and grade — no. (%)**

|                           |            |            |
|---------------------------|------------|------------|
| Endometrioid, grade 1     | 87 (23.5)  | 79 (21.6)  |
| Endometrioid, grade 2     | 103 (27.8) | 118 (32.2) |
| Endometrioid, grade 3     | 64 (17.3)  | 61 (16.7)  |
| Serous                    | 66 (17.8)  | 65 (17.8)  |
| Clear cell                | 10 (2.7)   | 12 (3.3)   |
| Mixed epithelial or other | 40 (10.8)  | 31 (8.5)   |


**Gross residual disease — no. (%)**

|         |            |            |
|---------|------------|------------|
| Absent  | 360 (97.3) | 359 (98.1) |
| Present | 10 (2.7)   | 7 (1.9)    |


Median BMI (range)¶ 32.0 (11.2–65.3) 32.9 (18–60.2)

**BMI category — no. (%)**

|                             |            |            |
|-----------------------------|------------|------------|
| Normal or underweight       | 72 (19.5)  | 71 (19.4)  |
| Overweight                  | 84 (22.7)  | 81 (22.1)  |
| Obesity class I, II, or III | 214 (57.8) | 214 (58.5) |



**C Distant Recurrence**



**No. at Risk**  
Chemoradiotherapy  
Chemotherapy only


**Figure 1. Relapse**  
Tick marks indicate

| No. at Risk       |     |     |     |     |     |    |    |
|-------------------|-----|-----|-----|-----|-----|----|----|
| Chemoradiotherapy | 370 | 302 | 249 | 175 | 111 | 51 | 20 |
| Chemotherapy only | 366 | 316 | 263 | 186 | 132 | 62 | 22 |

**Table 2.** Acute Adverse Events.

| Adverse Event                 | Chemoradiotherapy (N = 346) |           | Chemotherapy Only (N = 361) |           | P Value* |
|-------------------------------|-----------------------------|-----------|-----------------------------|-----------|----------|
|                               | Any Grade                   | Grade 3–5 | Any Grade                   | Grade 3–5 |          |
|                               | <i>percent of patients</i>  |           |                             |           |          |
| Constitutional symptom        | 87                          | 6         | 80                          | 2         | 0.004    |
| Fatigue                       | 85                          | 5         | 75                          | 2         | <0.001   |
| Cardiac event                 | 16                          | 3         | 19                          | 4         | 0.71     |
| Endocrine event               | 11                          | 1         | 11                          | 0         | 0.46     |
| Gastrointestinal event        | 90                          | 13        | 79                          | 4         | <0.001   |
| Renal or genitourinary event  | 33                          | 2         | 11                          | 1         | <0.001   |
| Blood or bone marrow event    | 96                          | 40        | 90                          | 52        | 0.01     |
| Infection                     | 23                          | 4         | 20                          | 5         | 0.18     |
| Lymphatic event               | 17                          | <1        | 16                          | <1        | 0.74     |
| Musculoskeletal event         | 20                          | 3         | 13                          | 1         | 0.01     |
| Metabolic or laboratory event | 48                          | 15        | 44                          | 9         | 0.02     |
| Neurologic event              | 76                          | 7         | 80                          | 5         | 0.99     |
| Pulmonary event               | 31                          | 2         | 26                          | 1         | 0.83     |
| Pain                          | 70                          | 8         | 68                          | 5         | 0.04     |


# Diseño del Estudio KEYNOTE-B21



**Table 1. Demographics and baseline disease characteristics in the ITT population**

|                                     | <b>Pembrolizumab +<br/>chemotherapy<br/>(n = 545)</b> | <b>Placebo +<br/>chemotherapy<br/>(n = 550)</b> |
|-------------------------------------|-------------------------------------------------------|-------------------------------------------------|
| Age, median (range),<br>years       | 62.0 (29.0-95.0)                                      | 62.0 (27.0-89.0)                                |
| <65                                 | 318 (58)                                              | 335 (61)                                        |
| ≥65                                 | 227 (42)                                              | 215 (39)                                        |
| Race                                |                                                       |                                                 |
| White                               | 315 (58)                                              | 362 (66)                                        |
| Asian                               | 189 (35)                                              | 157 (29)                                        |
| Black or<br>African American        | 11 (2)                                                | 13 (2)                                          |
| American Indian<br>or Alaska Native | 2 (<1)                                                | 3 (<1)                                          |
| Multiple                            | 23 (4)                                                | 10 (2)                                          |
| Missing                             | 5 (<1)                                                | 5 (<1)                                          |
| Region                              |                                                       |                                                 |
| Western Europe                      | 146 (27)                                              | 165 (30)                                        |
| North America                       | 94 (17)                                               | 119 (22)                                        |
| Rest of the world                   | 305 (56)                                              | 266 (48)                                        |
| ECOG performance<br>status          |                                                       |                                                 |
| 0                                   | 409 (75)                                              | 416 (76)                                        |
| 1                                   | 136 (25)                                              | 134 (24)                                        |
| Lymph node<br>dissection            |                                                       |                                                 |
| Yes                                 | 483 (89)                                              | 502 (91)                                        |
| No                                  | 62 (11)                                               | 48 (9)                                          |

|                             |          |          |
|-----------------------------|----------|----------|
| FIGO (2009) stage           |          |          |
| IA                          | 83 (15)  | 94 (17)  |
| IB                          | 63 (12)  | 50 (9)   |
| II                          | 40 (7)   | 41 (7)   |
| IIIA                        | 109 (20) | 94 (17)  |
| IIIB                        | 20 (4)   | 19 (3)   |
| IIIC1                       | 144 (26) | 169 (31) |
| IIIC2                       | 78 (14)  | 81 (15)  |
| IVA                         | 6 (1)    | 1 (<1)   |
| IVB <sup>a</sup>            | 2 (<1)   | 1 (<1)   |
| Histology                   |          |          |
| Endometrioid                | 297 (54) | 297 (54) |
| Non-endometrioid            | 248 (46) | 253 (46) |
| MMR status                  |          |          |
| pMMR                        | 404 (74) | 410 (75) |
| dMMR                        | 141 (26) | 140 (25) |
| Planned radiotherapy        |          |          |
| EBRT with cisplatin         | 94 (17)  | 95 (17)  |
| EBRT without cisplatin      | 256 (47) | 246 (45) |
| Brachytherapy only          | 49 (9)   | 52 (9)   |
| No EBRT or<br>brachytherapy | 146 (27) | 157 (29) |
| Lymph node<br>involvement   |          |          |
| No                          | 300 (55) | 284 (52) |
| Yes                         | 223 (41) | 250 (45) |
| Not evaluable               | 22 (4)   | 16 (3)   |



# HR: 1.02

(IC 95% 0.79 - 1.32),  $p = 0.570$

**Tasa DFS a 2 años:**

75% (Pembrolizumab)  
76% (Placebo)


**Eventos:** 22% en ambos brazos.

**OS:** Datos inmaduros (solo 10% de eventos totales).

**Conclusión:** En una población no seleccionada, la combinación adyuvante no altera la trayectoria de recurrencia en el cáncer de endometrio de alto riesgo.

# Eficacia en dMMR: Beneficio Clínico Transformador

Pembrolizumab reduce drásticamente el riesgo de recurrencia en tumores dMMR (n=281).



## HR: 0.31

(IC 95% 0.14 - 0.69)


**Mediana DFS:** No alcanzada en ningún brazo.

### Nota de Contexto:

Esta magnitud de beneficio (HR 0.31) es casi idéntica a la observada en enfermedad avanzada (KEYNOTE-868 HR 0.30), confirmando la profunda sensibilidad de la biología dMMR a la inmunoterapia temprana.

# Eficacia en pMMR: Ausencia de Beneficio

En tumores proficientes (n=814), la inmunoterapia adyuvante no altera el pronóstico.



## HR: 1.20

(IC 95% 0.91 - 1.57)

**Eventos:** 27% (Pembro) vs 23% (Placebo).

### Conclusión:

En el escenario de enfermedad microscópica (adyuvancia), la combinación de QT + PD-1 no logra inducir la respuesta inmunitaria necesaria en fenotipos tumorales intrínsecamente menos inmunogénicos.

Cualquier AE Grado  $\geq 3$


**71% (Pembro)**  
**vs 63% (Placebo)**

Discontinuación de cualquier  
fármaco por AE


**24% vs 16%**

Muertes relacionadas al tratamiento  
**0 en ambos brazos**

Eventos Más Frecuentes (Cualquier Grado  $>40\%$ )



|                                                | Pembro | Placebo |
|------------------------------------------------|--------|---------|
| Incidencia General de irAEs                    | 42%    | 24%     |
| irAEs Grado $\geq 3$                           | 10%    | 4%      |
| irAEs Clave (Incidencias Notables)             |        |         |
| Hipotiroidismo<br>*Mayoritariamente Grados 1/2 | 23%*   | 4%      |
| Hipertiroidismo                                | 12%    | 3%      |
| Reacciones a la infusión                       | 8%     | 14%     |
| Reacciones cutáneas severas                    | 5%     | <1%     |



**Stage I-IVA EC: adjuvant therapy for high-intermediate risk and high-risk patients<sup>a</sup>**

**High-intermediate risk**

High-intermediate risk,  
pN0 after lymph node staging

High-intermediate risk, without  
lymph-node staging

For patients with stage IA and IB with substantial LVSI  
For patients with stage IB G3  
For patients with stage II G1 with substantial LVSI  
For patients with stage II G2-G3 (dMMR or NSMP)

- Adjuvant EBRT is recommended [I, A]
- Adding (concomitant and/or sequential) ChT to EBRT could be considered, especially for G3 and/or substantial LVSI [II, C]

**High risk**

All stages and all histologies with p53-abn and myometrial invasion

All stages with serous or undifferentiated carcinoma including carcinosarcoma with myometrial invasion

All Stage III and IVA with no residual tumour, regardless of histology and regardless of molecular subtype

- Adjuvant EBRT + concurrent ChT [I, A]
- Sequential ChT and RT [I, B]
- ChT alone [I, B]

**Desescalamiento de RT en  
pacientes dMMR?**

# Criterios de adyuvancia



## Bajo riesgo

No se recomienda adyuvancia

## Intermedio

Braquiterapia o vigilancia.



## Intermedio alto

RT externa

## Alto riesgo

Quimiorradioterapia concurrente (IA) –  
Quimiorradioterapia secuencial – QT sola



# MANAGEMENT OF PATIENTS WITH DOUBLE CLASSIFIER ENDOMETRIAL CANCER

Alexandra Leary, MD, PhD

Deputy Director Dept of Medical Oncology

Institut Gustave Roussy


GINECO, France






# CLINICAL BEHAVIOR OF DOUBLE CLASSIFIERS

**POLEm/p53abn** behave like POLEm,  
not **p53abn**



**MMRd/p53abn** behave like MMRd,  
not **p53abn**





# MULTIPLE CLASSIFIERS

What we know regarding genomic profile and prognosis


**POLEm** 'trumps' **MMRd** and **p53abn** in terms of genomic profile and prognosis

**MMRd** 'trumps' **p53abn** in terms of genomic profile and prognosis

Double classifiers should be classified as:



# BUT LINES BETWEEN MOLECULAR CLASSES GET EVEN MORE CONFUSING...




Adapted from Vermij et al; Histopath 2019


\*Pathogenic *POLE* exonuclease domain mutations (EDM) as per León-Castillo et al. (J Pathol)

# PROGNOSIS OF ER-NEG NSMP

Within the NSMP subclass, ER-neg stand out



Within molecular classified EC, ER-neg NSMP behave like p53abn...



# Conclusiones

- ❖ El riesgo es lo más importante
- ❖ La clasificación molecular como pilar del manejo del cancer de endometrio
- ❖ Inmunoterapia en localizados??



Enfermedad avanzada

# GOG 209

**TAP vs TC**

**Phase III study**

- Endometrial ca.
- Surgical stage III / IV or Recurrent
- Measurable disease
- ER, PR status

**Randomize**

## Regimen I

**Doxorubicin** 45 mg / m<sup>2</sup>  
**CDDP** 50 mg / m<sup>2</sup>  
day 1


**Paclitaxel** 160 mg / m<sup>2</sup>  
day 2

G-CSF


## Regimen II

**Paclitaxel** 175 mg / m<sup>2</sup>  
**CBDCA** AUC = 6  
day 1

Ongoing with GOG Japan (JGOG)



13 (TC) vs 14 meses (TAP). HR 1.032, 90% CI 0.93 to 1.15



37 vs 41 meses. HR 1.002, 95% CI 0.88 to 1.15

# NRG-GY018 (NCT03914612)

## Key Eligibility Criteria

- Measurable stage III/IVA or measurable/nonmeasurable stage IVB or recurrent endometrial cancer
- Pathology report showing results of institutional MMR IHC testing
- ECOG PS 0, 1, or 2
- No prior chemo except prior adjuvant chemo if completed  $\geq 12$  mo before study

## Stratification Factors

- dMMR vs pMMR
- ECOG PS (0 or 1 vs 2)
- Prior adjuvant chemo (yes vs no)

N = 816  
(591 pMMR,  
225 dMMR)

R  
1:1

**Arm 1**  
Placebo IV Q3W +  
Paclitaxel 175 mg/m<sup>2</sup> IV Q3W +  
Carboplatin AUC 5 IV Q3W  
for 6 cycles

**Arm 1**  
Placebo IV Q6W  
for up to 14 additional  
cycles

**Arm 2**  
Pembrolizumab 200 mg IV Q3W +  
Paclitaxel 175 mg/m<sup>2</sup> IV Q3W +  
Carboplatin AUC 5 IV Q3W  
for 6 cycles

**Arm 2**  
Pembrolizumab  
400 mg IV Q6W  
for up to 14 additional  
cycles

## Endpoints

- **Primary:** PFS per RECIST v1.1 by investigator in pMMR and dMMR populations
- **Secondary:** Safety, ORR/DOR per RECIST v1.1 by BICR or investigator by treatment arm and MMR IHC status, OS in pMMR and dMMR populations, PRO/QoL in pMMR population, and concordance of institutional vs central MMR IHC testing results




ANNUAL MEETING  
ON WOMEN'S CANCER  
TAMPA, FL - 2019

BICR, blinded independent central review; dMMR, mismatch repair deficient; DOR, duration of response; ECOG PS, Eastern Cooperative Oncology Group performance status; IHC, immunohistochemistry; ORR, objective response rate; OS, overall survival; PFS, progression-free survival; pMMR, mismatch repair proficient; PRO, patient-reported outcomes; QoL, quality of life; RECIST, Response Evaluation Criteria in Solid Tumors.

**Table 1** *Characteristic of the Patients at Racalino* ☆


|                                      |            |            |            |            |            |            |          |
|--------------------------------------|------------|------------|------------|------------|------------|------------|----------|
| Histologic analysis — no. (%)        |            |            |            |            |            |            |          |
| Adenocarcinoma, NOS§                 | 12 (10.7)  | 14 (12.4)  | 26 (11.6)  | 24 (8.2)   | 33 (11.2)  | 57 (9.7)   |          |
| Clear cell                           | 1 (0.9)    | 0          | 1 (0.4)    | 17 (5.8)   | 20 (6.8)   | 37 (6.3)   |          |
| Dedifferentiated or undifferentiated | 4 (3.6)    | 4 (3.5)    | 8 (3.6)    | 7 (2.4)    | 6 (2.0)    | 13 (2.2)   |          |
| Endometrioid                         |            |            |            |            |            |            |          |
| G1                                   | 21 (18.8)  | 35 (31.0)  | 56 (24.9)  | 54 (18.4)  | 46 (15.6)  | 100 (17.0) |          |
| G2                                   | 52 (46.4)  | 41 (36.3)  | 93 (41.3)  | 51 (17.4)  | 58 (19.7)  | 109 (18.5) |          |
| G3                                   | 15 (13.4)  | 16 (14.2)  | 31 (13.8)  | 53 (18.1)  | 42 (14.2)  | 95 (16.2)  |          |
| Mixed epithelial                     | 3 (2.7)    | 2 (1.8)    | 5 (2.2)    | 6 (2.0)    | 11 (3.7)   | 17 (2.9)   |          |
| Serous                               | 4 (3.6)    | 1 (0.9)    | 5 (2.2)    | 78 (26.6)  | 72 (24.4)  | 150 (25.5) |          |
| Pending                              | 0          | 0          | 0          | 3 (1.0)    | 7 (2.4)    | 10 (1.7)   |          |
| Previous therapy                     |            |            |            |            |            |            |          |
| Chemotherapy — no. (%)               |            |            |            |            |            |            |          |
| Yes                                  | 5 (4.5)    | 8 (7.1)    | 13 (5.8)   | 72 (24.6)  | 77 (26.1)  | 149 (25.3) |          |
| No                                   | 107 (95.5) | 105 (92.9) | 212 (94.2) | 221 (75.4) | 218 (73.9) | 439 (74.7) |          |
| Radiotherapy — no. (%)               |            |            |            |            |            |            |          |
| Yes                                  | 41 (36.6)  | 55 (48.7)  | 96 (42.7)  | 114 (38.9) | 119 (40.3) | 233 (39.6) |          |
| No                                   | 71 (63.4)  | 58 (51.3)  | 129 (57.3) | 179 (61.1) | 176 (59.7) | 355 (60.4) |          |
| Surgery — no. (%)                    |            |            |            |            |            |            |          |
| Yes                                  | 98 (87.5)  | 105 (92.9) | 203 (90.2) | 261 (89.1) | 245 (83.1) | 506 (86.1) |          |
| No                                   | 14 (12.5)  | 8 (7.1)    | 22 (9.8)   | 29 (9.9)   | 46 (15.6)  | 75 (12.8)  |          |
| Missing data                         | 0          | 0          | 0          | 3 (1.0)    | 4 (1.4)    | 7 (1.2)    |          |
|                                      | 2          | 1 (0.9)    | 3 (4.4)    | 6 (2.7)    | 9 (3.1)    | 9 (3.1)    | 10 (3.1) |

# SLP en la Cohorte dMMR: Una Separación Temprana y Sostenida



Mediana de seguimiento de 12 meses (Corte NEJM 2023).

# SLP en la Cohorte pMMR: Eficacia en el Fenotipo de Mayor Riesgo



# Supervivencia Global dMMR (ASCO 2026): El Beneficio Inquebrantable

**80% vivos a los 48  
meses (Pembrolizumab)**

**60% vivos a los 48  
meses (Placebo)**


**HR 0.53 (IC 95%, 0.32-0.89)**

## El Efecto Crossover

Esta ventaja en supervivencia (HR 0.53) se mantuvo intacta a pesar de que  $\geq 55\%$  (n=62) de los pacientes en el brazo placebo recibieron ICI post-estudio.

# Supervivencia Global pMMR (ASCO 2026):


## 11.8 Meses de Ventaja



Ventaja clínica sostenida a pesar de que  $\geq 57\%$  ( $n=167$ ) del brazo placebo fue rescatado con ICI (mediana de duración de tratamiento rescate: 6.2 meses).

# ENGOT-EN6-NSGO/GOG-3031/RUBY (NCT03981796)

Phase 3, randomized, double-blind, multicenter study of dostarlimab plus carboplatin-paclitaxel versus placebo plus carboplatin/paclitaxel in patients with primary advanced or recurrent EC




On-study imaging assessments are to be performed Q6W (±7 days) from the randomization date until Week 25 (Cycle 8), followed by Q9W (±7 days) until Week 52. Subsequent tumor imaging is to be performed every 12 weeks (±7 days) until radiographic PD is documented by investigator assessment per RECIST v1.1 followed by one additional imaging 4-6 weeks later, or subsequent anticancer therapy is started, whichever occurs first. Thereafter, scans may be performed per standard of care.

<sup>a</sup>Mixed histology containing at least 10% carcinosarcoma, clear cell, or serous histology. <sup>b</sup>Patients were randomized based on either local or central MMR/MSI testing results. Central testing was used if local results were not available. For local determination of MMR/MSI status, IHC, next generation sequencing, and polymerase chain reaction assays were accepted. For central determination of MMR/MSI status IHC per Ventana MMR RxDx panel was used. <sup>c</sup>Treatment ends after 3 years, PD, toxicity, withdrawal of consent, investigator's decision, or death, whichever occurs first. Continued treatment with dostarlimab or placebo beyond 3 years may be considered following discussion between the Sponsor and the Investigator. AUC, area under the plasma or serum concentration-time curve; BICR, blinded independent central review; DCR, disease control rate; DOR, duration of response; EC, endometrial cancer; ECOG PS, Eastern Cooperative Oncology Group performance status; EOT, end of treatment; HRQOL, health-related quality of life; IHC, immunohistochemistry; INV, investigator assessment; MMR, mismatch repair; MSI, microsatellite instability; ORR, objective response rate; OS, overall survival; PD, progressive disease; PFS, progression-free survival; PRO, patient-reported outcome; R, randomization; RECIST v1.1, Response Evaluation Criteria in Solid Tumors version 1.1.

**Table 1.** Demographic and Clinical Characteristics of the

| Characteristic                                  |  | Histologic type — no. (%) |              |                                                                               |                       |
|-------------------------------------------------|--|---------------------------|--------------|-------------------------------------------------------------------------------|-----------------------|
|                                                 |  | Carcinosarcoma            | Endometrioid | Mixed carcinoma ≥10% of carcinosarcoma, clear-cell, or serous histologic type | Serous adenocarcinoma |
|                                                 |  | 4 (8)                     | 44 (83)      | 2 (4)                                                                         | 1 (2)                 |
|                                                 |  | 1 (2)                     | 56 (86)      | 4 (6)                                                                         | 1 (2)                 |
|                                                 |  | 25 (10.2)                 | 134 (54.7)   | 10 (4.1)                                                                      | 50 (20.4)             |
|                                                 |  | 19 (7.6)                  | 136 (54.6)   | 9 (3.6)                                                                       | 52 (20.9)             |
| Age                                             |  |                           |              |                                                                               |                       |
| Median (range) — yr                             |  |                           |              |                                                                               |                       |
| ≥65 Yr — no. (%)                                |  |                           |              |                                                                               |                       |
| Race or ethnic group — no. (%) †                |  |                           |              |                                                                               |                       |
| White                                           |  |                           |              |                                                                               |                       |
| Black                                           |  |                           |              |                                                                               |                       |
| Asian                                           |  |                           |              |                                                                               |                       |
| American Indian or Alaska Native                |  |                           |              |                                                                               |                       |
| Native Hawaiian or other Pacific Islander       |  |                           |              |                                                                               |                       |
| Unknown or not reported                         |  |                           |              |                                                                               |                       |
| ECOG performance category — no./total no. (%) ‡ |  |                           |              |                                                                               |                       |
| 0                                               |  |                           |              |                                                                               |                       |
| 1                                               |  |                           |              |                                                                               |                       |
| FIGO stage at diagnosis — no. (%) §             |  |                           |              |                                                                               |                       |
| I                                               |  |                           |              |                                                                               |                       |
| II                                              |  |                           |              |                                                                               |                       |
| III                                             |  |                           |              |                                                                               |                       |
| IV                                              |  |                           |              |                                                                               |                       |
| Unknown                                         |  |                           |              |                                                                               |                       |

## B Overall Population



### No. at Risk


|                   |     |     |     |     |     |     |    |    |    |    |    |    |    |    |    |    |   |   |   |   |
|-------------------|-----|-----|-----|-----|-----|-----|----|----|----|----|----|----|----|----|----|----|---|---|---|---|
| Dostarlimab group | 245 | 220 | 197 | 157 | 130 | 105 | 94 | 90 | 84 | 78 | 66 | 52 | 34 | 23 | 22 | 12 | 2 | 0 |   |   |
| Placebo group     | 249 | 219 | 200 | 144 | 103 | 74  | 59 | 57 | 48 | 42 | 39 | 32 | 20 | 14 | 13 | 5  | 2 | 1 | 1 | 0 |

### No. of Events

|                   |   |    |    |    |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-------------------|---|----|----|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Dostarlimab group | 0 | 12 | 25 | 55 | 80  | 103 | 110 | 113 | 118 | 122 | 127 | 128 | 131 | 132 | 132 | 133 | 134 | 135 |     |     |
| Placebo group     | 0 | 14 | 29 | 77 | 115 | 141 | 155 | 157 | 166 | 170 | 170 | 172 | 175 | 176 | 176 | 177 | 177 | 177 | 177 | 177 |

# 1. ITT

## Primary endpoint




No. at risk(events)


|                  |        |        |        |         |         |         |         |         |         |         |         |          |          |          |          |          |         |         |         |         |         |         |        |        |        |
|------------------|--------|--------|--------|---------|---------|---------|---------|---------|---------|---------|---------|----------|----------|----------|----------|----------|---------|---------|---------|---------|---------|---------|--------|--------|--------|
| Dostarlimab + CP | 245(0) | 239(3) | 232(9) | 223(16) | 211(27) | 201(37) | 198(40) | 188(48) | 184(51) | 181(54) | 175(60) | 168(67)  | 164(71)  | 154(80)  | 148(86)  | 137(93)  | 118(98) | 95(102) | 70(104) | 52(107) | 37(107) | 17(108) | 6(108) | 2(109) | 0(109) |
| Placebo + CP     | 249(0) | 244(3) | 239(8) | 228(19) | 223(23) | 210(36) | 197(47) | 181(63) | 168(74) | 156(84) | 143(97) | 135(105) | 127(111) | 122(116) | 117(120) | 112(123) | 96(127) | 78(131) | 53(134) | 39(139) | 22(142) | 13(143) | 4(144) | 2(144) | 0(144) |



## 2. dMMR:



### 3. pMMR



No. at risk(events)


Dostarlimab + CP 192(0) 187(2) 182(6) 175(11) 165(21) 156(30) 153(33) 144(41) 140(44) 137(47) 131(53) 125(59) 122(62) 113(71) 106(77) 96(84) 84(88) 67(91) 51(93) 38(96) 29(96) 11(96) 4(96) 1(97) 0(97)

Placebo + CP 184(0) 181(1) 177(5) 169(12) 167(14) 155(26) 146(34) 133(47) 125(54) 115(63) 104(74) 98(80) 93(84) 89(88) 86(91) 81(94) 73(95) 59(98) 41(101) 28(106) 15(108) 7(109) 3(109) 2(109) 0(109)

<sup>a</sup>Overall survival in the dMMR/MSS-H and MMRp/MSS populations was a prespecified exploratory endpoint. <sup>b</sup>Median expected duration of follow-up; range 31.2-49.5 months. CP, carboplatin-paclitaxel; HR, hazard ratio; MMRp, mismatch repair proficient; MSS, microsatellite stable; NE, not estimable; OS, overall survival.



Scan for slides



FIGO stage in newly diagnosed patients<sup>c</sup>

|     |           |           |            |    |
|-----|-----------|-----------|------------|----|
| I   | 1 (0.4)   | 0         | 0          | 1) |
| II  | 0         | 0         | 1 (0.4)    |    |
| III | 12 (5.0)  | 17 (7.1)  | 12 (5.0)   |    |
| IV  | 99 (41.4) | 96 (40.3) | 101 (41.9) |    |

Histology type,<sup>d</sup> No. (%)

|                   |            |            |            |
|-------------------|------------|------------|------------|
| Endometrioid      | 152 (63.6) | 141 (59.2) | 139 (57.7) |
| Serous            | 42 (17.6)  | 58 (24.4)  | 54 (22.4)  |
| Carcinosarcoma    | 18 (7.5)   | 12 (5.0)   | 21 (8.7)   |
| Mixed, epithelial | 9 (3.8)    | 9 (3.8)    | 11 (4.6)   |
| Clear cell        | 8 (3.3)    | 4 (1.7)    | 7 (2.9)    |
| Undifferentiated  | 5 (2.1)    | 4 (1.7)    | 3 (1.2)    |
| Mucinous          | 0          | 1 (0.4)    | 0          |
| Other             | 5 (2.1)    | 9 (3.8)    | 6 (2.5)    |

MMR status,<sup>ae</sup> No. (%)

|            |            |            |            |
|------------|------------|------------|------------|
| Proficient | 191 (79.9) | 192 (80.7) | 192 (79.7) |
| Deficient  | 48 (20.1)  | 46 (19.3)  | 49 (20.3)  |

HRRm status,<sup>f</sup> No. (%)

|          |            |            |            |
|----------|------------|------------|------------|
| HRRm     | 39 (16.3)  | 26 (10.9)  | 32 (13.3)  |
| Non-HRRm | 141 (59.0) | 138 (58.0) | 132 (54.8) |
| Unknown  | 59 (24.7)  | 74 (31.1)  | 77 (32.0)  |

PD-L1 expression,<sup>g</sup> No. (%)


|          |            |            |            |
|----------|------------|------------|------------|
| Positive | 150 (62.8) | 170 (71.4) | 163 (67.6) |
| Negative | 82 (34.3)  | 61 (25.6)  | 75 (31.1)  |
| Unknown  | 7 (2.9)    | 7 (2.9)    | 3 (1.2)    |

Previous chemotherapy, No. (%)

|     |           |           |           |
|-----|-----------|-----------|-----------|
| Yes | 54 (22.6) | 51 (21.4) | 51 (21.2) |
|-----|-----------|-----------|-----------|

(continued on following page)


A



No. at risk:

|                           | 0   | 3   | 6   | 9   | 12  | 15 | 18 | 21 | 24 | 27 | 30 | 33 |
|---------------------------|-----|-----|-----|-----|-----|----|----|----|----|----|----|----|
| Durvalumab + olaparib arm | 239 | 214 | 198 | 169 | 139 | 95 | 51 | 30 | 16 | 7  | 3  | 0  |
| Durvalumab arm            | 238 | 211 | 188 | 138 | 105 | 69 | 45 | 26 | 13 | 5  | 0  | 0  |
| Control arm               | 241 | 213 | 184 | 125 | 86  | 45 | 26 | 10 | 3  | 1  | 1  | 0  |


B




No. at risk:

|                           | 0   | 3   | 6   | 9   | 12  | 15  | 18  | 21 | 24 | 27 | 30 | 33 | 36 |
|---------------------------|-----|-----|-----|-----|-----|-----|-----|----|----|----|----|----|----|
| Durvalumab + olaparib arm | 239 | 233 | 227 | 208 | 202 | 152 | 109 | 77 | 38 | 18 | 8  | 2  | 0  |
| Durvalumab arm            | 238 | 227 | 221 | 205 | 192 | 147 | 105 | 64 | 34 | 17 | 6  | 0  | 0  |
| Control arm               | 241 | 229 | 215 | 201 | 185 | 138 | 104 | 69 | 35 | 15 | 4  | 0  | 0  |

ITT

**A**

**B**

**TABLE 2. PFS in the ITT Population and by MMR and PD-L1 Subgroup Status**



| Subgroup                                   | Durvalumab + Olaparib Arm             | Durvalumab Arm                       | Control Arm         |
|--------------------------------------------|---------------------------------------|--------------------------------------|---------------------|
| ITT                                        | n = 239                               | n = 238                              | n = 241             |
| Progression events or death, No. (%)       | 126 (52.7)                            | 139 (58.4)                           | 173 (71.8)          |
| PFS, months, median (95% CI) <sup>a</sup>  | 15.1 (12.6 to 20.7)                   | 10.2 (9.7 to 14.7)                   | 9.6 (9.0 to 9.9)    |
| HR (95% CI) v control arm <sup>b</sup>     | 0.55 (0.43 to 0.69); <i>P</i> < .0001 | 0.71 (0.57 to 0.89); <i>P</i> = .003 |                     |
| HR (95% CI) v durvalumab arm <sup>b</sup>  | 0.78 (0.61 to 0.99)                   |                                      |                     |
| 6-month PFS rate, % (95% CI) <sup>a</sup>  | 83.9 (78.6 to 88.0)                   | 83.8 (78.4 to 88.0)                  | 82.5 (76.9 to 86.8) |
| 12-month PFS rate, % (95% CI) <sup>a</sup> | 61.5 (54.9 to 67.4)                   | 48.5 (41.8 to 54.9)                  | 41.1 (34.6 to 47.5) |
| 18-month PFS rate, % (95% CI) <sup>a</sup> | 46.3 (39.2 to 53.0)                   | 37.8 (31.0 to 44.5)                  | 21.7 (16.0 to 27.9) |
| dMMR                                       | n = 48                                | n = 46                               | n = 49              |
| Progression events or death, No. (%)       | 18 (37.5)                             | 15 (32.6)                            | 25 (51.0)           |
| PFS, months, median (95% CI) <sup>a</sup>  | 31.8 (12.4 to NR)                     | NR (NR to NR)                        | 7.0 (6.7 to 14.8)   |
| HR (95% CI) v control arm <sup>c</sup>     | 0.41 (0.21 to 0.75)                   | 0.42 (0.22 to 0.80)                  |                     |
| HR (95% CI) v durvalumab arm <sup>c</sup>  | 0.97 (0.49 to 1.98)                   |                                      |                     |
| 6-month PFS rate, % (95% CI) <sup>a</sup>  | 87.2 (73.8 to 94.1)                   | 90.6 (76.9 to 96.4)                  | 73.1 (56.6 to 84.2) |
| 12-month PFS rate, % (95% CI) <sup>a</sup> | 70.0 (54.7 to 81.0)                   | 67.9 (51.1 to 80.0)                  | 43.3 (27.3 to 58.3) |
| 18-month PFS rate, % (95% CI) <sup>a</sup> | 62.7 (46.9 to 75.0)                   | 67.9 (51.1 to 80.0)                  | 31.7 (16.7 to 47.9) |

| pMMR                                       | n = 191             | n = 192             | n = 192             |
|--------------------------------------------|---------------------|---------------------|---------------------|
| Progression events or death, No. (%)       | 108 (56.5)          | 124 (64.6)          | 148 (77.1)          |
| PFS, months, median (95% CI) <sup>a</sup>  | 15.0 (12.4 to 18.0) | 9.9 (9.4 to 12.5)   | 9.7 (9.2 to 10.1)   |
| HR (95% CI) v control arm <sup>c</sup>     | 0.57 (0.44 to 0.73) | 0.77 (0.60 to 0.97) |                     |
| HR (95% CI) v durvalumab arm <sup>c</sup>  | 0.76 (0.59 to 0.99) |                     |                     |
| 6-month PFS rate, % (95% CI) <sup>a</sup>  | 83.1 (77.0 to 87.7) | 82.4 (76.1 to 87.1) | 84.4 (78.4 to 88.9) |
| 12-month PFS rate, % (95% CI) <sup>a</sup> | 59.4 (52.0 to 66.0) | 44.4 (37.1 to 51.4) | 40.8 (33.6 to 47.8) |
| 18-month PFS rate, % (95% CI) <sup>a</sup> | 42.0 (34.1 to 49.6) | 31.3 (24.2 to 38.6) | 20.0 (14.1 to 26.7) |
| PD-L1-positive <sup>d</sup>                | n = 150             | n = 170             | n = 163             |
| Progression events or death, No. (%)       | 68 (45.3)           | 97 (57.1)           | 114 (69.9)          |
| PFS, months, median (95% CI) <sup>a</sup>  | 20.8 (15.1 to NR)   | 11.3 (9.7 to 15.4)  | 9.5 (7.9 to 9.9)    |
| HR (95% CI) v control arm <sup>c</sup>     | 0.42 (0.31 to 0.57) | 0.63 (0.48 to 0.83) |                     |
| HR (95% CI) v durvalumab arm <sup>c</sup>  | 0.67 (0.49 to 0.91) |                     |                     |
| 6-month PFS rate, % (95% CI) <sup>a</sup>  | 85.2 (78.3 to 90.0) | 89.0 (83.1 to 92.9) | 81.7 (74.6 to 87.0) |
| 12-month PFS rate, % (95% CI) <sup>a</sup> | 67.3 (59.0 to 74.2) | 48.8 (40.8 to 56.3) | 38.6 (30.7 to 46.4) |
| 18-month PFS rate, % (95% CI) <sup>a</sup> | 54.8 (45.7 to 63.0) | 40.2 (32.1 to 48.2) | 21.5 (14.7 to 29.3) |

TABLE 3. Summary of AEs

| AE, No. (%)                                           | Overall (chemotherapy phase + maintenance phase) |                             |                          | Maintenance Phase                      |                             |                          |
|-------------------------------------------------------|--------------------------------------------------|-----------------------------|--------------------------|----------------------------------------|-----------------------------|--------------------------|
|                                                       | Durvalumab + Olaparib Arm<br>(n = 238)           | Durvalumab Arm<br>(n = 235) | Control Arm<br>(n = 236) | Durvalumab + Olaparib Arm<br>(n = 192) | Durvalumab Arm<br>(n = 183) | Control Arm<br>(n = 169) |
| Any-grade AE <sup>a</sup>                             | 237 (99.6)                                       | 232 (98.7)                  | 236 (100)                | 184 (95.8)                             | 158 (86.3)                  | 143 (84.6)               |
| Anemia <sup>b</sup>                                   | 147 (61.8)                                       | 112 (47.7)                  | 128 (54.2)               | 70 (36.5)                              | 16 (8.7)                    | 17 (10.1)                |
| Nausea                                                | 130 (54.6)                                       | 96 (40.9)                   | 105 (44.5)               | 79 (41.1)                              | 22 (12.0)                   | 25 (14.8)                |
| Fatigue or asthenia                                   | 129 (54.2)                                       | 101 (43.0)                  | 105 (44.5)               | 62 (32.3)                              | 19 (10.4)                   | 21 (12.4)                |
| Alopecia                                              | 121 (50.8)                                       | 118 (50.2)                  | 118 (50.0)               | 5 (2.6)                                | 2 (1.1)                     | 1 (0.6)                  |
| Neutropenia <sup>c</sup>                              | 99 (41.6)                                        | 84 (35.7)                   | 98 (41.5)                | 34 (17.7)                              | 13 (7.1)                    | 7 (4.1)                  |
| Constipation                                          | 78 (32.8)                                        | 64 (27.2)                   | 81 (34.3)                | 13 (6.8)                               | 13 (7.1)                    | 9 (5.3)                  |
| Thrombocytopenia <sup>d</sup>                         | 71 (29.8)                                        | 66 (28.1)                   | 52 (22.0)                | 27 (14.1)                              | 6 (3.3)                     | 9 (5.3)                  |
| Diarrhea                                              | 67 (28.2)                                        | 74 (31.5)                   | 66 (28.0)                | 34 (17.7)                              | 28 (15.3)                   | 20 (11.8)                |
| Vomiting                                              | 61 (25.6)                                        | 49 (20.9)                   | 43 (18.2)                | 39 (20.3)                              | 13 (7.1)                    | 16 (9.5)                 |
| Neuropathy peripheral                                 | 60 (25.2)                                        | 61 (26.0)                   | 66 (28.0)                | 12 (6.3)                               | 5 (2.7)                     | 5 (3.0)                  |
| Peripheral sensory neuropathy                         | 60 (25.2)                                        | 60 (25.5)                   | 66 (28.0)                | 3 (1.6)                                | 6 (3.3)                     | 2 (1.2)                  |
| Arthralgia                                            | 58 (24.4)                                        | 71 (30.2)                   | 58 (24.6)                | 22 (11.5)                              | 34 (18.6)                   | 16 (9.5)                 |
| Decreased appetite                                    | 55 (23.1)                                        | 42 (17.9)                   | 46 (19.5)                | 28 (14.6)                              | 9 (4.9)                     | 6 (3.6)                  |
| Leukopenia <sup>e</sup>                               | 48 (20.2)                                        | 40 (17.0)                   | 45 (19.1)                | 19 (9.9)                               | 7 (3.8)                     | 9 (5.3)                  |
| Urinary tract infection                               | 48 (20.2)                                        | 33 (14.0)                   | 50 (21.2)                | 25 (13.0)                              | 14 (7.7)                    | 23 (13.6)                |
| Any grade $\geq 3$ AE <sup>f</sup>                    | 160 (67.2)                                       | 129 (54.9)                  | 133 (56.4)               | 79 (41.1)                              | 30 (16.4)                   | 28 (16.6)                |
| Neutropenia <sup>c</sup>                              | 64 (26.9)                                        | 51 (21.7)                   | 55 (23.3)                | 12 (6.3)                               | 1 (0.5)                     | 1 (0.6)                  |
| Anemia <sup>b</sup>                                   | 56 (23.5)                                        | 37 (15.7)                   | 34 (14.4)                | 36 (18.8)                              | 0                           | 1 (0.6)                  |
| Leukopenia <sup>e</sup>                               | 15 (6.3)                                         | 11 (4.7)                    | 13 (5.5)                 | 2 (1.0)                                | 1 (0.5)                     | 0                        |
| Thrombocytopenia <sup>d</sup>                         | 14 (5.9)                                         | 16 (6.8)                    | 11 (4.7)                 | 1 (0.5)                                | 1 (0.5)                     | 0                        |
| Fatigue or asthenia                                   | 12 (5.0)                                         | 8 (3.4)                     | 7 (3.0)                  | 4 (2.1)                                | 1 (0.5)                     | 0                        |
| AEs of special interest to olaparib                   | 14 (5.9)                                         | 5 (2.1)                     | 4 (1.7)                  | 9 (4.7)                                | 4 (2.2)                     | 2 (1.2)                  |
| MDS/AML <sup>g</sup>                                  | 0                                                | 0                           | 0                        | 0                                      | 0                           | 0                        |
| New primary malignancies <sup>h</sup>                 | 2 (0.8)                                          | 1 (0.4)                     | 3 (1.3)                  | 1 (0.5)                                | 1 (0.5)                     | 2 (1.2)                  |
| Pneumonitis <sup>i</sup>                              | 12 (5.0)                                         | 4 (1.7)                     | 1 (0.4)                  | 8 (4.2)                                | 3 (1.6)                     | 0                        |
| Immune-mediated AEs <sup>j</sup>                      | 56 (23.5)                                        | 66 (28.1)                   | 16 (6.8)                 | 27 (14.1)                              | 27 (14.8)                   | 6 (3.6)                  |
| AEs leading to discontinuation of any study treatment | 58 (24.4)                                        | 49 (20.9)                   | 44 (18.6)                | 27 (14.1)                              | 11 (6.0)                    | 7 (4.1)                  |
| Durvalumab/placebo                                    | 22 (9.2)                                         | 26 (11.1)                   | 19 (8.1)                 | 16 (8.3)                               | 9 (4.9)                     | 4 (2.4)                  |
| Olaparib/placebo                                      | 21 (8.8)                                         | 11 (4.7)                    | 5 (2.1)                  | 21 (10.9)                              | 10 (5.5)                    | 5 (3.0)                  |
| Chemotherapy                                          | 31 (13.0)                                        | 31 (13.2)                   | 32 (13.6)                | 1 (0.5)                                | 2 (1.1)                     | 1 (0.6)                  |

# Randomized Phase II Trial of Carboplatin-Paclitaxel Versus Carboplatin-Paclitaxel-Trastuzumab in Uterine Serous Carcinomas That Overexpress Human Epidermal Growth Factor Receptor 2/neu



## RECURRENT DISEASE<sup>i,j</sup>

### First-Line Therapy for Recurrent Disease<sup>c,k</sup>

#### **Preferred**

- Carboplatin/Paclitaxel + Pembrolizumab (except for carcinosarcoma) (category 1)<sup>d,e,l,3</sup>
- Carboplatin/Paclitaxel + Dostarlimab-gxly (category 1)<sup>d,l,5</sup>
- Carboplatin/Paclitaxel + Durvalumab (for dMMR only) (category 1)<sup>d,l,6</sup>
- Carboplatin/Paclitaxel + Trastuzumab  
(for HER2-positive uterine serous carcinoma or carcinosarcoma)<sup>d,h,7</sup>
- Carboplatin/Paclitaxel (category 1 for carcinosarcoma)<sup>m,10</sup>

#### **Other Recommended**

- Carboplatin/Docetaxel<sup>n</sup>
- Carboplatin/Paclitaxel + Bevacizumab<sup>d,8,9</sup>

#### **Useful in Certain Circumstances**

- MMR-proficient (pMMR) tumors
  - ▶ Carboplatin/Paclitaxel/Durvalumab + Durvalumab/Olaparib (category 2B)<sup>d,11</sup>
- Biomarker-directed therapy: after prior platinum-based therapy including neoadjuvant and adjuvant
  - ▶ pMMR tumors
    - ◊ Lenvatinib + Pembrolizumab (category 1)<sup>e,12,13</sup>
  - ▶ TMB-high (TMB-H) tumors<sup>o</sup>
    - ◊ Pembrolizumab<sup>e,14</sup>
  - ▶ MSI-H/dMMR tumors<sup>p</sup>
    - ◊ Pembrolizumab<sup>e,15</sup>
    - ◊ Dostarlimab-gxly<sup>16</sup>

# 2das líneas

| RECURRENT DISEASE <sup>i,j</sup>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Second-Line or Subsequent Therapy <sup>c,k</sup>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| <b>Other Recommended</b> <ul style="list-style-type: none"> <li>• Cisplatin/Doxorubicin<sup>17</sup></li> <li>• Cisplatin/Doxorubicin/Paclitaxel<sup>q,17</sup></li> <li>• Cisplatin/Gemcitabine<sup>18</sup></li> <li>• Cisplatin</li> <li>• Carboplatin</li> <li>• Doxorubicin</li> <li>• Liposomal Doxorubicin</li> <li>• Paclitaxel<sup>19</sup></li> <li>• Albumin-bound Paclitaxel<sup>r</sup></li> <li>• Topotecan</li> <li>• Bevacizumab<sup>s,20</sup></li> <li>• Temsirolimus<sup>21</sup></li> <li>• Cabozantinib</li> <li>• Lenvatinib<sup>22</sup></li> <li>• Gemcitabine<sup>23</sup></li> <li>• Docetaxel (category 2B)</li> <li>• Ifosfamide (for carcinosarcoma)</li> <li>• Ifosfamide/Paclitaxel (for carcinosarcoma)<sup>24</sup></li> <li>• Cisplatin/Ifosfamide (for carcinosarcoma)</li> </ul> | <b>Useful in Certain Circumstances (Biomarker-directed therapy)</b> <ul style="list-style-type: none"> <li>• pMMR tumors <ul style="list-style-type: none"> <li>▶ Lenvatinib + Pembrolizumab (category 1)<sup>e,12,13</sup></li> <li>▶ Axitinib + Avelumab<sup>25</sup></li> </ul> </li> <li>• TMB-H tumors<sup>o</sup> <ul style="list-style-type: none"> <li>▶ Pembrolizumab<sup>e,14</sup></li> <li>▶ Ipilimumab + Nivolumab<sup>26</sup></li> </ul> </li> <li>• MSI-H/dMMR tumors<sup>p</sup> <ul style="list-style-type: none"> <li>▶ Pembrolizumab<sup>e,15</sup></li> <li>▶ Dostarlimab-gxly<sup>16</sup></li> <li>▶ Avelumab</li> <li>▶ Nivolumab<sup>t,27</sup></li> <li>▶ Tislelizumab-jsgr<sup>28</sup></li> <li>▶ Retifanlimab-dlwr (category 2B)<sup>29</sup></li> </ul> </li> <li>• HER2-positive tumors (IHC 3+ or 2+) <ul style="list-style-type: none"> <li>▶ Fam-trastuzumab deruxtecan-nxki<sup>30</sup></li> </ul> </li> <li>• HER2-positive tumors (IHC 1+, 2+ or 3+) <ul style="list-style-type: none"> <li>▶ Fam-trastuzumab deruxtecan-nxki (for carcinosarcoma)<sup>31</sup></li> </ul> </li> <li>• <i>NTRK</i> gene fusion-positive tumors <ul style="list-style-type: none"> <li>▶ Larotrectinib</li> <li>▶ Entrectinib</li> <li>▶ Repotrectinib<sup>u,32</sup></li> </ul> </li> <li>• RET gene fusion-positive tumors <ul style="list-style-type: none"> <li>▶ Selpercatinib</li> </ul> </li> </ul> |

# KeyNote-755 Trial

## Diseño

### Key eligibility criteria:


- Advanced endometrial carcinoma
- Previously treated with at least 1 prior platinum-based chemotherapy regimen in any setting, including neoadjuvant and adjuvant

### Key exclusion criteria:

- Endometrial sarcoma, including carcinosarcoma
- Active autoimmune disease
- A medical condition that required immunosuppression


### Stratification factors:

- ECOG PS
- Geographic region
- History of pelvic radiation




**Table 1.** Demographic and Disease Characteristics of All the Trial Patients at Baseline.\*


| Characteristic                                      | Lenvatinib plus Pembrolizumab<br>(N = 411) | Chemotherapy<br>(N = 416) |
|-----------------------------------------------------|--------------------------------------------|---------------------------|
| Age                                                 |                                            |                           |
| Median (range) — yr                                 | 64 (30–82)                                 | 65 (35–86)                |
| <65 yr — no. (%)                                    | 206 (50.1)                                 | 204 (49.0)                |
| Race — no. (%)†                                     |                                            |                           |
| White                                               | 261 (63.5)                                 | 246 (59.1)                |
| Black                                               | 17 (4.1)                                   | 14 (3.4)                  |
| Asian                                               | 85 (20.7)                                  | 92 (22.1)                 |
| Geographic region — no. (%)‡                        |                                            |                           |
| Region 1                                            | 234 (56.9)                                 | 240 (57.7)                |
| Region 2                                            | 177 (43.1)                                 | 176 (42.3)                |
| MMR status — no. (%)                                |                                            |                           |
| pMMR                                                | 346 (84.2)                                 | 351 (84.4)                |
| dMMR                                                | 65 (15.8)                                  | 65 (15.6)                 |
| ECOG performance-status score — no. (%)§            |                                            |                           |
| 0                                                   | 246 (59.9)                                 | 241 (57.9)                |
| 1                                                   | 164 (39.9)                                 | 175 (42.1)                |
| History of pelvic irradiation — no. (%)             | 174 (42.3)                                 | 186 (44.7)                |
| Histologic features at initial diagnosis — no. (%)¶ |                                            |                           |
| Endometrioid carcinoma                              | 243 (59.1)                                 | 254 (61.1)                |
| High grade                                          | 94 (22.9)                                  | 90 (21.6)                 |
| Low grade                                           | 59 (14.4)                                  | 54 (13.0)                 |
| Not specified                                       | 90 (21.9)                                  | 110 (26.4)                |
| Serous carcinoma                                    | 103 (25.1)                                 | 115 (27.6)                |
| Clear-cell carcinoma                                | 30 (7.3)                                   | 17 (4.1)                  |
| Mixed features                                      | 22 (5.4)                                   | 16 (3.8)                  |

**A pMMR Population****No. at Risk**

|                           |     |     |     |     |    |    |    |    |   |   |
|---------------------------|-----|-----|-----|-----|----|----|----|----|---|---|
| Lenvatinib+ pembrolizumab | 346 | 264 | 165 | 112 | 60 | 39 | 30 | 12 | 5 | 0 |
| Chemotherapy              | 351 | 177 | 83  | 37  | 15 | 8  | 3  | 1  | 1 | 0 |

**B All Patients****No. at Risk**

|                           |     |     |     |     |    |    |    |    |   |   |
|---------------------------|-----|-----|-----|-----|----|----|----|----|---|---|
| Lenvatinib+ pembrolizumab | 411 | 316 | 202 | 144 | 86 | 56 | 43 | 17 | 6 | 0 |
| Chemotherapy              | 416 | 214 | 95  | 42  | 18 | 10 | 4  | 1  | 1 | 0 |

**A pMMR Population**

Median  
Overall  
Survival  
(95% CI)  
*mo*


Lenvatinib+  
Pembrolizumab 17.4 (14.2–19.9)

Chemotherapy 12.0 (10.8–13.3)

Hazard ratio for death,  
0.68 (95% CI, 0.56–0.84)  
 $P < 0.001$

**No. at Risk**

|                          |     |     |     |     |     |     |    |    |   |   |
|--------------------------|-----|-----|-----|-----|-----|-----|----|----|---|---|
| Lenvatinib+pembrolizumab | 346 | 322 | 285 | 232 | 160 | 109 | 62 | 28 | 5 | 0 |
| Chemotherapy             | 351 | 319 | 262 | 201 | 120 | 70  | 33 | 11 | 3 | 0 |

**B All Patients**

Median  
Overall  
Survival  
(95% CI)  
*mo*

Lenvatinib+  
Pembrolizumab 18.3 (15.2–20.5)

Chemotherapy 11.4 (10.5–12.9)

Hazard ratio for death,  
0.62 (95% CI, 0.51–0.75)  
 $P < 0.001$

**No. at Risk**

|                          |     |     |     |     |     |     |    |    |   |   |
|--------------------------|-----|-----|-----|-----|-----|-----|----|----|---|---|
| Lenvatinib+pembrolizumab | 411 | 383 | 337 | 282 | 198 | 136 | 81 | 40 | 7 | 0 |
| Chemotherapy             | 416 | 373 | 300 | 228 | 138 | 80  | 40 | 11 | 3 | 0 |

**Table 3.** Adverse Events of Any Cause with an Incidence of 25% or More among All the Patients in Either Treatment Group, According to Preferred Term.

| Event                   | Lenvatinib plus Pembrolizumab<br>(N = 406) |                  | Chemotherapy<br>(N = 388) |                  |
|-------------------------|--------------------------------------------|------------------|---------------------------|------------------|
|                         | Any Grade                                  | Grade $\geq 3^*$ | Any Grade                 | Grade $\geq 3^*$ |
| Any adverse event       | 405 (99.8)                                 | 361 (88.9)       | 386 (99.5)                | 282 (72.7)       |
| Hypertension†           | 260 (64.0)                                 | 154 (37.9)       | 20 (5.2)                  | 9 (2.3)          |
| Hypothyroidism†‡        | 233 (57.4)                                 | 5 (1.2)          | 3 (0.8)                   | 0                |
| Diarrhea                | 220 (54.2)                                 | 31 (7.6)         | 78 (20.1)                 | 8 (2.1)          |
| Nausea                  | 201 (49.5)                                 | 14 (3.4)         | 179 (46.1)                | 5 (1.3)          |
| Decreased appetite      | 182 (44.8)                                 | 32 (7.9)         | 82 (21.1)                 | 2 (0.5)          |
| Vomiting                | 149 (36.7)                                 | 11 (2.7)         | 81 (20.9)                 | 9 (2.3)          |
| Weight decrease         | 138 (34.0)                                 | 42 (10.3)        | 22 (5.7)                  | 1 (0.3)          |
| Fatigue                 | 134 (33.0)                                 | 21 (5.2)         | 107 (27.6)                | 12 (3.1)         |
| Arthralgia              | 124 (30.5)                                 | 7 (1.7)          | 31 (8.0)                  | 0                |
| Proteinuria†            | 117 (28.8)                                 | 22 (5.4)         | 11 (2.8)                  | 1 (0.3)          |
| Anemia                  | 106 (26.1)                                 | 25 (6.2)         | 189 (48.7)                | 57 (14.7)        |
| Constipation            | 105 (25.9)                                 | 3 (0.7)          | 96 (24.7)                 | 2 (0.5)          |
| Urinary tract infection | 104 (25.6)                                 | 16 (3.9)         | 39 (10.1)                 | 4 (1.0)          |
| Neutropenia             | 30 (7.4)                                   | 7 (1.7)          | 131 (33.8)                | 100 (25.8)       |
| Alopecia                | 22 (5.4)                                   | 0                | 120 (30.9)                | 2 (0.5)          |

# Merck Announces TroFuse-005 Trial Evaluating Sacituzumab Tirumotecan (Sac-TMT) Met Primary Endpoints of Overall Survival (OS) and Progression- Free Survival (PFS) in Certain Patients With Advanced or Recurrent Endometrial Cancer







Muchas gracias