

project proposal:

Effect of homeostatic regulation on dynamics of recurrent neuronal networks

Tom Tetzlaff

t.tetzlaff@fz-juelich.de

Institute of Neuroscience and Medicine (INM-6), Jülich Research Centre and JARA

<http://www.csn.fz-juelich.de>

EITN fall school, Paris, 20.09.2023

https://github.com/tomtetzlaff/2023_eitnfallschool



This presentation is provided under the terms of the Creative Commons Attribution-ShareAlike License 4.0.

Outline

Background

Questions and hypotheses

Proposed work plan

Background

- both in nature and in Computational Neuroscience, neuronal network dynamics can become unstable, leading either
 - to an explosion of network activity or other pathological states, or
 - to dying out of network activity
- often accompanied by a loss of network functionality

Background

- both in nature and in Computational Neuroscience, neuronal network dynamics can become unstable, leading either
 - to an explosion of network activity or other pathological states, or
 - to dying out of network activityoften accompanied by a loss of network functionality
- typical causes:
 - strong excitatory feedback:
 - firing activity ↑
 - ↪ synaptic input ↑
 - ↪ firing activity ↑
 - ...

Background

- both in nature and in Computational Neuroscience, neuronal network dynamics can become unstable, leading either
 - to an explosion of network activity or other pathological states, or
 - to dying out of network activityoften accompanied by a loss of network functionality
- typical causes:
 - strong excitatory feedback:
 - firing activity ↑
 - ↪ synaptic input ↑
 - ↪ firing activity ↑
 - ...
 - correlation-based plasticity dynamics, such as spike-timing-dependent plasticity (STDP):
 - correlations in firing activity of a pair neurons ↑
 - ↪ synaptic coupling between them ↑
 - ↪ correlations in firing activity ↑
 - ...

Background


- both in nature and in Computational Neuroscience, neuronal network dynamics can become unstable, leading either
 - to an explosion of network activity or other pathological states, or
 - to dying out of network activityoften accompanied by a loss of network functionality
- typical causes:
 - strong excitatory feedback:
 - firing activity \uparrow
 - \curvearrowright synaptic input \uparrow
 - \curvearrowright firing activity \uparrow
 - ...
 - correlation-based plasticity dynamics, such as spike-timing-dependent plasticity (STDP):
 - correlations in firing activity of a pair neurons \uparrow
 - \curvearrowright synaptic coupling between them \uparrow
 - \curvearrowright correlations in firing activity \uparrow
 - ...
- typical countermeasures
 - Computational Neuroscience: laborious parameter tuning and/or unrealistic assumptions
 - (healthy) brains: **homeostatic regulation**

Background

Known forms of homeostasis in biological neuronal networks

- intrinsic plasticity (hours–days): regulation of cell-intrinsic excitability (e.g., thresholds; “slow adaptation”) (Naudé et al. 2012)

Intrinsic plasticity




(Watt and Desai 2010)

Background

Known forms of homeostasis in biological neuronal networks


- intrinsic plasticity (**hours–days**): regulation of cell-intrinsic excitability (e.g., thresholds; “slow adaptation”) (Naudé et al. 2012)
- short-term plasticity (**10-1000ms**): (transient) regulation of postsynaptic response amplitude (“synaptic weight”) based on presynaptic activity level (Tsodyks and Markram 1997)

Intrinsic plasticity



(Watt and Desai 2010)

Short-term plasticity



(Markram et al. 2015)


Background

Known forms of homeostasis in biological neuronal networks

- intrinsic plasticity (**hours–days**): regulation of cell-intrinsic excitability (e.g., thresholds; “slow adaptation”) (Naudé et al. 2012)
- short-term plasticity (**10-1000ms**): (transient) regulation of postsynaptic response amplitude (“synaptic weight”) based on presynaptic activity level (Tsodyks and Markram 1997)
- inhibitory plasticity (**minutes–hours**): regulation of strength of inhibitory connections controlled by pre- and postsynaptic activity (spike-timing-dependent plasticity)


(Hennequin, Agnes, and Vogels 2017)

Intrinsic plasticity



(Watt and Desai 2010)

Short-term plasticity




(Markram et al. 2015)

Background

Known forms of homeostasis in biological neuronal networks


- **intrinsic plasticity (hours–days)**: regulation of cell-intrinsic excitability (e.g., thresholds; “slow adaptation”) (Naudé et al. 2012)
- **short-term plasticity (10-1000ms)**: (transient) regulation of postsynaptic response amplitude (“synaptic weight”) based on presynaptic activity level (Tsodyks and Markram 1997)
- **inhibitory plasticity (minutes–hours)**: regulation of strength of inhibitory connections controlled by pre- and postsynaptic activity (spike-timing-dependent plasticity) (Hennequin, Agnes, and Vogels 2017)
- **synaptic scaling (days)**: regulation of synaptic weights based on postsynaptic activity level (Abbott and Nelson 2000; Tetzlaff et al. 2011)

Intrinsic plasticity



(Watt and Desai 2010)

Short-term plasticity




(Markram et al. 2015)

Background

Known forms of homeostasis in biological neuronal networks


- **intrinsic plasticity (hours–days)**: regulation of cell-intrinsic excitability (e.g., thresholds; “slow adaptation”) (Naudé et al. 2012)
- **short-term plasticity (10-1000ms)**: (transient) regulation of postsynaptic response amplitude (“synaptic weight”) based on presynaptic activity level (Tsodyks and Markram 1997)
- **inhibitory plasticity (minutes–hours)**: regulation of strength of inhibitory connections controlled by pre- and postsynaptic activity (spike-timing-dependent plasticity) (Hennequin, Agnes, and Vogels 2017)
- **synaptic scaling (days)**: regulation of synaptic weights based on postsynaptic activity level (Abbott and Nelson 2000; Tetzlaff et al. 2011)
- **synaptic normalization (hours–days)**: long-term synaptic plasticity under the constraint that the total amount of synaptic resources per neuron (e.g. postsynaptic receptors) is constant (Hartmann et al. 2015)

Intrinsic plasticity



(Watt and Desai 2010)

Short-term plasticity




(Markram et al. 2015)

Background

Known forms of homeostasis in biological neuronal networks


- **intrinsic plasticity (hours–days)**: regulation of cell-intrinsic excitability (e.g., thresholds; “slow adaptation”) (Naudé et al. 2012)
- **short-term plasticity (10-1000ms)**: (transient) regulation of postsynaptic response amplitude (“synaptic weight”) based on presynaptic activity level (Tsodyks and Markram 1997)
- **inhibitory plasticity (minutes–hours)**: regulation of strength of inhibitory connections controlled by pre- and postsynaptic activity (spike-timing-dependent plasticity) (Hennequin, Agnes, and Vogels 2017)
- **synaptic scaling (days)**: regulation of synaptic weights based on postsynaptic activity level (Abbott and Nelson 2000; Tetzlaff et al. 2011)
- **synaptic normalization (hours–days)**: long-term synaptic plasticity under the constraint that the total amount of synaptic resources per neuron (e.g. postsynaptic receptors) is constant (Hartmann et al. 2015)
- **structural plasticity (hours–days)**: removal or formation of connections controlled by postsynaptic activity level (“synaptic turnover”) (Butz and Ooyen 2013; Gallinaro and Rotter 2018; Gallinaro, Gašparović, and Rotter 2022)

Intrinsic plasticity




(Watt and Desai 2010)

Short-term plasticity



(Markram et al. 2015)

Structural plasticity




(Fu and Zuo 2011)

Background

Known forms of homeostasis in biological neuronal networks


- **intrinsic plasticity (hours–days)**: regulation of cell-intrinsic excitability (e.g., thresholds; “slow adaptation”) (Naudé et al. 2012)
- **short-term plasticity (10-1000ms)**: (transient) regulation of postsynaptic response amplitude (“synaptic weight”) based on presynaptic activity level (Tsodyks and Markram 1997)
- **inhibitory plasticity (minutes–hours)**: regulation of strength of inhibitory connections controlled by pre- and postsynaptic activity (spike-timing-dependent plasticity) (Hennequin, Agnes, and Vogels 2017)
- **synaptic scaling (days)**: regulation of synaptic weights based on postsynaptic activity level (Abbott and Nelson 2000; Tetzlaff et al. 2011)
- **synaptic normalization (hours–days)**: long-term synaptic plasticity under the constraint that the total amount of synaptic resources per neuron (e.g. postsynaptic receptors) is constant (Hartmann et al. 2015)
- **structural plasticity (hours–days)**: removal or formation of connections controlled by postsynaptic activity level (“synaptic turnover”) (Butz and Ooyen 2013; Gallinaro and Rotter 2018; Gallinaro, Gašparović, and Rotter 2022)
- ... and certainly many more

Intrinsic plasticity




(Watt and Desai 2010)

Short-term plasticity



(Markram et al. 2015)

Structural plasticity



(Fu and Zuo 2011)

Questions and hypotheses

- 1) What biologically plausible mechanisms can effectively stabilize healthy recurrent network dynamics (low rates, asynchronous firing)?


Questions and hypotheses

- 1) What biologically plausible mechanisms can effectively stabilize healthy recurrent network dynamics (low rates, asynchronous firing)?
- 2) How does the presence of such mechanisms alter network dynamics, such as the dynamics in response to perturbations?

Proposed work plan

- 1 implement a simple recurrent neuronal network model as presented during the NEST tutorial

(see NEST example: `balanced_random_network.py`)





Proposed work plan

- 1 implement a simple recurrent neuronal network model as presented during the NEST tutorial

(see NEST example: `balanced_random_network.py`)

- 2 equip the model with a classical form of spike-timing-dependent plasticity in the connections between excitatory neurons

(such as proposed by Morrison, Aertsen, and Diesmann (2007); see NEST synapse model `stdp_pl_synapse*`)



(Morrison, Aertsen, and Diesmann 2007)

Proposed work plan



- 1 implement a simple recurrent neuronal network model as presented during the NEST tutorial

(see NEST example: `balanced_random_network.py`)


- 2 equip the model with a classical form of spike-timing-dependent plasticity in the connections between excitatory neurons

(such as proposed by Morrison, Aertsen, and Diesmann (2007); see NEST synapse model `stdp_pl_synapse*`)

- 3 observe how the network activity evolves into an epileptic high-activity state for high levels of synaptic potentiation (learning rate)



(Morrison, Aertsen, and Diesmann 2007)



(Morrison, Aertsen, and Diesmann 2007)

Proposed work plan

- 1 implement a simple recurrent neuronal network model as presented during the NEST tutorial



(see NEST example: `balanced_random_network.py`)

- 2 equip the model with a classical form of spike-timing-dependent plasticity in the connections between excitatory neurons


(such as proposed by Morrison, Aertsen, and Diesmann (2007); see NEST synapse model `stdp_pl_synapse*`)

- 3 observe how the network activity evolves into an epileptic high-activity state for high levels of synaptic potentiation (learning rate)

- 4 extend the model with one (or several) of the mentioned forms of homeostasis



(Morrison, Aertsen, and Diesmann 2007)



(Morrison, Aertsen, and Diesmann 2007)

Proposed work plan

- 1 implement a simple recurrent neuronal network model as presented during the NEST tutorial

(see NEST example: `balanced_random_network.py`)



- 2 equip the model with a classical form of spike-timing-dependent plasticity in the connections between excitatory neurons

(such as proposed by Morrison, Aertsen, and Diesmann (2007); see NEST synapse model `stdp_pl_synapse*`)


- 3 observe how the network activity evolves into an epileptic high-activity state for high levels of synaptic potentiation (learning rate)

- 4 extend the model with one (or several) of the mentioned forms of homeostasis

- 5 investigate if the destabilization of network activity by strong synaptic potentiation can be prevented by the implemented form of homeostasis



(Morrison, Aertsen, and Diesmann 2007)



(Morrison, Aertsen, and Diesmann 2007)

Proposed work plan

- 1 implement a simple recurrent neuronal network model as presented during the NEST tutorial

(see NEST example: `balanced_random_network.py`)

- 2 equip the model with a classical form of spike-timing-dependent plasticity in the connections between excitatory neurons

(such as proposed by Morrison, Aertsen, and Diesmann (2007); see NEST synapse model `stdp_pl_synapse*`)



- 3 observe how the network activity evolves into an epileptic high-activity state for high levels of synaptic potentiation (learning rate)

- 4 extend the model with one (or several) of the mentioned forms of homeostasis


- 5 investigate if the destabilization of network activity by strong synaptic potentiation can be prevented by the implemented form of homeostasis

- 6 investigate how the implemented form of homeostasis affects network dynamics in the absence or presence of external perturbations (such as average firing rates, level of correlations in firing activity, power spectra of population activity, sensitivity to perturbations)

(Bachmann et al. 2020)



(Morrison, Aertsen, and Diesmann 2007)



(Morrison, Aertsen, and Diesmann 2007)

Thanks

References I

- Abbott, L. F. and S. B. Nelson (2000). "Synaptic plasticity: taming the beast". In: **Nature Neuroscience** 3, pp. 1178–1183. DOI: 10.1038/81453.
- Bachmann, Claudia et al. (2020). "Firing rate homeostasis counteracts changes in stability of recurrent neural networks caused by synapse loss in Alzheimer's disease". In: **PLoS Computational Biology** 16.8, e1007790. DOI: 10.1371/journal.pcbi.1007790.
- Butz, Markus and Arjen van Ooyen (2013). "A simple rule for dendritic spine and axonal bouton formation can account for cortical reorganization after focal retinal lesions". In: **PLoS Computational Biology** 9.10, e1003259. DOI: 10.1371/journal.pcbi.1003259.
- Fu, Min and Yi Zuo (2011). "Experience-dependent structural plasticity in the cortex". In: **Trends in Neuroscience** 34.4, pp. 177–187.
- Gallinaro, Júlia V., Nebojša Gašparović, and Stefan Rotter (2022). "Homeostatic control of synaptic rewiring in recurrent networks induces the formation of stable memory engrams". In: **PLoS Computational Biology** 18.2, e1009836. DOI: 10.1371/journal.pcbi.1009836.
- Gallinaro, Júlia V. and Stefan Rotter (2018). "Associative properties of structural plasticity based on firing rate homeostasis in recurrent neuronal networks". In: **Scientific Reports** 8.1. DOI: 10.1038/s41598-018-22077-3.
- Hartmann, Christoph et al. (2015). "Where's the noise? Key features of spontaneous activity and neural variability arise through learning in a deterministic network". In: **PLoS Computational Biology** 11.12, e1004640.
- Hennequin, Guillaume, Everton J Agnes, and Tim P Vogels (2017). "Inhibitory plasticity: balance, control, and codependence". In: **Annual review of Neuroscience** 40, pp. 557–579. DOI: 10.1146/annurev-neuro-072116-031005.
- Markram, Henry et al. (2015). "Reconstruction and simulation of neocortical microcircuitry". In: **Cell** 163.2, pp. 456–492. DOI: 10.1016/j.cell.2015.09.029.

References II

- Morrison, A., A. Aertsen, and M. Diesmann (2007). "Spike-Timing Dependent Plasticity in Balanced Random Networks". In: **Neural Computation** 19, pp. 1437–1467. DOI: 10.1162/neco.2007.19.6.1437.
- Naudé, Jérémie et al. (2012). "A theory of rate coding control by intrinsic plasticity effects". In: **PLoS Computational Biology** 8.1, e1002349.
- Tetzlaff, Christian et al. (2011). "Synaptic scaling in combination with many generic plasticity mechanisms stabilizes circuit connectivity". In: **Frontiers in Computational Neuroscience** 5, p. 47.
- Tsodyks, Misha V. and Henry Markram (1997). "The neural code between neocortical pyramidal neurons depends on neurotransmitter release probability". In: **Proceedings of the National Academy of Sciences USA** 94, pp. 719–723.
- Watt, Alanna J and Niraj S Desai (2010). "Homeostatic plasticity and STDP: keeping a neuron's cool in a fluctuating world". In: **Frontiers in Synaptic Neuroscience** 2, p. 1486.