

THE FOURIER TRANSFORM AND THE FFT ALGORITHM

Steffen Haug

THE FOURIER TRANSFORM

$$\hat{f}(\xi) = \int_{\mathbb{R}} f(t) e^{-2\pi i \xi t} dt \quad (1)$$

THE FOURIER TRANSFORM

$$\hat{f}(\xi) = \int_{\mathbb{R}} f(t) e^{-2\pi i \xi t} dt \quad (1)$$

Analytic evaluation generally not feasible.

$$\hat{f}(\xi) = \int_{\mathbb{R}} f(t) e^{-2\pi i \xi t} dt$$

Most engineering applications use its discrete counterpart – the DFT:

$$X_k = \sum_{n=0}^{N-1} x_n e^{-\frac{2\pi i k n}{N}} \quad (2)$$

$$X_k = \sum_{n=0}^{N-1} x_n e^{-\frac{2\pi i k n}{N}}$$

We want a way to compute this *efficiently*.

PRELIMINARIES

Eulers formula and the roots of unity

EULERS IDENTITY


$$e^{i\pi} = -1 \tag{3}$$

$$e^{i\pi} = -1$$


Just a special case of $e^{i\vartheta}$ when $\vartheta = \pi$.

$$e^{i\vartheta} = \cos \vartheta + i \sin \vartheta \tag{4}$$

$$e^{i\vartheta} = \cos \vartheta + i \sin \vartheta$$




$$e^{i\pi} = -1$$



THE ROOTS OF UNITY


$$z^N = 1$$

$$z^N = 1$$



SOLUTIONS: $e^{\frac{2\pi in}{N}}$ for n in $0, \dots, N - 1$.

IMPORTANT POINT



When N is even, every root has a “friend” with the opposite sign.

THE KEY IDEA

Decomposing the DFT

SIMPLIFYING ASSUMPTION

N is a power of two

Let $W = e^{\frac{-2\pi i}{N}}$ (notation from Cooley-Tukey).
We can restate (4) as

$$X_k = \sum_n x_n W^{nk} \tag{5}$$

GAINING SPEED

Reusing intermediate results

REFERENCES

- ▶ Cooley & Tukey: