

# Visualisierung von Daten

Erstellen Sie einen Lösungsvorschlag für eine andere Art der Visualisierung

**Estimates of relative survival rates, by cancer site<sup>20</sup>**

	% survival rates and their standard errors							
	5 year		10 year		15 year		20 year	
Prostate	98.8	0.4	95.2	0.9	87.1	1.7	81.1	3.0
Thyroid	96.0	0.8	95.8	1.2	94.0	1.6	95.4	2.1
Testis	94.7	1.1	94.0	1.3	91.1	1.8	88.2	2.3
Melanomas	89.0	0.8	86.7	1.1	83.5	1.5	82.8	1.9
Breast	86.4	0.4	78.3	0.6	71.3	0.7	65.0	1.0
Hodgkin's disease	85.1	1.7	79.8	2.0	73.8	2.4	67.1	2.8
Corpus uteri, uterus	84.3	1.0	83.2	1.3	80.8	1.7	79.2	2.0
Urinary, bladder	82.1	1.0	76.2	1.4	70.3	1.9	67.9	2.4
Cervix, uteri	70.5	1.6	64.1	1.8	62.8	2.1	60.0	2.4
Larynx	68.8	2.1	56.7	2.5	45.8	2.8	37.8	3.1
Rectum	62.6	1.2	55.2	1.4	51.8	1.8	49.2	2.3
Kidney, renal pelvis	61.8	1.3	54.4	1.6	49.8	2.0	47.3	2.6
Colon	61.7	0.8	55.4	1.0	53.9	1.2	52.3	1.6
Non-Hodgkin's	57.8	1.0	46.3	1.2	38.3	1.4	34.3	1.7
Oral cavity, pharynx	56.7	1.3	44.2	1.4	37.5	1.6	33.0	1.8
Ovary	55.0	1.3	49.3	1.6	49.9	1.9	49.6	2.4
Leukemia	42.5	1.2	32.4	1.3	29.7	1.5	26.2	1.7
Brain, nervous system	32.0	1.4	29.2	1.5	27.6	1.6	26.1	1.9
Multiple myeloma	29.5	1.6	12.7	1.5	7.0	1.3	4.8	1.5
Stomach	23.8	1.3	19.4	1.4	19.0	1.7	14.9	1.9
Lung and bronchus	15.0	0.4	10.6	0.4	8.1	0.4	6.5	0.4
Esophagus	14.2	1.4	7.9	1.3	7.7	1.6	5.4	2.0
Liver, bile duct	7.5	1.1	5.8	1.2	6.3	1.5	7.6	2.0
Pancreas	4.0	0.5	3.0	1.5	2.7	0.6	2.7	0.8

<sup>20</sup> Redesigned table based on Hermann Brenner, "Long-term survival rates of cancer patients achieved by the end of the 20th century: a period analysis," *The Lancet*, 360 (October 12, 2002), 1131-1135. Brenner recalculates survival rates from data collected by the U.S. National Cancer Institute, 1973-1998, from the Surveillance, Epidemiology, and End Results Program.