

# Computing Distance and Midpoint from Coordinates

## Answer Key

### High School Geometry

#### Quick Answers

---

- |         |                |                   |
|---------|----------------|-------------------|
| 1. 5    | 4. -1, 1       | 7. 4, 1, 5        |
| 2. 5, 2 | 5. $6\sqrt{2}$ | 8. 4, 3.5, 4, 3.5 |
| 3. 13   | 6. 3, 5        |                   |
- 

#### Curriculum

---

**Level:** High School Geometry

HSG-GPE.B.4–B.7 – *problems 1, 2, 3, 4, 5, 6, 7, 8*

*Difficulty 1–4 of 5 across 11 tagged checks.*

---

#### Common wrong answers

---

1. *If they got 7: added the horizontal and vertical legs instead of using the Pythagorean form*
  3. *If they got 17: added the horizontal and vertical legs instead of using the Pythagorean form*
  5. *If they got 12: added the two legs instead of using the Pythagorean form*
- 

#### 1. Distance from $A(1, 2)$ to $B(4, 6)$ .

**Solution:** The horizontal leg is  $4 - 1 = 3$  and the vertical leg is  $6 - 2 = 4$ , so the segment is the hypotenuse of a 3–4 right triangle.

$$\begin{aligned} AB &= \sqrt{(4 - 1)^2 + (6 - 2)^2} \\ &= \sqrt{3^2 + 4^2} = \sqrt{9 + 16} = \sqrt{25} \end{aligned}$$

$$\boxed{AB = 5}$$

#### 2. Midpoint of $A(2, -3)$ and $B(8, 7)$ .

**Solution:** Average each coordinate separately — the  $x$ 's with the  $x$ 's and the  $y$ 's with the  $y$ 's.

$$M = \left( \frac{2 + 8}{2}, \frac{-3 + 7}{2} \right) = \left( \frac{10}{2}, \frac{4}{2} \right)$$

$$\boxed{M = (5, 2)}$$

Sanity check: 5 is halfway between 2 and 8, and 2 is halfway between  $-3$  and 7.

**3. Distance from  $A(-3, 1)$  to  $B(2, 13)$ .**

**Solution:** Subtracting a negative adds: the horizontal leg is  $2 - (-3) = 5$ , and the vertical leg is  $13 - 1 = 12$ .

$$\begin{aligned} AB &= \sqrt{(2 - (-3))^2 + (13 - 1)^2} \\ &= \sqrt{5^2 + 12^2} = \sqrt{25 + 144} = \sqrt{169} \end{aligned}$$

$$\boxed{AB = 13}$$

This is the 5-12-13 triple.

*Watch for:* an answer of 17 — that is  $5+12$ , adding the legs instead of using the Pythagorean relationship between them.

**4. Midpoint of  $A(-5, 4)$  and  $B(3, -2)$ .**

**Solution:**

$$M = \left( \frac{-5 + 3}{2}, \frac{4 + (-2)}{2} \right) = \left( \frac{-2}{2}, \frac{2}{2} \right)$$

$$\boxed{M = (-1, 1)}$$

A negative  $x$  with a positive  $y$  places the midpoint in Quadrant II.

**5. Exact distance from  $A(0, 0)$  to  $B(6, 6)$ .**

**Solution:** Both legs measure 6, so this is an isosceles right triangle and the exact answer is a radical, not a decimal.

$$\begin{aligned} AB &= \sqrt{(6 - 0)^2 + (6 - 0)^2} = \sqrt{36 + 36} = \sqrt{72} \\ &= \sqrt{36 \cdot 2} = \sqrt{36} \sqrt{2} \end{aligned}$$

$$\boxed{AB = 6\sqrt{2}}$$

(That is about 8.49, but the problem asked for exact radical form.)

**6. Find  $B$  given  $M(3, 5)$  and  $A(-1, 2)$ .**

**Solution:** The midpoint formula is two averaging equations. Write each one and solve for the missing coordinate.

$$\begin{aligned} \frac{-1 + x_B}{2} &= 3 \implies -1 + x_B = 6 \implies x_B = 7 \\ \frac{2 + y_B}{2} &= 5 \implies 2 + y_B = 10 \implies y_B = 8 \end{aligned}$$

$$\boxed{B = (7, 8)}$$

Check by recomputing the midpoint of  $A(-1, 2)$  and  $B(7, 8)$ :  $\left( \frac{-1 + 7}{2}, \frac{2 + 8}{2} \right)$ , which gives the given midpoint  $\boxed{M = (3, 5)}$  ✓

**7. Median of triangle  $PQR$ :**  $P(1, 1)$ ,  $Q(7, 1)$ ,  $R(4, 6)$ .

(a) **Midpoint of  $\overline{PQ}$ .** Both endpoints have  $y = 1$ , so the segment is horizontal and only the  $x$ -coordinate needs averaging.

$$M = \left( \frac{1+7}{2}, \frac{1+1}{2} \right) = \left( \frac{8}{2}, \frac{2}{2} \right)$$

$$\boxed{M = (4, 1)}$$

(b) **Length of the median  $\overline{RM}$ .** Now use the distance formula on  $R(4, 6)$  and  $M(4, 1)$ . Both have  $x = 4$ , so the horizontal leg is 0 and the segment is vertical.

$$RM = \sqrt{(4-4)^2 + (1-6)^2} = \sqrt{0+25} = \sqrt{25}$$

$$\boxed{RM = 5}$$

The median from  $R$  is also the triangle's altitude here, because  $\overline{PQ}$  is horizontal and  $\overline{RM}$  is vertical.

**8. Coordinate proof: diagonals of  $ABCD$  with  $A(0, 0)$ ,  $B(6, 2)$ ,  $C(8, 7)$ ,  $D(2, 5)$ .**

**Solution — computation.** Diagonal  $\overline{AC}$  joins  $A(0, 0)$  to  $C(8, 7)$ :

$$\left( \frac{0+8}{2}, \frac{0+7}{2} \right) = \boxed{(4, 3.5)}$$

Diagonal  $\overline{BD}$  joins  $B(6, 2)$  to  $D(2, 5)$ :

$$\left( \frac{6+2}{2}, \frac{2+5}{2} \right) = \boxed{(4, 3.5)}$$

**Model proof (open response — grade the reasoning).** The two diagonals have the same midpoint,  $(4, 3.5)$ . A midpoint of a segment is by definition the point that divides it into two congruent halves, so that shared point cuts  $\overline{AC}$  into two equal parts and cuts  $\overline{BD}$  into two equal parts. Because the point lies on both diagonals at once, it is exactly where they cross. Therefore each diagonal passes through the other's midpoint — that is, the diagonals bisect each other.

*Look for:* both midpoints computed and stated to be equal; the word *bisect* tied to the definition of midpoint; the observation that a common midpoint is the intersection point. A student who computes only one midpoint has not proved anything.