


Genome assembly

Mark Stenglein




SMBE Regional Workshop
on Computational Biology
in Todos Santos, Mexico



Colorado State University Todos Santos Center
April 8-12, 2019

Genome assembly is the process of *attempting* to reconstruct a genome sequence

An assembly is only a “putative reconstruction” of the genome sequence [Miller, Koren, Sutton (2010)]



Baker M (2012) Nat Methods




Keith Bradnam, UC Davis

Genome assembly paper exercise

Your job is to assemble the ‘genome’ from which the ‘reads’ you’ve been given derive.

Rules/info:


- Like real sequencing data, these reads contain errors.
The error rate is ~2%
- These are single-end 11-base reads
- The average coverage is ~6x
- You’re not allowed to google the answer
- Also: the answer is in the slides: don’t cheat!
- You can use your computers (i.e. word processors or text editors) or paper and whatever strategy you want to do the assembly...



Genome assembly paper exercise

“Jinn (Arabic), also romanized as djinn … are supernatural creatures in early Arabian and later Islamic mythology and theology.”

<https://en.wikipedia.org/wiki/Jinn>




Conclusion: assembly is not trivial!

In this exercise, the ‘genome’ was only 65 positions long, and its alphabet contained 26 ‘bases’ (more information rich)

the human *haploid* genome is 3 Gb

Eukaryotic genomes can have billions of bases and there are only 4 bases (less information)




Bolzer et al (2005) PLoS Biol

Some of the reasons that assembly is difficult

1) Genomes are chock full of repetitive sequences

Alu sequences in the human genome
1 million copies, ~10% of the mass

2) Reads contain errors



Bolzer et al (2005) PLoS Biol

`_gew_kjinns`

`get_djinns_`

`l_get_djinn`

3) Uneven coverage, including possibly no coverage for particular regions (e.g. GC-rich regions)

4) Even with fast computers, it's still computationally difficult

5) Since you don't know what the 'answer' is, it can be difficult to assess whether your assembly is 'good' or not

6) Polyploidy means you are effectively assembling >1 closely related, but not identical, genome


7) Not to mention annotation, which can be as hard as assembly!

De novo assembly is like doing a jigsaw puzzle without the picture on the box



Images, metaphor: *Keith Bradnam, UC Davis*

Reads are assembled into contigs, contigs into scaffolds,
and scaffolds into chromosomes or genomes






These “contigs” could be scaffolded


Sometimes even ‘complete’ assemblies contain gaps

human Y chromosome from GRCh38.p12





Long reads are very useful for scaffolding genome assembly

A real life example: *Pseudomonas aeruginosa* WGS



Long reads have **terrible** error rates






But they're long!


a 242,000 bp read!

It is possible to use the consensus of the reads to perform long read only assembly




Long read only assembly w/ canu assembler produced (complete?) genome in 1 contig, but it still contained errors


assembly |
related
reference
genomes




The Illumina only assembly: most of genome covered but in 52 contigs




Repetitive sequences disrupt short-read assemblies



An rRNA contig assembled, but it sucked up the reads from all the rRNA loci



The hybrid assembly looks pretty good



A key question: How do you know if your assembly is any good?

- Size of the assembly: does it match estimates from other means?
- Size of the contigs/scaffolds: are they reasonably long?
- Are the expected ‘core genes’ present in the assembly?
- What fraction of reads map to the assembly?
- Does the assembly contain sequences of contaminating organisms?
- Is the assembly consistent with independently derived data? (optical mapping, transcriptome sequencing, genomes of related organisms?)

For what purpose do you need the assembly?

These questions apply to assemblies in databases too.


Mini exercise

Batrachochytrium dendrobatidis
cause of chytridiomycosis in amphibians



image: Gewin V. (2008) PLoS Biology

Visit the pages for the 2 assemblies.
Which is better?



NCBI Taxonomy Browser

Search for: Batrachochytrium dendrobatidis

Display: 3 levels using filter: none

Nucleotide Nucleotide EST Nucleotide GSS Protein Structure
 Domains GEO Datasets UniGene PubMed Central Gene
 Assembly LinkOut BLAST TRACE Host
 Bio Systems Clone DB dbVar GEO Profiles PubChem BioAssay

Lineage (full): root; cellular organisms; Eukaryota; Opisthokonta; Fungi; Chytridiomycota; Batrachochytrium

- [Batrachochytrium dendrobatidis](#) 2 Click on organism name to get more information
- [Batrachochytrium dendrobatidis JAM81](#) 1
- [Batrachochytrium dendrobatidis JEL423](#) 1

a common assembly metric:

N50: a measure of the average size of contigs & scaffolds

BD_JEL423

Organism name: [Batrachochytrium dendrobatidis JEL423 \(chytrids\)](#)

Infraspecific name: Strain: JEL423

BioSample: [SAMN02953669](#)

Submitter: Broad Institute

Date: 2006/10/25

Assembly level: Scaffold

Genome representation: full

GenBank assembly accession: GCA_000149865.1 (latest)

RefSeq assembly accession: n/a

RefSeq assembly and GenBank assembly identical: n/a

WGS Project: [AATT01](#)

Assembly method: Unknown v. Unknown

Genome coverage: 7.4x

Sequencing technology: Unknown

IDs: 172958 [UID] 172958 [GenBank]

History ([Show revision history](#))

Comment

Annotation was added to the contigs and scaffolds in May 2016. Please be aware that the annotation is preliminary and subject to curation.

Global statistics

Total sequence length	23,897,668
Total assembly gap length	319,935
Gaps between scaffolds	0
Number of scaffolds	70
Scaffold N50	1,707,251
Scaffold L50	5
Number of contigs	351
Contig N50	221,037
Contig L50	32
Total number of chromosomes and plasmids	1

v1.0

Organism name: [Batrachochytrium dendrobatidis JAM81 \(chytrids\)](#)

Infraspecific name: Strain: JAM81

BioSample: [SAMN02746048](#)

Submitter: US DOE Joint Genome Institute (JGI-PGF)

Date: 2011/04/12

Assembly level: Scaffold

Genome representation: full

RefSeq category: representative genome

GenBank assembly accession: GCA_000203795.1 (latest)

RefSeq assembly accession: GCF_000203795.1 (latest)

RefSeq assembly and GenBank assembly identical: yes

WGS Project: [ADAR01](#)

Assembly method: ARACHNE v. 20070201

Genome coverage: 8.74x

Sequencing technology: Sanger

IDs: 266258 [UID] 266258 [GenBank] 895298 [RefSeq]

History ([Show revision history](#))

Comment

URL -- <http://genome.jgi-psf.org/Batde5>

JGI Project ID: 4001669

The strain JAM81 was provided by Jessica Morgan, Animal Research Institute, Moorooka, Australia

The DNA was provided by Erica Rosenblum, University of Idaho, Moscow, ID (rosenblum@uidaho.edu)

Contacts: Igor Grigoriev, DOE Joint Genome Institute (ivgrigoriev@lbl.gov)

Jason ... [more](#)

Global statistics

Total sequence length	24,315,081
Total assembly gap length	184,767
Gaps between scaffolds	0
Number of scaffolds	127
Scaffold N50	1,484,462
Scaffold L50	6
Number of contigs	510
Contig N50	318,114
Contig L50	24
Total number of chromosomes and plasmids	0

I'm painting a somewhat bleak picture, but don't be too intimidated:
genome sequencing and assembly *is* possible.

Not all assembly problems are equally difficult!

tiny ssDNA genome





image: viralzone

bacterial genomes ~5 Mbp



Nakazawa et al (2009) Genome Research

Loblolly pine (*Pinus taeda*)
22 Gbp genome!



image: Univ of Alabama

Reading what others have done is a great way to figure out what you could do

MOLECULAR ECOLOGY
RESOURCES

Molecular Ecology Resources (2016) 16, 314–324

doi: 10.1111/1755-0998.12443

The *de novo* genome assembly and annotation of a female domestic dromedary of North African origin

ROBERT R. FITAK,^{*1} ELMIRA MOHANDESAN,^{*} JUKKA CORANDER[†] and PAMELA A. BURGER^{*}

**Institut für Populationsgenetik, Vetmeduni Vienna, Veterinärplatz 1, Vienna 1210, Austria, †Department of Mathematics and Statistics, University of Helsinki, Helsinki, FIN-0014, Finland*

You could call these ‘bioinformatics protocols’

mina, USA). Preprocessing of the sequence reads included the removal of adapter sequences and removal of reads with >10% uncalled bases and/or >50% of bases with a Phred-scaled quality score <4. After preprocessing, all 100-bp (paired-end) and 50-bp (mate-pair) reads were retained as the set of ‘raw’ reads. We trimmed the 3' end of all raw reads using a modified Mott algorithm in POPOOLATION v1.2.2 (Kofler *et al.* 2011) to a minimum quality score of 20 and a minimum length threshold of 50 bp and 30 bp for the paired-end and mate-pair reads, respectively.

We corrected the trimmed, paired-end reads for substitution sequencing errors using QUAKE v0.3.5 (Kelley *et al.* 2010). Salzberg *et al.* (2012) showed previously that the error correction of sequencing reads can greatly improve the *de novo* assembly of genomes, including genomes assembled using the program ABYSS (Simpson *et al.* 2009). QUAKE uses the distributions of infrequent and abundant *k*-mers to model the nucleotide error rates and subsequently corrects substitution errors. As input to QUAKE and again after error correction, we counted the frequency of 20-mers in the paired-end reads using DSK v1.6066 (Rizk *et al.* 2013). To estimate genome size, we divided the total number of error-free 20-mers by their peak coverage depth.

We assembled the genome using the trimmed and error-corrected paired-end reads with ABYSS v1.3.6. To determine the optimal *k*-mer length, we repeated the assembly using *k* = 40–88 in 8-bp increments. All scaffolding steps were performed using the trimmed mate-pair reads also in ABYSS, and only scaffolds longer than

Read and synthesize a bunch of these like you would ‘wet lab’ protocols

4. Sequence assembly

All cleaned sequences were assembled using the Newbler Assembler (25) v2.6 (build version 20110517_1502) with the following parameters “-scaffold -het -large -cpu 3 -siod -noinfo”. Our decision to use Newbler was influenced by the large proportion of 454 sequences used and the ability for Newbler to handle multiple data, which allowed BACends, Illumina, and 454 data to be combined. Assemblies were run on a 16-processor node with 256 GB of RAM. Our current assembly consists of 43,234 contigs with an average size of 15,456 bp (min= 436 bp; max=287,935 bp), an N50 size of 29,456 bp, and an N50 count of 6,448. Scaffolding by virtue of the cleaned paired-end reads resulted in 5,745 scaffolds, with an average size of 123 kb (min= 1,732 bp; max= 15.98 Mb), an N50 size of 4.93 Mb, and an N50 count of 50. Based on the N90 statistics, 0.00% of our assembled sequence resides within 155 scaffolds, each of which is 1.16 Mb.

Chamala et al (2016) Science

Bioinformatics protocols are analogous to any lab protocol

mina, USA). Preprocessing of the sequence reads included the removal of adapter sequences and removal of reads with >10% uncalled bases and/or >50% of bases with a Phred-scaled quality score <4. After preprocessing, all 100-bp (paired-end) and 50-bp (mate-pair) reads were retained as the set of ‘raw’ reads. We trimmed the 3’ end of all raw reads using a modified Mott algorithm in POPOOLATION v1.2.2 (Kofler *et al.* 2011) to a minimum quality score of 20 and a minimum length threshold of 50 bp and 30 bp for the paired-end and mate-pair reads, respectively.

We corrected the trimmed, paired-end reads for substitution sequencing errors using QUAKE v0.3.5 (Kelley *et al.* 2010). Salzberg *et al.* (2012) showed previously that the error correction of sequencing reads can greatly improve the *de novo* assembly of genomes, including genomes assembled using the program ABYSS (Simpson *et al.* 2009). QUAKE uses the distributions of infrequent and abundant k-mers to model the nucleotide error rates and subsequently corrects substitution errors. As input to QUAKE and again after error correction, we counted the frequency of 20-mers in the paired-end reads using DSK v1.6066 (Rizk *et al.* 2013). To estimate genome size, we divided the total number of error-free 20-mers by their peak coverage depth.

We assembled the genome using the trimmed and error-corrected paired-end reads with ABYSS v1.3.6. To determine the optimal k-mer length, we repeated the assembly using $k = 40\text{--}88$ in 8-bp increments. All scaffolding steps were performed using the trimmed mate-pair reads also in ABYSS, and only scaffolds longer than

Cells were analyzed using a Cell Lab Quanta SC flow cytometer (Beckman Coulter). CD14-positive cells were stained with CD14-FITC (Miltenyi Biotec). Cells were incubated with propidium iodide to assess cell viability.

Immunoblotting and antibodies. Cells were harvested and total protein extracted in a buffer containing 25 mM HEPES (pH 7.4), 10% glycerol, 150 mM NaCl, 0.5% Triton X-100, 1 mM EDTA, 1 mM MgCl₂, 1 mM ZnCl₂, and protease inhibitors. The extracts were clarified by centrifugation for 10 minutes at 20,800g at 4°C. The extracted proteins (15 µg) were fractionated by SDS-PAGE, transferred to a polyvinylidene difluoride membrane (Millipore), and probed with an anti-A3A polyclonal antiserum, an anti-GFP monoclonal antibody (Clontech), or an anti-eEF1alpha monoclonal antibody (Upstate). The anti-A3A polyclonal serum was generated by immunizing a rabbit with a peptide corresponding to A3A residues 171-199 (CPFQPWDGLEEHSQALSGRLRAILQNQGN) mixed with TiterMax Gold adjuvant (Sigma). Primary antibodies were detected by incubation with fluorescently labeled secondary antibodies and imaging on an Odyssey imaging device (LI-COR Biosciences).

DNA cytidine deaminase activity assays. PBMC or transfected HEK-293T cell lysates were prepared as above for immunoblotting. The deaminase activity in the lysates was determined using a FRET-based assay essentially as described⁵⁹. Briefly, serial dilutions of lysates were incubated for 2 h at 37°C with a DNA oligonucleotide 5’-(6-FAM)-AAA-TTCTAA-TAG-ATA-ATG-TGA-(TAMRA). FRET occurs between the fluorophores, decreasing FAM fluorescence. If

Questions?




Image: Keith Bradnam, UC Davis