
Appendix A

1

Citation: . . *Journal Not Specified* **2024**, *1*,
0. <https://doi.org/>

Received:

Revised:

Accepted:

Published:

Copyright: © 2024 by the author.
Submitted to *Journal Not Specified*
for possible open access publication
under the terms and conditions
of the Creative Commons Attri-
bution (CC BY) license ([https://
creativecommons.org/licenses/by/
4.0/](https://creativecommons.org/licenses/by/4.0/)).

Table S1. Replacing 15/day of total meat, red meat and processed meat with legumes and hazard ratios and 95% confidence intervals for hepatocellular carcinoma and intrahepatic cholangiocarcinoma.

15 g/day of legumes replacing:	Model 1 ¹	Model 2 ²
	HR (95% CI)	HR (95% CI)
Hepatocellular carcinoma		
Total red meat	1.01 (0.93-1.10)	1.06 (0.97-1.16)
Unprocessed red meat	1.01 (0.92-1.11)	1.05 (0.96-1.15)
Processed red meat	1.01 (0.88-1.16)	1.09 (0.95-1.26)
Intrahepatic cholangiocarcinoma		
Total red meat	0.94 (0.87-1.02)	0.97 (0.90-1.05)
Unprocessed red meat	0.92 (0.85-1.00)	0.95 (0.87-1.03)
Processed red meat	1.02 (0.89-1.17)	1.07 (0.93-1.22)

¹Multivariate Cox proportional hazards regression model adjusted for age (as underlying timescale), other food groups, and total food intake.
²Further adjusted for sex, educational level, Townsend deprivation index, living alone, physical activity, smoking, alcohol intake, and waist circumference.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

2
3
4