


# Introduction to the Biosphere-Atmosphere system

Lecture Autumn 2024  
**Part VI**

Steffen M. Noe

# Solar radiation



# What is the solar radiation?

Solar radiation is the electromagnetic energy emitted by the sun.


Solar energy can be expressed as “solar irradiance” which is the power per unit area ( $\text{Wm}^{-2}$ )

Annual average solar radiation arriving on top of the atmosphere is  $1361 \text{ Wm}^{-2}$ .

That's what we call the solar constant!

# What are we looking for

## Spectrum of Solar Radiation (Earth)



# Atmospheric absorption

By passing the atmosphere, the solar radiation is attenuated.


Attenuation happens by absorption and scattering.

Scattered light still contributes to the energy budget.

The maximum solar irradiance reaching to the ground at sea level is approx. 1000 Wm<sup>-2</sup>.

# Global horizontal irradiation

Europe



Average annual sum (4/2004 - 3/2010)



< 700    900    1100    1300    1500    1700    1900 >    kWh/m<sup>2</sup>

0    250    500 km

© 2011 GeoModel Solar s.r.o.

**solarGIS**

<http://solargis.info>


How much area of solar panel would be needed for your home?

My home needs approximately. 5700 kWh per year!  
the location predicts about 1000 kWh/m<sup>2</sup> production

I should be good with ~6 m<sup>2</sup> solar panels

but only if everything is in perfect angles, sun tracked etc!

More realistic calculations would get approx. 30-32 m<sup>2</sup>



# What is sunlight?


Sunlight is the portion of the solar radiation that consist of ultra violet (UV), visible, and infra red (IR) radiation.

It's the same as we already had!  
The solar radiation.

But, we experience sunlight as the combination of bright light and radiant heat.



How does the solar system determines long-term dynamics in the Earth system?



**Positive feedbacks:** Increase in albedo, lower temperature lead to more snow/ice cover and that reflects sunlight which lowers temperature. Reduction of forests further increases albedo.

**Negative feedbacks:** Ice sheets lead to erosion and less land is available for ice sheets to grow on which reduces albedo. Cold dry air because less evaporation occurs and ice sheet growth stops.

*Next glacial period will not occur due to high CO<sub>2</sub> concentrations!*


# Ice age on hold

## Determining the natural length of the current interglacial

P. C. Tzedakis<sup>1\*</sup>, J. E. T. Channell<sup>2</sup>, D. A. Hodell<sup>3</sup>, H. F. Kleiven<sup>4,5</sup> and L. C. Skinner<sup>3</sup>


No glacial inception is projected to occur at the current atmospheric CO<sub>2</sub> concentrations of 390 ppmv (ref. 1). Indeed, model experiments suggest that in the current orbital configuration—which is characterized by a weak minimum in summer insolation—glacial inception would require CO<sub>2</sub> concentrations below preindustrial levels of 280 ppmv (refs 2–4). However, the precise CO<sub>2</sub> threshold<sup>4–6</sup> as well as the timing of the hypothetical next glaciation<sup>7</sup> remain unclear. Past interglacials can be used to draw analogies with the present, provided their duration is known. Here we propose that the minimum age of a glacial inception is constrained by the onset of bipolar-seesaw climate variability, which requires ice-sheets large enough to produce iceberg discharges that disrupt the ocean circulation. We identify the bipolar seesaw in ice-core and North Atlantic marine records by the appearance of a distinct phasing of interhemispheric climate and hydrographic changes and ice-raftered debris. The glacial inception during Marine Isotope sub-Stage 19c, a close analogue for the present interglacial, occurred near the summer insolation minimum, suggesting that the interglacial was not prolonged by subdued radiative forcing<sup>7</sup>. Assuming that ice growth mainly responds to insolation and CO<sub>2</sub> forcing, this analogy suggests that the end of the current interglacial would occur within the next 1500 years, if atmospheric CO<sub>2</sub> concentrations did not exceed 240 ± 5 ppmv.

The notion that the Holocene (or Marine Isotope Stage 1, MIS1), already 11.6 thousand years (kyr) old, may be drawing to a close has been based on the observation that the duration of recent interglacials was approximately half a precession cycle (~11 kyr; ref. 8). However, uncertainty over an imminent hypothetical glaciation arises from the current subdued amplitude of insolation variations as a result of low orbital eccentricity (Fig. 1). It has thus been proposed that at times of weak eccentricity–precession forcing, obliquity is the dominant astronomical parameter driving ice-volume changes, leading to extended interglacial duration of approximately half an obliquity cycle (~21 kyr; ref. 9). In this view, the next glacial inception would occur near the obliquity minimum ~10 kyr from now<sup>7</sup>.



**Figure 1 | Astronomical parameters 100 kyr after present—800 kyr BP and palaeoclimatic records 0–800 kyr BP.** **a**, Eccentricity<sup>29</sup>; **b**, precession index, plotted on an inverse vertical axis<sup>29</sup>; **c**, obliquity<sup>29</sup>; **d**, 21 June insolation 65° N (ref. 29); **e**,  $\delta^{18}\text{O}_{\text{benthic}}$  record from the LR04 stack<sup>28</sup>; **f**, atmospheric CO<sub>2</sub> concentration in Antarctic ice cores<sup>12</sup>; **g**, atmospheric CH<sub>4</sub> concentration in the Antarctic EDC ice core<sup>13</sup>; **h**,  $\delta\text{D}$  composition of ice in the EDC ice core<sup>18</sup>. Marine Isotopic Stages and sub-Stages corresponding to interglacials are indicated. Ages in parentheses denote years after present. The dashed line indicates the current 21 June insolation level at 65° N.

# Vostok ice cores since 1970'ties used to assess Paleoclimate




Isotopically light water evaporates from the ocean and returns via rivers: the system is in balance


Glaciers expand, forming a new reservoir of isotopically light water on the land: sea level drops and the ocean becomes isotopically heavy

## How to assess temperature from ice cores?


- Air bubbles that contain  $CO_2$  and methane
- Oxygen isotopes  $H_2O^{16}$  (99.7%),  $H_2O^{18}$  (0.2%) and  $HDO^{16}$  (0.03%)
- Layers with no bubbles -> meltwater!




# Daily and seasonal rhythms




# Yearly change of solar radiation



A major cause of changing day length is the rotational angle



# Day-night overview






# Implications of seasonal behaviour of ecosystems



# Radiation measurements at SMEAR Estonia since summer 2015

incoming radiation




reflected radiation

Summer to Winter changes radiation by a factor of  $\sim 10!$



Incoming diffuse radiation is  $\sim 50\%$  of direct radiation

Surface reflected radiation is about 1/10 of the possible incoming radiation




Temperature follows the seasonal and daily sunlight cycle






Water vapour and rel. humidity follow the temperature and by that the solar cycle



## Action of seasonal cycle driven processes change the state of the atmosphere



Ozone is driven by solar cycle, NOx and SO<sub>2</sub> are driven by human activities



— 30m  
— 110m

# Theory of the seasonal behaviour

Atmos. Chem. Phys., 17, 15045–15053, 2017  
<https://doi.org/10.5194/acp-17-15045-2017>  
© Author(s) 2017. This work is distributed under  
the Creative Commons Attribution 4.0 License.



Atmospheric  
Chemistry  
and Physics  
Open Access  
EGU


## Annual cycle of Scots pine photosynthesis

Pertti Hari<sup>1</sup>, Veli-Matti Kerminen<sup>2</sup>, Liisa Kulmala<sup>1</sup>, Markku Kulmala<sup>2</sup>, Steffen Noe<sup>3</sup>, Tuukka Petäjä<sup>2</sup>,  
Anni Vanhatalo<sup>1</sup>, and Jaana Bäck<sup>1</sup>





$$\frac{dS}{dt} = \text{Max} \left\{ 0, a_1 (T + T_f) \right\} - a_2 S - a_3 \text{Max} (T_f - T) I. \quad (5)$$

Equation (5) defines the state of the photosynthetic machinery at any moment  $t$  when temperature and solar radiation records are available.

# Seasonal behaviour of the photosynthesis process





# Primary production dynamics measured at SMEAR Estonia



Seasonal cycle determines times when forests are sinks or sources for CO<sub>2</sub>

30m flux measurements are more sensitive to spatial stand heterogeneity (clear cut areas, different age structure, species composition).




$$\text{NEE} = \text{GPP} - \text{RE}$$

# Day-night overview




Daily courses of ecosystem processes are directed by the sunlight



Measured and predicted CO<sub>2</sub> exchange during two days.



A. early in the spring (May 8)  
and B in midsummer (July 18)  
as function of time

# Changes in radiation by cloud cover modulate the daily incoming radiation




Measured and predicted leaf CO<sub>2</sub> exchange  
(a) during a day of intermittent cloudiness (August 5),  
(b) during a cloudy day (July 22), and  
(c) during a sunny day when the stomata close partially (July 7) in Finnish Lapland, 68N.


# Different solar angles due to geological location




# Interplay of daily and seasonal cycle on GPP at different ecosystems



# Processes between the forest and the atmosphere




# Processes between the forest and the atmosphere



# Photosynthesis

General: Takes up carbon ( $\text{CO}_2$ ) from the atmosphere and releases oxygen ( $\text{O}_2$ ) to the atmosphere.

Basic needs:  
Sunlight for energy.  
Water for oxidation.




# Photosynthesis

It is split into light and dark reactions!

It produces sugars that are subsequently used in cellular respiration.



We can see on a leaf/needle scale the "net photosynthesis"  $A$ , is the difference between gross photosynthesis  $P$  and the respiration  $R$ .

$$A = P - R$$



# Photosynthesis

Respiration is also split into the photorespiration and the dark (cellular aerobic) respiration.




Uses energy but does not form sugars.

Use of sugar to produce energy and release CO<sub>2</sub>.

# Photosynthesis


We need gradients in water vapour, oxygen and carbon dioxide to maintain transport of the molecules needed for the process and exchange with the environment.



Another important gradient:


The light energy is also used to build up a proton gradient over the thylakoid membrane inside the chloroplasts.

# Photosynthesis on larger scales



Projected net primary production changes on global scale.

# Photosynthesis on larger scales



# Photosynthesis on larger scales

**Gross primary production GPP:**


The total amount of carbon fixed in the process of photosynthesis.

**Autotrophic respiration Ra:**

Carbon used in plant internal metabolism. It's about half the carbon fixed.

**Net primary production NPP:**

Net production of organic carbon by plants.  $NPP = GPP - Ra$



**Net ecosystem exchange NEE or net ecosystem production NPP:**

Net primary production minus the heterotrophic respiration.  $NEE = NEP = NPP - Rh$

**Heterotrophic respiration Rh:**

Heterotrophic respiration refers to carbon lost by organisms other than plants in ecosystem.


**Net biome exchange NBE or net biome production NBP:**

Change in carbon stocks including carbon losses by episodic natural or anthropogenic disturbances.  $NBE = NEE - Ld$


**Disturbance loss Ld:**

Episodic loss of carbon by disturbances (storms, timber production,...)


# Carbon balance and fluxes




# Processes between the forest and the atmosphere



# Soil efflux process




# Soil respiration and rainfall / soil humidity




**Figure 6** - Effect of rainfall events on the soil  $\text{CO}_2$  efflux emission. The precipitation did not change the mean  $\text{CO}_2$  emission, but the emission had a slight effect on the soil respiration, followed by soil temperature decrease for the SHF forest from Campina Reserve. In days of year (DOY) from 249-255 of 2007.

# Processes between the forest and the atmosphere



# Potential Evapotranspiration



Data taken from: Willmott and Matsuura (2001)

**Atlas of the Biosphere**  
Center for Sustainability and the Global Environment  
University of Wisconsin - Madison

## Some more numbers...

Most of the evaporation of moisture comes from the oceans, lakes and rivers! ~60%


Few is directly sublimated from ice caps, snow and glaciers.

Plant's evapotranspiration makes about 40% annual (summer only 50%, global land cover 30%)

Forests evapotranspiration is double than that of crops or water bodies!

Soils contain only a tiny fraction of the global water resource (~0.01%)

# Water cycle




**Fig. 14.3** The global water cycle, showing approximate magnitudes of the major pools (1,000 km<sup>3</sup>; boxes) and fluxes (1,000 km<sup>3</sup> year<sup>-1</sup>; arrows). Most water is in the ocean, ice, and groundwater, where it is not directly

accessible to terrestrial organisms. The major water fluxes are precipitation, evapotranspiration, and runoff. Modified from Carpenter and Biggs (2009)


# Active and passive processes

Land surface ~ 30%

- Rain/temp. forest ~ 14%
- Boreal forest (Taiga) ~ 17%
- Agricultural land ~ 39%
- Deserts ~ 30%



**active** control on water vapour and carbon by plants --> **FORCING**




Ocean surface ~ 70%



**passive** control on water vapour and carbon --> **FEEDBACK**

# Processes between the forest and the atmosphere



# Hydrocarbons and volatile organic compounds

Fossil fuels

Hydrocarbon = 'hydro(gen)' (H) + 'carbon' (C)

Alcohol (ethanol)

terpenes

benzene

methane

formaldehyde

Peroxy Acetyl Nitrate (PAN)

pinic acid

$\beta$ -carotene

Vitamin A


pesticides

$\alpha$ -pinene

Organic compound = hydrocarbon (+ further species: nitrogen (N), oxygen (O), sulphur (S) etc).

Occurrence: in living matter and the exchanged gases

# Sources of VOC





Source: modified from <http://www.defra.gov.uk/environment/airquality/publications/airqual-climatechange/pdf/chapter02.pdf>



# What is volatility?

From thermodynamics:

volatility is a measure of the tendency  
to vaporize!



# Anthropogenic VOC



# Sources of anthropogenic VOC



*Solvents*




*traffic*



*Solvents e.g. in:  
nail polish,  
perfumes,  
printer cart-  
ridges, tapes,  
advertisement prod.*




*Oil platforms*



# Anthropogenic alkanes

*Saturated hydrocarbons ( $C_xH_y$ ) excluding methane, emitted from all sources shown.*




## Atmospheric fate

reacts only with OH, but slow ( $k_{OH}(198K)=2.55 \times 10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ )

Long lived (days - months) ethane = 2.7 months, propane = 28 days

# Global distribution of surface ethane and propane



von Kuhlmann, 2002

*The simulation clearly shows the hot spots at metropolitan areas, industries, harbours, oil refineries and the transportation pattern of the long lived gases*

# More anthropogenic VOC


The gases you inhale in cities behind a vehicle.



Benzene



Xylene



Toluene




- reaction with OH (and O<sub>3</sub> for styrene)
- short live times (1h-2d)
- local air pollution
- aerosol formation

*Emitted by fossil fuel combustion and biomass burning*



umarsiddiqi.com

# Anthropogenic water soluble VOC



*formic acid  
ethanol (wine, beer etc)  
formaldehyde ( $\text{HCHO}$ )  
acetone*

Atmospheric chemical lifetime

long, more than 3 days up to months!  
(Exception Aldehydes, some hours)

Main sink are clouds and wash out!

Long range transport!



# Global biogenic VOC = BVOC



## Biogenic Volatile Organic Compounds: Annual Global Total Emission > 1.5 Gt




NCAR

Formic acid, acetic acid, ethane, toluene, camphene, terpinolene,  $\alpha$ -terpinolene,  $\alpha$ -thujene, cineole, ocimene,  $\gamma$ -terpinene, bornyl acetate, camphor, piperitone, linalool, tricyclene: 0.04 to 0.2% each

$\beta$ -pinene, d-carene, hexenal, hexenol, hexenyl-acetate, propene, formaldehyde, hexanal, butanone, sabinene, limonene, methyl butenol, butene,  $\beta$ -carophylene,  $\beta$ -phellandrene, p-cymene, myrcene: 0.2 to 1% each

Acetaldehyde, acetone, ethene, ethanol,  $\alpha$ -pinene: 1 to 7% each



Alex Guenther

iLEAPS meeting- September 29, 2003

# BVOC vs anthropogenic VOC

90 Tg year<sup>-1</sup>




760 Tg year<sup>-1</sup>




These numbers come from 2014!


# Breakdown of global anthropogenic sources



# Anthropogenic VOC emission per region



# Non-methane VOC emission density



## Legend


### NMVOC Total 2000

0 – 0.1 Gg NMVOC per cell
0.1 – 1
1 – 2
2 – 10
10 – 50
50 – 100
100 – 2,000 Gg NMVOC per cell

Units = Gigagrams of NMVOC  
(1 Gigagram =  $10^9$  grams = 1,000,000,000 grams)

Methane (~572 Tg/yr) is a source that needs to be handled separately!



# Emission sources




# What is a BVOC?



*Different plants emit so called  
Volatile organic compounds = VOC  
as they are originated from biological  
organism  
they are called  
Biogenic volatile organic compound = BVOC*



# The role of BVOC in atmospheric processes




-volatiles are originated mostly from natural sources (~650 GT/yr vs ~50 GT/yr man made, 2000)


-tropospheric ozone and hydroxyl radicals formed by sunlight and the processes above

-nitrogen oxides are mostly originated from burning processes, large fraction man-made

# BVOC fate after emission



# Short overview on aerosol effects



*Environmental importance: health (respiration), visibility, radiative balance, cloud formation, heterogeneous reactions, delivery of nutrients...*

D.J. Jacob

# Impact of plant originated BVOC emissions on visibility



(c) U.S. forest service (<http://webcam.srs.fs.fed.us/tools/winhaze/>)

If 88% are emitted from natural sources...

Why should we be concerned on that 12% fraction we emit ?



Natural



vs.

Anthropogenic

Natural sources are spread over the whole planet as the global forest cover indicates.

Anthropogenic sources are “concentrated” in areas where humans live. They form together with other pollutants unhealthy conditions.

# Lecture