

# Introduction to the Biosphere-Atmosphere system

Lecture Autumn 2024

**Part III**

Steffen M. Noe

# The concept of Biosphere

Concept introduced by geologist Eduard Suess in 1875.

One thing seems to be foreign on this large celestial body consisting of spheres, namely, organic life. But this life is limited to a determined zone at the surface of the lithosphere. The plant, whose deep roots plunge into the soil to feed, and which at the same time rises into the air to breathe, is a good illustration of organic life in the region of interaction between the upper sphere and the lithosphere, and on the surface of continents it is possible to single out an independent biosphere.



# The concept of Biosphere

Eduard Suess concept is of geological origin!

It already determines the typical features which we accept also today.

The Earth system is described in spheres.

The biosphere links the soil, on top of the lithosphere, with the atmosphere.

The concept of Suess ignores the oceans.


# The concept of Biosphere

Vladimir Ivanovich Vernadsky

The concept was refined to the modern form we usually accept today by him. But he is not much recognised in the west.

In his book "The Biosphere" (1926) he hypothesised that the life is the geological force that shapes the Earth.

In this theory, the principles of both life and cognition are essential features of the Earth's evolution, and must have been implicit in the earth all along.



It's a systems within systems approach!


# Excursion: Gaia hypothesis

Developed by James Lovelock (chemist) and Lynn Margulis (microbiologist) in the 1970'ies

The Gaia hypothesis or principle proposes that the Earth forms a synergistic, self-regulating complex system that maintains the climatic and biogeochemical conditions on Earth in a preferred homeostasis.

Originally it was proposed as Earth feedback system.


Sometimes, the Earth as a whole is seen as the "Biosphere".



# Excursion: homeostasis


Any self-regulating process by which biological systems tend to maintain stability while adjusting to conditions that are optimal for survival. If homeostasis is successful, life continues; if unsuccessful, disaster or death ensues. The stability attained is actually a dynamic equilibrium, in which continuous change occurs yet relatively uniform conditions prevail.

Source: Encyclopaedia Britannica



# Excursion: dynamic equilibrium or steady state

Exchange of matter or energy in an open system in a way that the “net” amount of the substance remains of the same amount.



## Some definitions...

The **biosphere** also known as the ecosphere, is the worldwide sum of all ecosystems.

Source Wikipedia

The part of the world in which life can exist.  
Living organisms together with their environment

Source: Merriam Webster


The regions of the surface and atmosphere of the earth or another planet occupied by living organisms.

Source: Oxford dictionary

# The most comprehensive definition

**Biosphere**, relatively thin life-supporting stratum of Earth's surface, extending from a few kilometres into the atmosphere to the deep-sea vents of the ocean. The biosphere is a global ecosystem composed of living organisms (biota) and the abiotic (nonliving) factors from which they derive energy and nutrients.

Source: Encyclopædia Britannica



# The concept of Biosphere revisited

If we take it together:

The Earth system is seen as spheres.

The Earth is a system of systems (complex adaptive system).


The Biosphere is one sphere.

The Biosphere is a system of systems (complex adaptive system)

The building blocks of the Biosphere are the ecosystems.

# How many Biospheres exist?

We call the Earth – Biosphere 1



Biosphere 2, constructed 1987-1991, as the largest artificial closed system created. Its purpose was to demonstrate the viability of a closed system to maintain human life in space.



However, an earlier experiment called BIOS-3 (1965) in Russia, Krasnoyarsk, is still used to do research on space travel and recycling.


# Biota

A 'biota' is the total **collection** of organisms of a geographic region or a time period, from local geographic scales and instantaneous temporal scales all the way up to whole-planet and whole-timescale spatiotemporal scales. The biotas of the Earth make up the biosphere.


Source: Wikipedia

Because biota are defined as collections of things, they must not have a systems character!

If they have interconnections and purpose, we'd rather name them Ecosystem.



# Biome



Terrestrial biomes according the Köppen scheme

# The biome concept

A **biome** is a formation of plants and animals that have common characteristics due to similar climates and can be found over a range of continents. Spanning continents, biomes are distinct biological communities that have formed in response to a shared physical climate.

Source: Wikipedia

The biome concept unites already the climatic influence on the living organisms.


The physical climate led to common characteristics given similar climatic conditions.

The interactions between biosphere and environment are not yet explicitly named but recognised.

# Sorting the puzzle

Living communities and their nonliving environment are **inseparably interrelated and constantly interact** upon each other. For convenience, any segment of the landscape that includes the biotic and abiotic components is called an ecosystem.

Source: Encyclopædia Britannica




# Primary producers

All life on Earth depends ultimately upon green plants, as well as upon water.

Plants are called primary producers. They utilise the energy of the sunlight to produce biomass by the process of photosynthesis.



# Consumers




# Decomposers


Decomposers are organisms that utilise the biomass produced by primary producers and decompose it in order to utilise the stored energy within this biomass.




# Example of a marine food web



# Energy flow in the biosphere



# Matter flow in the biosphere



# Major equation of the biosphere

$$\text{NPP} = \text{GPP} - \text{Re}$$


NPP is the net primary production

GPP is the gross primary production

Re is the ecosystem respiration

NPP is the rate of net chemical energy production in an ecosystem, which is the difference at which rate chemical energy is produced (GPP) and the rate at which chemical energy is lost to the ecosystem by respiration (Re).


# Feedbacks in the biosphere-atmosphere system




# Forest ecosystems



# Processes between the forest and the atmosphere



# Example: Carbon cycle



# Example: Carbon cycle

## Carbon ( $\text{CO}_2$ ) Budget





Figure 5.12 in IPCC, 2021: Chapter 5. In: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Canadell, J.G., P.M.S. Monteiro, M.H. Costa, L. Cotrim da Cunha, P.M. Cox, A.V. Eliseev, S. Henson, M. Ishii, S. Jaccard, C. Koven, A. Lohila, P.K. Patra, S. Piao, J. Rogelj, S. Syampungani, S. Zaehle, and K. Zickfeld, 2021: Global Carbon and other Biogeochemical Cycles and Feedbacks. In Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 673–816, doi: 10.1017/9781009157896.007 .]


# Measured rise of atmospheric CO<sub>2</sub> since the 60'ies



Measured rise in atmospheric CO<sub>2</sub> since the 60ties

# Carbon and economic growth

Global CO<sub>2</sub> emissions growth has generally resumed quickly from global crises.  
Emission intensity has steadily declined but not sufficiently to offset economic growth.





Economic activity is measured in purchasing power parity (PPP) terms in 2010 US dollars.

Source: [Friedlingstein et al 2021](#); [Global Carbon Project 2021](#)


# Where the carbon goes...

## Fate of Anthropogenic CO<sub>2</sub> Emissions (2010)




# Change in carbon dynamic

## ***RESEARCH: Sources and sinks of carbon***




Sources: CDIAC NOAA-ESRL; Houghton et al, 2012; Giglio et al 2013; Le Quéré et al 2015; Global Carbon Budget 2015

# Where the carbon goes... (last year estimates)



Source: [Friedlingstein et al 2020](#); [Global Carbon Budget 2020](#)

# Where the carbon goes... (recent year estimates)









Budget Imbalance:

(the difference between estimated sources & sinks)

3%  
-1.0 GtCO<sub>2</sub>/yr



# Changes in the sources/sinks over time

The sinks have continued to grow with increasing emissions, but climate change will affect carbon cycle processes in a way that will exacerbate the increase of CO<sub>2</sub> in the atmosphere



The budget imbalance is the total emissions minus the estimated growth in the atmosphere, land and ocean.  
It reflects the limits of our understanding of the carbon cycle.  
Source: [Friedlingstein et al 2021](#); [Global Carbon Project 2021](#)


# Forest uptake is increasing



Fernandez-Martinez et al. 2017, Sci.Rep.

# Role of forests in the climate system

CO<sub>2</sub> induced terrestrial climate feedback



# Aerosol effect on the climate feedback system

## Direct effect of aerosols on solar radiation and gross primary production in boreal and hemiboreal forests

Ekaterina Ezhova<sup>iD1</sup>, Ilona Ylivinkka<sup>iD1</sup>, Joel Kuusk<sup>2</sup>, Kaupo Komasaare<sup>3</sup>, Marko Vana<sup>3</sup>, Alisa Krasnova<sup>4</sup>, Steffen Noe<sup>iD4</sup>, Mikhail Arshinov<sup>iD5</sup>, Boris Belan<sup>iD5</sup>, Sung-Bin Park<sup>6</sup>, Jošt Valentin Lavrič<sup>iD6</sup>, Martin Heimann<sup>iD1,6</sup>, Tuukka Petäjä<sup>iD1</sup>, Timo Vesala<sup>1,7</sup>, Ivan Mammarella<sup>1</sup>, Pasi Kolari<sup>1</sup>, Jaana Bäck<sup>iD7</sup>, Üllar Rannik<sup>1</sup>, Veli-Matti Kerminen<sup>iD1</sup>, and Markku Kulmala<sup>iD1</sup>

<sup>1</sup>Institute for Atmospheric and Earth System Research/Physics, Faculty of Science, University of Helsinki, P.O. Box 64, 00014 Helsinki, Finland

<sup>2</sup>Tartu Observatory, Faculty of Science and Technology, University of Tartu, Tõravere, Nõo Parish, 61602 Tartu, Estonia


<sup>3</sup>Institute of Physics, Faculty of Science and Technology, University of Tartu, 50411, Tartu, Estonia

<sup>4</sup>Department of Plant Physiology, Institute of Agricultural and Environmental Sciences, Estonian University of Life Sciences, 51006 Tartu, Estonia

<sup>5</sup>V.E. Zuev Institute of Atmospheric Optics of Siberian Branch of the Russian Academy of Sciences, 634055 Tomsk, Russia

<sup>6</sup>Max Planck Institute for Biogeochemistry, 07745 Jena, Germany

<sup>7</sup>Institute for Atmospheric and Earth System Research/Forest Sciences, Faculty of Science, University of Helsinki, P.O. Box 64, 00014 Helsinki, Finland



Received: 09 Jul 2018 – Discussion started: 23 Jul 2018 – Revised: 12 Nov 2018 – Accepted: 22 Nov 2018 – Published: 17 Dec 2018






Figure 8. Estimated GPP dependences on  $R_d/R_g$  for all the sites (obtained as  $\text{GPP} = \text{LUE} \cdot \text{PAR}$  using the coefficients for PAR and LUE dependences on  $R_d/R_g$  reported in Table 4).

# How to measure biosphere-atmosphere interactions



***The goal is to measure energy and mass fluxes in a continuous and comprehensive way.***

Linking processes between atmosphere and forest ecosystem to assess their impact on climate.

Info: <http://smear.emu.ee> and <http://www.atm.helsinki.fi/SMEAR/>

# The SMEAR measurement network




## The Järvselja forest site:

heterogenous hemiboreal mixed forest (~55% conifers, ~45% deciduous) main species are Norway Spruce, Scots Pine, Silver Birch, Downy Birch

**Total area 11 000 ha, ~6 000 ha forest,  
~4 500 ha managed**



# SMEAR Estonia working hypothesis



✓ continuous

✓ episodes


✓ continuous (two locations)

# SMEAR Estonia

Järvselja/Apna 130 m tower


## Current measurements

- Eddy covariance at 30 and 70 m
- Gases CO<sub>2</sub>, H<sub>2</sub>O, Methane, Ozone, NO, NO<sub>2</sub>, SO<sub>2</sub> at 30, 50, 70, 90, 110m
- Turbulence and temperature at 30, 50, 70, 90, 110m
- Radiation 40m
- Meteorology (W, T, P, rH, rain)
- Soil fluxes
- Dendometers
- Above and below ground biomass
- Aerosol and air ion measurements
- Scaffolds 30m in Pine and Birch dominated forest




# Monthly CO<sub>2</sub> mixing ratio

Comparison between NOAA Mauna Loa and SMEAR Estonia




Monthly average CO<sub>2</sub> mixing ratio in ppmv

# SMEAR Estonia daily averages




Daily average CO<sub>2</sub> mixing ratio in ppmv




Daily average carbon uptake, GPP  
in  $\mu\text{mol m}^{-2} \text{s}^{-1}$

# Carbon fluxes at SMEAR Estonia



# Monthly CO<sub>2</sub> flux



# Lecture