

# Elbow MRI Dictation Cheat Sheet - Expanded Findings & Impressions

Copy-ready language. Replace bracketed text. Use as a workstation reference, not a substitute for surgical correlation or local report style.

## Core Measurements

| Parameter             | Normal / Threshold                                                         |
|-----------------------|----------------------------------------------------------------------------|
| UCL thickness         | Normal < 4 mm; thickened > 4 mm suggests chronic injury.                   |
| Ulnar nerve CSA       | Abnormal > 8-10 mm <sup>2</sup> at cubital tunnel.                         |
| Capitellar OCD sizing | Report AP x transverse x depth in mm; note fluid undermining.              |
| Radiocapitellar line  | Line through center of radial shaft should bisect capitellum on all views. |
| Carrying angle        | Normal ~5-15 degrees valgus; assess on coronal images.                     |

## Modified Outerbridge Chondromalacia Grading - Copy-Ready

Use this for any articular surface. Replace [surface/compartments] with the specific location: patellar median ridge, medial femoral condyle weightbearing surface, anterosuperior acetabulum, radiocapitellar joint, tibiotalar dome, first MTP joint, etc.

| Grade | MRI shorthand                                                                    |
|-------|----------------------------------------------------------------------------------|
| 0     | Normal cartilage.                                                                |
| 1     | Signal heterogeneity/softening or swelling; surface intact.                      |
| 2     | Superficial fissuring/fraying/partial-thickness defect <50% cartilage depth.     |
| 3     | Deep fissuring/partial-thickness defect >50% cartilage depth, not exposing bone. |
| 4     | Full-thickness cartilage loss with exposed subchondral bone, often edema/cysts.  |

### Grade 0 - normal cartilage

**F:** Articular cartilage of the [surface/compartments] is preserved in thickness and signal without focal fissuring, delamination, or subchondral marrow abnormality.

**I:** No chondral defect of the [surface/compartments].

### Grade 1 - chondral signal change / softening

**F:** Mild focal cartilage signal heterogeneity/softening of the [surface/compartments] with preserved cartilage thickness and intact articular surface. No subchondral marrow edema.

**I:** Grade 1 chondromalacia of the [surface/compartments].

### Grade 2 - superficial partial-thickness chondral loss

**F:** Superficial chondral fissuring/fraying of the [surface/compartments] involving less than 50% of cartilage thickness. No full-thickness defect or subchondral marrow edema.

**I:** Grade 2 chondromalacia of the [surface/compartments].

### Grade 3 - deep partial-thickness chondral loss

**F:** Deep partial-thickness chondral fissuring/defect of the [surface/compartments] involving greater than 50% of cartilage thickness without exposed subchondral bone. [Mild adjacent subchondral marrow edema.]

**I:** Grade 3 chondromalacia of the [surface/compartments].

### Grade 4 - full-thickness chondral loss

**F:** Full-thickness cartilage loss/fissuring of the [surface/compartments] measuring [X] x [Y] mm with exposed subchondral bone and [subchondral marrow edema/cystic change]. [Small unstable chondral flap/loose body if present.]

**I:** Grade 4 chondromalacia/full-thickness chondral defect of the [surface/compartments].

## Elbow MRI Tells

- Lateral epicondylitis: extensor carpi radialis brevis (ECRB) is the most commonly affected tendon at the common extensor origin.
- Thrower's triad: UCL injury + posteromedial olecranon osteophyte + radiocapitellar overload chondral changes.
- Posterolateral rotatory instability (PLRI): LUCL tear is the key lesion; look for pivot shift pattern on MRI and associated radial head subluxation.
- Terrible triad: elbow dislocation + radial head fracture + coronoid fracture.
- Capitellar OCD: T2 fluid undermining the fragment indicates instability.
- Distal biceps: intact lacertus fibrosus limits proximal retraction even with complete tendon rupture.
- Cubital tunnel: elbow flexion increases ulnar nerve pressure; check for subluxation anterior to the medial epicondyle on axial images.

## Normal / Negative Templates

### Normal elbow

**F:** The elbow joint is normally aligned. Common extensor and common flexor tendon origins demonstrate homogeneous low signal and normal morphology. The distal biceps tendon inserts normally at the radial tuberosity with normal fibrillar architecture. The distal triceps tendon inserts normally at the olecranon. The anterior bundle of the ulnar collateral ligament, radial collateral ligament, lateral ulnar collateral ligament, and annular ligament are intact. Articular cartilage of the radiocapitellar, ulnotrochlear, and proximal radioulnar articulations is preserved. The ulnar nerve is normal in caliber and signal within the cubital tunnel. No joint effusion, intra-articular body, or periarticular soft tissue abnormality.

**I:** Normal elbow MRI. No tendon tear, ligament injury, chondral defect, or nerve abnormality.

### Normal UCL

**F:** The anterior bundle of the ulnar collateral ligament is intact from humeral origin to sublime tubercle insertion with homogeneous low signal intensity, normal thickness (< 4 mm), and smooth margins. No periligamentous edema or medial soft tissue abnormality.

**I:** Intact ulnar collateral ligament. No sprain or tear.

## Useful Caveats

- Pseudodeflect of the capitellum: normal cortical irregularity/flattening at the bare area of the posterior capitellum; do not mistake for OCD. Located more posteriorly than true OCD, which favors the anterolateral capitellum.
- UCL T-sign on MR arthrography: contrast undermining the deep fibers of the UCL at the sublime tubercle indicates partial undersurface tear.
- Magic angle artifact: intermediate signal in tendons on short TE sequences when fibers oriented 55 degrees to B0; common at tendon insertions around the elbow.
- Post-surgical UCL reconstruction: expect graft signal heterogeneity and tunnel edema in the early postoperative period; diagnose graft failure when there is complete graft discontinuity or fluid-filled tunnels without graft visualization.

## Lateral Tendons

### Common Extensor Tendon - Normal

**F:** The common extensor tendon origin at the lateral epicondyle demonstrates homogeneous low signal intensity on all sequences with normal morphology and no peritendinous edema.

**I:** Normal common extensor tendon origin. No lateral epicondylitis.

### Common Extensor Tendinosis

**F:** The common extensor tendon origin is thickened with intermediate T2 signal and mild peritendinous edema. No discrete fiber disruption or fluid-signal cleft. The extensor carpi radialis brevis (ECRB) component is [preferentially involved].

**I:** Common extensor tendinosis/lateral epicondylitis without tear.

### Common Extensor Partial-Thickness Tear - Low Grade

**F:** Partial-thickness fiber disruption of the common extensor tendon origin with fluid-signal intensity within the tendon substance involving less than 50% of the tendon cross-section. Surrounding peritendinous edema and lateral epicondyle cortical irregularity. Remaining fibers are intact.

**I:** Low-grade partial-thickness tear of the common extensor tendon origin, lateral epicondylitis.

### Common Extensor Partial-Thickness Tear - High Grade

**F:** High-grade near-complete disruption of the common extensor tendon origin with extensive fluid-signal intensity replacing greater than 50% of the tendon substance. Marked peritendinous edema and lateral soft tissue swelling. A thin rim of intact fibers remains [deep/superficial].

**I:** High-grade partial-thickness tear of the common extensor tendon origin.

### Common Extensor Full-Thickness Tear

**F:** Full-thickness discontinuity of the common extensor tendon origin from the lateral epicondyle with [X] mm tendon retraction and fluid-filled gap at the tear site. Lateral soft tissue edema and [lateral collateral ligament complex is intact/involved].

**I:** Full-thickness tear of the common extensor tendon origin with [X] mm retraction. [Evaluate for concomitant lateral collateral ligament injury.]

## Medial Tendons

### Common Flexor Tendon - Normal

**F:** The common flexor-pronator tendon origin at the medial epicondyle demonstrates homogeneous low signal intensity with normal morphology. No medial soft tissue edema.

**I:** Normal common flexor tendon origin. No medial epicondylitis.

### Common Flexor Tendinosis

**F:** The common flexor-pronator tendon origin is thickened with intermediate T2 signal and medial peritendinous edema. Tendon continuity is preserved without discrete fiber disruption.

**I:** Common flexor tendinosis/medial epicondylitis without tear.

### Common Flexor Partial-Thickness Tear

**F:** Partial-thickness disruption of the common flexor-pronator tendon origin with fluid-signal intensity within the tendon substance. Medial soft tissue edema and medial epicondyle cortical irregularity. [Ulnar collateral ligament is intact/sprained.]

**I:** Partial-thickness tear of the common flexor-pronator tendon origin. [Evaluate for concomitant UCL injury.]

### Common Flexor Full-Thickness Tear

**F:** Full-thickness discontinuity of the common flexor-pronator tendon origin at the medial epicondyle with [X] mm retraction and fluid/hematoma at the tear site. Medial soft tissue edema and swelling. [UCL is intact/torn.]

**I:** Full-thickness tear of the common flexor-pronator tendon origin with retraction. [Comment on UCL status.]

## Distal Biceps Tendon

### Distal Biceps - Normal

**F:** The distal biceps tendon demonstrates normal fibrillar low-signal architecture on all sequences with normal caliber and smooth insertion at the radial tuberosity. No bicipitoradial bursal fluid.

**I:** Normal distal biceps tendon. No tendinosis, tear, or bursitis.

### Distal Biceps Tendinosis / Bicipitoradial Bursitis

**F:** The distal biceps tendon is mildly thickened with intermediate T2 signal at or near the radial tuberosity insertion. Fluid distends the bicipitoradial bursa. Tendon continuity is preserved.

**I:** Distal biceps tendinosis with bicipitoradial bursitis. No tear.

### Distal Biceps Partial-Thickness Tear

**F:** Partial thinning and irregularity of the distal biceps tendon at the radial tuberosity insertion with focal fluid-signal intensity within the tendon substance. Bicipitoradial bursal fluid is present. [Remaining fibers are intact but attenuated.]

**I:** Partial-thickness tear of the distal biceps tendon at the radial tuberosity.

### Distal Biceps Complete Tear

**F:** Complete discontinuity of the distal biceps tendon at the radial tuberosity with [X] cm proximal retraction of the tendon stump and surrounding hemorrhage/edema. The bicipitoradial bursa is distended with complex fluid. The lacertus fibrosus is [intact, limiting retraction / torn, allowing greater retraction].

**I:** Complete distal biceps tendon rupture with [X] cm retraction. Lacertus fibrosus is [intact/torn].

## Triceps Tendon

### Triceps Tendon - Normal

**F:** The distal triceps tendon demonstrates homogeneous low signal intensity with normal morphology and smooth insertion at the olecranon. No posterior soft tissue edema.

**I:** Normal distal triceps tendon.

### Triceps Tendinosis

**F:** The distal triceps tendon is thickened with intermediate T2 signal at the olecranon insertion. No discrete fiber disruption. Mild posterior soft tissue edema.

**I:** Distal triceps tendinosis at the olecranon insertion. No tear.

### Triceps Partial-Thickness Tear

**F:** Partial-thickness disruption of the distal triceps tendon at the olecranon insertion with fluid-signal intensity in the tendon substance and posterior peritendinous edema. Remaining fibers are continuous.

**I:** Partial-thickness tear of the distal triceps tendon.

### Triceps Full-Thickness Tear

**F:** Full-thickness discontinuity of the distal triceps tendon at the olecranon insertion with [X] mm proximal retraction and fluid-filled gap. Posterior soft tissue edema and [olecranon cortical irregularity/avulsion fragment]. [Fleck sign on lateral radiograph if available.]

**I:** Full-thickness distal triceps tendon tear with retraction. [Olecranon cortical avulsion.]

## UCL / Medial Ligaments

### UCL - Normal

**F:** The anterior bundle of the ulnar collateral ligament is homogeneous low signal on all pulse sequences with smooth margins extending from the undersurface of the medial epicondyle to the sublime tubercle. Thickness is normal (< 4 mm). No periligamentous edema.

**I:** Intact ulnar collateral ligament.

### UCL Sprain - Grade 1

**F:** The anterior bundle of the ulnar collateral ligament demonstrates T2 hyperintense edema along its course with preserved fiber continuity and normal thickness. Mild medial soft tissue edema.

**I:** Grade 1 UCL sprain. No tear.

### UCL Partial Tear - Proximal Attachment

**F:** The anterior bundle of the ulnar collateral ligament is thickened with partial-thickness fiber disruption and fluid-signal intensity at the humeral/proximal attachment. Remaining deep/superficial fibers are intact. Medial soft tissue edema. [Common flexor-pronator mass demonstrates reactive signal.]

**I:** Partial-thickness tear of the UCL at the proximal (humeral) attachment.

### UCL Partial Tear - Distal Attachment

**F:** The anterior bundle of the ulnar collateral ligament is thickened with partial-thickness fiber disruption at the distal/sublime tubercle attachment. Fluid signal undermines the deep fibers at the ulnar insertion. Medial periligamentous edema. [T-sign present on arthrographic images, if applicable.]

**I:** Partial-thickness tear of the UCL at the distal (sublime tubercle) attachment.

### UCL Complete Tear - Humeral Attachment

**F:** Complete discontinuity of the anterior bundle of the ulnar collateral ligament at the humeral origin with fluid-filled gap and retraction of the proximal ligament stump. Marked medial soft tissue edema. [Common flexor-pronator tendon demonstrates edema/partial tearing.]

**I:** Complete tear of the anterior bundle UCL at the humeral origin. [Comment on flexor-pronator status.]

### UCL Complete Tear - Sublime Tubercle Attachment

**F:** Complete discontinuity of the anterior bundle of the ulnar collateral ligament at the sublime tubercle insertion with fluid-filled gap and medial joint space widening. Medial soft tissue edema. [Ulnar nerve demonstrates reactive T2 hyperintensity.]

**I:** Complete tear of the anterior bundle UCL at the sublime tubercle. [Comment on ulnar nerve status.]

### UCL Chronic Changes

**F:** The ulnar collateral ligament is diffusely thickened (measuring [X] mm) with heterogeneous signal and periligamentous calcification/ossification. No acute fluid-signal cleft. [Medial epicondyle traction spur. Posteromedial olecranon osteophytes.]

**I:** Chronic UCL insufficiency/thickening with periligamentous calcification. [Correlate for valgus instability.]

## Lateral Collateral Complex

### Lateral Collateral Complex - Normal

**F:** The radial collateral ligament (RCL), lateral ulnar collateral ligament (LUCL), and annular ligament are intact with homogeneous low signal intensity and normal morphology. No lateral soft tissue edema.

**I:** Intact lateral collateral ligament complex.

### RCL Sprain

**F:** The radial collateral ligament demonstrates T2 hyperintense edema with preserved fiber continuity. LUCL and annular ligament are intact. Mild lateral soft tissue edema.

**I:** RCL sprain. No tear. LUCL and annular ligament intact.

### LUCL Tear

**F:** Discontinuity of the lateral ulnar collateral ligament with fluid-signal cleft and posterolateral capsular edema. Edema extends to the supinator crest. [RCL is intact/sprained. Annular ligament is intact/thickened.]

**I:** LUCL tear; evaluate for posterolateral rotatory instability (PLRI). [Comment on RCL and annular ligament status.]

### Annular Ligament Injury

**F:** The annular ligament is thickened and edematous with loss of normal low-signal morphology. [Mild proximal radioulnar joint fluid.] RCL and LUCL are [intact/injured].

**I:** Annular ligament sprain/thickening. [Comment on remainder of lateral collateral complex.]

## Cartilage / OCD

### Capitellar Chondral Softening

**F:** Focal cartilage signal heterogeneity/softening of the anterolateral capitellum with preserved cartilage surface and thickness. No subchondral marrow edema or loose body.

**I:** Grade 1 chondromalacia of the capitellum. No OCD lesion.

### Capitellar OCD - Stable

**F:** Osteochondral lesion of the anterolateral capitellum measuring [X] x [Y] x [Z] mm with overlying chondral irregularity and subchondral marrow edema/cystic change. No fluid undermining the fragment on T2-weighted images. No displaced loose body.

**I:** Capitellar osteochondritis dissecans, likely stable. No fluid undermining or displaced fragment.

### Capitellar OCD - Unstable

**F:** Osteochondral lesion of the anterolateral capitellum measuring [X] x [Y] x [Z] mm with overlying chondral irregularity, subchondral marrow edema, and T2 fluid signal undermining/surrounding the fragment. [Fragment is partially detached/displaced.] [Intra-articular loose body in the [olecranon/coronoid/radiocapitellar] recess.]

**I:** Unstable capitellar osteochondritis dissecans with fluid undermining the fragment. [Loose body.] Surgical consultation recommended.

### Trochlear Chondral Loss

**F:** Chondral fissuring/thinning of the trochlea with [focal/diffuse] partial-thickness cartilage loss and subchondral cystic change. [Mild subchondral marrow edema.]

**I:** [Grade 2/3/4] chondromalacia of the trochlea.

### Posteromedial Impingement

**F:** Posteromedial olecranon osteophyte with opposing trochlear chondral irregularity/thinning and mild synovitis. [Loose body in the posterior compartment.] [UCL is thickened/partially torn, consistent with chronic valgus overload.]

**I:** Posteromedial impingement with olecranon osteophyte and trochlear chondral loss, compatible with valgus extension overload syndrome. [Comment on UCL status.]

## Nerves

### Ulnar Nerve - Normal

**F:** The ulnar nerve is normal in caliber and signal within the cubital tunnel. No perineural edema, compressive mass, or subluxation. Cubital tunnel retinaculum is intact.

**I:** Normal ulnar nerve. No cubital tunnel syndrome.

### Ulnar Neuritis

**F:** The ulnar nerve within the cubital tunnel is enlarged (cross-sectional area [X] mm<sup>2</sup>) and T2 hyperintense with surrounding perineural edema. No compressive mass or ganglion cyst. Nerve is [normally positioned posterior to the medial epicondyle / subluxed anterior to the medial epicondyle].

**I:** Ulnar neuritis/cubital tunnel syndrome. [Comment on nerve subluxation if present.]

### Ulnar Nerve Subluxation

**F:** The ulnar nerve is subluxed anterior to the medial epicondyle on axial images with T2 hyperintensity and mild enlargement. [Cubital tunnel retinaculum is attenuated/absent.] No compressive mass.

**I:** Ulnar nerve subluxation with neuritis, anterior to the medial epicondyle.

### Ulnar Nerve - Mass Compression

**F:** A [T2 hyperintense/heterogeneous] mass within the cubital tunnel measuring [X] x [Y] x [Z] mm displaces and compresses the ulnar nerve, which is enlarged and T2 hyperintense proximal to the mass. [Mass characteristics: well-circumscribed/lobulated/enhancing.]

**I:** Cubital tunnel mass [ganglion cyst / schwannoma / other] with secondary ulnar nerve compression and neuritis.

### Radial Nerve / PIN - Normal

**F:** The radial nerve and posterior interosseous nerve (PIN) are normal in caliber and signal. No enlargement at the arcade of Frohse. Supinator and extensor muscle signal is normal.

**I:** Normal radial nerve/PIN. No radial tunnel syndrome.

### PIN Compression - Arcade of Frohse

**F:** The posterior interosseous nerve is enlarged and T2 hyperintense at the level of the arcade of Frohse / proximal edge of the supinator muscle. Edema within the supinator and [extensor] muscles is present, consistent with denervation. No compressive mass.

**I:** PIN entrapment at the arcade of Frohse with denervation signal in the supinator [and extensor muscles]. Correlate for radial tunnel syndrome.

### PIN Compression - Bicipitoradial Bursa

**F:** Distended bicipitoradial bursa with fluid/synovial thickening compresses the posterior interosseous nerve, which is enlarged and T2 hyperintense. [Denervation edema in the supinator and extensor musculature.]

**I:** Bicipitoradial bursitis with secondary PIN compression and denervation.

### Denervation Pattern - Radial Nerve Territory

**F:** Diffuse T2 hyperintense edema within the supinator and extensor musculature of the posterior forearm compartment without focal mass lesion, consistent with subacute denervation. [T1 signal is normal / demonstrates fatty atrophy indicating chronicity.]

**I:** Denervation pattern in the posterior interosseous nerve distribution. [Subacute if edema only / chronic if fatty replacement.] Correlate for radial nerve or PIN lesion.

## Effusion / Bursae / Loose Bodies

### Joint Effusion - Trace

**F:** Trace elbow joint effusion. No synovial thickening or intra-articular body.

**I:** Trace elbow joint effusion, nonspecific.

### Joint Effusion - Small with Synovitis

**F:** Small elbow joint effusion with mild synovial thickening. No intra-articular loose body.

**I:** Small elbow joint effusion with synovitis.

### Joint Effusion - Moderate with Loose Body

**F:** Moderate elbow joint effusion with synovial thickening and a [X] mm intra-articular osteochondral/ossific body in the [olecranon fossa / coronoid fossa / radiocapitellar recess]. [Additional bodies in the [location].]

**I:** Moderate elbow joint effusion with intra-articular loose body in the [location]. [Correlate for mechanical symptoms.]

### Olecranon Bursitis

**F:** Fluid distension and synovial thickening of the olecranon bursa measuring [X] x [Y] x [Z] mm overlying the posterior olecranon. [No internal complexity/debris.] [Complex fluid with internal septations/debris raising concern for septic bursitis.] No underlying triceps tendon tear.

**I:** Olecranon bursitis. [If complex: clinical correlation for septic bursitis recommended.]

### Bicipitoradial Bursitis

**F:** Fluid distension of the bicipitoradial bursa between the distal biceps tendon and radial tuberosity measuring [X] x [Y] mm. [Distal biceps tendon is intact/tendinotic/partially torn.] [PIN is displaced/compressed.]

**I:** Bicipitoradial bursitis. [Comment on biceps tendon and PIN status.]



### Multiple Loose Bodies / Synovial Chondromatosis

**F:** Moderate-to-large elbow joint effusion with numerous small intra-articular ossific/chondral bodies distributed throughout the olecranon fossa, coronoid fossa, and radiocapitellar recess. Synovial thickening/proliferation is present. [Secondary chondral irregularity of the [capitellum/trochlea].]

**I:** Multiple intra-articular loose bodies with synovitis, consistent with synovial chondromatosis. [Comment on secondary chondral damage.]

### Elbow Arthritis / Degenerative Changes

**F:** Degenerative changes of the elbow joint with marginal osteophytes at the [olecranon tip / coronoid process / radiocapitellar articulation], chondral thinning of the [ulnotrochlear / radiocapitellar] articulation, and small joint effusion. [Olecranon fossa/coronoid fossa loose body.] No acute fracture.

**I:** [Mild/moderate/severe] elbow osteoarthritis. [Loose body.]

### Occult Fracture / Bone Marrow Edema

**F:** Bone marrow edema pattern within the [radial head / coronoid process / capitellum / olecranon] on fluid-sensitive sequences with [preserved / irregular] overlying cortex. No displaced fracture fragment. Joint effusion with [posterior fat pad sign on lateral radiograph, if available].

**I:** Bone marrow edema of the [location] compatible with occult/nondisplaced fracture. Clinical and radiographic correlation recommended.

### Search Pattern

- **1. Check history, indication, and priors.**
- **2. Check adequacy, technique, and limitations.**
- **3. Examine localizers** for incidental pathology.
- **4. Check for effusion/collection** on fluid-weighted sequences:
  - Joint effusion, synovial thickening.
  - Bicipitoradial bursa (reactive to biceps tendon tear).
  - Olecranon bursa.
  - Correlate with enhancement if available.
- **5. Assess bone cortex and marrow:**
  - Cortical breaks/fractures, osteophytes, loose bodies, heterotopic ossification.
  - Transverse trochlear ridge pseudodefekt (expected finding).
  - Marrow replacement; hematopoietic marrow distribution.
  - OCD at capitellum and trochlea (fluid signal, cysts, high-intensity line through cartilage).
  - Marrow edema at epicondyles as clue to overlying cartilage defects.
  - Radiocapitellar alignment.
- **6. Assess cartilage:** PD and fluid-sensitive sequences; ulnotrochlear, radiocarpal, proximal radioulnar articulations; subchondral marrow changes.
- **7. Assess tendons:**
  - Continuity, position/size, signal on PD and fluid-sensitive.
  - Follow from muscle to myotendinous junction to tendon to bony attachment.
  - Medial compartment: common flexor tendon.
  - Lateral compartment: common extensor tendon.
  - Anterior compartment: biceps and brachialis.
  - Posterior compartment: triceps and anconeus (accessory anconeus replacing cubital tunnel retinaculum).
- **8. Assess ligaments:**
  - At least two projections (coronal + axial); fluid signal or contrast interposition (arthrogram).
  - Thickening, attenuation, abnormal intrasubstance signal, periligamentous edema, fiber discontinuity.
  - UCL (anterior bundle most commonly injured, T-sign = partial tear).
  - RCL; LUCL; annular ligament (axials, elbow dislocation setting).
- **9. Assess subcutaneous tissues and musculature:** T1 precontrast, fluid-sensitive correlation.
- **10. Assess nerves:**
  - Denervation changes first.
  - Ulnar nerve (best seen at elbow — osseous/alignment variations, osteophytes, synovial hypertrophy, space-occupying lesions).
  - Median and radial nerves (difficult to delineate from adjacent muscle, look for mass lesions along expected courses).
  - If nerve abnormality is primary, look for secondary denervation changes.
- **11. Double check** marrow signal, muscles, and subcutaneous tissues.
- **12. Proofread report.**

### Infection Search Pattern

- **1. Marrow/bone changes:**
- Marrow replacement (darker than muscle/disc on T1; geographic pattern more predictive than reticular/periarticular).
- Marrow edema (isolated finding raises concern in diabetics/high-risk).
- Bone erosion (loss of dark T1/T2 cortical line).
- **2. Surrounding inflammatory changes:** fascia, musculature, subcutaneous fat edema/enhancement.
- **3. Susceptibility suggesting air:** in collections, soft tissues, along fascial planes, in bone.
- **4. Abnormal enhancement:**
- Rim-enhancing collection = abscess.
- Superficial lesion without enhancement = ulcer.
- Enhancement along fascial planes = infectious/inflammatory fasciitis.
- Non-enhancement of normal structures = ischemia.
- Other patterns of enhancement may not be specific.
- **5. Differentiating osteomyelitis from neuropathic joint:**
- Joint effusion with marrow replacement on both sides of joint.
- Fluid collections/abscesses (periprosthetic, subperiosteal, intraosseous/Brodie's abscess).
- Sinus tract connecting abnormal marrow signal to skin surface.
- Disappearance of subchondral cysts on sequential imaging.
- Progressive marrow/soft tissue changes greater than expected on sequential imaging.

### Source Notes

Built from the original elbow MRI cheat sheet, standard MSK MRI reporting conventions, review-level radiology references for UCL/lateral collateral injuries, capitellar OCD staging, cubital tunnel syndrome, and PIN entrapment. Expanded to match the depth of the shoulder cheat sheet with graded findings for all major structures. Use surgical correlation and local report conventions as needed.