

```
space_time_rotation.wxm

GNU General Public License 2020 Stephen Athel Abbott.

An application document for Geometric Algebra using wxMaxima

Ref: The Survey, paragraph 4.1.2

Use of G(1,3) to rotate spacetime!

Initialization

(%i57) ext:["wxm"]$
      file_type_maxima:append(ext,file_type_maxima)$
      batchload("initialize_fns")$

the pseudoscalar and its inverse
the lowest useable dimension pseudoscalar should be {e1,e2} i.e. Plen = 2
e.g. for four dimensions edit Pseudos:{e1,e2,e3}$ to Pseudos:{e1,e2,e3,e4}$

(%i1) Pseudos:{e1,e2,e3,e4}$
      Pvar:listofvars(Pseudos)$
      Plen:length(Pvar)$
      I:Pseudos$
      ni:(Plen-1)*Plen/2$
      Ii:(-1)^ni*I$
      kill(ni)$
      ldisplay(Pvar)$

      (%t8) Pvar=[e1,e2,e3,e4]

(%i9) batchload("initialize_lsts")$

      (%t9) lstblds=[[{e1},{e2},{e3},{e4}],[{e1,e2},{e1,e3},{e1,e4},{e2,e3},{e2,e4},{
e3,e4}],[{e1,e2,e3},{e1,e2,e4},{e1,e3,e4},{e2,e3,e4}],[{e1,e2,e3,e4}]]
      (%t10) allblds=[{e1},{e2},{e3},{e4},{e1,e2},{e1,e3},{e1,e4},{e2,e3},{e2,e4},{e3,
e4},{e1,e2,e3},{e1,e2,e4},{e1,e3,e4},{e2,e3,e4},{e1,e2,e3,e4}]
      (%t11) invblds=[{e1},{e2},{e3},{e4},-{e1,e2},-{e1,e3},-{e1,e4},-{e2,e3},-{e2,e4}
,-{e3,e4},-{e1,e2,e3},-{e1,e2,e4},-{e1,e3,e4},-{e2,e3,e4},{e1,e2,e3,e4}]

end of Initialization
```

floating point print (display) precision

```
(%i12) fpprintprec:6$
      ratprint:false$
      ldisplay(fpprintprec,fpprec,ratprint)$

      (%t14) fpprintprec=6
      (%t15) fpprec=16
      (%t16) ratprint=false
```

The Survey, para.4.1.2

show the spacetime gammas required for the imitation of G(1,3), where, to avoid the use of gamma_zero, we have used the fourth axis, e4, for the time axis and the intrinsic maxima imaginary, %i, for the space axes

```
(%i17) g1:%i*{e1}$
      g2:%i*{e2}$
      g3:%i*{e3}$
      g4:{e4}$
```

the spacetime coordinate vector using the gammas

```
(%i21) x:x1*g1+x2*g2+x3*g3+t*g4;

      (%o21) %i*{e3}*x3+%i*{e2}*x2+%i*{e1}*x1+{e4}*t
```

spacetime vector rotation for a simple velocity, vel*vhat (= vel*g1)

```
(%i22) vel:0.8$
      alpha:atanh(vel)$
      ahalf:alpha/2;

      (%o24) 0.5493
```

form the rotation bivector

```
(%i25) vhat:g1$
      B:vhat&*g4*ahalf$
      ev(% ,numer,expand);

      (%o27) 0.5493*%i*{e1,e4}
```

form the left and right exponential multipliers

```
(%i28) mvexp(-B,13)$
      lexp:ev(% ,numer,expand);

      (%o29) 1.1547-0.5774*%i*{e1,e4}

(%i30) mvexp(+B,13)$
      rexp:ev(% ,numer,expand);

      (%o31) 0.5774*%i*{e1,e4}+1.1547
```

apply the rotation to the spacetime coordinate vector and find the rotated spacetime vector

```
(%i32) xbar:lexp&*x&*rexp$
      ev(% ,numer,expand)$
      collectterms(% ,%i,e1,e4);

      (%o34) %i*(1.0*{e3}*x3+1.0*{e2}*x2)+%i*{e1}*(1.66667*x1-1.33333*t)+{e4}*
(1.66667*t-1.33333*x1)
```