



Bat roost dynamics

Stephen Gregory¹, Philip Briggs², Karen Haysom² & Franck Courchamp¹

¹Ecologie, Systématique & Evolution, Université Paris-Sud XI ²Bat Conservation Trust, 8 Battersea Park Road, London

July 16 2009

Background and questions

- Why bats?
- Questions

Background and questions


- Why bats?
- Questions

Roost dynamics

- Analyses
- Species results

Background and questions

Why bats?



Bat behaviours

More than 1000 species of bat Worldwide

Bat behaviours

- More than 1000 species of bat Worldwide
- Most are sociable

Bat behaviours

- More than 1000 species of bat Worldwide
- Most are sociable
- Bat social behaviours include:
 - grouping for thermoregulation
 - babysitting or creching
 - o alturistic feeding
 - o information exchange

Bat behaviours


- More than 1000 species of bat Worldwide
- Most are sociable
- Bat social behaviours include:
 - o grouping for thermoregulation
 - babysitting or creching
 - alturistic feeding
 - o information exchange
- Social behaviours require a minimum number of participants

Bat behaviours

- More than 1000 species of bat Worldwide
- Most are sociable
- Bat social behaviours include:
 - o grouping for thermoregulation
 - babysitting or creching
 - alturistic feeding
 - information exchange
- Social behaviours require a minimum number of participants


Bats may exhibit Allee effects

An Allee effect is...




An Allee effect is...

 a positive relationship between 'fitness' & population size




An Allee effect is...

- a positive relationship between 'fitness' & population size
- 'fitness' either individual (e.g. survival) or group (population growth rate)




Allee effects act to...



Allee effects act to...


increase 'fitness' while a population is growing



Allee effects act to...

increase 'fitness' while a population is growing

BUT




Allee effects act to...

increase 'fitness' while a population is growing

BUT

 decrease 'fitness' while a population is declining




Allee effects act to...

increase 'fitness' while a population is growing

BUT

 decrease 'fitness' while a population is declining

Allee effects may drive declining populations to extinction



UK National Bat Monitoring Programme (NBMP)

Collaborators







UK National Bat Monitoring Programme (NBMP)

National Bat Monitoring Programme initiated in 1996

Collaborators







UK National Bat Monitoring Programme (NBMP)

- National Bat Monitoring Programme initiated in 1996
- Monitors national population trends of 11 UK bat species

Collaborators







UK National Bat Monitoring Programme (NBMP)

- National Bat Monitoring Programme initiated in 1996
- Monitors national population trends of 11 UK bat species
- Over 2000 volunteers supported by network of 'bat groups'

Collaborators







UK National Bat Monitoring Programme (NBMP)

- National Bat Monitoring Programme initiated in 1996
- Monitors national population trends of 11 UK bat species
- Over 2000 volunteers supported by network of 'bat groups'
- Includes data for some species monitored since 1977

Collaborators







UK National Bat Monitoring Programme (NBMP)

- National Bat Monitoring Programme initiated in 1996
- Monitors national population trends of 11 UK bat species
- Over 2000 volunteers supported by network of 'bat groups'
- Includes data for some species monitored since 1977
- NBMP uses 3 core methods; I analyse 'Maternity roost counts'

Collaborators







Maternity roost counts

Collaborators









Maternity roost counts

Maternity roosts are places where bats gather to breed

Collaborators









Maternity roost counts

- Maternity roosts are places where bats gather to breed
- Usually in buildings (e.g. homes) but also trees, etc

Collaborators









Maternity roost counts

- Maternity roosts are places where bats gather to breed
- Usually in buildings (e.g. homes) but also trees, etc
- Breeding bats show habitual behaviour facilitating survey

Collaborators









Maternity roost counts

- Maternity roosts are places where bats gather to breed
- Usually in buildings (e.g. homes) but also trees, etc
- Breeding bats show habitual behaviour facilitating survey
- Volunteers count bats on 2 evenings during breeding period

Collaborators









Maternity roost counts

- Maternity roosts are places where bats gather to breed
- Usually in buildings (e.g. homes) but also trees, etc
- Breeding bats show habitual behaviour facilitating survey
- Volunteers count bats on 2 evenings during breeding period
- Counts not undertaken in extreme weather

Collaborators









Maternity roost counts

- Maternity roosts are places where bats gather to breed
- Usually in buildings (e.g. homes) but also trees, etc
- Breeding bats show habitual behaviour facilitating survey
- Volunteers count bats on 2 evenings during breeding period
- Counts not undertaken in extreme weather
- 8 species are monitored using Maternity roost counts

Collaborators









Background and questions

- Why bats?
- Questions

Questions

Questions

1) Are bat counts affected by weather?

Questions

1) Are bat counts affected by weather?

2) Do bats exhibit an Allee effect?

Talk outline

Background and questions

- Why bats?
- Questions

Roost dynamics

Analyses

UK rainfall this week in 1977 Weather data Thanks: ENSEMBLES and ECA&D

Weather data

 Used ENSEMBLES gridded weather data for Europe http://eca.knmi.nl/

Thanks: ENSEMBLES and ECA&D

UK rainfall this week in 1977

Weather data

- Used ENSEMBLES gridded weather data for Europe http://eca.knmi.nl/
- Extracted daily UK temperature & rain using ncdf and spatstat in R

Thanks: ENSEMBLES and ECA&D

UK rainfall this week in 1977

Smoothing counts for weather variation

Smoothing counts for weather variation

Used weather variables to smooth counts at each roost

Smoothing counts for weather variation

- Used weather variables to smooth counts at each roost
- Compared models of varying complexity using AIC:

Smoothing counts for weather variation

- Used weather variables to smooth counts at each roost
- Compared models of varying complexity using AIC:

Complexity	Model
Highest	
\	
Lowest	

Smoothing counts for weather variation

- Used weather variables to smooth counts at each roost
- Compared models of varying complexity using AIC:

Complexity	Model		
Highest	saturated Generalized Additive Model (GAM) with automatic smooth		
	↓ all combinations of weather covariables		
	non-linear vs linear terms		
Lowest	Generalized Linear Model (GLM) with no weather covariables		

Weather results

Weather results

Saturated GAMs had lowest deviance for all species

Weather results

- Saturated GAMs had lowest deviance for all species
- Temperature and rain affect bat counts non-linearly

Weather results

- Saturated GAMs had lowest deviance for all species
- Temperature and rain affect bat counts non-linearly

Smoothed counts were extracted for the Allee effect analysis

Allee effect analysis method

Allee effect analysis method

Followed simple empirical method of Tobin et al. 2007

Thanks: Patrick Tohin

Allee effect analysis method

Followed simple empirical method of Tobin et al. 2007

For each roost, compare count in year t and t+1 to see if it increased



Thanks: Patrick Tobin

Allee effect analysis method

Followed simple empirical method of Tobin et al. 2007

For each roost, compare count in year t and t+1 to see if it increased



Plot the proportion of roosts increasing against their sqrt(size) in year t

Thanks: Patrick Tohin

Allee effect analysis method

Followed simple empirical method of Tobin et al. 2007

For each roost, compare count in year t and t+1 to see if it increased


 \Rightarrow

Plot the proportion of roosts increasing against their sqrt(size) in year t

 Fit models representing [i] random walk (RW), [ii] negative density dependence (DD), and [iii] an Allee effect (AE) and measure support

Thanks: Patrick Tohin

Roost dynamics models



Talk outline


Background and questions


- Why bats?
- Questions

Roost dynamics

- Analyses
- Species results
































Daubenton's bat (Myotis daubentonii)









Natterer's bat (Myotis nattereri)









Noctule bat (Nyctalus noctula)









Common pipistrelle (Pipistrellus pipistrellus)








Soprano pipistrelle (Pipistrellus pygmaeus)








Brown long-eared bat (Plecotus auritus)









Lesser horseshoe bat (Rhinolophus hipposideros)







Allee effect results

Allee effect results

Some species do (others do not)

Allee effect results

- Some species do (others do not)
- An Allee effect may drive roost abandonment in these species

Allee effect results

- Some species do (others do not)
- An Allee effect may drive roost abandonment in these species
- These species use the same roost year-round

Allee effect results

- Some species do (others do not)
- An Allee effect may drive roost abandonment in these species
- These species use the same roost year-round

Management recommendation

Species exhibiting Allee effects should be subject to careful/supplemental monitoring



Questions

stephen.gregory@u-psud.fr



SG acknowledges the British Ecological Society for a travel grant

Model AIC_w

Table: Model AICc weights (AIC_w)

Species	AIC_w		
	RW	DD	AE
Serotine	0.001	0.882	0.117
Daubenton's	0.096	0.363	0.541
Natterer's	0.000	0.684	0.316
Noctule	0.008	0.983	0.009
Common pipistrelle	0.000	0.795	0.205
Soprano pipistrelle	0.000	0.117	0.883
Long-eared	0.001	0.870	0.129
Lesser horseshoe	0.000	0.053	0.947