



Wylfe Grayling Long-Term Study

Stephen Gregory, Richard Cove, Anton Ibbotson



Biosketch: Stephen D. Gregory



- Education:
 - BSc Zoology (University of Swansea)
 - MSc Ecology (University of Oxford)
 - PhD Ecology & Statistics (University of Paris)
- Projects:
 - Foraging behaviour of Lesser horseshoe bats, Swansea
 - Food competition between native & black rats, Galapagos
 - Demographic consequences of sociality in animals
 - Effects of climate and habitat change on Orangutan, Borneo
- Current:
 - Drivers of salmon population dynamics, UK and France
 - Grayling population dynamics on the Wylfe, UK



In 1996...



In 1996...



Anton Ibbotson



In 1996...

Wylfe Grayling Long-Term Study begins!



Anton Ibbotson



In 1996...



Me!



Wylfe Grayling Long-Term Study



Wylfe Grayling LTS: longest grayling dataset

- 1996 – 2013: 18 years!



Wylfe Grayling LTS: longest grayling dataset

- 1996 – 2013: 18 years!
- 9760+ records from 8463 individual grayling
 - Length
 - Scales



Wylfe Grayling LTS: longest grayling dataset

- 1996 – 2013: 18 years!
- 9760+ records from 8463 individual grayling
 - Length
 - Scales
- River flow
 - Stockton Park
 - Norton Bavant



Wylfe Grayling LTS: individual-based data

- Individual marks



Wylfe Grayling LTS: individual-based data



- Individual marks



Wylfe Grayling LTS: individual-based data

- Individual marks





Long data sets needed to study changes in
fish population size and their causes



Grayling: an indicator of river health

- Salmonid ~ similar needs to salmon & trout

Grayling: an indicator of river health

- Salmonid ~ similar needs to salmon & trout
- Most sensitive to environmental change:

	Temperature (oC)		
	Salmon	Trout	Grayling
Lower	0 – 6	0 – 4	0 – 4
Optimum	6 – 20	4 – 19	4 – 18
Upper	20 – 34	19 – 30	18 – 25

Grayling: an indicator of river health

- Salmonid ~ similar needs to salmon & trout
- Most sensitive to environmental change:

	Temperature (oC)		
	Salmon	Trout	Grayling
Lower	0 – 6	0 – 4	0 – 4
Optimum	20 – 34	19 – 30	18 – 25
Upper	6 – 20	4 – 19	4 – 18

- Most sensitive reproductive strategy

Grayling reproduction: a risky strategy



Grayling reproduction: a risky strategy

- Spawn in April – May: weather dependent

Grayling reproduction: a risky strategy

- Spawn in April – May: weather dependent
- Eggs shallow in redd:

	Salmon	Trout	Grayling
Depth (cm)	10 – 15	10 – 15	0 – 5

Average redd characteristics



Measure	River Pollon	River Suran
Mean bottom velocity	37.2 cm/s	33.7 cm/s
Selected depth	10 – 40 cm	20 – 30 cm



Grayling reproduction: a risky strategy

- Spawn in April – May: weather dependent
- Eggs shallow in redd:

	Salmon	Trout	Grayling
Depth (cm)	10 – 15	10 – 15	0 – 5

- Shallow redd ~ risk of “egg washout” in floods

UK flooding





UK flooding

- 4 of 5 UK wettest recorded years since 2000
 - (2000 [1], 2012 [2], 2008 [4], 2002 [5])

UK flooding

- 4 of 5 UK wettest recorded years since 2000
– (2000 [1], 2012 [2], 2008 [4], 2002 [5])
- January 2014 = England's wettest in c.250 years

UK flooding

- 4 of 5 UK wettest recorded years since 2000
– (2000 [1], 2012 [2], 2008 [4], 2002 [5])
- January 2014 = England's wettest in c.250 years

High rainfall -> High river flows

Wylfe January 2014



Frome



Frome: 2012



Questions




Questions

How might flooding affect grayling populations?

- Shared patterns in flow and grayling counts?






Wylfe average daily flow




Grayling population:

all ages







all ages



all ages





Grayling population: separate ages




Grayling population:

0+ only







0+ only




0+ only







0+ only



April 2000



er of grayling in first shock



April 2000 - Record rainfall

A consistently wet April has led to records being broken throughout the UK


Introduction

Record rainfall

A consistently wet April has led to records being broken throughout the UK.

According to the England and Wales precipitation series (an index which begins in 1766) the total was 143 mm (using best available data/estimates on 2 May 2000) making it the wettest April since records began. Previous highest April totals were in 1782 (139 mm), 1818 (136 mm) and 1998 (133 mm). The chart below shows the series April 1766-2000.


England & Wales Precipitation, April 1766-2000




1999 – 2000


125 -> 26 =
80% decrease in 0+






0+ only







0+ only



April 2012

Number of grayling in first shock



Record rainfall - April to July 2012

From April to July 2012 the UK experienced a period of exceptionally wet weather, breaking previous rainfall records and resulting in several significant flood events.

The wet weather affected all of England, Wales and eastern Scotland. April, June, and the period April to July were each the wettest on record in the England & Wales precipitation series from 1766, while for the UK overall, summer 2012 (June, July and August) was the wettest since 1912. The record rainfall brought the [2010-12 England and Wales drought](#) to an abrupt end. In contrast to the wet weather elsewhere, the far north-west of Scotland saw well below-average rainfall from March to October 2012.

The persistent wet weather was due to a shift in the jet stream to a much more southerly track than normal, bringing a succession of Atlantic low pressure systems and associated fronts across the southern half of the UK.

Impacts

After the drought, the wet weather was initially very welcome, bringing much-needed rain for farmers and growers. However, before long it brought new problems.

Waterlogging made access to land difficult, reduced yields and caused some crops to rot. Various flood incidents through the period caused widespread problems, particularly to the transport network. Surface water flooding and debris closed main roads. Railway lines were blocked by flooding and landslips. Birmingham airport diverted inbound flights in late June.

2011 – 2012

118 -> 7 =
94% decrease in 0+




“Provisional” observations




- Patterns: not cause and effect
 - Needs closer examination, e.g., 2005, 2007
- Based on counts, not population estimates
- Alternative factors:
 - Temperature
 - Habitat loss
 - etc



Length: poor 0+ growth 2013?



Length: poor 0+ growth 2013?



Questions

How might flooding affect grayling populations?

- Shared patterns in flow and grayling counts?

How might flooding affect grayling in the future?

- Forecast from observed population patterns




Procedure: “matrix population model”

$$\begin{pmatrix} N_{t+l_1} \\ N_{t+l_2} \\ N_{t+l_3} \end{pmatrix} = \begin{pmatrix} F_1 & F_2 & F_3 \\ S_1 & 0 & 0 \\ 0 & S_2 & 0 \end{pmatrix} \begin{pmatrix} N_{t_1} \\ N_{t_2} \\ N_{t_3} \end{pmatrix} .$$

1. Use observed data to calculate age-specific:
 - Death rate
 - Birth rate






Procedure: “matrix population model”

$$\begin{pmatrix} N_{t+l_1} \\ N_{t+l_2} \\ N_{t+l_3} \end{pmatrix} = \begin{pmatrix} F_1 & F_2 & F_3 \\ S_1 & 0 & 0 \\ 0 & S_2 & 0 \end{pmatrix} \begin{pmatrix} N_{t_1} \\ N_{t_2} \\ N_{t_3} \end{pmatrix} .$$

1. Use observed data to calculate age-specific:
 - Death rate
 - Birth rate
2. Competition for food among 0+ individuals





Procedure: matrix population model

$$\begin{pmatrix} N_{t+l_1} \\ N_{t+l_2} \\ N_{t+l_3} \end{pmatrix} = \begin{pmatrix} F_1 & F_2 & F_3 \\ S_1 & 0 & 0 \\ 0 & S_2 & 0 \end{pmatrix} \begin{pmatrix} N_{t_1} \\ N_{t_2} \\ N_{t_3} \end{pmatrix} .$$

1. Use observed data to calculate age-specific:
 - Mortalities
 - Fecundities
2. Competition for food among 0+ individuals
3. Calculate “population projection matrix”




Procedure: matrix population model

$$\begin{pmatrix} N_{t+l_1} \\ N_{t+l_2} \\ N_{t+l_3} \end{pmatrix} = \begin{pmatrix} F_1 & F_2 & F_3 \\ S_1 & 0 & 0 \\ 0 & S_2 & 0 \end{pmatrix} \begin{pmatrix} N_{t_1} \\ N_{t_2} \\ N_{t_3} \end{pmatrix} .$$


1. Use observed data to calculate age-specific:
 - Mortalities
 - Fecundities
2. Competition for food among 0+ individuals
3. Calculate “population projection matrix”
4. Forecast future population changes




Population forecast without floods



Population forecast with floods



Population forecast with floods



What does this mean?

- Floods *could* cause population decline
- Juvenile (0+) fish will be most affected
- Sensitivity analysis of model parameters:
 - Identify information gaps
 - Guide additional data collection



What can be done?

- Consider management options
 - Habitat alterations
 - Woody debris
- Test possible implications of management
- Catchment-based water management
- Extend investigation to other species?




Next steps



- Compare effects of flow to other factors:
 - river temperature
 - etc
- Grayling population & climate change
- Individual analyses

Thank you



The Grayling Research Trust





The Grayling Society


Promoting Awareness, Conservation & Angling for Grayling, Worldwide



Observed population change: '09 – '13



Condition factor



the temperature
of environment on

quite rare

erations
(sometimes caused by flow

ashout”
ing population dynamics

ent effects

- Habitat changes

- Generalising the investigations to other species, e.g., trout

