Phys040B: Formulae

SI LEADER: Stephen Iota (siota001@ucr.edu)

Course: Physics 40B (Spring 2019), Prof. Barsukov

LAST UPDATED: April 25, 2019

0 General relationships

$$\mathbf{F}_{\text{net}} = m \frac{\mathrm{d}^2 \mathbf{x}}{\mathrm{d}t^2} \qquad \qquad \mathbf{F} = -\nabla U \qquad \qquad U = -\int \mathbf{F} dx$$

$$K = \frac{p^2}{2m} \qquad \qquad a_{\phi} = v^2/r \qquad \qquad W = -\Delta U$$

$$\omega = 2\pi f \qquad \qquad T = 1/f$$

1 Gravitation

$$\mathbf{F} = \frac{Gm_1m_2}{r^2} \hat{r} \qquad U = -\frac{Gm_1m_2}{r}$$

$$v = \sqrt{GM/r} \qquad T^2 = \frac{4\pi^2}{GM}r^3$$

2 Fluids

$$P = \frac{F}{A}$$

$$P = P_0 + \rho g d$$

$$F_1 = \frac{A_1}{A_2} F_2 + \rho g h A_1$$

$$F_B = \rho_f V_f g$$

$$v_1 A_1 = v_2 A_2$$

$$P + 1/2 \rho v^2 + \rho g h = \text{constant}$$

3 Oscillations

$$x(t) = A\cos wt + \phi$$
 $\omega = \sqrt{k/m}$ $\mathbf{F} = -k\mathbf{x}$ $v_{\text{max}} = A\omega$ $T = 2\pi\sqrt{m/k}$ $U = 1/2kx^2$