
Analysis I

Theorems & Lemmas

Stephan Oseghale

Mathematics Bachelor
ETH Zürich
November 16, 2025

Contents

1	Sequences of Real Numbers	1
1.1	Convergence of Sequences	1
1.2	Convergent Subsequences and Accumulation Points	1
1.3	Addition, Multiplication and Inequalities	2

1 Sequences of Real Numbers

1.1 Convergence of Sequences

Definition 1.1: Sequences

A **sequence** is a function $a : \mathbb{N} \rightarrow \mathbb{R}$. The image $a(n)$ of $n \in \mathbb{N}$ is also written as a_n and is called the n -th element of a . Instead of $a : \mathbb{N} \rightarrow \mathbb{R}$ one often writes $(a_n)_{n \in \mathbb{N}}, (a_n)_{n=0}^{\infty}, (a_n)_{n \geq 0}$.

Definition 1.2: (Eventually) Constant Sequences

A sequence $(x_n)_{n=0}^{\infty}$ is **constant** if $x_n = x_m \forall n, m \in \mathbb{N}$. It is **eventually constant** if there exists $N \in \mathbb{N}$ such that $x_n = x_m \forall n, m \geq N$.

Definition 1.3: Convergence of Sequences

Let $(x_n)_{n=0}^{\infty}$ be a sequence in \mathbb{R} . We say that $(x_n)_{n=0}^{\infty}$ **converges** (or is **convergent**) if $\exists A \in \mathbb{R}$ such that

$$\forall \varepsilon > 0 \exists N \in \mathbb{N} : |x_n - A| < \varepsilon \quad \forall n \geq N.$$

In this case we write

$$\lim_{n \rightarrow \infty} x_n = A \tag{1.1}$$

and call A the **limit** of $(x_n)_{n=0}^{\infty}$.

Lemma 1.4: Uniqueness of the Limit

A convergent sequence $(x_n)_{n=0}^{\infty}$ has exactly one limit.

Proof. Let $A, B \in \mathbb{R}$ be limits of $(x_n)_{n=0}^{\infty}$. Fix $\varepsilon > 0$. Then there exists $N_A, N_B \in \mathbb{N}$ such that $|x_n - A| < \varepsilon$ for all $n \geq N_A$ and $|x_n - B| < \varepsilon$ for all $n \geq N_B$. We define $N := \max\{N_A, N_B\}$. Then it holds that

$$|A - B| \leq |A - x_N| + |x_N - B| < \varepsilon + \varepsilon = 2\varepsilon.$$

Since $\varepsilon > 0$ is arbitrary, it follows that $A = B$. □

1.2 Convergent Subsequences and Accumulation Points

Definition 1.5: Subsequences

Let $(x_n)_{n=0}^{\infty}$ be a sequence. A **subsequence** is of the form $(x_{n_k})_{k=0}^{\infty}$, where $(n_k)_{k=0}^{\infty}$ is a strictly increasing sequence of non-negative integers, i.e., $n_{k+1} > n_k \forall k \in \mathbb{N}$.

Remark 1.6. *Since $n_{k+1} > n_k$ for all $k \in \mathbb{N}$ it follows by induction that $n_k \geq k$ for all $k \in \mathbb{N}$.*

Proof. For $k = 0$ we have that $n_0 \geq 0$, because $(n_k)_{k=0}^{\infty}$ is a sequence of non-negative integers. So the condition is fulfilled. For the inductive step we want to show that the condition holds for $k + 1$ under the assumption that the condition is true for k . Because $(n_k)_{k=0}^{\infty}$ is also a strictly increasing sequence, we have that $n_{k+1} > n_k \geq k$. Additionally since $n_k \in \mathbb{N}$, we have that $n_{k+1} \geq n_k + 1$. So it follows that $n_{k+1} \geq n_k + 1 \geq k + 1$, which proves the condition for $k + 1$. □

Lemma 1.7: Subsequences of Convergent Sequences are Convergent

Let $(x_n)_{n=0}^{\infty}$ be a sequence converging to $A \in \mathbb{R}$. Then every subsequence $(x_{n_k})_{k=0}^{\infty}$ also converges to A .

Proof. Let $(x_n)_{n=0}^{\infty}$ be a sequence converging to $A \in \mathbb{R}$. Fix $\varepsilon > 0$. Since $(x_n)_{n=0}^{\infty}$ converges to A , there exists $N \in \mathbb{N}$ such that $|x_n - A| < \varepsilon \forall n \geq N$. As by Remark 1.6 we know that $n_k \geq k$ for all $k \in \mathbb{N}$. Therefore for all $k \geq N$ it holds that $|x_{n_k} - A| < \varepsilon$. \square

Definition 1.8: Accumulation Points of Sequences

Let $(x_n)_{n=0}^{\infty}$ be a sequence in \mathbb{R} . A point $A \in \mathbb{R}$ is an **accumulation point** of $(x_n)_{n=0}^{\infty}$ if

$$\forall \varepsilon > 0 \forall N \in \mathbb{N} \exists n \geq N : |x_n - A| < \varepsilon.$$

Proposition 1.9: Subsequences and Accumulation Points

Let $(x_n)_{n=0}^{\infty}$ be a sequence in \mathbb{R} . A point A is an accumulation point of $(x_n)_{n=0}^{\infty}$ if and only if there exists a convergent subsequence of $(x_n)_{n=0}^{\infty}$ with limit A .

Corollary 1.10: Infinitely Many Terms Near an Accumulation Point

If $A \in \mathbb{R}$ is an accumulation point of $(x_n)_{n=0}^{\infty}$, then for every $\varepsilon > 0$ there are infinitely many n with $x_n \in (A - \varepsilon, A + \varepsilon)$.

Proof. By Proposition 1.9, there exists a subsequence $(x_{n_k})_{k=0}^{\infty}$ with $\lim_{k \rightarrow \infty} = A$. Hence for every $\varepsilon > 0$ there exists K such that $x_{n_k} \in (A - \varepsilon, A + \varepsilon)$ for all $k \geq K$, providing infinitely many elements of the sequence inside the interval $(A - \varepsilon, A + \varepsilon)$. \square

Corollary 1.11: Accumulation Points of Convergent Sequences

convergent sequence has exactly one accumulation point, namely its limit.

1.3 Addition, Multiplication and Inequalities

Proposition 1.12: Limits and Operations

Let $(x_n)_{n=0}^{\infty}$ and $(y_n)_{n=0}^{\infty}$ be sequences converging to $A, B \in \mathbb{R}$ respectively. Then:

1. The sequence $(x_n + y_n)_{n=0}^{\infty}$ converges to $A + B$.
2. The sequence $(x_n y_n)_{n=0}^{\infty}$ converges to AB .
3. Given $\alpha \in \mathbb{R}$, the sequence $(\alpha x_n)_{n=0}^{\infty}$ converges to αA .
4. Suppose $x_n \neq 0$ for all $n \in \mathbb{N}$ and $A \neq 0$. Then the sequence $(x_n^{-1})_{n=0}^{\infty}$ converges to A^{-1} .

Proposition 1.13: Limits and Inequalities

Let $(x_n)_{n=0}^{\infty}$ and $(y_n)_{n=0}^{\infty}$ be sequences converging to $A, B \in \mathbb{R}$ respectively.

1. If $A < B$, then there exists $N \in \mathbb{N}$ such that $x_n < y_n$ for all $n \geq N$.
2. If there exists $N \in \mathbb{N}$ such that $x_n \leq y_n$ for all $n \geq N$, then $A \leq B$.

Remark 1.14. In Proposition 1.13 even if we assume that $x_n < y_n$ for all $n \in \mathbb{N}$, we cannot conclude that $A < B$. for example take

$$x_n = \frac{1}{n}, \quad y_n = \frac{1}{n}.$$

Then we have that $x_n < y_n$ for all $n \in \mathbb{N}$ but $A = B = 0$.

Lemma 1.15: Sandwich Lemma

Let $(x_n)_{n=0}^{\infty}, (y_n)_{n=0}^{\infty}, (z_n)_{n=0}^{\infty}$ be sequences such that for some $N \in \mathbb{N}$, we have that

$$x_n \leq y_n \leq z_n \quad \forall n \geq N.$$

Suppose that both $(x_n)_{n=0}^{\infty}$ and $(z_n)_{n=0}^{\infty}$ converge to the same limit. Then $(y_n)_{n=0}^{\infty}$ also converges, and we have that

$$\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} y_n = \lim_{n \rightarrow \infty} z_n.$$

Proof. Let $(x_n)_{n=0}^{\infty}, (y_n)_{n=0}^{\infty}, (z_n)_{n=0}^{\infty}$ be sequences such that for some $N_0 \in \mathbb{N}$, we have that

$$x_n \leq y_n \leq z_n \quad \forall n \geq N_0.$$

Additionally suppose that $(x_n)_{n=0}^{\infty}$ and $(z_n)_{n=0}^{\infty}$ converge to $A \in \mathbb{R}$. Fix $\varepsilon > 0$. Since $(x_n)_{n=0}^{\infty}, (z_n)_{n=0}^{\infty}$ converge to A there exists $N_x, N_z \in \mathbb{N}$ such that

$$\begin{aligned} A - \varepsilon &< x_n < A + \varepsilon \quad \forall n \geq N_x \\ A - \varepsilon &< z_n < A + \varepsilon \quad \forall n \geq N_z. \end{aligned}$$

So we choose $N := \max\{N_0, N_x, N_z\}$. Then we have that

$$A - \varepsilon < x_n \leq y_n \leq z_n < A + \varepsilon \quad \forall n \geq N,$$

which shows that $\lim_{n \rightarrow \infty} y_n = A$. □