
Analysis I

Theorems & Lemmas

Stephan Oseghale

Mathematics Bachelor
ETH Zürich
November 24, 2025

Contents

1	Functions	1
2	The Real Numbers	2
2.1	Groups, Rings, Fields	2
2.2	Order Relation	4
2.3	Ordered Fields	5
2.4	Completeness Axiom	8
2.5	Intervals	8
2.6	Complex Numbers	9
2.7	Consequences of Completeness	12
3	Sequences of Real Numbers	12
3.1	Convergence of Sequences	12
3.2	Convergent Subsequences and Accumulation Points	13
3.3	Addition, Multiplication and Inequalities	14
3.4	Superior and Inferior Limits	16
3.5	Cauchy Sequences	18
3.6	Improper Limits	19
3.7	Sequences of Complex Numbers	20

1 Functions

Definition 1.1: Functions/Maps/Transformations

A **function** f from a set X to a set Y is an assignment of an element of Y to each element of X . The element $y \in Y$ to which $x \in X$ is assigned to is denoted $f(x)$. We write $f : X \rightarrow Y$ and sometimes also speak of a **map**, **mapping** or a **transformation**. The set X is the **domain** and the set Y is the **codomain**. We refer to the set X as **domain** or **domain of definition**, and the set Y as **domain of values** or **codomain**. The set

$$\{(x, f(x)) \mid x \in X\} \subseteq X \times Y$$

is called the **graph** of f . In the context of a function $f : X \rightarrow Y$, an element of the domain of definition is also called **argument**, and an element $y = f(x) \in Y$ assumed by the function, is also called **value** of the function. If $f : X \rightarrow Y$ is a function, one also writes

$$\begin{aligned} f : X &\rightarrow Y \\ x &\mapsto f(x), \end{aligned}$$

where $f(x)$ could be a concrete formula. We pronounce ' \mapsto ' as 'is mapped to'. Two functions $f_1 : X_1 \rightarrow Y_1$ and $f_2 : X_2 \rightarrow Y_2$ are said to be equal if $X_1 = X_2$, $Y_1 = Y_2$ and $f_1(x) = f_2(x) \quad \forall x \in X_1$.

Definition 1.2: Injective, Surjective and Bijective Functions

Let $f : X \rightarrow Y$ be a function. We call f :

1. **injective** (or an **injection**) if

$$\forall x_1, x_2 \in X : x_1 \neq x_2 \Rightarrow f(x_1) \neq f(x_2);$$

2. **surjective** (or a **surjection**) if

$$\forall y \in Y \exists x \in X : f(x) = y;$$

3. **bijective** (or a **bijection**) if f is both injective and surjective.

Thus, a function $f : X \rightarrow Y$ is *not* injective if there exists distinct $x_1 \neq x_2 \in X$ such that $f(x_1) = f(x_2)$, and *not* surjective if there exists $y \in Y$ such that $f(x) \neq y$ for all $x \in X$.

Definition 1.3: Image and Preimage of a Function

For $f : X \rightarrow Y$ and $A \subseteq X$, define the **image** of A under the function f as

$$f(A) := \{y \in Y \mid \exists x \in X : f(x) = y\}.$$

For $B \subseteq Y$, define the **preimage** of B under the function f as

$$f^{-1}(B) := \{x \in X \mid f(x) \in B\}.$$

Remark 1.4. *Saying that $f : X \rightarrow Y$ is surjective is equivalent to $f(X) = Y$. Equivalently, f is surjective if $f^{-1}(\{y\}) \neq \emptyset$ for all $y \in Y$.*

2 The Real Numbers

2.1 Groups, Rings, Fields

Definition 2.1: Groups

A **group** is a non-empty set G together with a rule (called an *operation*) denoted by $\star : G \times G \rightarrow G$ that combines any two elements of G into another element of G . This operation must satisfy three conditions:

- **Associativity:** No matter how you place parentheses, the result is the same for all $a, b, c \in G$,

$$(a \star b) \star c = a \star (b \star c).$$

- **Neutral element:** There is a special element $e \in G$ such that combining it with any $a \in G$ leaves a unchanged, i.e.,

$$\forall a \in G : a \star e = e \star a = a.$$

- **Inverse element:** Every $a \in G$ has a 'partner' $a^{-1} \in G$ that 'cancels it out', giving the neutral element, i.e.,

$$a \star a^{-1} = a^{-1} \star a = e.$$

Note that, in general, one does not require that $a \star b = b \star a$. If the order of the operation does not matter, i.e., $a \star b = b \star a$ for all $a, b \in G$, the group is called **commutative** or **abelian**.

Lemma 2.2: Basic Properties of Groups

Let (G, \star) be a group. Then:

1. The neutral element is unique.
2. The inverse of an element is unique.
3. The inverse of the inverse of an element is the element itself, namely $(a^{-1})^{-1} = a$ for all $a \in G$.

Proof. 1. Assume that, in addition to $e \in G$, we have a second element e' with the property that $e' \star a = a \star e' = a$ for all elements $a \in G$. Then, we can choose $a = e$ to obtain

$$e \star e' = e.$$

Similarly, since e is a neutral element, we have

$$e \star e' = e'.$$

Combining the two identities, we get

$$e = e \star e' = e'.$$

This proves that $e' = e$, so we speak of *the* neutral element of a group.

2. Assume that for an element $a \in G$, there exists two elements $b, c \in G$ that are both the inverse of a , namely

$$a \star b = b \star a = e, \quad a \star c = c \star a = e.$$

Then, using associativity, we observe that

$$b = b \star e = b \star (a \star c) = (b \star a) \star c = e \star c = c.$$

This proves that the inverse of an element a is unique, so we can speak of *the* inverse element, and the notation a^{-1} makes sense.

3. Since $a \star a^{-1} = e$, we deduce that a is the inverse element of a^{-1} , thus

$$(a^{-1})^{-1} = a. \quad (2.1)$$

□

Groups capture the idea of combining elements with a single operation. But to describe the arithmetic of numbers more faithfully, we also need a second operation (as we do with addition and multiplication). This leads us to the notion of *rings* and *fields*.

Definition 2.3: Rings and Fields

A **ring** is a non-empty set R in which we can both 'add' and 'multiply' elements with two operations '+' and '·'. Also, these two operations are compatible with each other. More precisely:

- $(R, +)$ is a **commutative group**, with neutral element denoted 0.
- Multiplication \cdot is **associative**, has a **neutral element** (usually written as 1), and **distributes over addition**, i.e.,

$$a \cdot (b + c) = a \cdot b + a \cdot c, \quad (b + c) \cdot a = b \cdot a + c \cdot a \quad \forall a, b, c \in R.$$

If multiplication is also commutative, we call $(R, +, \cdot)$ a **commutative ring**. Note that, unlike addition, we do not require that every element has an inverse for multiplication. A **field** is a special kind of commutative ring, i.e. every non-zero element has an inverse for multiplication. In other words, if $(R, +, \cdot)$ is a commutative ring, then $(R, +, \cdot)$ is a field if $R \setminus \{0\}$ forms a commutative group under multiplication. Traditionally, we use the letter F to denote a field. We also write $F^* = F \setminus \{0\}$ for the set of all invertible elements of F .

Lemma 2.4: Basic Properties of Fields

Let $(F, +, \cdot)$ be a field and let $a, b \in F$. Then:

1. $0 \cdot a = a \cdot 0 = 0$.
2. $a \cdot (-b) = -(a \cdot b) = (-a) \cdot b$. In particular $(-1) \cdot a = -a$.
3. $(-a) \cdot (-b) = a \cdot b$. In particular, $(-a)^{-1} = -(a^{-1})$ whenever $a \neq 0$.

Proof. 1. Since 0 is the neutral element for the addition, we have $0+0=0$. Hence, using distributivity,

we get

$$0 \cdot a = (0 + 0) \cdot a = (0 \cdot a) + (0 \cdot a).$$

Adding $-0 \cdot a$ (i.e., the inverse of $0 \cdot a$), we deduce that $0 \cdot a = 0$. The case of $a \cdot 0$ is analogous.

2. By the distributive law,

$$a \cdot b + a \cdot (-b) = a \cdot (b + (-b)) = a \cdot 0 = 0.$$

So $a \cdot (-b)$ is the additive inverse of $a \cdot b$, i.e., $-(a \cdot b) = a \cdot (-b)$. Taking $b = 1$ gives $-a = (-1) \cdot a$. The validity of $(-a) \cdot b = -(a \cdot b)$ follows by exchanging a and b in the argument above.

3. By 2. we know that $-(a \cdot b) = a \cdot (-b)$. Hence, recalling Equation 2.1,

$$a \cdot b = -(a \cdot (-b)).$$

On the other hand, applying 2. with $(-b)$ instead of b , we also have

$$-(a \cdot (-b)) = (-a) \cdot (-b).$$

Combining the two identities above, we conclude that $(-a) \cdot (-b) = a \cdot b$. Finally, taking $b = a^{-1}$ yields $(-a) \cdot (-a^{-1}) = a \cdot a^{-1} = 1$, which gives the second assertion. \square

2.2 Order Relation

Definition 2.5: Cartesian Product

Let X and Y be two sets. The **cartesian product** $X \times Y$ is the set of ordered pairs of elements of X and Y , i.e.,

$$X \times Y := \{(x, y) \mid x \in X, y \in Y\}.$$

Definition 2.6: Subsets

Let P and Q be sets. Then

- P is a **subset** of Q , written $P \subset Q$ (or $P \subseteq Q$), if every element of P also belongs to Q .
- P is a **proper subset** of Q , written $P \subsetneq Q$, if P is a subset of Q but $P \neq Q$.
- We write $P \not\subseteq Q$ if P is not a subset of Q .

Definition 2.7: Relations

Let X be a set. A **relation** on X is a subset $\mathcal{R} \subseteq X \times X$, that is, a collection of ordered pairs of elements of X . If $(x, y) \in \mathcal{R}$ we write $x\mathcal{R}y$. Common symbols for relations include $<, \leq, \sim, \equiv, \cong$. If \sim is a relation on X , we write $x \not\sim y$ if $x \sim y$ does not hold. A relation \sim may have the following properties:

1. **Reflexive:** if $x \sim x \quad \forall x \in X$.
2. **Transitive:** if $x \sim y$ and $y \sim z$, then $x \sim z$.
3. **Symmetric:** if $x \sim y$, then $y \sim x$.
4. **Antisymmetric:** if $x \sim y$ and $y \sim x$, then $x = y$.

A relation is an **equivalence relation** if it is reflexive, transitive and symmetric. It is an **order relation** if it is reflexive, transitive and antisymmetric.

2.3 Ordered Fields**Definition 2.8: Ordered Field**

Let F be a field, and let \leq be an order relation on F . We call (F, \leq) , or simply F , an **ordered field** if the following hold:

1. **Linearity of order:** for all $x, y \in F$, at least one of $x \leq y$ or $y \leq x$ holds.
2. **Compatibility with addition:** for all $x, y, z \in F$,

$$x \leq y \Rightarrow x + z \leq y + z.$$

3. **Compatibility with multiplication:** for all $x, y \in F$,

$$0 \leq x \wedge 0 \leq y \Rightarrow 0 \leq x \cdot y.$$

Lemma 2.9: Ordered Field: Basic Consequences

Let (F, \leq) be an ordered field, and let $x, y, z, w \in F$. Then:

- (a) (Trichotomy) Either $x < y$, or $x = y$, or $x > y$.
- (b) If $x < y$ and $y \leq z$, then $x < z$. (Analogously, $x \leq y$ and $y < z$ imply $x < z$.)
- (c) (Addition of inequalities) If $x \leq y$ and $z \leq w$, then $x + z \leq y + w$. (Analogously, $x < z$ and $z \leq w$ imply $x + z < y + w$.)
- (d) $x \leq y$ if and only if $0 \leq y - x$.
- (e) $x \leq 0$ if and only if $0 \leq -x$.
- (f) $x^2 \geq 0$, and $x^2 > 0$ if $x \neq 0$.
- (g) $0 < 1$.
- (h) If $0 \leq x$ and $y \leq z$, then $xy \leq xz$.
- (i) If $x \leq 0$ and $y \leq z$, then $xy \geq xz$.
- (j) If $0 < x \leq y$, then $0 < y^{-1} \leq x^{-1}$.
- (k) If $0 \leq x \leq y$ and $0 \leq z \leq w$, then $0 \leq xz \leq yw$.
- (l) If $x + y \leq x + z$, then $y \leq z$.
- (m) If $xy \leq xz$ and $x > 0$, then $y \leq z$.

Lemma 2.10: Integers and Rationals Inside an Ordered Field

Let (F, \leq) be an ordered field, and denote by 0 and 1 the neutral elements for addition and multiplication, respectively. Then:

- (i) The elements $\dots, -2, -1, 0, 1, 2, \dots$ defined by

$$2 = 1 + 1, \quad 3 = 2 + 1, \dots, \quad -n = (-1) \cdot n$$

are all distinct and satisfy

$$\dots < -2 < -1 < 0 < 1 < 2 < 3 < \dots$$

We denote this set of elements by \mathbb{Z} , and we call them 'integers'

- (ii) Every fraction pq^{-1} with $p, q \in \mathbb{Z}$, $q \neq 0$, lies in F and the set of all such elements is denoted by \mathbb{Q} . Also,

$$\mathbb{Z} \subsetneq \mathbb{Q} \subseteq F.$$

Proof. (i) By Lemma 2.9(g), we have that $0 < 1$. Then Lemma 2.9(c) yields $0 < 1 < 2 < 3 < \dots$, and taking negatives gives $\dots < -2 < -1 < 0$. Hence all these elements are distinct.

(ii) For $q \neq 0$, q is invertible in F ; define $\frac{p}{q} = pq^{-1}$. The set of such fractions is a field contained in F , which we denote by \mathbb{Q} .

To show that \mathbb{Q} strictly contains \mathbb{Z} , consider $\frac{1}{2}$ (the inverse of 2). Since $2 > 1$, it follows from

Lemma 2.9(j) that $0 < \frac{1}{2} < 1$, so $\frac{1}{2} \notin \mathbb{Z}$. □

Definition 2.11: Absolute Value and Sign

Let (F, \leq) be an ordered field.

- The **absolute value** (or **modulus**) is the function $|\cdot| : F \rightarrow F$ defined by

$$|x| = \begin{cases} x, & x \geq 0, \\ -x, & x < 0. \end{cases}$$

- The **sign** is the function $\text{sgn} : F \rightarrow \{-1, 0, 1\}$ defined by

$$\text{sgn}(x) = \begin{cases} -1, & x < 0, \\ 0, & x = 0, \\ 1, & x > 0. \end{cases}$$

Lemma 2.12: Absolute Value and Sign: Basic Properties

Let (F, \leq) be an ordered field and let $x, y \in F$. Then:

- (a) $x = \text{sgn}(x)|x|$, $|-x| = |x|$, $\text{sgn}(-x) = -\text{sgn}(x)$.
- (b) $|x| \geq 0$, and $|x| = 0$ if and only if $x = 0$ (by Trichotomy Lemma ??).
- (c) (Multiplicativity) $\text{sgn}(xy) = \text{sgn}(x)\text{sgn}(y)$ and $|xy| = |x||y|$.
- (d) If $x \neq 0$, then $|x^{-1}| = |x|^{-1}$.
- (e) $|x| \leq y$ iff $-y \leq x \leq y$.
- (f) $|x| < y$ iff $-y < x < y$.
- (g) (Triangle inequality) $|x + y| \leq |x| + |y|$.
- (h) (Inverse triangle inequality) $||x| - |y|| \leq |x - y|$.

Proof. (g) Thanks to (e) we have $-|x| \leq x \leq |x|$ and $-|y| \leq y \leq |y|$. Adding these two inequalities we get

$$-(|x| + |y|) \leq x + y \leq |x| + |y|.$$

Applying (e) again yields the result.

- (h) From (g) we have $|x| \leq |x - y| + |y|$, therefore

$$|x| - |y| \leq |x - y|.$$

Exchanging the roles of x and y , we also have $|y| - |x| \leq |y - x| = |x - y|$. Combining these two inequalities yields

$$-|x - y| \leq |x| - |y| \leq |x - y|,$$

and the result follows by applying (e) again. □

2.4 Completeness Axiom

Definition 2.13: Completeness Axiom

Let (K, \leq) be an ordered field. We say that (K, \leq) is **complete** (or a **completely ordered field**) if the following statement holds:

Let X, Y be non-empty subsets of K such that $x \leq y$ for all $x \in X$ and $y \in Y$. Then there exists $c \in K$ lying between X and Y , in the sense that $x \leq c \leq y$ for all $x \in X$ and $y \in Y$.

The statement above is called the **completeness axiom**.

Definition 2.14: Real Numbers

We call **the field of real numbers**, any completely ordered field and denote it by \mathbb{R} .

2.5 Intervals

Definition 2.15: Intervals

Let $a, b \in \mathbb{R}$. We define:

- The **closed interval**

$$[a, b] := \{x \in \mathbb{R} \mid a \leq x \leq b\};$$

- The **open interval**

$$(a, b) := \{x \in \mathbb{R} \mid a < x < b\};$$

- The **half-open intervals**

$$[a, b) := \{x \in \mathbb{R} \mid a \leq x < b\} \quad \text{and} \quad (a, b] := \{x \in \mathbb{R} \mid a < x \leq b\};$$

- The **unbounded closed intervals**

$$[a, \infty) := \{x \in \mathbb{R} \mid a \leq x\} \quad \text{and} \quad (-\infty, b] := \{x \in \mathbb{R} \mid x \leq b\};$$

- The **unbounded open intervals**

$$(a, \infty) := \{x \in \mathbb{R} \mid a < x\} \quad \text{and} \quad (-\infty, b) := \{x \in \mathbb{R} \mid x < b\};$$

Definition 2.16: Set Operations

Let P, Q be sets. The **intersection** $P \cap Q$, the **union** $P \cup Q$, the **relative complement** $P \setminus Q$ and the **symmetric difference** $P \triangle Q$ are defined by

$$\begin{aligned} P \cap Q &= \{x \mid x \in P \text{ and } x \in Q\}, \\ P \cup Q &= \{x \mid x \in P \text{ or } x \in Q\}, \\ P \setminus Q &= \{x \mid x \in P \text{ and } x \notin Q\}, \\ P \triangle Q &= (P \setminus Q) \cup (Q \setminus P) = (P \cup Q) \setminus (P \cap Q). \end{aligned}$$

Definition 2.17: Union and Intersection of several Sets

Let \mathcal{A} be a family of sets (i.e., a set whose elements are sets). We define the **union** and **intersection** of the sets in \mathcal{A} as

$$\bigcup_{A \in \mathcal{A}} A = \{x \mid \exists A \in \mathcal{A} : x \in A\}, \quad \bigcap_{A \in \mathcal{A}} A = \{x \mid \forall A \in \mathcal{A} : x \in A\}.$$

If $\mathcal{A} = \{A_1, A_2, \dots\}$, we also write

$$\bigcup_{i=1}^{\infty} A_i = \{x \mid \exists i \geq 1 : x \in A_i\}, \quad \bigcap_{i=1}^{\infty} A_i = \{x \mid \forall i \geq 1 : x \in A_i\}.$$

Definition 2.18: Neighborhoods

Let $x \in \mathbb{R}$. A **neighborhood** of x is a set containing an interval I such that $x \in I$. Given $\delta > 0$, the open interval $(x - \delta, x + \delta)$ is called the δ -**neighborhood** of x .

Definition 2.19: Open and Closed Sets

A subset $U \subseteq \mathbb{R}$ is called **open** in \mathbb{R} if for every $x \in U$ there exists open interval I such that $x \in I$ and $I \subseteq U$. A subset $F \subseteq \mathbb{R}$ is called **closed** in \mathbb{R} if its complement $\mathbb{R} \setminus F$ is open.

Remark 2.20. The sets \emptyset and \mathbb{R} are both open in \mathbb{R} . Hence, they are also closed since $\emptyset^c = \mathbb{R}$ and $\mathbb{R}^c = \emptyset$. We note that $\mathbb{Q} \subseteq \mathbb{R}$ and $[a, b) \subseteq \mathbb{R}$ are neither open nor closed.

Remark 2.21. Let \mathcal{U} be a family of open sets, and \mathcal{F} be a family of closed subsets of \mathbb{R} . Then the union and intersection

$$\bigcup_{U \in \mathcal{U}} U, \quad \bigcap_{F \in \mathcal{F}} F$$

are open and closed, respectively.

2.6 Complex Numbers

Starting from the field of real numbers \mathbb{R} , we define the set of **complex numbers** as

$$\mathbb{C} = \mathbb{R}^2 = \{(x, y) \mid x, y \in \mathbb{R}\}.$$

We denote the elements $z = (x, y) \in \mathbb{C}$ in the form $z = x + iy$, where i is the **imaginary unit**. Here $x \in \mathbb{R}$ is the **real part** of z , written as $x = \operatorname{Re}(z)$, and $y \in \mathbb{R}$ is the **imaginary part**, written

as $y = \text{Im}(z)$. Elements with $\text{Im}(z) = 0$ are called **real**, while those with $\text{Re}(z) = 0$ are **purely imaginary**. Via the injective map $\mathbb{R} \ni x \mapsto x + i \cdot 0 \in \mathbb{C}$, we identify \mathbb{R} with the subset of real numbers inside \mathbb{C} .

As you may expect from previous knowledge, we want to satisfy $i^2 = -1$. To achieve this, we define addition and multiplication on \mathbb{C} so that it becomes a field. Additionally, we want these operations to coincide with the usual addition and multiplication when considering real numbers.

Since $i^2 = -1$, using commutativity and distributivity we get

$$(x_1 + iy_1)(x_2 + iy_2) = x_1x_2 + ix_1y_2 + iy_2x_1 + i^2y_1y_2 = (x_1x_2 - y_1y_2) + i(x_1y_2 + y_1x_2).$$

This motivates the following definition

Definition 2.22: Addition and Multiplication on \mathbb{C}

On $\mathbb{C} = \mathbb{R} \times \mathbb{R}$ we define **addition** and **multiplication** as follows:

$$\begin{aligned} (x_1, y_1) + (x_2, y_2) &= (x_1 + x_2, y_1 + y_2), \\ (x_1, y_1) \cdot (x_2, y_2) &= (x_1x_2 - y_1y_2, x_1y_2 + x_2y_1). \end{aligned}$$

Proposition 2.23: \mathbb{C} is a Field

With the operation of Definition 2.22, together with the zero element $(0, 0)$ and the unit element $(1, 0)$, the set \mathbb{C} is a field.

Definition 2.24: Complex Conjugation

For $z = x + iy \in \mathbb{C}$ we define its **conjugate** as $\bar{z} = x - iy$. The mapping $\mathbb{C} \ni z \mapsto \bar{z} \in \mathbb{C}$ is called **complex conjugation**.

Lemma 2.25: Properties of Complex Conjugation

For all $z, w \in \mathbb{C}$:

- (i) $z\bar{z} = x^2 + y^2 \in \mathbb{R}_{\geq 0}$. In particular, $z\bar{z} = 0$ if and only if $z = 0$.
- (ii) $\overline{z + w} = \bar{z} + \bar{w}$.
- (iii) $\overline{z\bar{w}} = \bar{z}w$.

Proof. Property (i) follows from the fact that, for $z = x + iy$, $(x + iy)(x - iy) = x^2 + y^2$. Also, $x^2 + y^2 = 0$ if and only if $x + iy = 0$. Properties (ii) and (iii) follow from a direct computation, writing $z = x_1 + iy_1$ and $w = x_2 + iy_2$, which yields

$$\begin{aligned} \overline{z + w} &= \overline{(x_1 + x_2) + i(y_1 + y_2)} = (x_1 + x_2) - i(y_1 + y_2) = (x_1 - iy_1) + (x_2 - iy_2) = \bar{z} + \bar{w}, \\ \overline{z \cdot w} &= \overline{(x_1x_2 - y_1y_2) + i(x_1y_2 + x_2y_1)} = (x_1x_2 - y_1y_2) - i(x_1y_2 + x_2y_1) \\ &= (x_1 - iy_1) \cdot (x_2 - iy_2) = \bar{z} \cdot \bar{w}. \quad \square \end{aligned}$$

Definition 2.26: Absolute Value

The **absolute value** (or **norm**) on \mathbb{C} is the map $|\cdot| : \mathbb{C} \rightarrow \mathbb{R}$ given by

$$|z| = \sqrt{z\bar{z}} = \sqrt{x^2 + y^2}, \quad z = x + iy \in \mathbb{C}.$$

Lemma 2.27: Cauchy-Schwarz Inequality

If $z = x_1 + iy_1$, and $w = x_2 + iy_2$, then

$$x_1x_2 + y_1y_2 \leq |z||w|. \quad (2.2)$$

Proof. We observe that

$$\begin{aligned} |z|^2|w|^2 - (x_1x_2 + y_1y_2)^2 &= (x_1^2 + y_1^2)(x_2^2 + y_2^2) - (x_1x_2 + y_1y_2)^2 \\ &= x_1^2x_2^2 + y_1^2y_2^2 + y_1^2x_2^2 + x_1^2y_2^2 - (x_1^2x_2^2 + y_1^2y_2^2 + 2x_1x_2y_1y_2) \\ &= y_1^2x_2^2 + x_1^2y_2^2 - 2x_1x_2y_1y_2 \\ &= (y_1x_2 - x_1y_2)^2 \geq 0. \end{aligned}$$

This proves that $(x_1x_2 + y_1y_2)^2 \leq |z|^2|w|^2$, so it follows that

$$|x_1x_2 + y_1y_2| \leq |z||w|.$$

Since $x \leq |x|$ for all $x \in \mathbb{R}$, we obtain Equation 2.2. □

Proposition 2.28: Triangle Inequality

For all $z, w \in \mathbb{C}$, one has

$$|z + w| \leq |z| + |w|.$$

Proof. For $z = x_1 + iy_1$ and $w = x_2 + iy_2$, using Lemma 2.27, we have

$$\begin{aligned} |z + w|^2 &= (x_1 + x_2)^2 + (y_1 + y_2)^2 \\ &= |z|^2 + |w|^2 + 2(x_1x_2 + y_1y_2) \\ &\leq |z|^2 + |w|^2 + 2|z||w| = (|z| + |w|)^2. \end{aligned}$$

Taking square roots proves the result. □

Definition 2.29: Circular Disks

For $z \in \mathbb{C}$ and $r > 0$, we define the **open disk** with radius $r > 0$ around z as

$$B(z, r) := \{w \in \mathbb{C} \mid |z - w| < r\},$$

and the **closed disk** with radius $r > 0$ around z as

$$\overline{B(z, r)} := \{w \in \mathbb{C} \mid |z - w| \leq r\}.$$

In other words, the open disk $B(z, r)$ is the set of points at distance strictly less than r from z . We note that this definition is compatible with the one of neighborhoods in \mathbb{R} : if $x \in \mathbb{R}$ and $r > 0$,

then

$$B(x, r) \cap \mathbb{R} = (x - r, x + r).$$

Definition 2.30: Open and Closed Sets

A set $U \subseteq \mathbb{C}$ is **open** if for every $z \in U$ there exists $r > 0$ such that $B(z, r) \subseteq U$. A set $C \subseteq \mathbb{C}$ is **closed** if its complement $\mathbb{C} \setminus C$ is open.

2.7 Consequences of Completeness

3 Sequences of Real Numbers

3.1 Convergence of Sequences

Definition 3.1: Sequences

A **sequence** is a function $a : \mathbb{N} \rightarrow \mathbb{R}$. The image $a(n)$ of $n \in \mathbb{N}$ is also written as a_n and is called the n -th element of a . Instead of $a : \mathbb{N} \rightarrow \mathbb{R}$ one often writes $(a_n)_{n \in \mathbb{N}}$, $(a_n)_{n=0}^{\infty}$, $(a_n)_{n \geq 0}$.

Definition 3.2: (Eventually) Constant Sequences

A sequence $(x_n)_{n=0}^{\infty}$ is **constant** if $x_n = x_m \forall n, m \in \mathbb{N}$. It is **eventually constant** if there exists $N \in \mathbb{N}$ such that $x_n = x_m \forall n, m \geq N$.

Definition 3.3: Convergence of Sequences

Let $(x_n)_{n=0}^{\infty}$ be a sequence in \mathbb{R} . We say that $(x_n)_{n=0}^{\infty}$ **converges** (or is **convergent**) if $\exists A \in \mathbb{R}$ such that

$$\forall \varepsilon > 0 \exists N \in \mathbb{N} : |x_n - A| < \varepsilon \quad \forall n \geq N.$$

In this case we write

$$\lim_{n \rightarrow \infty} x_n = A \tag{3.1}$$

and call A the **limit** of $(x_n)_{n=0}^{\infty}$.

Lemma 3.4: Uniqueness of the Limit

A convergent sequence $(x_n)_{n=0}^{\infty}$ has exactly one limit.

Proof. Let $A, B \in \mathbb{R}$ be limits of $(x_n)_{n=0}^{\infty}$. Fix $\varepsilon > 0$. Then there exists $N_A, N_B \in \mathbb{N}$ such that $|x_n - A| < \varepsilon$ for all $n \geq N_A$ and $|x_n - B| < \varepsilon$ for all $n \geq N_B$. We define $N := \max\{N_A, N_B\}$. Then it holds that

$$|A - B| \leq |A - x_N| + |x_N - B| < \varepsilon + \varepsilon = 2\varepsilon.$$

Since $\varepsilon > 0$ is arbitrary, it follows that $A = B$. □

3.2 Convergent Subsequences and Accumulation Points

Definition 3.5: Subsequences

Let $(x_n)_{n=0}^{\infty}$ be a sequence. A **subsequence** is of the form $(x_{n_k})_{k=0}^{\infty}$, where $(n_k)_{k=0}^{\infty}$ is a strictly increasing sequence of non-negative integers, i.e., $n_{k+1} > n_k \forall k \in \mathbb{N}$.

Remark 3.6. Since $n_{k+1} > n_k$ for all $k \in \mathbb{N}$ it follows by induction that $n_k \geq k$ for all $k \in \mathbb{N}$.

Proof. For $k = 0$ we have that $n_0 \geq 0$, because $(n_k)_{k=0}^{\infty}$ is a sequence of non-negative integers. So the condition is fulfilled. For the inductive step we want to show that the condition holds for $k + 1$ under the assumption that the condition is true for k . Because $(n_k)_{k=0}^{\infty}$ is also a strictly increasing sequence, we have that $n_{k+1} > n_k \geq k$. Additionally since $n_k \in \mathbb{N}$, we have that $n_{k+1} \geq n_k + 1$. So it follows that $n_{k+1} \geq n_k + 1 \geq k + 1$, which proves the condition for $k + 1$. \square

Lemma 3.7: Subsequences of Convergent Sequences are Convergent

Let $(x_n)_{n=0}^{\infty}$ be a sequence converging to $A \in \mathbb{R}$. Then every subsequence $(x_{n_k})_{k=0}^{\infty}$ also converges to A .

Proof. Let $(x_n)_{n=0}^{\infty}$ be a sequence converging to $A \in \mathbb{R}$. Fix $\varepsilon > 0$. Since $(x_n)_{n=0}^{\infty}$ converges to A , there exists $N \in \mathbb{N}$ such that $|x_n - A| < \varepsilon \forall n \geq N$. As by Remark 3.6 we know that $n_k \geq k$ for all $k \in \mathbb{N}$. Therefore for all $k \geq N$ it holds that $|x_{n_k} - A| < \varepsilon$. \square

Definition 3.8: Accumulation Points of Sequences

Let $(x_n)_{n=0}^{\infty}$ be a sequence in \mathbb{R} . A point $A \in \mathbb{R}$ is an **accumulation point** of $(x_n)_{n=0}^{\infty}$ if

$$\forall \varepsilon > 0 \forall N \in \mathbb{N} \exists n \geq N : |x_n - A| < \varepsilon.$$

Proposition 3.9: Subsequences and Accumulation Points

Let $(x_n)_{n=0}^{\infty}$ be a sequence in \mathbb{R} . A point A is an accumulation point of $(x_n)_{n=0}^{\infty}$ if and only if there exists a convergent subsequence of $(x_n)_{n=0}^{\infty}$ with limit A .

Proof. First assume that $A \in \mathbb{R}$ is an accumulation point of $(x_n)_{n=0}^{\infty}$. We construct $(n_k)_{k \geq 0}$ recursively:

- first, apply the definition of accumulation point with $N = 1$ and $\varepsilon = 1 = 2^0$ to find $n_0 \geq 1$ with $|x_{n_0} - A| \leq 2^0$,
- second, apply the definition the definition of accumulation point with $N = n_0 + 1$ and $\varepsilon = 2^{-1}$ to find $n_1 \geq n_0 + 1$ with $|x_{n_1} - A| \leq 2^{-1}$,
- more in general given n_{k-1} , we apply the definition of accumulation point with $N = n_{k-1} + 1$ and $\varepsilon = 2^{-k}$ to find $n_k \geq n_{k-1} + 1$ with $|x_{n_k} - A| \leq 2^{-k}$.

Now given $\varepsilon > 0$ choose N such that $2^{-N} < \varepsilon$. Then for all $k \geq N$ we have that

$$|x_{n_k} - A| \leq 2^{-k} \leq 2^{-N} < \varepsilon,$$

so $\lim_{k \rightarrow \infty} x_{n_k} = A$.

Conversely, assume that there exists a subsequence $(x_{n_k})_{k=0}^{\infty}$ converging to A . Fix $\varepsilon > 0$ and $N \in \mathbb{N}$. Since $\lim_{k \rightarrow \infty} x_{n_k} = A$, there exists N_0 such that $|x_{n_k} - A| < \varepsilon$ for all $k \geq N_0$. Hence if we

choose $k = \max\{N_0, N\}$, because $n_k \geq n$ (recall Remark 3.6) we have that $n_k \geq N$ and $|x_{n_k} - A| < \varepsilon$. Thus A is an accumulation point. \square

Corollary 3.10: Infinitely Many Terms Near an Accumulation Point

If $A \in \mathbb{R}$ is an accumulation point of $(x_n)_{n=0}^\infty$, then for every $\varepsilon > 0$ there are infinitely many n with $x_n \in (A - \varepsilon, A + \varepsilon)$.

Proof. By Proposition 3.9, there exists a subsequence $(x_{n_k})_{k=0}^\infty$ with $\lim_{k \rightarrow \infty} x_{n_k} = A$. Hence for every $\varepsilon > 0$ there exists K such that $x_{n_k} \in (A - \varepsilon, A + \varepsilon)$ for all $k \geq K$, providing infinitely many elements of the sequence inside the interval $(A - \varepsilon, A + \varepsilon)$. \square

Corollary 3.11: Accumulation Points of Convergent Sequences

convergent sequence has exactly one accumulation point, namely its limit.

3.3 Addition, Multiplication and Inequalities

Proposition 3.12: Limits and Operations

Let $(x_n)_{n=0}^\infty$ and $(y_n)_{n=0}^\infty$ be sequences converging to $A, B \in \mathbb{R}$ respectively. Then:

1. *The sequence $(x_n + y_n)_{n=0}^\infty$ converges to $A + B$.*
2. *The sequence $(x_n y_n)_{n=0}^\infty$ converges to AB .*
3. *Given $\alpha \in \mathbb{R}$, the sequence $(\alpha x_n)_{n=0}^\infty$ converges to αA .*
4. *Suppose $x_n \neq 0$ for all $n \in \mathbb{N}$ and $A \neq 0$. Then the sequence $(x_n^{-1})_{n=0}^\infty$ converges to A^{-1} .*

Proposition 3.13: Limits and Inequalities

Let $(x_n)_{n=0}^\infty$ and $(y_n)_{n=0}^\infty$ be sequences converging to $A, B \in \mathbb{R}$ respectively.

1. *If $A < B$, then there exists $N \in \mathbb{N}$ such that $x_n < y_n$ for all $n \geq N$.*
2. *If there exists $N \in \mathbb{N}$ such that $x_n \leq y_n$ for all $n \geq N$, then $A \leq B$.*

Remark 3.14. *In Proposition 3.13 even if we assume that $x_n < y_n$ for all $n \in \mathbb{N}$, we cannot conclude that $A < B$. for example take*

$$x_n = \frac{1}{n}, \quad y_n = \frac{1}{n}.$$

Then we have that $x_n < y_n$ for all $n \in \mathbb{N}$ but $A = B = 0$.

Lemma 3.15: Sandwich Lemma

Let $(x_n)_{n=0}^\infty, (y_n)_{n=0}^\infty, (z_n)_{n=0}^\infty$ be sequences such that for some $N \in \mathbb{N}$, we have that

$$x_n \leq y_n \leq z_n \quad \forall n \geq N.$$

Suppose that both $(x_n)_{n=0}^\infty$ and $(z_n)_{n=0}^\infty$ converge to the same limit. Then $(y_n)_{n=0}^\infty$ also converges, and we have that

$$\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} y_n = \lim_{n \rightarrow \infty} z_n.$$

Proof. Let $(x_n)_{n=0}^\infty, (y_n)_{n=0}^\infty, (z_n)_{n=0}^\infty$ be sequences such that for some $N_0 \in \mathbb{N}$, we have that

$$x_n \leq y_n \leq z_n \quad \forall n \geq N_0.$$

Additionally suppose that $(x_n)_{n=0}^\infty$ and $(z_n)_{n=0}^\infty$ converge to $A \in \mathbb{R}$. Fix $\varepsilon > 0$. Since $(x_n)_{n=0}^\infty, (z_n)_{n=0}^\infty$ converge to A there exists $N_x, N_z \in \mathbb{N}$ such that

$$\begin{aligned} A - \varepsilon < x_n < A + \varepsilon & \quad \forall n \geq N_x \\ A - \varepsilon < z_n < A + \varepsilon & \quad \forall n \geq N_z. \end{aligned}$$

So we choose $N := \max\{N_0, N_x, N_z\}$. Then we have that

$$A - \varepsilon < x_n \leq y_n \leq z_n < A + \varepsilon \quad \forall n \geq N,$$

which shows that $\lim_{n \rightarrow \infty} y_n = A$. □

Definition 3.16: Bounded Sequences

A sequence $(x_n)_{n=0}^\infty$ is called **bounded** if there exists a real number $M \geq 0$ such that

$$|x_n| \leq M \quad \forall n \in \mathbb{N}.$$

Lemma 3.17: Convergent Sequences are Bounded

Every convergent sequence is bounded.

Proof. Let $(x_n)_{n=0}^\infty$ be a sequence converging to $A \in \mathbb{R}$. Let $\varepsilon = 1$. Then, by convergence of $(x_n)_{n=0}^\infty$, there exists N such that $|x_n - A| \leq 1$ for all $n \geq N$. So we have that

$$|x_n| = |x_n - A + A| \leq |x_n - A| + |A| \leq 1 + |A| \quad \forall n \geq N.$$

We choose

$$M = \max(|x_0|, |x_1|, \dots, |x_{N-1}|, 1 + |A|).$$

Then $|x_n| \leq M$ for all $n \in \mathbb{N}$ as desired. □

Definition 3.18: Monotone Sequences

A sequence $(x_n)_{n=0}^\infty$ is called:

- **(monotonically) increasing** if $m > n \Rightarrow x_m \geq x_n$,
- **strictly (monotonically) increasing** if $m > n \Rightarrow x_m > x_n$,
- **(monotonically) decreasing** if $m > n \Rightarrow x_m \leq x_n$,
- **strictly (monotonically) decreasing** if $m > n \Rightarrow x_m < x_n$.

If a sequence is decreasing or increasing we call it monotone. If a sequence is strictly increasing or strictly decreasing then we call it strictly monotone.

Remark 3.19. *An equivalent formulation of monotone sequences can be given using only successive terms:*

- $(x_n)_{n=0}^\infty$ is increasing if $x_{n+1} \geq x_n$ for all n ,
- $(x_n)_{n=0}^\infty$ is strictly increasing if $x_{n+1} > x_n$ for all n ,
- $(x_n)_{n=0}^\infty$ is decreasing if $x_{n+1} \leq x_n$ for all n ,
- $(x_n)_{n=0}^\infty$ is strictly decreasing if $x_{n+1} < x_n$ for all n .

Theorem 3.20: Convergence of Monotone Sequences

A monotone sequence $(x_n)_{n=0}^\infty$ converges if and only if it is bounded. More precisely, let $X = \{x_n \mid n \in \mathbb{N}\}$ denote the set of points in the sequence.

- If $(x_n)_{n=0}^\infty$ is increasing, then $\lim_{n \rightarrow \infty} x_n = \sup(X)$,
- if $(x_n)_{n=0}^\infty$ decreasing, then $\lim_{n \rightarrow \infty} x_n = \inf(X)$.

Proof. If $(x_n)_{n=0}^\infty$ converges Lemma 3.17 says that it's bounded.

Conversely, let $(x_n)_{n=0}^\infty$ be a bounded monotone sequence. Wlog assume that $(x_n)_{n=0}^\infty$ is increasing (otherwise consider $(-x_n)_{n=0}^\infty$). Since $(x_n)_{n=0}^\infty$ is bounded from above, the set $X = \{x_n \mid n \in \mathbb{N}\}$ has a supremum, that we'll call $A = \sup(X)$.

By definition of A :

- (i) $x_n \leq A \quad \forall n \in \mathbb{N}$,
- (ii) $\forall \varepsilon > 0$ there exists $N \in \mathbb{N}$ such that $x_N > A - \varepsilon$.

Then, for all $n \geq N$ using (ii) and monotonicity, we have that $x_n \geq x_N > A - \varepsilon$. Then using (i), we conclude that

$$A - \varepsilon < x_n < A + \varepsilon \quad \forall n \geq N.$$

□

3.4 Superior and Inferior Limits

Let $(x_n)_{n=0}^\infty$ be a bounded sequence. To study its behavior for large n it is useful to look at its tails

$$X_{\geq n} = \{x_k \mid k \geq n\} \subseteq \mathbb{R}.$$

The concept of limits can be restated using the tails of a sequence, i.e., the sequence $(x_n)_{n=0}^\infty$ converges to $A \in \mathbb{R}$ if and only if, for every $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that $X_N \subseteq (A - \varepsilon, A + \varepsilon)$.

However, since not every sequence has a limit we now introduce a related notion (the **superior** and **inferior limits**), which always exist for bounded sequences.

For each $n \in \mathbb{N}$, define

$$s_n = \sup(X_{\geq n}) = \sup_{k \geq n} x_k, \quad i_n = \inf(X_{\geq n}) = \inf_{k \geq n} x_k.$$

Since $X_{\geq m} \subset X_{\geq n}$, whenever $m > n$, we have that

$$i_n \leq i_m \leq s_m \leq s_n \quad \forall m > n.$$

Thus, $(s_n)_{n=0}^\infty$ is a monotonically decreasing sequence, while $(i_n)_{n=0}^\infty$ is a monotonically increasing sequence. Moreover, since $(x_n)_{n=0}^\infty$ is bounded both $(s_n)_{n=0}^\infty$ and $(i_n)_{n=0}^\infty$ are bounded as well. Hence

by Theorem 3.20, both sequences converge. Their limits will be called the *superior* and the *inferior limit* of $(x_n)_{n=0}^{\infty}$ respectively.

Note that, since $x_n \in X_{\geq n}$, we have that

$$i_n \leq x_n \leq s_n \quad \forall n \in \mathbb{N}. \quad (3.2)$$

Definition 3.21: Superior and Inferior Limits

Let $(x_n)_{n=0}^{\infty}$ be a bounded sequence in \mathbb{R} . The numbers

$$\limsup_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} \left(\sup_{k \geq n} x_k \right), \quad \liminf_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} \left(\inf_{k \geq n} x_k \right)$$

are called the **superior** and **inferior limit** of $(x_n)_{n=0}^{\infty}$ respectively. From Equation 3.2 and Proposition 3.13, we have

$$\liminf_{n \rightarrow \infty} x_n \leq \limsup_{n \rightarrow \infty} x_n.$$

Lemma 3.22: Convergence and Superior/Inferior Limits

A bounded sequence $(x_n)_{n=0}^{\infty}$ in \mathbb{R} converges if and only if

$$\limsup_{n \rightarrow \infty} x_n = \liminf_{n \rightarrow \infty} x_n.$$

Proof. For every $n \in \mathbb{N}$, define

$$i_n = \inf_{k \geq n} x_k, \quad s_n = \sup_{k \geq n} x_k,$$

and set

$$I = \lim_{n \rightarrow \infty} i_n = \liminf_{n \rightarrow \infty} x_n, \quad S = \lim_{n \rightarrow \infty} s_n = \limsup_{n \rightarrow \infty} x_n.$$

First suppose that $I = S$. Since $i_n \leq x_n \leq s_n$ (see Equation 3.2), the Sandwich Lemma 3.15 implies that the sequence $(x_n)_{n=0}^{\infty}$ converges, and its limit equals $I = S$.

Conversely, assume that $(x_n)_{n=0}^{\infty}$ converges to $A \in \mathbb{R}$. Given $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$A - \varepsilon < x_n < A + \varepsilon \quad \forall n \geq N.$$

Then for all $n \geq N$, the same inequalities holds for i_n and s_n , i.e.,

$$A - \varepsilon \leq i_n \leq s_n \leq A + \varepsilon.$$

Taking limits and using Proposition 3.13, we obtain

$$A - \varepsilon \leq I \leq S \leq A + \varepsilon.$$

Since $\varepsilon > 0$ is arbitrary, it follows that $A = I = S$, which proves the result. \square

Theorem 3.23: Superior and Inferior Limits are Accumulation Points

Let $(x_n)_{n=0}^{\infty}$ be a bounded sequence and let $A = \limsup_{n \rightarrow \infty} x_n$. Then A is an accumulation point of $(x_n)_{n=0}^{\infty}$, and for every $\varepsilon > 0$ the following hold:

1. only finitely many elements satisfy $x_n \geq A + \varepsilon$;
2. infinitely many elements satisfy $A - \varepsilon < x_n < A + \varepsilon$.

An analogous statement holds for the inferior limit.

Proof. Since the sequence $(s_n)_{n=0}^{\infty}$ is monotonically decreasing and converges to A , given $\varepsilon > 0$, there exists $N_0 \in \mathbb{N}$ such that

$$A \leq s_n < A + \varepsilon \quad \forall n \geq N_0. \quad (3.3)$$

We first prove that A is an accumulation point.

Fix $N \in \mathbb{N}$ and set $N_1 = \max\{N, N_0\}$. Since $s_{N_1} = \sup_{k \geq N_1} x_k$, there exists $n_1 \geq N_1 \geq N_0$ such that

$$s_{N_1} - \varepsilon < x_{n_1} \leq s_{N_1}.$$

Thus, combining this bound with Equation 3.3 we obtain

$$A - \varepsilon < s_{N_1} - \varepsilon < x_{n_1} \leq s_{N_1} < A + \varepsilon.$$

This construct shows that for any $\varepsilon > 0$ and any $N \in \mathbb{N}$, there exists $n_1 \geq N$ such that $A - \varepsilon < x_{n_1} < A + \varepsilon$. Thus A is an accumulation point for $(x_n)_{n=0}^{\infty}$.

We now prove 1. and 2.. From Equation 3.3 we have $x_n < A + \varepsilon$ for all $n \geq N_0$, so only finitely many terms satisfy $x_n \geq A + \varepsilon$. This shows 1..

Also since A is an accumulation point, it follows from Corollary 3.10 that infinitely many terms of the sequence lie within any interval $(A - \varepsilon, A + \varepsilon)$. \square

Corollary 3.24: Bounded Sequences have Convergent Subsequences

Every bounded sequence has at least one accumulation point and therefore possesses a convergent subsequence.

Proof. By Theorem 3.23, the number

$$A = \limsup_{n \rightarrow \infty} x_n$$

is always an accumulation point of $(x_n)_{n=0}^{\infty}$. Moreover, by Proposition 3.9, every accumulation point is the limit of a convergent subsequence. Hence every bounded sequence admits at least one convergent subsequence. \square

3.5 Cauchy Sequences

Definition 3.25: Cauchy Sequences

A sequence $(x_n)_{n=0}^{\infty}$ is called a **Cauchy sequence** if for every $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that

$$|x_n - x_m| < \varepsilon \quad \forall n, m \geq N.$$

Lemma 3.26: Cauchy Sequences are Bounded*Every Cauchy sequence is bounded.*

Proof. By definition, there exists $N \in \mathbb{N}$ such that

$$|x_n - x_N| \leq 1 \quad \forall n \geq N.$$

Hence, for $n \geq N$, we have $|x_n| \leq 1 + |x_N|$. Now, define

$$M = \max\{|x_0|, |x_1|, \dots, |x_{N-1}|, 1 + |x_N|\}.$$

Then, $|x_n| \leq M$ for all $n \in \mathbb{N}$, so $(x_n)_{n=0}^\infty$ is bounded. \square

Theorem 3.27: Convergence and Cauchy Sequences*A sequence $(x_n)_{n=0}^\infty$ of real numbers converges if and only if it is a Cauchy sequence.*

Proof. Suppose first that $(x_n)_{n=0}^\infty$ converges to some $A \in \mathbb{R}$, and let us prove that $(x_n)_{n=0}^\infty$ is a Cauchy sequence.

Given $\varepsilon > 0$, choose $N \in \mathbb{N}$ such that

$$|x_n - A| < \frac{\varepsilon}{2} \quad \forall n \geq N.$$

Then for all $n, m \geq N$, we have that

$$|x_n - x_m| \leq |x_n - A| + |x_m - A| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon,$$

hence $(x_n)_{n=0}^\infty$ is a Cauchy sequence.

Viceversa, let $(x_n)_{n=0}^\infty$ be a Cauchy sequence. Since it is bounded (by Lemma 3.26), Corollary 3.24 implies that there exists a subsequence $(x_{n_k})_{k=0}^\infty$ converging to some $A \in \mathbb{R}$. Given $\varepsilon > 0$, choose $N_0 \in \mathbb{N}$ such that

$$|x_n - x_m| < \frac{\varepsilon}{2} \quad \forall n, m \geq N_0,$$

and choose $N_1 \in \mathbb{N}$ such that

$$|x_{n_k} - A| < \frac{\varepsilon}{2} \quad \forall k \geq N_1.$$

Let $N = \max\{N_0, N_1\}$. Since $n_N \geq N$ (see Remark 3.6), for all $n \geq N$ we have

$$|x_n - A| \leq |x_n - x_{n_N}| + |x_{n_N} - A| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon.$$

Thus, $(x_n)_{n=0}^\infty$ converges to A . \square

3.6 Improper Limits

We now extend the notion of limit to allow the **improper limit values** $+\infty$ (often abbreviated as ∞) and $-\infty$.

Definition 3.28: Improper Limits

Let $(x_n)_{n=0}^{\infty}$ be a sequence in \mathbb{R} . We say $(x_n)_{n=0}^{\infty}$ **diverges to** $+\infty$, and we write

$$\lim_{n \rightarrow \infty} x_n = +\infty,$$

if for every $M > 0$ there exists $N \in \mathbb{N}$ such that $x_n > M$ for all $n \geq N$.

Similarly, $(x_n)_{n=0}^{\infty}$ **diverges to** $-\infty$ if for every $M > 0$ there exists $N \in \mathbb{N}$ such that $x_n < -M$ for all $n \geq N$. In both cases, we say that $(x_n)_{n=0}^{\infty}$ has an **improper limit**.

An unbounded sequence doesn't need to diverge to $+\infty$ or $-\infty$. For instance, the sequence $x_n = (-1)^n n$, is unbounded but neither diverges to $+\infty$ nor to $-\infty$.

The notion of improper limit allows us to extend the definitions of superior and inferior limits to *unbounded* sequences. If $(x_n)_{n=0}^{\infty}$ is not bounded from above, then

$$\sup_{k \geq n} x_k = +\infty \quad \forall n \in \mathbb{N},$$

and we write

$$\limsup_{n \rightarrow \infty} x_n = +\infty.$$

If $(x_n)_{n=0}^{\infty}$ is bounded from above but not from below, then we define

$$\limsup_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} (\sup_{k \geq n} x_k),$$

where the right-hand side is a real limit if the decreasing sequence $\sup_{k \geq n} x_k$ is bounded, and the improper limit $-\infty$ otherwise. The definition of the inferior limit extends analogously.

3.7 Sequences of Complex Numbers

Informally, a **sequence of complex numbers** is just like a sequence of real numbers, except that each term is a complex number instead of a real one. Thus, we study ordered lists (z_0, z_1, \dots) , where $z_n : \mathbb{N} \rightarrow \mathbb{C}$. As in the real case, we are mainly interested in their convergence, divergence and limit behavior.

To analyze sequences in \mathbb{C} , it is often sufficient to consider separately the corresponding sequences of real and imaginary parts in \mathbb{R} .

Definition 3.29: Sequences of Complex Numbers

A sequence of complex numbers $(z_n)_{n=0}^\infty$, where

$$z_n = x_n + iy_n,$$

is said to **converge** to a limit $A + iB \in \mathbb{C}$ if the two sequences of real numbers $(x_n)_{n=0}^\infty$ and $(y_n)_{n=0}^\infty$ converge to A and B , respectively. In this case, we write

$$\lim_{n \rightarrow \infty} z_n = A + iB.$$

We say that $(z_n)_{n=0}^\infty$ **diverges to ∞** if the sequence of moduli $(|z_n|)_{n=0}^\infty$ diverges to $+\infty$, i.e.,

$$\lim_{n \rightarrow \infty} |z_n| = \lim_{n \rightarrow \infty} \sqrt{x_n^2 + y_n^2} = +\infty.$$

Remark 3.30. *As for sequences of real numbers, one can consider subsequences of sequences \mathbb{C} . Given a strictly increasing sequence of non-negative integers $(n_k)_{k=0}^\infty$, the corresponding subsequence is*

$$(z_{n_k})_{k=0}^\infty = (x_{n_k} + iy_{n_k})_{n=0}^\infty.$$