

Étale cohomology

by Prof. Dr. Jens Franke

notes by Stefan Albrecht

University Bonn – winter term 2023/24

Contents

1	Motivation and basic definitions	2
1.1	Introduction and motivation	2
1.2	Flat morphisms	3

1 Motivation and basic definitions

1.1 Introduction and motivation

Problem: For varieties X over an algebraically closed field k (and hopefully more general schemes) define a cohomology theory $H^*(X)$ with properties similar to $H_{\text{sing}}^*(X(\mathbb{C})_{\text{ord. top. space}})$. Hopefully, there exists a Lefschitz fixed point formula

$$\#(\text{fixed points of } f \text{ with multiplicity}) = \sum_{i=0}^{2 \dim X} (-1)^i \text{Tr}(f^* | H^i(X)). \quad (\text{L})$$

The aim of Grothendieck was to apply this to a program proposed by Weil of studying the congruence zeta function of X by applying (L) to $f = F_X$ given by $[x_0, \dots, x_n] \mapsto [x_0^q, \dots, x_n^q]$, yielding

$$\#X(\mathbb{F}_q) = \sum_{i=0}^{2 \dim X} (-1)^i \text{Tr}(F_X^* | H^i(X)).$$

Counterexamples $H_{dR}^*(X) = \mathbb{H}^*(X_{\text{Zar}}, \mathcal{O}_X \rightarrow \Omega_X^1 \rightarrow \dots)$ (de Rham cohomology) is ok if the characteristic of k is zero but not in char p where it is unsuitable for Weil's program. Similarly, $H^*(X_{\text{Zar}}, \mathbb{Z})$ does not work: $\mathbb{Z}(X) \rightarrow \mathbb{Z}(V)$ is surjective when X is irreducible, implying vanishing higher sheaf cohomology.

Restrictions on the ring of coefficients: If X is a supersingular elliptic curve over $\overline{\mathbb{F}}_q$ then $H^1(X)$ ought to be two-dimensional, but $\text{End}(X) \otimes \mathbb{Q}$ is a quaternion algebra over \mathbb{Q} which is non-split precisely over \mathbb{Q}_p and \mathbb{R} , in which case it cannot act on a two-dimensional vector space. This excludes \mathbb{Q}_p and \mathbb{R} as the field of definition and hence also \mathbb{Q} and \mathbb{Z} .

Étale cohomology with coefficients $\mathbb{Z}/l^n\mathbb{Z}$, l a prime invertible in k . Then

$$H^*(X, \mathbb{Q}_l) := (\varprojlim H^*(X_{\text{ét}}, \mathbb{Z}/l^n\mathbb{Z})) \otimes_{\mathbb{Z}_l} \mathbb{Q}_l.$$

Deligné used this to show the Riemann hypothesis for congruence zeta function.

Other theories include Crystalline cohomology with coefficients in $W(\overline{\mathbb{F}}_q)$. Scholze has a way of working with \mathbb{Z}_p directly, using the pro-étale site, and a proposal to work with \mathbb{C} coefficients. But it is not clear how to do this.

Hence we will mostly study finite coefficients. If one works over \mathbb{C} , the exact exponential sequence $0 \rightarrow 2\pi i\mathbb{Z} \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_X^\times \rightarrow 0$ is important. and we want at least the exactness of

$$0 \rightarrow \mu_{l^n} \rightarrow \mathcal{O}_X^\times \xrightarrow{f \mapsto f^{l^n}} \mathcal{O}_X^\times \rightarrow 0. \quad (*)$$

Note that $\mu_{l^n} \cong \mathbb{Z}/l^n\mathbb{Z}$ non-canonically if $k = \bar{k}$ and l is invertible in k . Unfortunately, but not unexpectedly, this is not exact on X_{Zar} . If this were exact, one could hope to get some information from it provided that $H^1(C, \mathcal{O}_C^\times) \cong \mathbb{Z} \times \text{Jac}_C(k)$. The idea of Grothendieck was to enforce the exactness of $(*)$ by considering $V \rightarrow F(V)$ for étale morphisms $V \rightarrow X$ instead of only Zariski open subsets. Then, when $f \in \mathcal{O}_V^\times(V)$ one has an l^n -th root of f on $U = \{(x, \varphi) \mid x \in V, \varphi^{l^n} = f(x)\}$.

1.2 Flat morphisms

Definition 1. M is a *flat* A -module if $T \mapsto M \otimes_A T$ is exact or, equivalently, if $\mathrm{Tor}_p^A(M, T) = 0$ for all T and $p > 0$. An A -algebra B is flat if it is flat as an A -module.

Definition 2. For a morphism $f : X \rightarrow Y$ of schemes, f is called *flat* if it satisfies the following equivalent conditions:

- a) For all $x \in X$, $\mathcal{O}_{X,x}$ is a flat $\mathcal{O}_{Y,f(x)}$ -algebra.
- b) For affine open subsets $U \subseteq X, V \subseteq Y$ s.t. $f(U) \subseteq V$, $\mathcal{O}_X(U)$ is flat as an $\mathcal{O}_Y(V)$ -algebra.
- c) There are affine open subsets $U_i \subseteq X, V_i \subseteq Y$ s.t. $f(U_i) \subseteq V_i$, $\mathcal{O}_X(U_i)$ is a flat $\mathcal{O}_Y(V_i)$ -algebra and $X = \bigcup_{i \in I} U_i$.

Remark 1. a) See stacksproject 01U2

- b) Other literature: SGA1: Etale fundamental group, SGA4₁: Topoi, Grothendieck topology, SGA4₂: Etale topology, SGA4₃: Proper and smooth base change, SGA4₂¹: various stuff and Arcata – Introduction to étale cohomology by Deligne, SGA5: l -adic cohomology
Milne: Etale cohomology, Kiehl-Freitag: Etale cohomology and Weil conjectures
Matsumura: Commutative Algebra, Matsumura: Commutative Ring Theory

Let A be a ring, X quasi-compact and separated $\mathrm{Spec} A$ -scheme and \mathcal{M} a quasi-coherent \mathcal{O}_X -module. Then $H^*(X, \mathcal{M})$ can be calculated using $\check{H}(\mathcal{U}, -)$ for affine coverings. Hence, by the exactness of $- \otimes_A \tilde{A}$, this gives

Proposition 1. a) Let \tilde{A} be a flat A -algebra, then $H^*(\tilde{X}, \tilde{\mathcal{M}}) \cong H^*(X, \mathcal{M}) \otimes_A \tilde{A}$, where $\tilde{X} = X \times_{\mathrm{Spec} A} \mathrm{Spec} \tilde{A} \xrightarrow{p} X$ and $\tilde{\mathcal{M}} = p^* \mathcal{M}$.

- b) Let $f : X \rightarrow Y$ be a quasi-compact separated morphism and $g : \tilde{Y} \rightarrow Y$ a flat morphism, \mathcal{M} a quasi-coherent \mathcal{O}_X -module. Then $g^* R^* f_* \mathcal{M} \cong R^* \tilde{f}_* \tilde{g}^* \mathcal{M}$ where $\tilde{X} = X \times_Y \tilde{Y}$.

Remark 2. Base change results for étale cohomology are similar. We have b) if f is proper or if f is of finite type and g is smooth, and the sheaves are of torsion.

Definition 3. f is called *faithfully flat* if it is flat and surjective on points. \tilde{A} is a faithfully flat A -algebra if it is flat and $R \otimes_A \tilde{A} = 0$ implies $R = 0$.

Definition 4. ¹ Let $f : X \rightarrow Y$ be a morphism of schemes. A descent datum (of quasi-coherent sheaves of modules) for f is a quasi-coherent \mathcal{O}_X -module \mathcal{M} with an isomorphism $\mu : p_1^* \mathcal{M} \cong p_2^* \mathcal{M}$, where

$$\begin{array}{ccc} X \times_Y X \times_Y X & \xrightarrow[p_{23}]{p_{12}, p_{13}} & X \times_Y X \xrightarrow{p_1, p_2} X \\ & \searrow q_1, q_2, q_3 \nearrow & \\ & & \end{array}$$

¹ see tag 023A or SGA1, VI for fibred categories: descend data for X -schemes to Y -schemes and ample line bundles

are the different projections, and the diagram

$$\begin{array}{ccccc}
 q_1^* \mathcal{M} & \xlongequal{\quad} & p_{12}^* p_1^* \mathcal{M} & \xrightarrow[p_{12}^* \mu]{\cong} & p_{12}^* p_2^* \mathcal{M} & \xlongequal{\quad} & q_2^* \mathcal{M} \\
 & \searrow & \downarrow p_{13}^* \mu & & \downarrow p_{23}^* \mu & \swarrow & \\
 & p_{13}^* p_1^* \mathcal{M} & \xrightarrow[p_{13}^* \mu]{\cong} & p_{13}^* p_2^* \mathcal{M} & & p_{23}^* p_1^* \mathcal{M} & \xrightarrow[p_{23}^* \mu]{\cong} & p_{23}^* p_2^* \mathcal{M} \\
 & & & \searrow & \swarrow & & & \\
 & & & q_3^* \mathcal{M} & & & &
 \end{array}$$

must commute. A morphism of descent data is a morphism $\varphi : \mathcal{M} \rightarrow \widetilde{\mathcal{M}}$ compatible with μ and $\widetilde{\mu}$, i.e. $(p_2^* \varphi) \mu = \widetilde{\mu} (p_1^* \varphi)$

Remark 3. We have a functor

$$\mathrm{QCoh}(Y) \rightarrow \mathrm{Desc}_{\mathrm{QCoh}(X),f}, \quad \mathcal{N} \mapsto (f^* \mathcal{N}, \text{ the canonical iso } p_1^* f^* \mathcal{N} \cong p_2^* f^* \mathcal{N}).$$

One would like this to be an equivalence of categories. It has a right adjoint

$$(\mathcal{RM})(U) = \{m \in \mathcal{M}(f^{-1}U) \mid \mu p_1^* m = p_2^* m\}$$

Proposition 2 (stacks loc.cit., SGA1.VII.1, Milne). *If f is faithfully flat and quasi-compact, the above functor $\mathrm{QCoh}(Y) \rightarrow \mathrm{Desc}_{\mathrm{QCoh}(X),f}$ is an equivalence of categories.*

Proof. If f has a section, the inverse image along that section is an inverse functor. In general, base change with $f : X \rightarrow Y$ reduces to this situation, provided that f is separated, which is a situation one can reduce to. \square

Corollary 5. *If f is faithfully flat, $\mathcal{O}_Y(V) = \{\lambda \in \mathcal{O}_X(f^{-1}U) \mid p_1^* \lambda = p_2^* \lambda\}$.*

Remark 4. Both quasi-compactness and quasi-coherence in proposition 2 are needed. Consider $Y = \mathrm{Spec} R$, R a PID with $\mathrm{Spec} R$ infinite,

$$X = \coprod_{m \in \mathrm{mSpec}} \mathrm{Spec} R_m, \quad N_1 = \coprod_{m \in \mathrm{mSpec} R} R/m \rightarrow N_2 = \coprod_{m \in \mathrm{mSpec} R} R/m,$$

then it is easy to see that this inclusion does not split, but it splits canonically after applying $- \otimes_R R_m$, giving rise to a morphism of descent data which does not descend to a morphism $N_2 \rightarrow N_1$.

Definition 6. A morphism $i : X \rightarrow Y$ in a category \mathcal{A} is an effective monomorphism if for all objects T ,

$$\mathrm{Hom}_{\mathcal{A}}(T, X) \xrightarrow{\varphi \mapsto i\varphi} \{f \in \mathrm{Hom}_{\mathcal{A}}(T, Y) \mid \sigma f = \widetilde{\sigma} f \text{ for all } \sigma, \sigma' : Y \rightarrow S \text{ s.t. } \sigma i = \widetilde{\sigma} i\}$$

is bijective. $p : X \rightarrow Y$ is an effective epimorphism if it is an effective monomorphism in $\mathcal{A}^{\mathrm{op}}$, i.e.

$$\mathrm{Hom}_{\mathcal{A}}(Y, T) \xrightarrow[p \cong]{\varphi \mapsto \varphi p} \{f \in \mathrm{Hom}_{\mathcal{A}}(X, T) \mid f \sigma = f \widetilde{\sigma} \text{ for all } \sigma, \widetilde{\sigma} : S \rightarrow X \text{ s.t. } p \sigma = p \widetilde{\sigma}\}.$$

Remark 5. If $X \times_Y X$ exists, f being an effective epimorphism is equivalent to it being a coequalizer of $X \times_Y X \rightrightarrows X$.

Proposition 3 (SGA1.VIII.4 or stacks 023Q). *Every fpqc (quasi-compact faithfully flat) morphism of schemes is an effective epimorphism.*