

POST-RESUSCITATION CARE




IMMEDIATE TREATMENT

- Airway and breathing**
- Maintain SpO₂ 94 – 98%
 - Insert advanced airway
 - Waveform capnography
 - Ventilate lungs to normocapnia

- Circulation**
- 12-lead ECG
 - Obtain reliable intravenous access
 - Aim for SBP > 100 mmHg
 - Fluid (crystalloid) – restore normovolaemia
 - Intra-arterial blood pressure monitoring
 - Consider vasopressor/ inotrope to maintain SBP

- Control temperature**
- Constant temperature 32°C – 36°C
 - Sedation; control shivering

DIAGNOSIS



OPTIMISING RECOVERY

- ICU management**
- Temperature control: constant temperature 32°C – 36°C for ≥ 24h; prevent fever for at least 72h
 - Maintain normoxia and normocapnia; protective ventilation
 - Avoid hypotension
 - Echocardiography
 - Maintain normoglycaemia
 - Diagnose/treat seizures (EEG, sedation, anti-epileptic drugs)
 - Delay prognostication for at least 72h

Secondary prevention
e.g. ICD, screen for inherited disorders, risk factor management

Functional assessments
before hospital discharge

Structured follow up after
hospital discharge

Rehabilitation