Table of Distributions

stesiam

Διακριτές κατανομές

| Distribution Συμβολισμός | Parameters | f(x) | Value domain | F(x) | E(X) | Var(X) | $M_x(t)$ | $P_x(t)$ |
|--|---|--|--------------|--|-----------------|-----------------|--|--------------------|
| Bernoulli Bernoulli(p) | $p \in [0,1]$ | $\begin{cases} q, & \text{eáv k=0} \\ p, & \text{eáv k=1} \end{cases}$ | x = {0, 1} | $\begin{cases} 0, & \text{ean } k < 0 \\ q, & \text{ean } 0 < = k < 1 \\ 1, & \text{ean } k > = 1 \end{cases}$ | p | pq | $p(e^t + q)$ | (pt+q) |
| $\begin{array}{c} \textbf{Binomial} \\ B(n,p) \end{array}$ | $n \in \{0,1,\}$ $p \in [0,1]$ | $\binom{n}{k} p^x q^{n-x}$ | x = 0,1,,n | | np | npq | $p(e^t + q)^n$ | $(pt+q)^n$ |
| $\begin{array}{c} \textbf{Poisson} \\ Poisson(\lambda) \end{array}$ | $\lambda > 0$ | $e^{-\lambda} rac{\lambda^x}{x}$ | x = 0,1, | _ | λ | λ | $e^{\lambda(e^t-1)}$ | $e^{\lambda(t-1)}$ |
| $\begin{array}{c} \textbf{Geometric 0} \\ G_0(p) \end{array}$ | $p \in [0,1]$ | $q^x p$ | x = 0,1,2, | $1 - q^{x-1}$ | $\frac{q}{p}$ | $\frac{q}{p^2}$ | $\frac{p}{1-qe^t}$ | |
| $\begin{array}{c} \textbf{Geometric 1} \\ G_1(p) \end{array}$ | $p \in [0,1]$ | $q^{x-1}p$ | x = 1,2, | $1-q^x$ | $\frac{1}{p}$ | $\frac{q}{p^2}$ | $\begin{array}{c} \frac{pe^t}{1-qe^t},\\ t<-ln(q) \end{array}$ | |
| $\begin{array}{c} \textbf{Negative Binomial 0} \\ NB_0(r,p) \end{array}$ | | | | | | | | |
| $\begin{array}{c} \textbf{Negative Binomial 1} \\ NB_1(r,p) \end{array}$ | | | | | | | | |
| $\begin{array}{c} \textbf{Hypergeometric} \\ h(v,a,b) \end{array}$ | a,b,n: pos. integers $n <= a+b$ | $\frac{\binom{a}{x}\binom{b}{n-x}}{\binom{a+b}{n}}$ | x=0,1,2,n | | $\frac{a}{a+b}$ | | _ | _ |
| $\begin{array}{c} \textbf{Discrete Uniform} \\ U(a,b) \end{array}$ | a, b, n : pos. integers $n \le a + b$ | $\frac{\binom{a}{x}\binom{b}{n-x}}{\binom{a+b}{n}}$ | x=0,1,2,n | | $\frac{a+b}{2}$ | | _ | _ |

Συνεχείς κατανομές

| Distribution | Parameters | f(x) | Value domain | F(x) | E(X) | Var(X) | $M_x(t)$ |
|---|--|--|---------------------------|---|-----------------------------------|--|---|
| $\begin{array}{c} \textbf{Uniform} \\ U(a,b) \end{array}$ | $-\infty < a < b < +\infty$ | $\frac{1}{b-a}$ | $x \in [a, b]$ | $\begin{cases} 0, & x < a \\ \frac{x-a}{b-a}, & a \le x \le b \\ 1, & x > b \end{cases}$ | $\frac{a+b}{2}$ | $\frac{(b-a)^2}{12}$ | $\frac{e^{tb}-e^{ta}}{t(b-a)}$ |
| $\begin{array}{c} \textbf{Normal} \\ N(n,p) \end{array}$ | $n \in \{0,1,\}$ $p \in [0,1]$ | $\phi(x) = \frac{1}{\sigma\sqrt{2\pi}}e^{-\frac{(x-\mu)^2}{2\sigma^2}}$ | $x \in [-\infty, \infty]$ | | μ | σ^2 | $e^{\mu t + \frac{\sigma^2 t^2}{2}}$ |
| $\begin{array}{c} \textbf{Stndard Normal} \\ N(0,1) \end{array}$ | - | $\frac{1}{\sqrt{2\pi}}e^{-\frac{x^2}{2}}$ | x = 0,1, | | 0 | 1 | |
| Exponential $Exp(\lambda)$ | $\lambda > 0$ | $\lambda e^{-\lambda x}$ | $x \in [0, \infty]$ | $1 - e^{-\lambda x}$ | $\frac{1}{\lambda}$ | $\frac{1}{\lambda^2}$ | |
| $\begin{array}{c} \textbf{Gamma} \\ Gamma(k,\theta) \end{array}$ | $k > 0$ $\theta > 0$ | $\frac{x^{k-1}e^{-\frac{x}{\theta}}}{\theta^k\gamma(k)}$ | $x \in [0, \infty]$ | | $k\theta$ | $k\theta^2$ | $(1-\theta t)^{-k}, t<\tfrac{1}{\theta}$ |
| $\begin{array}{c} \textbf{Beta} \\ Beta(\alpha,\beta) \end{array}$ | $\begin{array}{c} \alpha > 0 \\ \beta > 0 \end{array}$ | $\frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(a,b)}$ | $x \in [0, 1]$ | | $rac{lpha}{lpha+eta}$ | $\frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$ | |
| Weibull $Wei(\lambda,k)$ | $\lambda \in (0, +\infty)$ $k \in (0, +\infty)$ | $\begin{cases} \frac{k}{\lambda} (\frac{x}{\lambda}) e^{-(\frac{x}{\lambda})^k}, & xheq0\\ 0, & x < 0 \end{cases}$ | $x \in [0, \infty]$ | $\begin{cases} 1 - e^{-\left(\frac{x}{\lambda}\right)}, & xheq0\\ 0, & x < 0 \end{cases}$ | $\lambda \Gamma(1 + \frac{1}{k})$ | | |
| Erlang $Erl(k, \lambda)$ εναλλ. $beta = \frac{1}{\lambda}$ | $k \in 1, 2, 3, \dots$ $\lambda \in (0, \infty)$ | $\frac{\lambda^k x^{k-1} e^{-\lambda x}}{(k-1)!}$ | $x \in [0, \infty]$ | | $rac{k}{\lambda}$ | $\frac{k}{\lambda^2}$ | $(1 - \frac{t}{\lambda})^{-k} \\ t < \lambda$ |
| $\begin{array}{c} \textbf{Pareto I} \\ Pareto(\alpha,\beta) \end{array}$ | a > 0 $b > 0$ | $\frac{\alpha\beta^{\alpha}}{(\beta+x)^{a+1}}$ | $x \in [0, +\infty]$ | | $\alpha > 1$ | $\frac{\alpha\beta^2}{(\alpha-1)^2(\alpha-2)}$ | - |
| Pareto II Pareto | a > 0 $b > 0$ | | $x \in [0, +\infty]$ | | | | - |
| $\begin{array}{c} \textbf{Lognormal} \\ log N(\mu, \sigma^2) \end{array}$ | $\mu \in (-\infty, +\infty)$ $\sigma > 0$ | $\frac{e^{-\frac{1}{2}(\frac{logx-\mu}{\sigma})}}{x\sigma\sqrt{2\pi}}$ | $x \in (0, +\infty)$ | Compl. expression | $e^{\mu + \frac{\sigma^2}{2}}$ | $(e^{\sigma^2}-1)(e^{2\mu+\sigma^2})$ | _ |