

The *Herschel*-ATLAS Data Release 2 - Paper 2: Catalogues of far-infrared and submillimetre sources in the fields at the south and north Galactic Poles

S.J. Maddox^{1,2*}, E. Valiante¹, P. Cigan¹, L. Dunne^{1,2},
M.W.L. Smith¹, S. Eales¹, S. Dye³, C. Furlanetto^{3,5}, E. Ibar⁴,
J. Millard¹, N. Bourne², H. Gomez¹ and R.J. Ivison⁶

¹*School of Physics and Astronomy, Cardiff University, The Parade, Cardiff CF24 3AA, UK*

²*Institute for Astronomy, The University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh, EH9 3HJ, UK*

³*School of Physics and Astronomy, University of Nottingham, University Park, Nottingham, NG7 2RD, UK*

⁴*ÁgInstituto de Física y Astronomía, Universidad de Valparaíso, Avda. Gran Bretaña 1111, Valparaíso, Chile.*

⁵*Departamento de Física, Universidade Federal do Rio Grande do Sul., Av. Bento Gonçalves, 9500, 91501-970, Porto Algres, RS Brazil*

⁶*European Southern Observatory, Karl-Schwarzschild-Strasse 2, 85748, Garching, Germany*

ABSTRACT

The *Herschel* Astrophysical Terahertz Large Area Survey (H-ATLAS) is a survey of 660 deg² with the PACS and SPIRE cameras in five photometric bands - 100, 160, 250, 350 and 500 μ m. This is the second of three papers describing the data release for the large fields at the south and north Galactic poles. In this paper we describe the catalogues of far-infrared and submillimetre sources for the SGP and NGP, which cover 285 deg² and 170 deg², respectively. The catalogues contain 118,986 sources for the NGP field and 183,784 sources for the SGP field detected at more than 4σ significance in any of the 250 μ m, 350 μ m or 500 μ m bands. We present photometry in all five bands for each source, including aperture photometry for sources known to be extended. The rms positional accuracy for the faintest sources is about 2.4 arc seconds in right ascension and declination. We describe a statistical analysis of the catalogues and discuss all the practical issues - completeness, reliability, flux boosting, accuracy of positions, accuracy of flux measurements - necessary to use the catalogues for astronomical projects.

Key words: methods: data analysis - catalogues - surveys - galaxies: statistics - cosmology: observations - submillimetre: galaxies

1 INTRODUCTION

This is the second of three papers describing the second major data release of the *Herschel* Astrophysical Terahertz Large Area Survey (the *Herschel* ATLAS or H-ATLAS), the largest single key project carried out in open time with the *Herschel* Space Observatory (Pilbratt et al. 2010). The H-ATLAS is a survey of approximately 660 deg² of sky in five photometric bands: 100, 160, 250, 350 and 500 μ m (Eales et al. 2010). Although the original goal of the survey was to study dust, and the newly formed stars hidden by dust, in galaxies in the nearby ($z < 0.4$) universe (Dunne et al. 2011, Eales et al. 2017), in practice the exceptional sensitivity of

Herschel, aided by the large negative k -correction at submillimetre wavelengths (Blain et al. 1993), has meant that the median redshift of the sources detected in the survey is ≈ 1 (Pearson et al. 2013), and our source catalogues include sources up to a redshift of at least 6.03 (Fudamoto et al. 2017; Zavala et al. 2017).

The five H-ATLAS fields were selected to be areas with relatively little emission from dust in the Milky Way, as judged from the IRAS 100 μ m images (Neugebauer 1984), and with a large amount of data in other wavebands. In 2010 for the Science Demonstration Phase (SDP) of *Herschel*, we released the data products for one 16 deg² field in the GAMA 9 hour field (Ibar et al. 2010; Pascale et al. 2011; Rigby et al. 2011; Smith et al. 2011). In our first large data release (DR1), we released the data products for three fields

* E-mail: maddoxs@cardiff.ac.uk

on the celestial equator centred at approximately 9, 12 and 15 hours (Valiante et al. 2016; Bourne et al. 2016), covering a total area of 161 deg^2 . These data products included the *Herschel* images in all five bands, a catalogue of the 120,230 sources detected on these images and of the 44,835 optical counterparts to these sources.

Our second data release is for the two larger fields at the north and south Galactic poles (NGP and SGP). The NGP field is centred approximately at a right ascension of $13^h 18^m$ and a declination of $+29^\circ 13'$ (J2000) and has an area of 180.1 deg^2 . The NGP field is a roughly square region (Fig. 1) and, among many other interesting known extragalactic objects, includes the Coma Cluster. The SGP field is centred approximately at a right ascension of $0^h 6^m$ and a declination of $-32^\circ 44'$ (J2000) and has an area of 317.6 deg^2 . The SGP field is elongated in right ascension (Fig. 2). Smith et al. (2017; S17) provide a comprehensive list of the multi-wavelength data that exists for these fields.

Our data release for these fields is described in three papers. In the first paper (Smith et al. 2017, S17), we describe the images of these fields, including a description of how these images can be used by the astronomical community for a variety of scientific projects. In this paper, we describe the production and properties of the catalogues of far-infrared and submillimetre sources detected on these images. A third paper (Furlanetto et al. 2017; F17) describes a search for the optical/near-infrared counterparts to the *Herschel* sources in the NGP field and the resulting multi-wavelength catalogue.

The arrangement of this paper as follows. Section 2 describes the detection of the sources. Section 3 describes the photometry of the sources. Section 4 describes the catalogues and their properties. Section 5 gives a summary of the paper.

The catalogues described in this paper can be obtained from the H-ATLAS website (h-atlas.org). Note that since we only search for sources where the images are made out of more than one individual *Herschel* observation, the area covered by the catalogues (170 deg^2 and 285 deg^2 for the NGP and SGP, respectively) is slightly smaller than the sizes of the images (180.1 deg^2 and 317.6 deg^2 , respectively - S17).

2 SOURCE DETECTION

2.1 The maps and background subtraction

A detailed description of the processing necessary to produce maps from the *Herschel* raw data is presented in S17. The resulting maps have pixel sizes 3, 4, 6, 8 and 12 arc seconds for 100, 160, 250, 350 and $500 \mu\text{m}$ respectively. The maps made with the PACS camera (100 and $160 \mu\text{m}$, Poglitsch et al. 2010) have units of Jy per pixel. The maps made with the SPIRE camera (250, 350 and $500 \mu\text{m}$, Griffin et al. 2010) have units of Jy per beam. The beam areas at 250, 350 and $500 \mu\text{m}$ are 469, 831 and 1804 square arc seconds, respectively. The noise on the images is a combination of instrumental noise and the confusion noise from sources that are too faint to be detected individually. S17 describes a detailed analysis of the noise properties of the images.

Before attempting to detect sources in the maps, we first subtracted a smoothly varying 'sky' level to remove the foreground emission from dust in our galaxy, so-called 'cirrus emission', and also the emission from




Figure 1. The coverage map for $250 \mu\text{m}$ observations of the NGP field. The map shows the number of samples from the bolometer timelines contributing to each map pixel, which ranges from 1 to 43, with the median value being 10. The range of the grayscale is from 0 samples (white) to 27 samples (black).

clustered extragalactic sources fainter than our detection limit. We used the `nebuliser` function, a programme produced by the Cambridge Astronomy Survey Unit to estimate and subtract the sky level on astronomical images (see <http://casu.ast.cam.ac.uk/surveys-projects/software-release/background-filtering>).

The choice of the filter scale used in `nebuliser` is quite critical, since it must be small enough for `nebuliser` to remove small-scale patches of cirrus emission but not so small that the flux from large galaxies is reduced. In practice, for the SPIRE maps we found that a median filter scale of 30 pixels (3 arc minutes in the $250 \mu\text{m}$ band) followed by a linear filter scale of 15 pixels was an acceptable combination.

We tested whether this filtering scale reduced the flux density of extended extragalactic sources by creating simulated maps, placing artificial extended sources on these maps, and then measuring the flux densities of these sources after the application of `nebuliser`. Since the nearby galaxies detected by *Herschel* are mostly spiral galaxies, we used exponential profiles for the artificial sources, convolving these with the SPIRE point-spread function. The surface-brightness limit on photographic plates is typically $\mu_B \approx 25 \text{ mag arcsec}^{-2}$, giving rise to the widely used D25 optical diameters for galaxies. Since these diameters are roughly equivalent to a distance of five scale lengths from the centre of a galaxy, we truncated the profiles of our artificial submillimetre sources at five scale lengths, which gave us diameters, roughly equivalent to D25 diameters in the optical waveband, that ranged from 24 to 192 arcsec. The simulations showed that significant flux is lost only for sources that have diameters larger than 3 arc minutes, and even for sources above this size, the flux loss is $\lesssim 10\%$.

We note that the application of `nebuliser` will change




Figure 2. The coverage map for $250\ \mu\text{m}$ observations of the SGP field. The map shows the number of samples from the bolometer timelines contributing to each map pixel, which ranges from 1 to 36, with the median value being 9. The range of the grayscale is from 0 samples (white) to 21 samples (black).

the clustering statistics of extragalactic sources. Apart from the foreground cirrus emission, **nebuliser** removes the background produced by the sources that are too faint to be detected individually. This background varies because of the clustering of these faint sources. A source catalogue made without any background subtraction will include more sources where the this background is high as a result of clusters of these faint sources, and so the clustering of the sources in this catalogue will be stronger than in a catalogue produced from an image in which this background emission has been removed. An investigation of the clustering in the H-ATLAS catalogues, which includes an analysis of the effect of the subtraction of this background, will be presented by Amvrosiadis et al. (in preparation).

For the PACS maps, the $1/f$ -noise from the instrument is much larger than for SPIRE, making the foreground cirrus emission and the background emission from faint galaxies difficult to detect. Since we could not clearly detect the foreground/background emission on smaller scales, we used a **nebuliser** scale of 5 arcminutes.

As the result of the application of **nebuliser**, the maps have a modal pixel value which is close to zero. For the SPIRE bands, the instrumental noise is low enough that the flux distribution of detected sources skews the pixel distribution to positive values so the mean is slightly positive (1.0, 1.0 and 0.6 mJy/beam at 250, 350 and 500 μm). The PACS detector is less sensitive and less stable than SPIRE, and so the instrumental noise dominates over the confusion noise and the pixel distribution is close to Gaussian; the mean of the **nebulised** PACS maps are very close to zero (0.016 and 0.016 MJy/sr, respectively, for the 100 and 160 μm maps).

2.2 Source Detection

In this section we describe the method used to find the sources on the images. Additional details are given in V16. Sources were detected using the MADX algorithm (Maddox et al in prep) applied to the SPIRE maps. MADX creates maps of the signal-to-noise ratio and identifies sources by finding peaks in the signal to noise. The detection and mea-

surement of fluxes is optimised by using a matched filter that is applied to both the signal map and the noise map.

The SPIRE instrumental noise maps are created from the number of detector passes and the estimated instrumental noise per pass, $\sigma_{\text{inst}}/\sqrt{N_{\text{sample}}}$, as described in S17 and V16.

Since the noise consists of both instrumental noise and confusion noise from the background of undetected sources, we follow the approach of Chapin et al (2011) to calculate the optimal matched filter in each of the three SPIRE bands. Details of estimation and form of the matched filter are discussed in V16. The resulting matched filters are slightly more compact than the corresponding PSFs, and have slightly negative regions outside the FWHM.

In the first step of the source detection, peak pixels which have values $> 2.5\sigma$ in the filtered 250 μm map are considered as potential sources. We use the 250- μm map since most sources have the highest signal-to-noise in this map. The source position is determined by fitting a Gaussian to the flux densities in the pixels surrounding the pixel containing the peak emission. As an initial estimate of the flux density of the source in each SPIRE band, MADX takes the flux density in the pixel closest to the 250- μm position.

The high source density on the SPIRE maps means that these flux estimates often contain contributions from neighbouring sources. To mitigate this effect, MADX uses the following procedure. In each band, MADX sorts the sources in order of decreasing flux density. The flux density of the brightest source is then more precisely estimated from a bicubic interpolation in the filtered map at the 250- μm position from the flux densities in the 3×3 pixels around the position. Using this flux estimate, a point source profile is then subtracted from the map at this position. The program then moves to the next brightest source and follows the same set of steps. One consequence of these steps is that some sources will now have 250- μm flux densities less than the original 2.5σ cut. Since most sources are brighter at 250 μm than at the two longer wavelengths, the estimates of the flux densities in the 350 μm and 500 μm bands are sometimes negative.




Figure 3. The distribution of instrumental and total noise for the 250 μm , 350 μm and 500 μm fluxes bands for the NGP and SGP fields. Green shows the instrumental noise and black the total noise for all pixels; red shows the instrumental noise and blue the total noise at the positions of all sources. The multiple peaks are the results of our tiling strategy. The main peak corresponds to the large fraction of the survey area that was covered by two individual *Herschel* observations (S17). The smaller peaks correspond to the smaller fraction of the survey area that was either covered by more than two observations or, in the case of one end of the SGP (S17), a single observation (the small peak at the right in the bottom panels).

3 PHOTOMETRY

3.1 For point sources

3.1.1 SPIRE

V16 carried out extensive simulations to determine the errors on the flux density estimates for point sources. They followed the simple procedure of injecting artificial sources of known flux density on to the real maps and then using MADX to estimate their flux densities (Section 2.2). They found that at 250 μm , the detection wavelength, the confusion noise varies as a function of source flux and gave a simple formula to approximate this:

$$\sigma_{\text{con}250}^2 = \min(0.0049, f_{250}/5.6)^2 + 0.00253^2 \quad (1)$$

They found that at 350 μm and 500 μm band the confusion noise is roughly constant, with $\sigma_{\text{con}350} = 0.00659$ and $\sigma_{\text{con}500} = 0.00662$.

We calculated the errors in the flux density estimates for each individual source, using these formulae to estimate the confusion noise and the maps of the instrumental noise (Section 2.2) to estimate the instrumental noise, then adding the confusion and instrumental noise in quadrature.

Our strategy of creating the H-ATLAS survey from overlapping tiles (S17) means that the instrumental noise varies over the maps. Figure 3 shows histograms of instrumental noise and total noise (instrumental noise plus confusion noise) for all pixels and at the positions of all sources. The multiple peaks are the results of our tiling strategy. The main peak corresponds to the large fraction of the survey area that was covered by two individual *Herschel* observations (S17). The smaller peaks correspond to the smaller fraction of the survey area that was either covered by more

than two observations or, in the case of one end of the SGP (S17), a single observation.

The variation of noise across the maps means that the 4σ flux-density limit varies over the fields, and hence the available area depends on the chosen flux limit. Figure 4 shows the relationship between area and flux-density limit for each of the H-ATLAS fields, including the GAMA fields.

3.1.2 PACS

As in V16, we used aperture photometry to estimate the flux densities in the two PACS bands. We did this for two reasons. First, the PACS PSF for our observing mode (fast-parallel scan mode) is not well determined near its peak (see V16 for an extensive discussion). Second, if we estimated the 100 and 160- μm flux densities at the 250- μm position, as we did for the 350 and 500- μm bands, we would be likely to significantly underestimate the flux density because of the higher resolution of the PACS maps.

V16 describe an extensive investigation of the optimum aperture size, and we follow them in using an aperture with a radius equal to the FWHM, which is 11.4 arc seconds for 100 μm and 13.7 arc seconds for 160 μm . Since the ‘sky’ level has already been subtracted with *nebuliser*, we made no correction for any residual sky level. To reduce any pixelisation effects, we divided each pixel into 16, assigning one sixteenth of the flux density in each sub-pixel, and added up the flux density in each sub-pixel within the aperture. Since only $\approx 10\%$ of the SPIRE sources were clearly detected on the PACS images, we centered the aperture on the 250- μm position.

We corrected the aperture flux densities to total flux densities using the table of the Encircled Energy Fraction (EEF) described in V16 and available at <http://www.h->




Figure 4. The relationship between area and 4σ flux-density limit for the H-ATLAS fields: NGP - black; SGP - blue; GAMA9 - magenta; GAMA12 - green; GAMA15 - cyan. The more sensitive areas correspond to the tile overlaps in each field. The westerly end of SGP has only a single SPIRE observation, which explains the kink at high flux densities in the blue line in the panels.

atlas.org/. We made a further correction to allow for the effect of the errors on the $250\text{-}\mu\text{m}$ positions, since any error in the position will lead to the small PACS apertures missing flux. V16 describe simulations of this effect, and we follow them in compensating for this effect by multiplying the flux densities by 1.1 and 1.05 at 100 and $160\text{ }\mu\text{m}$, respectively.

We describe how we estimated the errors on these flux estimates in the following section.

3.2 Extended sources

The approach in the previous section gives optimal flux density estimates for point sources, but will seriously underestimate the flux density of extended sources. As in V16, we used the optical sizes of optical counterparts to the *Herschel* sources to indicate which sources are likely to require aperture photometry rather than the methods described in the last section. We followed different methods for the NGP and the SGP because of the lack of a comprehensive identification analysis for the SGP.

3.2.1 The NGP

In the NGP, F17 carried out a search for optical counterparts to the *Herschel* sources on the r-band images of the Sloan Digital Sky Survey (SDSS) which was almost exactly the same as that carried out by Bourne et al. (2016) for the H-ATLAS GAMA fields. Our initial list of NGP sources that might require aperture photometry were the sources with optical identifications with reliability $R > 0.8$ from F17.

In our previous data release (V16) we calculated the sizes of our apertures from the SDSS parameter `isoA_r`, which was available in SDSS DR7. However, this parameter was not available in SDSS DR10, on which F17 based their analysis. After an investigation of the various size measurements available in DR10, we found that the parameter `petroR90_r`, the 90% Petrosian radius, met our needs since there is a simple scaling between it and `isoA_r`, with $\text{isoA}_r \approx 1.156 \text{ petroR90}_r$.

We considered that for H-ATLAS sources with optical counterparts with `petroR90_r` less than 8.6 arcsec (equivalent to the value of `isoA_r` of 10 arcsec used in V16) the source is still unlikely to be extended in the SPIRE bands, and for these H-ATLAS sources we preferred the flux densities in the SPIRE bands produced by MADX (Section 3.1.1). However, if the H-ATLAS source had an optical counterpart with `petroR90_r` greater than 8.6 arcsec, and for all PACS photometry, we measured aperture photometry. We calculated the radius of the aperture using the same formula as V16 (with `isoA_r` replaced by `petroR90_r`):

$$r_{\text{ap}} = \sqrt{\text{FWHM}^2 + (1.156 \text{ petroR90}_r)^2}, \quad (2)$$

where `FWHM` is the Full-Width Half-Maximum of the point-spread function for the passband being measured. As discussed above (Section 3.1.2), we also use aperture photometry in the PACS bands for sources without reliable optical counterparts, using an aperture with a radius equal to the FWHM.

After calculating the aperture using equation 2, we visually compared it with the $250\text{-}\mu\text{m}$ emission from the source,

since in some case the aperture is not well-matched to the 250- μm emission, either being too small, too large, with the wrong shape or including the flux from a neighbouring galaxy (see V16 for examples). In these cases, we chose a more appropriate aperture for the galaxy. We also visually inspected the 3000 sources with the brightest 250- μm flux densities from MADX in order to check whether there were any obvious additional extended sources. For these sources too, we chose appropriate apertures to include all of the emission. In total, for the NGP there are 74 of these ‘customised apertures’. The details of these customised apertures are given as part of the data release.

We calculated errors in the aperture flux densities from the results of the Monte-Carlo simulation of S17. S17 placed apertures randomly on the SGP and NGP maps in areas which are made from two individual observations ($N_{\text{scan}} = 2$), varying the aperture radii from approximately the beam size up to $100''$ in $2''$ intervals and using 3000 random positions for each aperture radius. They found that the error, σ_{ap} , depends on the radius the aperture as a double power-law:

$$\sigma_{\text{ap}}(\text{mJy}) = \begin{cases} Ar^\alpha & \text{if } r \leq 50'' \\ B(r - 50)^\beta + A50^\alpha & \text{for } r > 50'' \end{cases} \quad (3)$$

The constants A , B , α , and β are given in Table 3 of S17. We used this equation for the sources on parts of the images made from two observations. In parts of the images made from more than two observations the instrumental noise is less; for sources in these more sensitive parts of the images we used the extension of equation 3 derived by S17: equation 4 in S17 for SPIRE and equation 6 in S17 for PACS.

Finally, we only used the aperture flux density if it is significantly larger than the point-source estimate, if

$$F_{\text{ap}} - F_{\text{ps}} > \sqrt{\sigma_{\text{ap}}^2 - \sigma_{\text{ps}}^2}. \quad (4)$$

In summary, of the 118,986 sources in the NGP, we measured aperture flux densities at 250 μm for 889 sources.

3.2.2 The SGP

For the SGP area no SDSS data exists and we have not carried out the comprehensive identification analysis that we have carried out for the other four fields. Instead, we have carried out a rudimentary identification analysis using the 2MASS survey. We first found a 2MASS galaxy parameter that provides a useful estimate of the size of the galaxy. We found that the 2MASS parameter `sup_r_3sig` (super-coadd 3-sigma isophotal semi-major axis) has a simple scaling with the `isoA_r`: `isoA_r` ≈ 1.96 `sup_r_3sig`. We found all 2MASS galaxies in the SGP region with `sup_r_3sig` > 5.1 arcsec, equivalent to `isoA_r` = 10 arcsec. There are 6249 of these galaxies. We then found all H-ATLAS sources in the SGP within 5 arcsec of a 2MASS galaxy. There are 3444 of these sources, and a simple probability analysis shows that only 23 (0.7%) of these matches should not be physical associations of the H-ATLAS source and the 2MASS galaxy.

For these sources, we calculated the radius of the aperture to use for photometry using the relationship:

$$r_{\text{ap}} = \sqrt{\text{FWHM}^2 + (1.96 \text{ sup_r_3sig})^2}. \quad (5)$$

This is the same as equation 2, except for the change in the parameter used to estimate the size of the galaxy.

As for the NGP, we then visually compared the apertures with the 250- μm emission from the source, modifying the aperture when necessary (see above). We also visually inspected the 5000 sources with the brightest 250- μm flux densities from MADX in order to check whether there were any obvious additional extended sources. For these sources too, we chose appropriate apertures to include all of the emission. In total, for the SGP there are XXX customised apertures, for which the details are given as part of the data release.

In the case of the SGP, we centered the apertures on the 250- μm positions rather than on the optical positions. Since the ‘sky’ level on both the PACS and SPIRE images has already been subtracted with `nebuliser`, we made no correction for any residual sky level. To reduce any pixelisation effects, we divided each pixel into 16, assigning one sixteenth of the flux density in each sub-pixel, and added up the flux density in each sub-pixel within the aperture. We corrected the PACS flux densities to total flux densities using the table EEF described in V16 and available at <http://www.h-atlas.org/>. We corrected all the SPIRE aperture flux densities for the fraction of the PSF outside the aperture using a table of corrections determined from the best estimate of the SPIRE PSF (Griffin et al. 2013), which is provided as part of the data release (see V16 for more details).

We followed the same procedure for estimating the errors in the flux densities and for deciding whether to use the aperture flux density rather than the flux density from MADX as for the NGP. In summary, of the XXX sources in the SGP, we measured aperture flux densities at 250 μm for XXX sources.

3.3 Colour corrections and flux calibration

The width of the SPIRE filters means that both the size of the PSF and the power detected by SPIRE depend on the spectral energy distribution of the source. The SPIRE data-reduction pipeline and ultimately our flux densities are based on the assumption that the flux density of a source varies with frequency as ν^{-1} . If the user knows the SED of a source, the flux densities should be corrected using corrections from either Table 5.7 or 5.8 from the SPIRE handbook¹. It is important to apply these corrections, since they can be quite large: for a point sources with a typical dust spectrum ($T = 20\text{K}$, $\beta = 2$) the multiplicative correction is 0.96, 0.94 and 0.90 at 250, 350 and 500 μm , respectively.

As with SPIRE, the PACS flux densities are also based on the assumption that flux density of the source is proportional to ν^{-1} , and a correction is required for sources which follow a different SED. The required corrections are described in the PACS Colour-Correction document².

On top of all other errors, there is an additional error due to the uncertain photometric calibration of *Herschel*. As in V16, we assume conservative calibration errors of 5.5% for

¹ http://herschel.esac.esa.int/Docs/SPIRE/spire_handbook.pdf

² http://herschel.esac.esa.int/twiki/pub/Public/PacsCalibrationWeb/cc_report_v1.pdf

Table 1. Stars in H-ATLAS

Name	Position
EY Hya	08:46:21.4 +01:37:53
IN Hya	09:20:36.7 +00:10:53
NU Com	13:10:08.5 +24:36:02
19 PsA	22:42:22.3 -29:21:43
V PsA	22:55:19.9 -29:36:48
S Sci	00:15:22.4 -32:02:44
XY Sci	00:06:35.9 -32:35:38
eta Sci	00:27:55.9 -33:00:27
Y Sci	23:09:05.7 -30:08:04
HD 119617	13:43:35.2 +35:20:45
R Sci	01:26:58.2 -32:32:37
Fomalhaut	22:57:39.2 -29:37:22

the three SPIRE wavebands and 7% for PACS (see V16 for more details).

4 THE CATALOGUES

We included all sources in the catalogues if they were detected at at least 4σ in one or more of the three SPIRE bands: 250, 350 and 500 μm . We eliminated all sources from the original list of point sources produced by MADX if they fell within the aperture of an extended source. All of the H-ATLAS fields were observed at least twice, making it possible to search for moving sources such as asteroids. We found 9 asteroids in the GAMA fields (V16), eliminating these from the final catalogue. We carried out the same search for the NGP and SGP but found no moving objects.

The sources in the final catalogues are almost all extragalactic sources. We carried out a search for clusters of sources in all the H-ATLAS fields (Eales et al. in preparation). In the GAMA9 field, we found several clusters of sources that are likely to be clusters of prestellar cores, implying that the catalogue for this field is likely to contain a few tens of Galactic sources. However, we found no similar clusters in the other fields and the GAMA9 field is at a much lower Galactic latitude than the other fields. Prestellar cores are therefore likely to be a very minor contaminant to the catalogues for these fields. There are a few debris disks and AGB stars in the catalogues, and an incomplete list is given in Table 1. However, well over 99% of the sources are extragalactic. The extragalactic sources range from nearby galaxies, such as the spectacular galaxy, NGC 7793, in the centre of the SGP, one of the brightest galaxies in the nearby Sculptor group, to galaxies up to a redshift of ≈ 6 (Fudamoto et al. 2017).

4.1 The statistics of the catalogues

The catalogue for the NGP contains 118,986 sources, of which 112,074 were detected at $> 4\sigma$ at 250 μm , 46,872 at $> 4\sigma$ at 350 μm and 10,369 at $> 4\sigma$ at 500 μm . The effective sensitivity of the PACS images was much less, but the catalogues contain flux-density measurements at 100 μm and 160 μm for all the sources in the catalogue. 4,207 sources were detected at $> 3\sigma$ at 100 μm and 6,342 sources were detected at $> 3\sigma$ at 160 μm .




Figure 5. The cumulative number of sources as a function of signal-to-noise at 100 μm (black), 160 μm (blue), 250 μm (cyan), 350 μm (green) and 500 μm (red). The NGP area is shown in the top panel, and the SGP in bottom panel. The vertical dotted line shows the 4σ limit for the 250 μm selection. The other bands are truncated at 3σ .

The catalogue for the SGP contains 372596 sources, of which 354022 were detected at $> 4\sigma$ at 250 μm , 167904 at $> 4\sigma$ at 350 μm and 33336 at $> 4\sigma$ at 500 μm . 5978 sources were detected at $> 3\sigma$ at 100 μm and 11009 sources were detected at $> 3\sigma$ at 160 μm .

The cumulative number of sources as a function of signal-to-noise in the 5 bands is shown in Figure 5. The 250 μm band is the most sensitive, and so has the largest number detected sources. Of the PACS bands, the 160 μm band detects most sources above 3σ .

The observed number of sources as function of flux in the PACS and SPIRE bands is shown in Figure 6. Note that this is the observed flux in the catalogue, before any corrections are made for source SED (Section 3.3) or ‘flux boosting’ (Section 4.3), which are necessary before the flux densities are compared with model predictions.

4.2 Positional Accuracy

V16 carried out extensive simulations to investigate the accuracy of the H-ATLAS catalogues by injecting artificial




Figure 6. The cumulative number of sources as a function of flux at 100 μm (black), 160 μm (blue), 250 μm (cyan), 350 μm (green) and 500 μm (red). The NGP area is shown in the top panel, and the SGP in the bottom panel. The counts are plotted only above the limit of 3σ in each wave band.

sources on to the GAMA images, and then using MADX to detect the sources and measure their flux densities and positions. The results of these ‘in-out’ simulations apply to the NGP and SGP catalogues, which were produced using almost exactly the same methods.

We investigated the accuracy of the source positions in two ways: (1) by looking at the positional offsets between the *Herschel* sources and galaxies found on optical images; (2) from the in-out simulations. Bourne et al. (2016) and F17 describe the details of the first method, which takes account of the clustering of the galaxies in the optical catalogue and the PSF of the *Herschel* observations. We also use the positions of the galaxies to correct the astrometry of the individual *Herschel* observations, which ties the positions of the *Herschel* sources to the astrometric frame of the optical catalogue (S17).

In the case of the NGP, we applied this method using the galaxies found on the SDSS r-band images (F17), which thus ultimately ties the *Herschel* positions to the SDSS astrometric frame. In the case of the SGP, we used the galaxies found in the VLT Survey Telescope ATLAS (Shanks et al. 2015), which thus ultimately ties the astronomy in the

SGP to the astrometric frame of this survey. We find that the positional error, σ_{pos} , varies from 1.2'' to 2.4'' as the signal-to-noise in flux varies from 10 to 5, with a relationship between positional accuracy and flux density given by $\sigma_{\text{pos}} = 2.4(\text{SNR}/5)^{-0.84}$. This agrees well with the errors in the measured positions of the artificial sources in the in-out simulations (V16).

The mean positional errors as a function of position within the NGP and SGP fields are shown in Fig. 7. Though there are hints of systematic variations in different part of the fields, these are only ≈ 1 arcsec, less than the quoted pointing accuracy of *Herschel* of ≈ 2 arcsec (Pilbratt et al. 2010).

4.3 Purity, flux boosting and completeness

Based on Gaussian statistics, we expect $\approx 0.2\%$ of the sources in the catalogue to be spurious. However, V16 argue that this is likely to be a significant overestimate because our errors, while being good estimates of the errors on the flux measurements, are likely to underestimate the signal-to-noise of a detection.

A major problem in submillimetre surveys, where source confusion is usually an issue, is flux bias or ‘flux boosting’, in which the measured flux densities are systematically too high. V16 used the in-out simulations to quantify this effect in the H-ATLAS. Table 6 in V16 gives estimates of the flux bias as a function of flux density for all three SPIRE bands. The table shows that at the 4σ detection flux density, the measured flux densities are on average higher than the true flux densities by $\approx 20\%$, 5% and 4% at 250 μm , 350 μm and 500 μm , respectively. Astronomers interested in comparing the flux densities in the catalogue with the predictions of models should be aware of this effect. Table 6 in V16 can be used to correct the flux densities for this effect.

V16 also used the in-out simulations to estimate the completeness of the survey as a function of measured flux density in all three SPIRE bands. This is shown in Figure 21 of V16 and listed in Table 7 of V16. The completeness at 250 μm is $\approx 87\%$ at the 4σ detection limit of the survey.

5 SUMMARY

We have described the construction of the source catalogues from the *Herschel* survey of fields around the north and south Galactic poles. This survey which was carried out in five photometric bands - 100, 160, 250, 350 and 500 μm - was part of the *Herschel* Astrophysical Terahertz Large Area Survey (H-ATLAS), a survey of 660 deg^2 of the extragalactic sky. Our source catalogues cover 285 deg^2 around the SGP and 170 deg^2 around the NGP.

The catalogues contain 118,986 sources for the NGP field and 183,784 sources for the SGP field detected at more than 4σ significance in any of the 250 μm , 350 μm or 500 μm bands. We present photometry in all five bands for each source, including aperture photometry for sources known to be extended. We discuss all the practical issues - completeness, reliability, flux boosting, accuracy of positions, accuracy of flux measurements - necessary to use the catalogues for astronomical projects.




Figure 7. The mean positional errors in R.A. and Dec. averaged in areas $0.5 \times 0.5 \text{ deg}^2$ as a function of position on the sky for the NGP field and the SGP fields.

ACKNOWLEDGMENTS

PC, LD and SM acknowledge support from the European Research Council (ERC) in the form of Consolidator Grant COSMICDUST (ERC-2014-CoG-647939, PI H.L. Gomez). SJM LD and RJI acknowledge support from the ERC in the form of the Advanced Investigator Program, COSMICISM (ERC-2012-ADG 20120216, PI R.J. Ivison). EV and SAE acknowledge funding from the UK Science and Technology Facilities Council consolidated grant ST/K000926/1. MS and SAE have received funding from the European Union Seventh Framework Programme ([FP7/2007-2013] [FP7/2007-2011]) under grant agreement No. 607254.

Neugebauer, G. et al. 1984, *ApJ*, 278, L1
 Pascale, E., et al. 2011, *MNRAS*, 415, 911
 Pearson, E.A. et al. 2013, *MNRAS*, 435, 2753
 Pilbratt, G. L., et al. *A&A*, 518, L1
 Poglitsch, A., et al. 2010 *A&A*, 518, L2
 Rigby, E.E, et al. 2011, *MNRAS*, 415, 2336
 Shanks, T. et al. 2015, *MNRAS*, 451, 4238
 Smith, D.J.B. et al. 2011, *MNRAS*, 416, 857
 Smith, M.W.L. et al. 2017, *ApJ* suppl., in press (S17; Paper I)
 Valiante, E. et al. 2016, *MNRAS*, 462, 3146 (V16)
 Zavala, J.A. et al. 2017, *Nature Astronomy*, in press (arXiv: 1707.09022)

REFERENCES

Blain, A.W. and Longair, M.S. 1993, *MNRAS*, 264, 509
 Bourne, N. et al. 2016, *MNRAS*, 462, 1714
 Chapin, E.L. et al. 2011, *MNRAS*, 411, 505
 Dunne, L. et al. 2011, *MNRAS*, 417, 1510
 Eales, S., et al. 2010, *PASP*, 122, 499
 Eales, S., et al. 2017, *MNRAS*, in press (arXiv: 1710.01314)
 Fudamoto, Y. et al. 2017, *MNRAS*, 472, 2028
 Furlanetto, C. et al. 2017, *MNRAS*, submitted (F17; Paper III)
 Griffin, M.J. et al. 2010 *A&A*, 518, L3
 Griffin, M.J. et al. 2013, *MNRAS*, 434, 992
 Ibar, E., 2010, *MNRAS*, 409, 38