

THE ISOPERIMETRIC PROBLEM

Dissertation submitted at the University of Leicester
in partial fulfilment of the requirements for
the degree of Bachelor of Science of Mathematics

by

Steven Cheung
Department of Mathematics
University of Leicester

May 2024

Contents

Declaration	1
Abstract	2
Introduction	3
Historical Notes	3
Important Preliminaries	3
1 The Isoperimetric Theorems for 2D, 3D and nD Cases	5
1.1 2 Dimensional Case (Plane)	5
1.1.1 Unpacking Steiner's proof	5
1.2 3 Dimensional Case (Sphere)	5
1.3 n Dimensional Case (\mathbb{R}^n)	5
2 Manifolds	6

Declaration

All sentences or passages quoted in this project dissertation from other people's work have been specifically acknowledged by clear cross referencing to author, work and page(s). I understand that failure to do this amounts to plagiarism and will be considered grounds for failure in this module and the degree examination as a whole.

Name: Steven Cheung

Signed:

Date:

Abstract

In general, we want the maximum area whose boundary has a specific length.

For the 2-dimensional case.

For the 3-dimensional.

For the n -dimensional.

Manifolds?

Introduction

short intro to the problem

Historical Notes

Isoperimeter, isos which is ancient Greek for equal and perimetron for perimeter. The isoperimetric perimetric problem, even though not properly formalized, was already thought about. I'm talking about the legend of Dido. To begin, I'd like to quote Virgil

At last they landed, where from far your eyes May view the turrets of new
Carthage rise; There bought a space of ground, which Byrsa call'd, From the
bull's hide they first inclos'd and wall'd.

Virgil's version has it that Dido, daughter of the king of Tyre, fled home after her brother killed her husband. Dido ended up on the north coast of Africa, where she bargained to buy as much land as she could enclose with an oxhide. Thus, she cut the hide into thin strips, presumably met and solved enclosing the largest area with a given perimeter - the isoperimetric problem. Dido may have been clever enough and chose an area by the coast to exploit the shore as part of the perimeter. But this mostly spoils the purity of posed problem.

Since ancient Greece, people has long considered that of all the planes closed curve with equal perimeters, which bounds the greatest area? As they wanted to measure the size of islands by timing how long it takes to circumnavigate the entire island.

The Greek have pretty much solved it, by their standards. Zendorus was an ancient Greek mathematician from around 200B.C. to 120B.C.. Zendorus had proved that a circle has greater area than any polygon with the same perimeter. Most of his work was lost. Fortunately, parts of his work survived through references by Pappus and Theon of Alexandria.

Important Preliminaries

(to do: find a good book to give the proof of the Jordan Curve theorem or from papers) We will take for granted the Jordan Curve Theorem

Theorem 1 (Jordan Curve Theorem). *A simple closed curve in the plane divides the plane into two regions, one compact and one non-compact, and in the common boundary of both regions*

Note. *When we talk of the region bounded by a simple closed curve in the plane, we always mean the compact region*

— (to do: change the 3 def below to more rigorous ones, ideally from books)

Definition 1. *A closed curve, is a curve that changes direction but does not cross itself whilst changing direction.*

Definition 2. *A simple curve, is a curve that changes direction but does not cross itself whilst changing direction.*

The two definitions, above, are vital into understanding the main theorem.

Definition 3. *A function is bounded if $\exists M \in \mathbb{R}$ such that $|f(x)| \leq M$*

Definition 4. *A set is said to be compact if it is closed and bounded.*

(to do: Change above definition to the covers one, from topology. Add the Heine-Borel Theorem below)

The Isoperimetric Theorems for 2D, 3D and n D Cases

1.1 2 Dimensional Case (Plane)

Theorem 2. *Let C be a simple closed curve in the plane with length L and bounding a region of area A . Then $L^2 \leq 4\pi A$ with equality if and only if C is a circle.*

The circle therefore bounds the biggest area among all simple closed curves in the plane with a given length.

1.1.1 Unpacking Steiner's proof

The proof that I will be unpacking and taking a closer look at will be from the book (reference the book here), and is credited by them to Jakob Steiner.

Proof. We will begin the proof by assuming the solution exists

□

1.2 3 Dimensional Case (Sphere)

1.3 n Dimensional Case (\mathbb{R}^n)

Manifolds