

1 For  $n \geq 1$  let  $f_n: [1, 2] \rightarrow \mathbb{R}$  be defined as follows: for any  $x \in [1, 2]$ ,

$$f_1(x) = 0, \quad f_{n+1}(x) = \sqrt{x + f_n(x)}, \quad \forall n \geq 1.$$

Prove that  $\{f_n\}_{n \geq 1}$  converges uniformly to  $f(x) = \frac{1 + \sqrt{1 + 4x}}{2}$ .

**Solution** Fix  $x \in [1, 2]$ . Notice that 2 is an upper bound for  $f_n(x)$ . We will show this by induction.

Base step:

$$2 - f_1(x) = 2 > 0 \implies 2 \geq f_1(x)$$

Inductive step:

Suppose we have  $2 \geq f_n(x)$ . We wish to show that  $2 \geq f_{n+1}(x)$ .

$$2 - f_{n+1}(x) = 2 - \sqrt{x + f_n(x)} = \frac{4 - x - f_n(x)}{2 + \sqrt{x + f_n(x)}} \geq \frac{2 - f_n(x)}{2 + \sqrt{x + f_n(x)}} \geq 0 \implies 2 \geq f_{n+1}(x)$$

Hence the inductive step holds.

Next, we will show that  $f_n(x)$  is monotonically increasing.

$$\begin{aligned} f_{n+1}(x) - f_n(x) &= \sqrt{x + f_n(x)} - f_n(x) \\ &= \frac{x + f_n(x) - f_n^2(x)}{\sqrt{x + f_n(x)} + f_n(x)} \\ &\geq \frac{2 + f_n(x) - f_n^2(x)}{\sqrt{x + f_n(x)} + f_n(x)} \\ &= \frac{(2 - f_n(x))(1 + f_n(x))}{\sqrt{x + f_n(x)} + f_n(x)} \geq 0 \end{aligned}$$

Since  $f_n(x)$  is monotonically increasing and bounded above by 2, it converges pointwise to  $f(x)$ . Moreover, since  $f_n$  is continuous for all  $n \geq 1$  (each  $f_n$  where  $n \geq 2$  a composition of continuous functions)  $[1, 2]$  is compact, by Dini's theorem,  $f_n$  converges uniformly to  $f$  as  $n \rightarrow \infty$ .

As  $\lim_{n \rightarrow \infty} f_n(x)$  exists for all  $x \in [1, 2]$ ,  $\sqrt{x + f_n(x)}$  converges to  $\sqrt{x + f(x)}$  since it is a composition of continuous functions. Thus,

$$f_{n+1}(x) = \sqrt{x + f_n(x)} \implies f(x) = \sqrt{x + f(x)} \implies f(x) = \frac{1 + \sqrt{1 + 4x}}{2}$$

2 For  $n \geq 1$  let

$$f_n: [0, \infty) \rightarrow \mathbb{R}, \quad f_n(x) = \frac{nx^2 + 1}{nx + 1}.$$

Study the pointwise and uniform convergence of  $f_n$  on each of the intervals  $[0, \infty)$ ,  $(0, \infty)$ ,  $[1, \infty)$ .

**Solution**  $[0, \infty)$ :

When  $x = 0$ , we have  $f_n(0) = 1$ , so  $f_n(0) \xrightarrow{n \rightarrow \infty} f(0) = 1$ .

For  $x > 0$ , we have

$$f_n(x) = \frac{x^2 + \frac{1}{n}}{x + \frac{1}{n}} \xrightarrow{n \rightarrow \infty} f(x) = \frac{x^2}{x} = x$$

Thus,  $f(x)$  is not continuous, so  $f_n(x)$  cannot converge uniformly to  $f(x)$  on this interval.

$(0, 1)$ :

From the above calculation, we have that  $f_n(x) \xrightarrow{n \rightarrow \infty} f(x) = x$ . Moreover,

$$d(f_n, f) = \sup_{x \in [0, \infty)} |f_n(x) - f(x)| = \sup_{x \in [0, \infty)} \left| \frac{1-x}{nx+1} \right| \geq 1$$

Thus,  $f$  converges pointwise on  $x \in (0, \infty)$ , but it does not converge uniformly.

$[1, \infty)$ :

The calculation from the first case still holds, so we have that  $f(x) = x$ . Note that

$$(f_n(x) - x)' = -\frac{n+1}{(nx+1)^2} \leq 0$$

so  $f_n(x) - x$  is decreasing for all  $x$ . So, since  $f_n(x) - x \leq 0$ ,  $|f_n(x) - x|$  is increasing for all  $x \geq 1$ . Thus,

$$d(f_n, f) = \sup_{x \in [1, \infty)} |f_n(x) - x| = \lim_{x \rightarrow \infty} |f_n(x) - x| = \lim_{x \rightarrow \infty} \left| \frac{\frac{1}{x} - 1}{n + \frac{1}{x}} \right| = \frac{1}{n}$$

Then by the Archimedean principle, it is clear that  $f_n$  converges uniformly to  $f$  on  $[1, \infty)$ .

3 Given a metric space  $(X, d)$ , let  $C(X)$  denote the set of bounded and continuous functions  $f: X \rightarrow \mathbb{R}$ . For  $f, g \in C(X)$ , we define

$$d(f, g) = \sup_{x \in X} |f(x) - g(x)|.$$

- Prove that  $(C(X), d)$  is a metric space.
- Show that  $C(X)$  is complete, connected, but not compact.

**Solution** a.  $C(X)$  is clearly non-empty. One such function in  $C(X)$  is  $f(x) \equiv 0$ . Then we need to show that  $d$  is a metric.

$d(f, g) \geq 0$  clearly since it is the supremum of an absolute value.

$$d(f, g) = 0 \iff \sup_{x \in X} |f(x) - g(x)| = 0 \iff |f(x) - g(x)| = 0 \iff f(x) = g(x) \forall x \in X \iff f = g$$

$$d(f, g) = \sup_{x \in X} |f(x) - g(x)| = \sup_{x \in X} |g(x) - f(x)| = d(g, f)$$

$$d(f, g) = \sup_{x \in X} |f(x) - g(x)|$$

$$= \sup_{x \in X} |(f(x) - h(x)) + (h(x) - g(x))|$$

$$\leq \sup_{x \in X} |f(x) - h(x)| + \sup_{x \in X} |h(x) - g(x)| = d(f, h) + d(h, g)$$

- b. Let  $\{f_n\}_{n \geq 1}$  be a Cauchy sequence of functions in  $C(X)$ . Fix  $x \in X$ . Then  $\{f_n(x)\}_{n \geq 1}$  is Cauchy in  $\mathbb{R}$ . As  $\mathbb{R}$  is complete,  $f_n(x) \xrightarrow{n \rightarrow \infty} f(x) \in \mathbb{R}$ . This applies for all  $x$ , so hence,  $f_n$  converges pointwise to  $f$ . We now show that  $f$  belongs to  $C(X)$ .

To show that  $f$  is continuous, since all the  $f_n$  are continuous, it suffices to show that they converge uniformly to  $f$ .

Fix  $\varepsilon > 0$ . Then as  $\{f_n\}_{n \geq 1}$  is Cauchy, there exists  $N_1 \in \mathbb{N}$  such that for all  $n, m \geq N_1$ , we have

$$d(f_n, f_m) = \sup_{x \in X} |f_n(x) - f_m(x)| < \frac{\varepsilon}{100}.$$

for all  $x \in X$ . Taking  $m \rightarrow \infty$  yields

$$d(f_n, f) = \sup_{x \in X} |f_n(x) - f(x)| \leq \frac{\varepsilon}{100} < \varepsilon$$

for all  $x \in X$ . Thus,  $f_n \xrightarrow{n \rightarrow \infty} f$  uniformly, so  $f$  is continuous on  $X$ .

Since  $f_n \xrightarrow{n \rightarrow \infty} f$  uniformly, there exists  $N \in \mathbb{N}$  such for all  $n \geq N$ , we have that

$$d(f_n, f) = \sup_{x \in X} |f_n(x) - f(x)| < 1 \implies 1 - f_n(x) \leq f(x) \leq 1 + f_n(x)$$

for all  $x \in X$ . But  $f_n(x) \in C(X)$ , so there exists  $M$  such that  $|f_n(x)| \leq M$ . Thus,

$$1 - M \leq f(x) \leq 1 + M$$

so  $f$  is bounded. Hence,  $f \in C(X)$ , so  $(C(X), d)$  is complete.

We wish to show that  $C(X)$  is connected. Let  $f, g \in C(X)$ , and define

$$\gamma: [0, 1] \rightarrow C(X), \quad \gamma(t) = (1 - t)f + tg$$

Note that  $\gamma(0) = f$  and  $\gamma(1) = g$ . It suffices to show that  $\gamma$  is continuous on  $[0, 1]$ .

Note that for all  $t \in [0, 1]$ ,  $\gamma(t) \in C(X)$ . Indeed, it is continuous because it is a sum and product of continuous functions, and it is bounded because  $f$  and  $g$  are both bounded. We now wish to prove that  $\gamma$  is continuous.

As  $g$  and  $f$  are bounded, there exists  $M \in \mathbb{R}$  such that  $|f(x)| \leq \frac{M}{2}$  and  $|g(x)| \leq \frac{M}{2}$  for all  $x \in X$ .

Fix  $\varepsilon > 0$ ,  $t_0 \in [0, 1]$ , and choose  $\delta = \frac{\varepsilon}{2M}$ . Then if  $|t - t_0| < \delta$ , we have

$$\begin{aligned} d(\gamma(t), \gamma(t_0)) &= \sup_{x \in X} |\gamma(t)(x) - \gamma(t_0)(x)| \\ &= \sup_{x \in X} |(1 - t)f(x) + tg(x) - (1 - t_0)f(x) - t_0g(x)| \\ &= \sup_{x \in X} |f(x)(t_0 - t) + g(x)(t - t_0)| \\ &= |t - t_0| \sup_{x \in X} |g(x) - f(x)| \\ &\leq |t - t_0| \left( \sup_{x \in X} |g(x)| + \sup_{x \in X} |f(x)| \right) \\ &\leq \frac{\varepsilon}{2M} \cdot M < \varepsilon \end{aligned}$$

Thus,  $C(X)$  is path connected  $\implies C(X)$  is connected.

Lastly,  $C(X)$  is not compact. Consider the sequence  $\{f_n\}_{n \geq 1}$  with  $f_n \equiv n$ . The sequence clearly does not admit a convergent subsequence, so  $C(X)$  is not sequentially compact  $\implies C(X)$  is not compact.

4 Consider the subset of  $C([0, 1])$  defined as follows:

$$X = \{f: [0, 1] \rightarrow \mathbb{R} \mid f(0) = 0 \text{ and } |f(x) - f(y)| \leq |x - y|\}.$$

Prove that  $X$  is compact.

**Solution** We first show that  $X$  is uniformly bounded. Let  $f \in X$ . Then  $|f(x) - f(0)| = |f(x)| \leq |x - 0| \leq 1$  for all  $x \in [0, 1]$ . Thus,  $X$  is uniformly bounded.

We now show that  $X$  is equicontinuous. Fix  $\varepsilon > 0$  and choose  $\delta = \frac{\varepsilon}{2}$ . Then for all  $f \in X$  and all  $x, y \in [0, 1]$  with  $|x - y| < \delta$ ,

$$|f(x) - f(y)| \leq |x - y| \leq \frac{\varepsilon}{2} < \varepsilon$$

Thus,  $X$  is equicontinuous.

Hence, by Arzelà–Ascoli,  $X$  is sequentially compact  $\implies X$  is compact.

5 Let  $X$  and  $Y$  be two metric spaces and let  $f: X \rightarrow Y$  be a function. Assume  $A$  and  $B$  are open subsets of  $X$  such that  $f$  is continuous on  $A$  and  $f$  is continuous on  $B$ .

- Show that  $f$  is continuous on  $A \cup B$ .
- Is this result still true if  $A$  and  $B$  were both closed subsets of  $X$ ?
- Is the result true for the union of an infinite number of open sets?
- Is the result true for the union of an infinite number of closed sets?

**Solution** a. Let  $x_0 \in A \cup B$ . Assume without loss of generality that  $x_0 \in A$ .

Fix  $\varepsilon > 0$

Then as  $A$  is open, there exists  $\delta_1$  such that  $B_{\delta_1}^X(x_0) \subseteq A$ . As  $f$  is continuous on  $A$ , there exists  $\delta_2$  such that for all  $x \in B_{\delta_2}^X(x_0) \cap A$ , we have  $f(x) \in B_\varepsilon^Y(f(x_0))$ . Choose  $\delta = \min\{\delta_1, \delta_2\}$ . The  $B_\delta^X(x_0) \subseteq A$ , so for all  $x \in A \cup B$  such that  $x \in B_\delta^X(x_0)$ , we have  $f(x) \in B_\varepsilon^Y(f(x_0))$ .

We can apply the same argument, but with  $A$  and  $B$  switched. Hence,  $f$  is continuous on  $A \cup B$ .

- b. Yes. Let  $x_0 \in A \cup B$ . Assume without loss of generality that  $x_0 \in A$ .

Let  $\{x_n\}_{n \geq 1} \subseteq A \cup B$  be such that  $x_n \xrightarrow{n \rightarrow \infty} x_0$ . Let  $\{a_n\}_{n \geq 1}$ , where  $a_n$  is the subsequence containing all  $x_n$  such that  $x_n \in A$ . Similarly, let  $\{b_n\}_{n \geq 1}$  be the subsequence containing all  $x_n$  such that  $x_n \in B$ .

If  $\{b_n\}_{n \geq 1}$  has finitely many terms, then there exists  $N \in \mathbb{N}$  such that if  $n \geq N$ ,  $x_n \in A$ . Then  $f(x_n) \xrightarrow{n \rightarrow \infty} f(x_0)$ , since  $f$  is continuous on  $A$ .

If  $\{a_n\}_{n \geq 1}$  has finitely many terms, then we can apply the same argument as above with  $a_n$ ,  $b_n$  and  $A$ ,  $B$  switched.

If both subsequences have infinitely many terms, then we have that  $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n = x_0$  by uniqueness of limits, so  $x_0 \in A \cap B$ . Then as  $f$  is continuous on both  $A$  and  $B$ , we have that

$$\left. \begin{array}{l} a_n \xrightarrow{n \rightarrow \infty} x_0 \\ b_n \xrightarrow{n \rightarrow \infty} x_0 \end{array} \right\} \implies x_n \xrightarrow{n \rightarrow \infty} x_0$$

Hence  $f$  is continuous on  $A \cup B$ .

- c. Yes. We can apply the same argument:  $x_0$  must belong to one open set, say  $A$ , so we can fit an open ball in  $A$ . As  $f$  is continuous on  $A$ , we can make  $\delta$  small enough so that  $B_\delta^X(x_0) \subseteq A$  and  $f(B_\delta^X(x_0)) \subseteq B_\varepsilon^Y(f(x_0))$ .

- d. No. Consider  $f: \mathbb{R} \rightarrow \mathbb{R}$ , with  $f(0) = 0$  and  $f(x) = 1$  otherwise. Then  $f$  is continuous on  $\{0\}$  and  $[\frac{1}{n}, \infty)$  for all  $n \geq 1$ .

But  $\{0\} \cup \bigcup_{n=1}^{\infty} [\frac{1}{n}, \infty) = [0, \infty)$ , and  $f$  is clearly not continuous at  $x = 0$ .

- 6 Let  $f: [0, 1] \rightarrow \mathbb{R}$  be a function with Darboux's property such that for any  $y \in \mathbb{R}$ , the set  $f^{-1}(\{y\})$  is closed. Prove that  $f$  is continuous.

**Solution** Suppose  $f$  is discontinuous at  $x_0 \in [0, 1]$ . Then there exists  $\varepsilon_0 > 0$  such that for all  $\delta > 0$ , there exists  $x_\delta$  such that  $|x_0 - x_\delta| < \delta$  but  $|f(x_0) - f(x_\delta)| \geq \varepsilon_0$ . Assume without loss of generality that this implies

$$f(x_\delta) \geq \varepsilon_0 + f(x_0)$$

If we parametrize  $\delta$  as  $\frac{1}{n}$ , then we get a sequence  $\{x_n\}_{n \geq 1}$  with  $x_n \xrightarrow{n \rightarrow \infty} x_0$  and  $f(x_n) \geq \varepsilon_0 + f(x_0)$ .

As  $f(x_0) < \varepsilon_0 + f(x_0) \leq f(x_n)$ , then by the Darboux property, there exists  $y_n \in (x_n, x_0)$  (or  $y_n \in (x_0, x_n)$  if  $x_n > x_0$ ) such that  $f(y_n) = \varepsilon_0 + f(x_0)$ . Note that  $y_n \xrightarrow{n \rightarrow \infty} x_0$  also.

Since  $\{y_n\}_{n \geq 1} \subseteq f^{-1}(\{\varepsilon_0 + f(x_0)\})$  closed, we have that  $y_n \xrightarrow{n \rightarrow \infty} x_0 \in f^{-1}(\{\varepsilon_0 + f(x_0)\})$  by completeness and uniqueness of limits. But this implies that  $f(x_0) = \varepsilon_0 + f(x_0) = f(x_0)$ , which is a contradiction. Hence,  $f$  is continuous.

In the case that  $f(x_\delta) \leq -\varepsilon_0 + f(x_0)$ , the same argument applies, but with inequalities switched around.

- 7 Let  $f, g: [a, b] \rightarrow [a, b]$  be two continuous functions such that  $f \circ g = g \circ f$ . Show that there exists  $x_0 \in [a, b]$  such that  $f(x_0) = g(x_0)$ .

**Solution** Assume  $f(x) \neq g(x)$  for all  $x \in [a, b]$ . Assume without loss of generality that  $f(x) < g(x)$ .

By Brouwer's fixed point theorem, there exists  $x_0 \in [0, 1]$  such that  $f(x_0) = x_0$ . Then define

$$\begin{aligned} x_1 &= g(x_0) = g(f(x_0)) = f(g(x_0)) = f(x_1) \\ x_2 &= g(x_1) = g(f(x_1)) = f(g(x_1)) = f(x_2) \end{aligned}$$

Repeating this process inductively yields a sequence  $\{x_n\}_{n \geq 1}$  such that

$$\begin{aligned} x_{n+1} &= g(x_n) \\ x_n &= f(x_n) \end{aligned}$$

for all  $n \geq 1$ . Also, as  $g(x) > f(x)$ , we have that  $x_{n+1} = g(x_n) > f(x_n) = x_n$ . So  $x_n$  is monotonically increasing. Moreover, as each  $x_n \in [a, b]$ ,  $x_n$  is bounded above by  $a$ . Thus, the sequence must converge. Call its limit  $x$ . Since  $[a, b]$  is compact, we have by uniqueness of limits that  $x \in [a, b]$ . Since  $f$  and  $g$  are continuous on  $[a, b]$ , we have

$$\lim_{n \rightarrow \infty} g(x_n) = g(x) = f(x) = \lim_{n \rightarrow \infty} f(x_n)$$

as desired.

If  $g(x) < f(x)$ , then the argument would be the same, but  $x_n$  would be decreasing instead. However, it would be bounded below by  $a$  and we would get the same result.