



Math 673

# Multigrid Methods: A Mostly Matrix-Based Approach

## Chapter06: Multigrid and the Conforming Finite Element Method

Abner J. Salgado and Steven M. Wise

asalgad1@utk.edu swise1@utk.edu  
University of Tennessee

Fall 2024



# Chapter 06, Part 1 of 2

## Multigrid and the Conforming Finite Element Method



## Introduction

Let  $\Omega \subset \mathbb{R}^d$ ,  $d = 1, 2$  or  $3$ , be an open polyhedral domain. Often, we will also assume that  $\Omega$  is convex. The weak form of the model problem may be expressed as follows: given  $f \in L^2(\Omega)$ , find  $u \in H_0^1(\Omega)$  such that

$$a(u, v) = (f, v)_{L^2(\Omega)}, \quad \forall v \in H_0^1(\Omega), \quad (1)$$

where

$$a(u, v) := (\nabla u, \nabla v)_{L^2(\Omega)}. \quad (2)$$


The finite element approximation is based on this weak formulation. In this chapter, we will use the theory of the last chapter to prove that basic multigrid algorithms will converge when applied to the finite element approximation of the model problem. The path forward is simple. We need only check that the basic assumptions hold.



# Nested Families of Finite Element Spaces



Let  $\mathcal{T}_0$  be a conforming triangulation of  $\Omega$ . This means that there are no “hanging nodes”. We define  $\mathcal{T}_1$  to be the triangulation of  $\Omega$  that results from bisecting ( $d = 1$ ) or quadrisecting ( $d = 2$ ) simplices of the triangulation  $\mathcal{T}_0$ . See the two following two figures. The refinement of the tetrahedra in  $d = 3$  is more complicated, and we skip that case for the sake of brevity.



**Figure:** Bisecting the triangulation  $\mathcal{T}_0$  in 1D.




Figure: Quadrisectioning the triangulation  $\mathcal{T}_0$  in 2D.

For  $d = 2$ , we connect the edge midpoints. Observe that the daughter triangles are similar to the mother. Continuing, we can recursively define a nested family of the conforming triangulations, indexed by  $\ell$ ,

$$\mathcal{T}_0, \mathcal{T}_1, \mathcal{T}_2, \dots, \mathcal{T}_\ell, \dots, \mathcal{T}_L.$$




## Definition

Let  $\{\mathcal{T}_\ell\}_{\ell=0}^L$  be a nested family of triangulations of  $\Omega$ . For  $0 \leq \ell \leq L$ , define the **grid spacing**,  $h_\ell$ , via


$$h_\ell := \max_{K \in \mathcal{T}_\ell} \text{diam}(K).$$

Subordinate to  $\mathcal{T}_\ell$ , define the **finite element space**,  $V_\ell$ , via

$$V_\ell := \left\{ v_\ell \in C^0(\overline{\Omega}) \mid v_\ell|_K \in \mathbb{P}_1(K), \forall K \in \mathcal{T}_\ell, v_\ell|_{\partial\Omega} \equiv 0 \right\}.$$



$$\mathcal{T}_1 : n_1 = 13.$$



$$\mathcal{T}_0 : n_0 = 2$$

**Figure:** Quadrisecting a complicated triangulation  $\mathcal{T}_0$  in 2D. Interior vertices are marked with a filled dot.





## Definition

Let  $0 \leq \ell \leq L$ , and suppose  $V_\ell$  and  $\mathcal{T}_\ell$  are defined as above. Suppose that  $\{\mathbf{N}_{\ell,j}\}_{j=1}^{n_\ell}$  is the set of interior vertices of the triangulation  $\mathcal{T}_\ell$ . By  $\mathcal{B}_\ell$  we denote the **Lagrange nodal basis** for  $V_\ell$ , that is,

$$\mathcal{B}_\ell := \{\psi_{\ell,i}\}_{i=1}^{n_\ell}, \quad (3)$$

where  $\psi_{\ell,i} \in V_\ell$  is the unique function with the property that


$$\psi_{\ell,i}(\mathbf{N}_{\ell,j}) = \delta_{i,j}.$$

The functions  $\psi_{\ell,i}$  are called **hat functions**.


## Remark

*We will always assume that  $n_0 > 0$ . In other words, there is always at least interior vertex.*

See the next two figures for plots of hat functions in two dimensions.



**Figure:** Interior vertices  $N_{0,1}$  and  $N_{0,2}$  in the triangulation  $\mathcal{T}_0$ , and the basis function,  $\psi_{0,2}$ .



**Figure:** A uniform mesh in 2D and a Langrange nodal basis (hat) function.



The following is easily established.

### Proposition

*Let  $0 \leq \ell \leq L$ , and suppose  $V_\ell$  and  $\mathcal{T}_\ell$  are defined as above. Suppose that  $\mathcal{B}_\ell := \{\psi_{\ell,i}\}_{i=1}^{n_\ell}$  is as defined in (3). Then,  $\mathcal{B}_\ell$  is a bona fide basis of  $V_\ell$ .*

### Proof.

Exercise. □



## Lemma

For  $0 \leq \ell \leq L$ , with  $V_\ell$  and  $\mathcal{T}_\ell$  defined as above, we have

$$V_1 \subset V_2 \subset \cdots \subset V_\ell \subset V_L \subset H_0^1(\Omega),$$

and each  $V_\ell$  is a finite dimensional vector space with

$$\dim(V_\ell) = n_\ell.$$

Furthermore,

$$0 < n_0 < n_1 < n_2 < \cdots < n_\ell < \cdots < n_L < \infty,$$

where  $n_\ell$  is precisely the number of interior vertices of the triangulation  $\mathcal{T}_\ell$ .

Finally,

$$h_{\ell-1} = 2h_\ell, \quad 1 \leq \ell \leq L.$$

Proof.

Exercise.





## Lemma

Suppose that  $\mathcal{B}_\ell$  is the Lagrange nodal basis for  $V_\ell$ ,  $0 \leq \ell \leq L$ . Then, for every  $\ell$ , with  $1 \leq \ell \leq L$ , there exist unique numbers

$$p_{\ell-1,j,i} \in \mathbb{R}, \quad 1 \leq j \leq n_\ell, \quad 1 \leq i \leq n_{\ell-1},$$

with the property that

$$\psi_{\ell-1,i} = \sum_{j=1}^{n_\ell} p_{\ell-1,j,i} \psi_{\ell,j}, \quad (4)$$

for each  $1 \leq i \leq n_{\ell-1}$ .



## Proof.

Since  $V_{\ell-1}$  is a linear subspace of  $V_\ell$  and  $\mathcal{B}_\ell$  is a basis for the latter, for every  $\psi_{\ell-1,i} \in \mathcal{B}_{\ell-1} \subset V_{\ell-1} \subset V_\ell$ , there exists unique coefficients

$$p_{\ell-1,j,i} \in \mathbb{R}, \quad 1 \leq j \leq n_\ell,$$

such that

$$\psi_{\ell-1,i} = \sum_{j=1}^{n_\ell} p_{\ell-1,j,i} \psi_{\ell,j}.$$

Recall that basis representation are unique. □



## Definition

For  $0 \leq \ell \leq L$ , define the **prolongation matrix**,

$$P_{\ell-1} \in \mathbb{R}^{n_\ell \times n_{\ell-1}},$$

via

$$[P_{\ell-1}]_{i,j} := p_{\ell-1,i,j}. \quad (5)$$





## Lemma

*Let  $1 \leq \ell \leq L$ , and suppose  $v_{\ell-1} \in V_{\ell-1}$  is arbitrary. Suppose  $\mathbf{v}_{\ell-1}$  is the coordinate vector of  $v_{\ell-1}$  in the Lagrange nodal basis  $\mathcal{B}_{\ell-1}$  and  $\mathbf{v}_{\ell}$  is the coordinate vector of  $v_{\ell-1}$  in the basis  $\mathcal{B}_{\ell}$ . Then*

$$\mathbf{v}_{\ell} = \mathbf{P}_{\ell-1} \mathbf{v}_{\ell-1}.$$



## Proof.

We write, as usual,

$$[\mathbf{v}_{\ell-1}]_i = v_{\ell-1,i}, \quad 1 \leq i \leq n_{\ell-1},$$

and

$$[\mathbf{v}_{\ell}]_i = v_{\ell,i}, \quad 1 \leq i \leq n_{\ell}.$$

Thus

$$\begin{aligned} v_{\ell-1} &= \sum_{i=1}^{n_{\ell-1}} v_{\ell-1,i} \psi_{\ell-1,i} \\ &\stackrel{(4)}{=} \sum_{i=1}^{n_{\ell-1}} v_{\ell-1,i} \sum_{j=1}^{n_{\ell}} p_{\ell-1,j,i} \psi_{\ell,j} \\ &= \sum_{j=1}^{n_{\ell}} \left\{ \sum_{i=1}^{n_{\ell-1}} p_{\ell-1,j,i} v_{\ell-1,i} \right\} \psi_{\ell,j} \\ &= \sum_{j=1}^{n_{\ell}} [\mathbf{P}_{\ell-1} \mathbf{v}_{\ell-1}]_j \psi_{\ell,j}. \end{aligned}$$



## Proof (Cont.)

But

$$\mathbf{v}_{\ell-1} = \sum_{j=1}^{n_\ell} [\mathbf{v}_\ell]_j \psi_{\ell,j}.$$

Since basis representations are unique,

$$\mathbf{v}_\ell = \mathbf{P}_{\ell-1} \mathbf{v}_{\ell-1}.$$





## Definition

For  $1 \leq \ell \leq L$ , define the **restriction matrix** via

$$R_{\ell-1} := P_{\ell-1}^T \in \mathbb{R}^{n_{\ell-1} \times n_{\ell}}.$$



### Lemma

*For  $1 \leq \ell \leq L$  and suppose  $R_{\ell-1}$  and  $P_{\ell-1}$  are defined as above. Then*

$$\text{rank}(P_{\ell-1}) = \text{rank}(R_{\ell-1}) = n_{\ell-1}.$$



## Proof.

Suppose

$$P_{\ell-1} \mathbf{v}_{\ell-1} = \mathbf{0} \in \mathbb{R}^{n_\ell}.$$

This represents a linear combination of the  $n_{\ell-1}$  columns of  $P_{\ell-1}$ . Using the notation from the last lemma and its proof, we have

$$\mathbf{v}_\ell = \mathbf{0} \in \mathbb{R}^{n_\ell},$$

where  $\mathbf{v}_{\ell-1}$  and  $\mathbf{v}_\ell$  are coordinate vectors of some function  $v_{\ell-1} \in V_{\ell-1}$  in the bases  $\mathcal{B}_{\ell-1}$  and  $\mathcal{B}_\ell$ , respectively.

The only way that  $\mathbf{v}_\ell = \mathbf{0}$  is if  $v_{\ell-1} \equiv 0$  in  $V_{\ell-1}$ . But then  $\mathbf{v}_{\ell-1} = \mathbf{0}$ . Thus, the columns of  $P_{\ell-1}$  are linearly independent and

$$\text{rank}(P_{\ell-1}) = n_{\ell-1}.$$





# The Stiffness Matrices



## Definition

For  $\ell \geq 0$  and  $V_\ell, \mathcal{T}_\ell$  defined as above, we define the **stiffness matrices**  $A_\ell \in \mathbb{R}^{n_\ell \times n_\ell}$  via

$$[A_\ell]_{i,j} := a(\psi_{\ell,j}, \psi_{\ell,i})$$

for all  $1 \leq i, j \leq n_\ell$ , where  $a(\cdot, \cdot)$  is the energy inner product defined in (2).





We need the following well-known result to show that the stiffness matrix is positive definite.

### Theorem (Poincaré inequality)

*Suppose  $d = 1, 2$  or  $3$  and  $\Omega$  is an open polyhedral domain. There is a constant  $C_P > 0$ , depending only on the domain  $\Omega$ , such that*

$$C_P \|v\|_{L^2(\Omega)}^2 \leq a(v, v), \quad (6)$$

*for all  $v \in H_0^1(\Omega)$ .*



## Lemma

*The stiffness matrices  $A_\ell \in \mathbb{R}^{n_\ell \times n_\ell}$  are all SPD. Moreover, they satisfy the Galerkin condition (Assumption (G0)), that is, for  $1 \leq \ell \leq L$ ,*

$$A_{\ell-1} = R_{\ell-1} A_\ell P_{\ell-1}. \quad (7)$$



## Proof.

(Symmetry):

$$\begin{aligned}[A_\ell]_{i,j} &= a(\psi_{\ell,j}, \psi_{\ell,i}) \\ &= a(\psi_{\ell,i}, \psi_{\ell,j}) \\ &= [A_\ell]_{j,i}.\end{aligned}$$

(Positivity): Let  $\mathbf{v}_\ell \in \mathbb{R}^{n_\ell}$  be arbitrary and suppose  $v_\ell \in V_\ell$  is the unique function with coordinates  $\mathbf{v}_\ell$ . Then, using the Poincaré inequality,

$$\begin{aligned}0 \leq C_P \|\mathbf{v}_\ell\|_{L^2(\Omega)}^2 &\stackrel{(6)}{\leq} a(\mathbf{v}_\ell, \mathbf{v}_\ell) \\ &= a\left(\sum_{j=1}^{n_\ell} v_{\ell,j} \psi_{\ell,j}, \sum_{j=1}^{n_\ell} v_{\ell,i} \psi_{\ell,i}\right) \\ &= \sum_{i=1}^{n_\ell} \sum_{j=1}^{n_\ell} v_{\ell,i} a(\psi_{\ell,j}, \psi_{\ell,i}) v_{\ell,j} \\ &= \mathbf{v}_\ell^T A_\ell \mathbf{v}_\ell.\end{aligned}$$

But  $\mathbf{v}_\ell \equiv 0$  iff  $\mathbf{v}_\ell = \mathbf{0}$ . Hence  $A_\ell$  is SPD.



## Proof (Cont.)

(Galerkin condition): By definition

$$\begin{aligned}
 [A_{\ell-1}]_{i,j} &= a(\psi_{\ell-1,j}, \psi_{\ell-1,i}) \\
 &\stackrel{(4)}{=} a\left(\sum_{s=1}^{n_\ell} p_{\ell-1,s,j} \psi_{\ell,s}, \sum_{t=1}^{n_\ell} p_{\ell-1,t,i} \psi_{\ell,t}\right) \\
 &= \sum_{s=1}^{n_\ell} \sum_{t=1}^{n_\ell} p_{\ell-1,s,j} a(\psi_{\ell,s}, \psi_{\ell,t}) p_{\ell-1,t,i} \\
 &= \sum_{t=1}^{n_\ell} \sum_{s=1}^{n_\ell} p_{\ell-1,t,i} a(\psi_{\ell,s}, \psi_{\ell,t}) p_{\ell-1,s,j} \\
 &\stackrel{(5)}{=} \sum_{t=1}^{n_\ell} \sum_{s=1}^{n_\ell} [R_{\ell-1}]_{i,t} [A_\ell]_{t,s} [P_{\ell-1}]_{j,s}.
 \end{aligned}$$

Thus

$$A_{\ell-1} = R_{\ell-1} A_\ell P_{\ell-1}.$$





## Remark

*Observe that this last result also confirms the fact that  $P_{\ell-1}$  and  $R_{\ell-1}$  are of full rank. Otherwise,  $A_{\ell-1}$  could not be positive definite.*



# The Stiffness Matrices

Now, for our level- $\ell$  Euclidean inner products and norms, we use the now-familiar notation:

$$(\mathbf{u}_\ell, \mathbf{v}_\ell)_\ell := \mathbf{v}_\ell^T \mathbf{u}_\ell, \quad \forall \mathbf{u}_\ell, \mathbf{v}_\ell \in \mathbb{R}^{n_\ell},$$

and

$$\|\mathbf{u}_\ell\|_\ell := \sqrt{(\mathbf{u}_\ell, \mathbf{u}_\ell)_\ell} \quad \forall \mathbf{u}_\ell \in \mathbb{R}^{n_\ell}.$$

The stiffness-matrix induced inner products and norms are, recall,

$$(\mathbf{u}_\ell, \mathbf{v}_\ell)_{A_\ell} := (A_\ell \mathbf{u}_\ell, \mathbf{v}_\ell)_\ell, \quad \forall \mathbf{u}_\ell, \mathbf{v}_\ell \in \mathbb{R}^{n_\ell},$$

and

$$\|\mathbf{u}_\ell\|_{A_\ell} := \sqrt{(\mathbf{u}_\ell, \mathbf{u}_\ell)_{A_\ell}} \quad \forall \mathbf{u}_\ell \in \mathbb{R}^{n_\ell}.$$



Next we estimate the size of the condition number of  $A_\ell$ . To do this we need a couple of results whose proofs can be found in most good finite element books.

### Lemma

*Suppose  $d = 1, 2$  or  $3$  and  $0 \leq \ell \leq L$ . Let  $v_\ell \in V_\ell$  be arbitrary and assume that  $\mathbf{v}_\ell \in \mathbb{R}^{n_\ell}$  is its unique coordinate vector in the basis  $\mathcal{B}_\ell$ . Then, there are constants  $C_2 \geq C_1 > 0$ , both independent of  $\ell$  and  $v_\ell$ , such that*

$$C_1 h_\ell^d \|\mathbf{v}_\ell\|_\ell^2 \leq \|v_\ell\|_{L^2(\Omega)}^2 \leq C_2 h_\ell^d \|\mathbf{v}_\ell\|_\ell^2. \quad (8)$$



## Lemma

Suppose  $d = 1, 2$  or  $3$  and  $0 \leq \ell \leq L$ . There is a constant  $C_3 > 0$  independent of  $\ell \geq 0$  such that

$$a(v_\ell, v_\ell) \leq C_3 h_\ell^{-2} \|v_\ell\|_{L^2(\Omega)}^2. \quad (9)$$

for all  $v_\ell \in V_\ell$ . As a consequence of (8),

$$a(v_\ell, v_\ell) \leq C_2 C_3 h_\ell^{d-2} \|\mathbf{v}_\ell\|_\ell^2, \quad (10)$$

for all  $v_\ell \in V_\ell \xleftrightarrow{\mathcal{B}_\ell} \mathbf{v}_\ell \in \mathbb{R}^{n_\ell}$ .





## Remark

*These results require some conditions on the underlying family of conforming meshes. such as global quasi-uniformity and shape regularity, which hold thanks to our construction of the family  $\mathcal{T}_\ell$ ,  $\ell \geq 0$ .*

## Remark

*Estimate (8) is an example of finite-dimensional norm equivalence, and estimate (9) is called an **inverse inequality**.*



Proofs of the following facts can be found in the book by Braess.

### Lemma (Asymptotic sharpness)

*Let  $V_\ell$  and  $\mathcal{T}_\ell$  be defined as usual. There exists a family of non-trivial functions  $\tilde{v}_\ell \in V_\ell$  and a constant  $C_4 > 0$ , independent of  $\ell$  and  $\tilde{v}_\ell$ , such that, for every  $0 \leq \ell \leq L$*

$$C_4 h_\ell^{-2} \|\tilde{v}_\ell\|_{L^2(\Omega)}^2 \leq a(\tilde{v}_\ell, \tilde{v}_\ell). \quad (11)$$

*There exists a family of non-trivial functions  $\hat{v}_\ell \in V_\ell$  and a constant  $C_5 > 0$ , independent of  $\ell$  and  $\hat{v}_\ell$ , such that, for every  $0 \leq \ell \leq L$*

$$a(\hat{v}_\ell, \hat{v}_\ell) \leq C_5 \|\hat{v}_\ell\|_{L^2(\Omega)}^2. \quad (12)$$



## Theorem

*Let  $d = 1, 2$ , or  $3$  and  $V_\ell$  and  $\mathcal{T}_\ell$  be defined as usual. There exist constants  $C_7 \geq C_6 > 0$ , independent of  $\ell \geq 0$ , such that*

$$C_6 h_\ell^{-2} \leq \kappa_2(A_\ell) = \frac{\lambda_\ell^{(n_\ell)}}{\lambda_\ell^{(1)}} \leq C_7 h_\ell^{-2}. \quad (13)$$

*In particular, there are constant  $C_7^{(i)}, C_6^{(i)} > 0$  for  $i = 1, n_\ell$ , such that*

$$C_6^{(n_\ell)} h_\ell^{d-2} \leq \lambda_\ell^{(n_\ell)} \leq C_7^{(n_\ell)} h_\ell^{d-2}$$

*and*

$$C_7^{(1)} h_\ell^d \leq \lambda_\ell^{(1)} \leq C_6^{(1)} h_\ell^d.$$



## Proof.

First we recall some basis facts for the Rayleigh quotient for  $A_\ell$ :

$$R(\mathbf{v}_\ell) := \frac{\mathbf{v}_\ell^T A_\ell \mathbf{v}_\ell}{\mathbf{v}_\ell^T \mathbf{v}_\ell}.$$

The smallest and largest eigenvalues satisfy

$$\lambda_\ell^{(1)} = R\left(\mathbf{v}_\ell^{(1)}\right) = \min_{\mathbf{v}_\ell} R(\mathbf{v}_\ell) > 0$$

and

$$\lambda_\ell^{(n_\ell)} = R\left(\mathbf{v}_\ell^{(n_\ell)}\right) = \max_{\mathbf{v}_\ell} R(\mathbf{v}_\ell),$$

where

$$A_\ell \mathbf{v}_\ell^{(k)} = \lambda_\ell^{(k)} \mathbf{v}_\ell^{(k)}, \quad 1 \leq k \leq n_\ell.$$



## Proof (Cont.)

(Upper bound in (13)): As usual we use the correspondence

$$\mathbf{v}_\ell \in V_\ell \xleftrightarrow{\mathcal{B}_\ell} \mathbf{v}_\ell \in \mathbb{R}^{n_\ell}.$$

Then, for arbitrary  $\mathbf{v}_\ell \in V_\ell$ ,

$$\begin{aligned} R(\mathbf{v}_\ell) &:= \frac{\mathbf{v}_\ell^T \mathbf{A}_\ell \mathbf{v}_\ell}{\mathbf{v}_\ell^T \mathbf{v}_\ell} \\ &= \frac{a(\mathbf{v}_\ell, \mathbf{v}_\ell)}{\|\mathbf{v}_\ell\|_\ell^2} \\ &\stackrel{(10)}{\leq} \frac{C_2 C_3 h_\ell^{d-2} \|\mathbf{v}_\ell\|_\ell^2}{\|\mathbf{v}_\ell\|_\ell^2} \\ &=: C_7^{(n_\ell)} h_\ell^{d-2}. \end{aligned}$$

This implies

$$\lambda_\ell^{(n_\ell)} \leq C_7^{(n_\ell)} h_\ell^{d-2}.$$



## Proof (Cont.)

Similarly,

$$\begin{aligned}
 R(\mathbf{v}_\ell) &:= \frac{a(\mathbf{v}_\ell, \mathbf{v}_\ell)}{\|\mathbf{v}_\ell\|_\ell^2} \\
 &\stackrel{(6)}{\geq} \frac{C_P \|\mathbf{v}_\ell\|_{L^2(\Omega)}^2}{\|\mathbf{v}_\ell\|_\ell^2} \\
 &\stackrel{(8)}{\geq} \frac{C_P C_1 h_\ell^d \|\mathbf{v}_\ell\|_\ell^2}{\|\mathbf{v}_\ell\|_\ell^2} \\
 &=: C_7^{(1)} h_\ell^d.
 \end{aligned}$$

Therefore,

$$\lambda_\ell^{(1)} \geq C_7^{(1)} h_\ell^d.$$

We conclude that

$$\kappa_2(\mathbf{A}_\ell) = \frac{\lambda_\ell^{(n_\ell)}}{\lambda_\ell^{(1)}} \leq \frac{C_7^{(n_\ell)} h_\ell^{d-2}}{C_7^{(1)} h_\ell^d} =: C_7 h_\ell^{-2}$$



## Proof (Cont.)

(Lower bound in (13)): Next, it follows that

$$\begin{aligned}
 \lambda_\ell^{(n_\ell)} &= R\left(\mathbf{v}_\ell^{(n_\ell)}\right) \\
 &\geq R(\tilde{\mathbf{v}}_\ell) \\
 &= \frac{a(\tilde{\mathbf{v}}_\ell, \tilde{\mathbf{v}}_\ell)}{\|\tilde{\mathbf{v}}_\ell\|_\ell^2} \\
 &\stackrel{(11)}{\geq} \frac{C_4 h_\ell^{-2} \|\tilde{\mathbf{v}}_\ell\|_{L^2(\Omega)}^2}{\|\tilde{\mathbf{v}}_\ell\|_\ell^2} \\
 &\stackrel{(8)}{\geq} \frac{C_1 C_4 h_\ell^{d-2} \|\tilde{\mathbf{v}}_\ell\|_\ell^2}{\|\tilde{\mathbf{v}}_\ell\|_\ell^2} \\
 &=: C_6^{(n_\ell)} h_\ell^{d-2},
 \end{aligned}$$

where we use the correspondence

$$\tilde{\mathbf{v}}_\ell \in V_\ell \xleftrightarrow{\mathcal{B}_\ell} \tilde{\mathbf{v}}_\ell \in \mathbb{R}^{n_\ell}.$$



## Proof (Cont.)

Finally,

$$\begin{aligned}
 \lambda_{\ell}^{(1)} &= R\left(\mathbf{v}_{\ell}^{(1)}\right) \\
 &\leq R(\hat{\mathbf{v}}_{\ell}) \\
 &= \frac{a(\hat{\mathbf{v}}_{\ell}, \hat{\mathbf{v}}_{\ell})}{\|\hat{\mathbf{v}}_{\ell}\|_{\ell}^2} \\
 &\stackrel{(12)}{\leq} \frac{C_5 \|\hat{\mathbf{v}}_{\ell}\|_{L^2(\Omega)}^2}{\|\hat{\mathbf{v}}_{\ell}\|_{\ell}^2} \\
 &\stackrel{(8)}{\leq} \frac{C_2 C_5 h_{\ell}^d \|\hat{\mathbf{v}}_{\ell}\|_{\ell}^2}{\|\hat{\mathbf{v}}_{\ell}\|_{\ell}^2} \\
 &=: C_6^{(1)} h_{\ell}^d,
 \end{aligned}$$

where we used the correspondence

$$\hat{\mathbf{v}}_{\ell} \in V_{\ell} \xleftrightarrow{\mathcal{B}_{\ell}} \hat{\mathbf{v}}_{\ell} \in \mathbb{R}^{n_{\ell}}.$$





## Proof (Cont.)

We conclude that

$$\kappa_2(A_\ell) \geq \frac{C_6^{(n_\ell)} h_\ell^{d-2}}{C_6^{(1)} h_\ell^d} =: C_6 h_\ell^{-2}.$$

