

## Angler Satisfaction Survey Results as Indicators of Fisheries Changes: A Case Study

Journal:	<i>North American Journal of Fisheries Management</i>
Manuscript ID	Draft
Manuscript Type:	Article
Keywords:	creel survey, angler satisfaction, quantile regression, maximum likelihood, population estimation

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

Angler Satisfaction Survey Results as Indicators of  
Fisheries Changes: A Case Study

Steven H. Ranney\*

596 North Laurel Parkway

Bozeman, Montana 59718 USA

Suggested running head:

Angler Satisfaction Survey Results May Portend Fishery Health Changes

\*Email: [Steven.Ranney@gmail.com](mailto:Steven.Ranney@gmail.com)

## Abstract

**Objective:** I used long-term datasets to determine if angler complaints about fish size were evident in biological data and how complaints about angler crowding could serve as early indicators of fishery health.

**Methods:** I used standardized sampling data from 2004 through 2022 and evaluated fish weight-at-length with quantile regression compared to the reference year 2003. I also estimated population sizes for Brown Trout *Salmo trutta* and Rainbow Trout *Onchorhynchus mykiss* with maximum likelihood estimation, and use extant data related to angler demographics, catch-and-release mortality, catch rate, and summer season-length to predict changes to the Brown and Rainbow Trout populations with concomitant changes to the angler population.

**Result:** For all years except 2004, Brown and Rainbow Trout weight-at-length was significantly lower when compared to the reference population showing that angler complaints were reflected in the biological data. A significant linear model of estimated population size as a function of time predicts that Brown and Rainbow Trout populations could reach 0 at long-term sampling locations by 2025. Angler population size predictions and expected catch rates suggest that during summer season 2023, each Brown Trout was caught an estimated 1.8-4.7 times and Rainbow Trout an average of 2.4-7.4 times. As angler populations increase and fish populations decrease, the average number of times a fish is caught each summer will continue to increase and could lead to reduced spawning success through repeated handling stress.

**Conclusion:** Angler satisfaction survey results could provide early insight into fisheries management questions that may have otherwise been overlooked. When paired with

analysis of long-term data sets, angler satisfaction surveys may provide fisheries scientists and managers the ability to validate or repudiate angler concerns.

**Keywords:** creel survey, angler satisfaction, quantile regression, maximum likelihood, population estimation

<A>Impact statement

Angler satisfaction data from creel surveys may provide anecdotal evidence related to potential areas of concern for fishery health. This anecdotal evidence, in conjunction with analysis of standardized long-term data sets, could provide direction to fisheries managers as areas in which they should focus their research.

<A>Introduction

Creel surveys of angler populations are the primary means through which fisheries managers collect information on angler effort and catch rates (Malvestuto 1996; Pollock et al. 1994). Surveys of recreational anglers provide crucial information related to fishing effort, harvest, and released catch to fisheries managers (McCormick and Meyer 2017). These estimates are important for providing data upon which fisheries management decisions can be based. Angler effort and catch rates, for example, can be used to estimate the number of fish harvested and released in sport fisheries (Malvestuto 1996; Pollock et al. 1994). Creel surveys are often labor intensive and expensive to conduct (Jones and Pollock 2012; McCormick and Meyer 2017) and many

rely on angler-reported data to estimate the metrics used as the basis for fisheries management (McCormick et al. 2013).

In addition to effort, harvest, and released-catch information, recreational angler creel surveys may also be used to obtain biological fisheries data. Length and weight data, some of the most important information for fisheries management (Anderson and Neumann 1996; Neumann et al. 2012), can be collected on a subsample of the harvest (Jones and Pollock 2012). Disease status, fishing mortality, and other biological indicators (e.g., structures used for aging) may be collected simultaneously. Data on angler demographics, social, and economic variables may also be collected (Jones and Pollock 2012). Angler satisfaction data may also be collected from creel surveys, whether or not they are conducted in-person or through angler self-reporting via mail or online surveys. For cases where fisheries management agencies are beholden to—either directly or indirectly—their constituents (i.e., commercial or recreational anglers), if angler satisfaction drops, anglers may voice their opinions to their elected representatives or directly to fisheries managers. In either case, hearing from unsatisfied anglers, if a frequent occurrence, could be an indicator that something has gone awry in a fishery. Fisheries managers should take these complaints seriously and investigate, if there are data available.

Analysis of long-term datasets related to angler satisfaction may provide early insight into potential problem areas in a fishery. For example, decreases in angler-reported catch rates may be an indication of negative population-level issues in the fishery population. Similarly, decreases in angler satisfaction related to the size of fish caught may also imply a survival or growth issue in the fishery. Additionally, decreases

in angler satisfaction related to angler crowding could lead to questions related to increased harvest or catch-and-release mortality, given more anglers would logically lead to increased total catch. The Madison River in Montana serves as a good case study for using angler satisfaction as insight into fisheries changes given the frequency with which anglers are surveyed, the river's global popularity, and the availability of long-term datasets.

The Madison River in Montana is one of the most popular fisheries in the state [Montana Fish, Wildlife & Parks (MTFWP) 2017; Horton et al. 2018; MTFWP 2019; Skaar 2022; Lohrenz et al. 2023] and hosted over 300,000 angler days in 2020 (Lohrenz et al. 2023), the last year for which estimated effort is available. From its source in Yellowstone National Park (National Park Service 2023) through its confluence with the Gallatin and Jefferson Rivers at Missouri Headwaters State Park near Three Forks, Montana, the river is important to local economies to support tourism and agriculture as well as jobs in the commercial angling outfitting industry (Horton et al. 2018; Lohrenz et al. 2023). Because the river is a destination for trout anglers worldwide (Skaar 2022), a number of management strategies have been implemented since the 1950s to mitigate angler conflicts and crowding (Horton et al. 2018). The most numerous and frequently-targeted fish species on the Madison River include Brown Trout *Salmo trutta*, Rainbow Trout *Onchorhynchus mykiss*, and Mountain Whitefish *Prosopium williamsoni* (Lohrenz et al. 2023). Anglers on the upper Madison River (Figure 1) are responsible for approximately 58% of all angler trips on the Madison River (MTFWP 2017, 2019; League and Caball 2022).

The upper Madison River is managed with catch-and-release regulations for trout and use of artificial lures only (MTFWP 2023). Anglers  $\leq 14$  years old may take 1 trout daily, any size (MTFWP 2023). Additional regulations related to fishing from boats and vessels are in place for two sections of the upper Madison River. Although not a regulation, most anglers on the upper Madison River use fly-fishing gear (Roulson 2002; Horton et al. 2018). Of upper-river anglers, 73.2% are not residents of Montana and nearly 90% are male (Skaar 2022). Angling is the most popular activity by far on the upper Madison River; >90% of respondents of river-use recreational surveys at 11 of 13 public river access sites upstream of Ennis, Montana, were angling (Skaar 2022).

Angler-survey results from the Madison River indicate that angler satisfaction of the size of fish caught has been declining steadily over the last five decades (Horton et al. 2018). In the 1970s, 41.1% of anglers reported the size of fish caught on the Madison River as “excellent” and only 0.2% reported the size of fish caught as “poor” (Horton et al. 2018). By the 2010s, only 14.4% of respondents reported the size of trout caught as “excellent” while 5.3% reported the same metric as “poor” (Horton et al. 2018). Additionally, recent decreases in angler satisfaction related to angler crowding have been noted with several potential mitigation strategies having gone through internal and public review; Horton et al. (2018) provides a list of past actions undertaken to reduce angler conflicts and crowding since 1959 including float fishing closures, moratoriums on new outfitters, and different restrictions on fishing from vessels. Horton et al. (2018) noted that increased angler crowding may be suggestive of a concomitant

increase in catch-and-release angling mortality and that the increased mortality from extant levels of angling pressure may be more than the fish populations can withstand.

The goal of this paper is twofold: (1) determine if the decreases in the reported angler satisfaction related to the sizes of Brown and Rainbow Trout caught on the upper Madison River, Montana, are reflected in the long-term fisheries population monitoring data and (2) evaluate the trends in population size estimates of Brown and Rainbow Trout the upper Madison River, Montana in recent years and to estimate what may happen to those populations if there are no changes in management strategies to reduce angler crowding, a complaint of anglers on the upper Madison River. I use long-term standardized sampling data to compare fish weight-at-length through quantile regression, estimate fisheries population sizes and evaluate population over time, and use extant data related to angling pressure, summer-season length, fishing day length, angler trip length, and catch rates to simulate how the angling population may change over time, estimate the average number of times individual fish are caught in a summer season, and how estimated angler population changes could impact the Brown and Rainbow Trout populations over time.

## Methods

### Quantile Regression of Weight at Length

I used 20 years' worth of fisheries data collected by MTFWP at two upper Madison River long-term fisheries index sites, Varney and Pine Butte (Figure 1; MTFWP 2023). These locations were established as long-term monitoring locations on the upper Madison River in 1981 (Pine Butte) and 1967 (Varney). The Pine Butte location is 5.3



river km (rkm) long and the Varney location is 7.8 rkm long (MTFWP 2023). All sampling was completed during the months of September and October from 2003 through 2022. All sampling events were completed by electrofishing from drift boats with mobile anodes or rafts with boom-type anodes (MTFWP 2023). Although I retrieved 21 years' worth of population data from the FishMT portal (MTFWP 2023), for allometric growth parameter comparison, I used data from every third year, beginning with 2004, to reduce the complexity of the results while still providing long-term inferences and allowing the breadth of the data set to be used. As a result, I used 8 years' worth of data: 2003 (the "reference" population), 2004, 2007, 2010, 2013, 2016, 2019, 2022. Within species, I pooled data from the Varney and Pine Butte sampling areas and assigned equal weights to both populations regardless of contributed sample size (Cade et al. 2008; Ranney 2018b).

Angler-survey results indicate that anglers claim that the size of fish caught in the upper Madison River has been declining steadily since the 1970s (Horton et al. 2018). To evaluate how allometric growth patterns have changed since 2003, I evaluated changes in 75th quantile ( $\square = 0.75$ ) intercepts ( $\beta_0$ ) and slopes ( $\beta_1$ ) for both Brown and Rainbow Trout in the upper Madison River. I used methods described in Ranney (2018b) to compare yearly  $\beta_0$  and  $\beta_1$  values by regressing  $\log_{10}W$  as a function of  $\log_{10}TL$ . I used the linear quantile regression function `rq()` in the quantile regression package "quantreg" (Koenker 2021) in R version 4.3.2 (R Development Core Team 2023). Similar to Ranney (2018b), I used  $\square = 0.75$  because this is a fish with "above average" body weight at a given length. I used a single linear model (Cade et al. 2008, 2011; Ranney 2018b; Cade and Gilham 2024) to provide estimates for separate

181 population-level  $\beta_0$  and  $\beta_1$  values when compared to a reference population based on  
182 the methods described by Ranney (2018b). I used the 2003 sampling event as my  
183 “reference” population for two reasons: (1) it was a reasonably long time ago; and (2)  
184 the Brown and Rainbow Trout populations had recovered from the whirling disease  
185 impacts from the 1990s (M. Duncan, MTFWP, personal communication). I estimated the  
186 standard error (SE) of  $\beta_{0j}$  and  $\beta_{1j}$ , where  $j$  represented the individual yearly population  
187 intercept and slope values, respectively, by resampling, with replacement, 5,000  
188  $\log_{10}W$ - $\log_{10}TL$  pairs 1000 times. I calculated 95% confidence intervals (CI) around  
189 each  $\beta_{0j}$  and  $\beta_{1j}$  with the  $t$  statistic = 1.960.

190 I compared individual yearly population  $\beta_{0j}$  and  $\beta_{1j}$  to the 2003 reference  
191 population  $\beta_0$  and  $\beta_1$  by setting the reference population as the base level then setting  
192 the contrasts between the rest of the categorical factors (i.e., yearly population identifier  
193 and its interaction with the continuous predictor variable; Ranney 2018b). Similar to  
194 calculating the SE of the  $\beta_0$  and  $\beta_1$  for each population-level model, I estimated the SE  
195 of the differences in  $\beta_{0j}$  and  $\beta_{1j}$  from the reference population  $\beta_0$  and  $\beta_1$  with resampling.  
196 I resampled 5,000  $\log_{10}W$ - $\log_{10}TL$  pairs 1000 times and calculated the 95% CIs around  
197 the differences in  $\beta_0$  and  $\beta_1$  with the  $t$  statistic = 1.960. I compared quantiles of weight at  
198 specific length values for both Brown and Rainbow Trout by regressing  $\log_{10}W$  as a  
199 function of  $\log_{10}TL$  for both species. I then estimated the weight and 95% CIs of weight  
200 for the midpoints of the length categories (Gabelhouse 1984) for Brown Trout  
201 [midpoints: substock (SS) = 75mm; stock-quality (S-Q) = 190mm; quality-preferred (Q-  
202 P) = 265mm; preferred-memorable (P-M) = 340mm; memorable-trophy (M-T) = 420mm;  
203 Milewski and Brown 1994] and Rainbow Trout [midpoints: substock (SS) = 125mm;

stock-quality (S-Q) = 325mm; quality-preferred (Q-P) = 450mm; preferred-memorable (P-M) = 575mm; memorable-trophy (M-T) = 725mm; Simpkins and Hubert 1996]

## Population Estimates

I used annual sampling data from 2003 through 2022 to make population estimates. From 2003 through 2022, Montana FWP conducted annual mark-recapture sampling in the upper Madison River by capturing, marking, and releasing individuals during at least 3 sampling events at each location each year. During mark-recapture sampling events, marked fish were released unharmed and allowed to redistribute among unmarked fish. Mortalities that occurred from either a mark or recapture event were subtracted from the total number of fish. After redistribution of marked fish, MTFWP conducted 2-4 recapture runs at each sampling location an average of 10 d (SD = 2.5 d) later.

I estimated yearly population size ( $\hat{N}$ ) and 95% CI per rkm of Brown and Rainbow Trout at the Pine Butte and Varney sampling locations by 10 mm length class in the upper Madison River. I used a maximum likelihood estimator (MLE) to estimate population size. I calculated the likelihood function as

$$L(N|n, q) = \frac{N!}{n!(N-n)!} q^n (1-q)^{N-n}$$

where  $N$  is the population size,  $n$  is the total number of fish captured in the recapture sampling event,  $q$  is the probability of capture of fish in that 10 mm length class, and  $L$  is the likelihood of  $N$  given  $n$  and  $q$  (Hayes et al. 2007). I allowed  $N$  to vary and estimated  $\hat{N}$  from the value of  $N$  for which the likelihood was maximized. I modeled  $q$  from observed values by fitting a smoothing spline to each species:year:location combination

as a function of the observed proportion of fish recaptured to the total number of fish captured in the recapture event (Anderson 1995) by 10 mm length class because capture probability can change continuously with size. I constructed 95% CI around  $\hat{N}$  by determining values for the parameters that gave a log-likelihood value that was less than the maximum value of the log-likelihood by 3.841 (Hayes et al. 2007). I used a critical value of 3.841 because this is the 5% critical value for the  $\chi^2$  distribution with 1 degree of freedom (Hayes et al. 2007).

I evaluated 20-year population trends in Brown and Rainbow Trout at Pine Butte and Varney long-term monitoring locations by summing the  $\hat{N}$  from each 10 mm length class for each year:species:location combination. I removed  $\hat{N}$  values where  $\hat{q} < 0.01$  from these final estimates because poor capture efficiency overestimates the  $\hat{N}$  for these length classes. I calculated  $\hat{N}/\text{km}$  by dividing the total  $\hat{N}$  for each monitoring location and dividing by the total length of each monitoring location and regressed  $\Sigma \hat{N}$  as a function of year with ordinary least squares regression.

#### <B>Angler Population Simulations

I used data on  $n$  anglers/year data from Horton et al. (2018) for 2011, 2013, 2015, and 2016 to create a linear model to predict anglers/year on the upper Madison River through 2030. I validated this limited model with observed anglers/year for 2017, 2019, and 2020 (MTFWP 2017, 2019; League and Caball 2022). From the linear model, I extrapolated the number of annual anglers/year through 2030 ( $\pm$  95% prediction intervals) that would fish the upper Madison River. From the linear fit and prediction intervals, I simulated 7 years' worth of summer-season anglers, trips, effort, and catch

with the AnglerCreelSurveySimulation package (Ranney 2018a) in R version 4.3.2 (R Development Core Team 2023). The AnglerCreelSurveySimulation package allows fisheries managers to simulate a creel survey to better understand how a single access-point survey or bus route-type survey may best represent their fishery, including estimates of effort ( $\hat{E}$ ) and catch (including catch rate,  $\hat{R}$ ; Jones and Robson 1991, Jones and Pollock 2012). I used predicted output from the linear model of anglers/year and data from Horton et al. (2018) and Skaar (2022) to seed the angler population simulation with data related to  $n$  anglers, summer-season length (June 1 through August 30; 91 days), fishing day length (11 hours), catch rate (Brown trout = 0.38 fish/hr; Rainbow trout = 0.62 fish/hr), and average trip length (5.6 h). The AnglerCreelSurveySimulation package assigns angler start times with the uniform distribution and catch rates with the gamma distribution with the shape parameter ( $k$ ) equal to the mean catch rate and the rate parameter equal to 1 (Ranney 2018a).

Although AnglerCreelSurveySimulation (Ranney 2018a) is designed to allow fisheries managers to estimate effort and catch of an angler population with varying sampling probabilities (i.e.,  $0 \leq p \leq 1$ ), if the sampling probability is set to 1 and there is a single “access point” (e.g., the entirety of the upper Madison River), then all anglers can be “sampled” by the simulated “surveyor” upon completion of their trip. As a result,  $\hat{E}$  and  $\hat{R}$  would equal the “true” effort and catch by each angler. In using a simulated angler population, I can observe all of the simulated trips and calculate “true” catch and effort for multiple summer seasons.

I estimated future catch-and-release related mortality on the upper Madison River from these reasonable predictions of  $n$  anglers/year. I used mortality rates from Horton

et al. (2018) to calculate the number of Brown and Rainbow Trout that would succumb to catch-and-release mortality each summer from 2023 through 2030. Mortality rates used by MTFWP are 0.02 for Brown Trout and 0.08 for Rainbow Trout (Horton et al. 2018). I estimated the number of times that a fish will be caught in the summer season by dividing  $\hat{N}$  by  $\hat{R}$  from all summer-season anglers. All analyses were conducted in R version 4.3.2 (R Development Core Team 2023) and for all analyses, alpha was set to 0.05.

<A>Results

Long-term fall electrofishing samples from 2003 through 2022 yielded observations from 57,865 Brown Trout and 41,676 Rainbow Trout. Brown Trout captures averaged 2,893 (SD = 853) and Rainbow Trout captures averaged 2,084 (SD = 543) per year. Although Rainbow Trout X Westslope Cutthroat Trout *O. clarkii lewisi* hybrids and Westslope Cutthroat Trout observations existed in the dataset, together they represented < 0.001% of the total data and were removed from consideration.

<B>Quantile Regression of Weight at Length

The  $\square = 0.75$  quantile regression model for  $\beta_0$  and  $\beta_1$  estimates from the reference data set (year = 2023) for Brown Trout were -4.6404 and 2.8818, respectively, and for Rainbow Trout were -4.4300 and 2.7955, respectively (Table 1). The 95% CIs from the reference data set for Brown Trout were  $\beta_0 = -4.6859$  to  $-4.595$  and  $\beta_1 = 2.864$  to  $2.9002$  and for Rainbow Trout were  $\beta_0 = -4.4890$  to  $-4.3710$  and  $\beta_1 = 2.7717$  to  $2.8193$ . The lowest  $\beta_0$  for Brown Trout (-5.212) and Rainbow Trout (-5.1083) was during

2013 (Table 1). The lowest  $\beta_1$  for Brown Trout (2.8305) and Rainbow Trout (2.6933) was during 2004 (Table 1). For both Brown and Rainbow Trout, 2004 was the only year in which the intercept was higher and the slope was shallower than the reference population (Table 1). For all other years investigated, intercepts were lower and slopes were steeper than the reference population for both species (Table 1). There were no instances for either species in any year in which bootstrapped 95% CIs of 75th quantile  $\beta_0$  and  $\beta_1$  overlapped with the reference data set (Table 2).

Estimates and 95% CIs of weight at TL equal to the midpoints of length categories across all quantiles from  $\square = 0.05$  up to 0.95 by increments of 0.05 overlapped for both species for all years, especially at the upper length ranges for Brown Trout (TL = 420mm; Figure 2A) and for length ranges  $\geq 450$ mm for Rainbow Trout (Figure 2B). For Brown Trout, the 2004 population was heavier than the reference population at all quantiles (except for the lowest extremes; (Figure 2A). In the substock (SS) and stock-quality (S-Q) length categories, individuals in all years after 2004 weighed less than the reference population at all quantiles 0.05 through 0.95 (Figure 2A). Overlap in predictions of weight-at-length for years after 2004 in the lower quantiles ( $\square < 0.35$ ) for quality-preferred (Q-P) and preferred-memorable (P-M) length categories became more pronounced. Similar to Brown Trout, prediction of weight-at-length for Rainbow Trout in 2004 at all quantiles for SS and S-Q length categories were higher than the reference populations except at the extreme lowest quantiles (i.e.,  $< 0.20$ ) for SS (Figure 2B). For SS and S-Q length categories, predictions of weight-at-length for years  $> 2004$  are all lower than the reference population. In Q-P, P-M, and M-T length categories, the overlap of all quantiles and years  $\geq 2004$  was challenging to

distinguish and, in many cases, predictions of weight-at-length did not follow any specific trend. Notably, the 95% CIs at TL = 575 and 725 were much broader than those in lower length categories (Figure 2B).

### <B>Population Estimates

Annual fall sampling from established, long-term monitoring locations indicated that Brown Trout populations are declining while Rainbow Trout populations appear stable or are increasing (Figure 3). A linear model describing the Brown Trout  $\hat{N}$  decreases in the upper Madison River was significant at the Pine Butte ( $R^2 = 0.1572$ ;  $\beta_0 = 73596.74$ ;  $\beta_1 = -35.84$ ;  $p = 0.047$ ) and Varney long-term monitoring locations ( $R^2 = 0.2969$ ;  $\beta_0 = 65417.88$ ;  $\beta_1 = -31.87$ ;  $p = 0.0013$ ). Slopes for linear models fit to long-term sampling  $\hat{N}$  suggest that Rainbow Trout populations are stable or increasing at both Pine Butte and Varney, but the models fit to yearly data were not significant at  $\alpha = 0.05$  (Pine Butte:  $R^2 = 0.06165$ ;  $\beta_0 = 65654.24$ ;  $\beta_1 = -31.80$ ;  $p = 0.15$ ; Varney:  $R^2 = 0.1619$ ;  $\beta_0 = -66330.16$ ;  $\beta_1 = 33.43$ ;  $p = 0.055$ ; Figure 3).

### <B>Angler Population Simulations

Angling pressure (i.e., anglers/year) from 2011 through 2016 appeared linear in Horton et al. (2018), and the linear model fit to observed (2011, 2013, 2015) and estimated (2016) data was significant (adjusted  $R^2 = 0.96$ ;  $\beta_0 = -33991356$ ;  $\beta_1 = 16949$ ;  $p = 0.013$ ; Figure 4). Although “hold-out” data did not fall on the predicted line, these data did fall within the 95% prediction interval of the linear model (Figure 4). If the linear



model of angler pressure growth continues to hold into the future, angling pressure would be expected to increase year-over-year by roughly 5% (Figure 4; Table 3).

Similar to the expected linear increase in angling pressure, Brown and Rainbow Trout catch rates were also predicted to increase year-over-year by roughly 5% (Table 3) with a concomitant increase in hooking mortality from catch-and-release fishing (Table 3). Simulations of predicted angling pressure for 2023 on the upper Madison River during the summer season predicted a total catch of Brown Trout of 235,195 individuals ( $\pm 95\%$  prediction interval of  $\sim 69,000$ ; Table 3) and 395,059 Rainbow Trout ( $\pm 95\%$  prediction interval of  $\sim 114,000$ ; Table 3). At assumed levels of catch-and-release mortality (0.02 for Brown Trout, 0.08 for Rainbow Trout; Horton et al. 2018), catch and release mortality for Brown Trout in 2023 was predicted to be 4,704 individuals and 31,605 individuals for Rainbow Trout (Table 3). By 2030, if there is no change in the expected catch rates of Brown or Rainbow Trout, summer-season catch of Brown and Rainbow Trout would be predicted to increase to 328,666 and 552,697 individuals, respectively (Table 3). If 2023 predicted catch were distributed evenly across the 86.4 rkms of the upper Madison River, Brown Trout were estimated to have been caught an average of 1.8-4.7 times at Pine Butte and Varney and Rainbow Trout were estimated to have been caught an average of 2.4-7.4 times at Pine Butte and Varney. During the predicted summer season of 2030, at  $\hat{N}$  and with predicted growth in the angler population, Brown Trout were estimated to have been caught an average of 2.9-9.8 times at Pine Butte and Varney. During a predicted 2030 summer season, Rainbow Trout at Pine Butte and Varney were estimated to have been caught 3.8-7.7 times.

## 364 <A>Discussion

365 Anglers on heavily-pressured fisheries often have anecdotal evidence about the  
366 health of the fishery before fisheries managers may have the time or funding to  
367 investigate anglers' concerns. Here, angler survey responses related to their opinions  
368 about fish size are reflected in long-term fisheries monitoring data. Similarly, evidence  
369 suggests that as angler overcrowding continues, Brown and Rainbow Trout populations  
370 may continue to drop. Because Brown Trout populations have been declining on the  
371 upper Madison River at Pine Butte and Varney since at least 2003 and declines in  
372 Brown Trout populations are outpacing limited increases from Rainbow Trout, angler  
373 survey results from Madison River anglers could "tip off" managers to a potential  
374 upcoming problem with the health of the fishery. Using declines in angler satisfaction as  
375 a potential early-warning system to system health may be a useful tool for fisheries  
376 managers.

377 The deviation between the actual weights of fish in a population and an expected  
378 length-specific weight can indicate if biotic or abiotic conditions are beneficial or  
379 detrimental to that population (Ranney et al. 2008). The changes seen here in Brown  
380 and Rainbow Trout weight-at-length growth patterns are challenging to explain without a  
381 broader study related to additional system factors (e.g., prey availability, habitat  
382 availability, temperature regimes). Increased water temperatures unrelated to  
383 management of Hebgen Dam (i.e., from climate change) may be impacting growth and  
384 weight at length of Brown and Rainbow Trout (Vincent 1978; Jobling 1995; Wootton  
385 1996; Réalis-Doyelle et al. 2018) and in the early 2000s, the Madison River basin had  
386 been going through several years of continuous drought (MTFWP 2005) which may

have been driving the negative changes in Brown and Rainbow Trout weight at length seen here as early as 2007. I chose to use quantile regression to analyze weight at length in this analysis because it is particularly helpful when evaluating populations that are regularly monitored (Cade et al. 2008, 2011; Crane et al. 2015; Crane and Farrell 2017; Ranney 2018b; Cade and Gilham 2024), especially those that are trophy fisheries or may be valuable to local economies or as a source of recreation. Further, quantile regression estimates from linear models that contain relevant grouping factors is a comprehensive and rigorous analysis of changes in weight at length (Cade et al. 2008, 2011; Ranney 2018b; Neely et al. 2021; Cade and Gilham 2024).

I estimated population sizes at two specific long-term monitoring locations with a maximum likelihood estimator. Although these  $\hat{N}$  came from standardized sampling, estimating the population size of an open population without an encounter history matrix can be challenging (Hayes et al. 2007; Link and Barker 2010). Open populations include immigration, emmigration, mortality, and recruitment (Hayes et al. 2007; Link and Barker 2010). Assuming that anyone could estimate the abundance of an open population with precision without having an encounter history matrix of uniquely marked fish would be challenging. Tagging fish with unique identifiers and encouraging anglers to report catching tagged fish may help MTFWP make better estimates of the population of Brown and Rainbow Trout in the upper Madison River. Similarly, with uniquely-marked fish and reports of catch of these uniquely marked fish, managers could better estimate the recycling rate [i.e., total catch in a fishing season divided by the number of individuals caught at least once (Schill et al. 1986; Jones et al. 2022)] of their fishery (Jones et al. 2022). Without unique tagging, it's impossible to know how much catch

410 increased because of multiple recaptures. Indeed, beginning in Autumn 2023, MTFWP  
411 Began implementing a tagging study on the upper Madison River, including uniquely-  
412 identified tags. Calculating the recycling rates of a fishery may help managers better  
413 understand how catch is distributed across the fish population. For example, for a  
414 fisheries wide recycling rate of 2.0, each fish may have been caught twice or half of the  
415 fish were caught four times (Jones et al. 2022). For the case of Brown and Rainbow  
416 Trout in the upper Madison River, estimating  $\hat{N}$  with accuracy for the entire river is  
417 challenging, given the nature of open populations. However, if the estimated angling  
418 pressure in the upper Madison River during the 2023 summer season were evenly  
419 distributed across the entire 86.4 km of the river, each Brown Trout in the Pine Butte  
420 and Varney sampling areas would have been caught 1.8-4.7 and 2.6-4.8 times,  
421 respectively. These values increase as  $\hat{N}$  drops and the angler population increases if  
422 catch rate stays the same, an unlikely prospect.

423         The catch-and-release hooking mortality rates used herein are based on work  
424 from Boyd et al. (2010) in which cumulative stressors (including the number of times a  
425 fish was caught over the summer) and temperature regimes similar to the upper  
426 Madison River were not taken into account (Horton et al. 2018). “Fight time” [i.e., time to  
427 landing (Boyd et al. 2010)] and the handling stress of increased recycling rate should be  
428 considered a cumulative stressor that should be taken into account in catch-and-release  
429 hooking-mortality estimates (Schill et al. 1986). Further, handling stress increases  
430 cortisol levels in salmonids (Donaldson 1981; Barton and Iwama 1991) and may affect  
431 the timing (Schreck et al. 2001) and success of reproduction (Campbell et al. 1992). As  
432 a result, understanding how the cumulative stress associated with multiple capture and

handling events over the few months immediately preceding expected Brown Trout spawning affects reproduction success is important in comprehending the population-level consequences of those stressors (Schreck et al. 2001). Although acute stressors were shown to not affect spawning success of semelparous salmon on their spawning grounds, more chronic events (e.g., elevated water temperatures, multiple catch-and-release events in a season) demonstrate sustained elevation of stress hormones (McConnachie et al. 2012) which carries significant reproductive costs. Including estimates of these chronic stressors into future estimates of catch-and-release hooking mortality on the Madison River would likely increase the expected estimated mortality rate and may provide additional explanations for the decline in Brown Trout observed in these data.

Anecdotal evidence from local fishing guides on the upper Madison River suggests that the growth in the angler population is slowing (J. Seckinger, personal communication). While the trajectory of the increases may change (i.e., shallower slope or leveling off asymptotically in the model), the popularity of the Madison River as a global fishing destination is clear (Skaar 2022). Although temporary reductions in angling pressure have been observed in recent years (i.e., the 2020 COVID-19 global pandemic and poor air quality from drifting smoke from regional wildfires in 2021), the upward trajectory in angling pressure on the upper Madison River is likely to continue (Skaar 2022). Creel surveys focused on angler demographics and opinions reflect an increasing proportion of anglers expressing concern about the unacceptability of the numbers of people and boats on the Madison River (Skaar 2022). Regardless, the upper Madison River continues to draw global tourists (Skaar 2022) and as the

population of cities in and surrounding SW Montana continues to grow, likely so will the population of anglers on the upper Madison River. Continued increases in angling pressure will lead to increases in catch-and-release mortality. If predictions of Brown and Rainbow Trout  $\hat{N}$  from MLE and subsequent models hold, the lower boundary of the 95% prediction interval of  $\hat{N}$  reaches 0 in 2025 at Pine Butte and Varney for Brown Trout and 2025 for Rainbow Trout at Pine Butte. Given that each long-term sampling site is an open population, Brown and Rainbow Trout are not likely to be locally extirpated. However, the negative feedback loop of decreasing  $\hat{N}$  coupled with increased angling pressure is expected to lead to increased recycling rates which will lead to increased stress in Brown and Rainbow Trout in the upper Madison River which may lead to additional declines in reproductive success, leading to a continued decrease in  $\hat{N}$ . One notable caveat in the predictions of angler pressure and catch I have made is that I have assumed that the seed metrics I used (i.e., season length, fishing day length, mean trip length, and mean catch rate) were not likely to change over the course of the predictions. It is possible that one or more of these metrics could change in the future. One potential scenario is to assume that as Brown Trout populations decline while angler pressure increases, catch rate will decline concomitantly. Similarly, MTFWP may enact changes to fishing day length that limit the amount of time anglers can spend on the water. For example, MTFWP already limits fishing on temperature and drought-impacted streams with periodic closures (colloquially known as “hoot owl” restrictions, prohibiting angling between the hours of 1400 and 0000 the following morning).

478 Fisheries managers have several tools at their disposal to manage fisheries,  
479 including supplemental stocking to increase recruitment to the recreationally-angled  
480 population, supplemental feeding of exogenous resources to improve growth, and  
481 requiring “hookless fishing” in which anglers are required to use hookless flies,  
482 simulating the “thrill of the catch” while hooking mortality becomes effectively zero. In a  
483 large, economically-important river like the upper Madison River, developing options  
484 that are easier to implement is warranted, ideally implementing those that are simple  
485 and may impact one or more of the fisheries dynamic rate functions. Fisheries  
486 managers cannot control much related to water temperature and volume (only the latter  
487 of which can be influenced by managers on impounded rivers and even then, only to a  
488 certain extent), but changes to the angling population could be applied that may  
489 improve the fishery. Because the deleterious impacts of increased angling pressure on  
490 Brown and Rainbow Trout appear to be predominantly associated with handling stress  
491 (and subsequent hooking mortality), reducing fisheries mortality by limiting the number  
492 of anglers on the river is one simple option because reduction in the number of anglers  
493 on the river would reduce catch-and-release mortality. However, this option is likely to  
494 receive significant negative feedback from the angling public and outfitter/fishing guide  
495 community reliant on the river for income. Requiring the use of barbless hooks is an  
496 implementable solution that may reduce hooking mortality but also allow the angler  
497 population to grow. Whether or not the use of barbless hooks directly affects mortality is  
498 debatable (i.e., Thompson 1946; Hunsaker et al. 1970; Titus and Vanicek 1988; Turek  
499 and Brett 1997). More likely, barbless hooks reduce the average time from landing to  
500 release (Schaeffer and Hoffman 2002) thus reducing fish stressors. Coupled with using

501   barbless hooks, requiring all anglers to use a landing net while fishing the upper  
502   Madison River would eliminate the need for anglers to be handling fish at all and may  
503   reduce immunosuppression in landed fish.

504           Identifying the exact cause of the changes in weight at length of Brown and  
505   Rainbow Trout is beyond the scope of this analysis as is identifying the cause of  
506   declines of Brown Trout and Rainbow Trout and Pine Butte; however, given the  
507   increased angling pressure on the upper Madison River, it stands to reason that  
508   changes in water temperature regimes, increased recycling rate, and handling stress  
509   could be a starting point for future investigations. Indeed, environmental (e.g., water  
510   temperature) and mechanical (e.g., capture) stressors may lead to individuals making  
511   trade-offs between reproductive efforts, somatic growth, and survival (Schreck et al.  
512   2001) and decreased weights at nearly all fish lengths in all quantiles when compared to  
513   the reference population for Brown Trout may be reflective of these tradeoffs. Potential  
514   areas to evaluate include changes in river thermal regimes, anglers targeting larger-  
515   sized fish, and fishing during spawning season that potentially contribute to additional  
516   increased mortality of larger fish.

517           Managing fisheries that are recreationally and economically important can be  
518   challenging. Balancing the desires of anglers, fishing guides and outfitters with the need  
519   to manage the natural resource requires walking a fine line. I encourage fisheries  
520   managers to review angler satisfaction survey results related to their fisheries. If a  
521   recreational or commercially-important fishery has no data related to angler satisfaction,  
522   I encourage managers to establish regular, valid creel surveys to sample angler  
523   demographics, satisfaction, and other key metrics. A global destination for trout anglers,



fisheries managers on the upper Madison River have been in “uncharted waters” since 2018 with regards to catch and expected mortality (Horton et al. 2018). If there is evidence that a globally-recognized recreationally and economically-important fishery may be suffering a quality decline, including decreases in angler satisfaction in the size of fish they have caught over time (Horton et al. 2018), evidence of declines in estimates of weight-at-length and abundance (this paper), complaints of angler overcrowding (Horton et al. 2018; Skaar 2022), and evidence of angling mortality that exceeds the estimates of what a fishery can sustain (Horton et al. 2018), fishery managers should leverage the angler feedback they receive and look for supporting evidence in extant data sets. If evidence exists to support angler claims, identifying the fewest number of new regulations that have the largest positive impact on the fishery may be challenging but software exists for simulating management changes [e.g., Fisheries Analysis and Modeling Simulator (available through the American Fisheries Society), the R packages Fisheries Library in R (FLR), the fishmethods: Fishery Science Methods and Models package, etc.]. Still more challenging may be educating the angling public on why new regulations are required. Managing angler expectations will be important.

#### Acknowledgements

Thanks to M. Duncan and D. Schmetterling (MTFWP) for answering a number of my early questions about FWP reports and for help retrieving data related to long-term sampling on the upper Madison River, M. Wuellner (University of Nebraska - Kearney) whose review of an early draft of this manuscript provided significant improvement, and

547 X anonymous reviewers whose comments improved this manuscript. There is no  
548 conflict of interest declared in this article.

549

550 <A>Data Availability Statement

551 A repository that includes supplementary data and R code used in this  
552 manuscript is available at [https://github.com/stevenranney/madison\\_river\\_fisheries](https://github.com/stevenranney/madison_river_fisheries).

553

554 <A>Ethics Statement

555 There were no ethical guidelines applicable to this study.

## References

- Anderson, C. S. (1995). Measuring and correcting for size selection in electrofishing mark-recapture experiments. *Transactions of the American Fisheries Society*, 124, 663-676. [https://doi.org/10.1577/1548-8659\(1995\)124%3C0663:MACFSS%3E2.3.CO;2](https://doi.org/10.1577/1548-8659(1995)124%3C0663:MACFSS%3E2.3.CO;2)
- Anderson, R. O., & Neumann, R. M. (1996). Length, weight, and associated structural indices. In B. R. Murphy and D. W. Willis (Eds.), *Fisheries Techniques* (2nd ed., pp 447–482). American Fisheries Society.
- Barton, B. A., & Iwama, G. K.. (1991). Physiological changes in fish from stress in aquaculture with emphasis on the response and effects of corticosteroids. *Annual Review of Fish Diseases*, 1, 3-26. [https://doi.org/10.1016/0959-8030\(91\)90019-G](https://doi.org/10.1016/0959-8030(91)90019-G)
- Boyd, J. W., Guy, C. S., Horton, T. B., & Leathe, S. A. (2010). Effects of catch-and-release angling on salmonids at elevated water temperatures. *North American Journal of Fisheries Management*, 30, 898-907. <https://doi.org/10.1577/M09-107.1>
- Cade, B. S., & Gilham, A. T. (2024). Summarizing quantile estimates of weight at length to compare condition: Geographic and temporal variation in Arctic Grayling. *Transactions of the American Fisheries Society*, 00, 1-21. <https://doi.org/10.1002/tafs.10466>
- Cade, B. S., Terrell, J. W., & Neely, B. C. (2011). Estimating geographic variation in allometric growth and body condition of Blue Suckers with quantile regression. *Transactions of the American Fisheries Society*, 140, 1657-1669. <https://doi.org/10.1080/00028487.2011.641885>

- 578 Cade, B. S., Terrell, J. W., & Porath, M. T. (2008). Estimating fish body condition with  
579 quantile regression. *North American Journal of Fisheries Management*, 28, 349-359.  
580 <http://dx.doi.org/10.1577/M07-048.1>
- 581 Campbell, P. M., Pottinger, T. G., & Sumpter, J. P. (1992). Stress reduces the quality of  
582 gametes produced by Rainbow Trout. *Biology of Reproduction*, 47, 1140-1150.  
583 <https://doi.org/10.1095/biolreprod47.6.1140>
- 584 Crane, D. P., & Farrell, J. M. (2017). Trends in body condition of Smallmouth Bass and  
585 Northern Pike (1982–2013) following multiple ecological perturbations in the St.  
586 Lawrence River. *Canadian Journal of Fisheries and Aquatic Sciences*, 74, 1158–  
587 1172. <https://doi.org/10.1139/cjfas-2016-0160>
- 588 Crane, D. P., Farrell, J. M., Einhouse, D. W., Lantry, J. R., & Markham, J. L. (2015).  
589 Trends in body condition of native piscivores following invasion of Lakes Erie and  
590 Ontario by the Round Goby. *Freshwater Biology*, 60, 111–124.  
591 <https://doi.org/10.1111/fwb.12473>
- 592 Donaldson, E. M. (1981). The pituitary-interrenal axis as an indicator of stress in fish. In  
593 A. D. Pickering (Eds.), *Stress in Fish* (pp. 11-47). Academic Press, London.
- 594 Gabelhouse, D. W. Jr. (1984). A length-categorization system to assess fish stocks.  
595 *North American Journal of Fisheries Management*, 4, 273-285.  
596 [https://doi.org/10.1577/1548-8659\(1984\)4%3C273:ALSTAF%3E2.0.CO;2](https://doi.org/10.1577/1548-8659(1984)4%3C273:ALSTAF%3E2.0.CO;2)
- 597 Hayes, D. B., Bence, J. R., Kwak, T. J., & Thompson, B. E. (2007). Abundance,  
598 biomass, and production. In C. S. Guy and M. L. Brown (Eds), *Analysis and*  
599 *Interpretation of Freshwater Fisheries Data* (pp. 327-374). American Fisheries  
600 Society. <https://doi.org/10.47886/9781888569773>

- 601 Horton, T., Moser, D., Morris, C., Puls, A., Lohrenz, T., & Lewis, M. (2018). *Angler*  
602 *satisfaction, demographic, and creel surveys—Upper Madison River, 2015-2017.*  
603 *Report to Northwestern Energy, Helena.* Montana Fish, Wildlife & Parks.
- 604 Hunsaker, J., Marnell, L. F., & Sharpe, F. P. (1970). Hooking mortality of Yellowstone  
605 cutthroat trout. *Progressive Fish-Culturist*, 32, 231-235. [https://doi.org/10.1577/1548-](https://doi.org/10.1577/1548-8640(1970)32[231:HMOYCT]2.0.CO;2)  
606 [8640\(1970\)32\[231:HMOYCT\]2.0.CO;2](https://doi.org/10.1577/1548-8640(1970)32[231:HMOYCT]2.0.CO;2)
- 607 Jobling, M. (1995). *Environmental Biology of Fishes*. Chapman & Hall, London.
- 608 Jones, C. M., & Pollock, K. H. (2012). Recreational survey methods: estimation of effort,  
609 harvest, and released catch. In A. V. Zale, D. L. Parrish, and T. M. Sutton (Eds.),  
610 *Fisheries Techniques (3rd ed., pp 883-919)*. American Fisheries Society.  
611 <https://doi.org/10.47886/9781934874295>
- 612 Jones, C. M., & Robson, D. (1991). Improving precision in angler surveys: traditional  
613 access design versus bus route design. *American Fisheries Society Symposium*  
614 *12:177-188.*
- 615 Jones, T. S., Jones, M. I., Treml, M., & Heinrich, T. (2022). Estimating recycling of fish  
616 in catch-and-release fisheries. *Fisheries*, 47, 529-536.  
617 <https://doi.org/10.1002/fsh.10824>
- 618 Koenker, R. (2021). quantreg: Quantile Regression. R package version 5.85.  
619 <https://CRAN.R-project.org/package=quantreg>
- 620 League, C., & Caball, B. (2022). *Montana statewide angling pressure 2020, Helena.*  
621 *Montana Fish, Wildlife & Parks.*
- 622 Link, W. A., & Barker, R. J. (2010). *Bayesian Inference with Ecological Applications.*  
623 *Elsevier. London.*

- 624 Lohrenz, T., Duncan, M., Dukovcic, J., McCormak, M., & Jaeger, M. (2023). *Madison*  
625 *River Drainage 2188 Project Monitoring Report 2022, Helena*. Northwestern Energy,  
626 Environmental Division.
- 627 Malvestuto, S. P. (1996). Sampling the Recreational Creel. In B. R. Murphy and D. W.  
628 Willis (Eds.), *Fisheries Techniques* (2nd ed., pp. 591-623). American Fisheries  
629 Society.
- 630 McConnachie, S. H., Cook, K. V., Patterson, D. A., Gilmour, K. M., Hinch, S. G., Farrell,  
631 A. P., & Cooke, S. J. (2012). Consequences of acute stress and cortisol  
632 manipulation on the physiology, behavior, and reproductive outcome of female  
633 Pacific salmon on spawning grounds. *Hormones and Behavior*, 62, 67-76.
- 634 McCormick, J. L., & Meyer, K. A. (2017). Sample size estimation for on-site creel  
635 surveys. *North American Journal of Fisheries Management*, 37, 970-980.  
636 <https://doi.org/10.1080/02755947.2017.1342723>
- 637 McCormick, J. L., Quist, M. C., & Schill, D. J. (2013). Self-reporting bias in chinook  
638 salmon sport fisheries in Idaho: implications for roving creel surveys. *North American*  
639 *Journal of Fisheries Management*, 33, 723-731.  
640 <https://doi.org/10.1080/02755947.2013.808293>
- 641 Milewski, C. L., & Brown, M. L. (1994). Proposed standard weight ( $W_s$ ) equation and  
642 length category standards for stream-dwelling brown trout. *Journal of Freshwater*  
643 *Ecology*, 9, 111-116. <https://doi.org/10.1080/02705060.1994.9664437>
- 644 Montana Fish, Wildlife & Parks. (2005). *Annual FWP drought summary, Helena*.  
645 Montana Fish, Wildlife & Parks.

- 646 Montana Fish, Wildlife & Parks. (2017). *Montana statewide angling pressure 2017*  
647 *summary report, Helena*. Montana Fish, Wildlife & Parks.
- 648 Montana Fish, Wildlife & Parks. (2019). *Montana statewide angling pressure 2019*  
649 *summary report, Helena*. Montana Fish, Wildlife & Parks.
- 650 Montana Fish, Wildlife & Parks. (2023). *FishMT survey and inventory data, Madison*  
651 *River, Helena*. Accessed 18 October 2023. Available:  
652 [myfwp.mt.gov/fishMT/reports/surveyreport](https://myfwp.mt.gov/fishMT/reports/surveyreport)
- 653 Montana Fish, Wildlife & Parks. (2023). *2023 Fishing: Montana Fishing Regulations,*  
654 *Helena*. Montana Fish, Wildlife & Parks.
- 655 National Park Service. (2023). *The Madison River near West Yellowstone, Montana.*  
656 National Park Service. Accessed 2023-11-05. Available  
657 <https://www.nps.gov/articles/000/madison-river-west-yellowstone.htm>
- 658 Neely, B. C., Lynott, S. T., & Koch, J. D. (2021). Comparison of growth, body weight,  
659 and contribution to recreational creel of Channel Catfish and Channel Catfish x Blue  
660 Catfish hybrids in two public put-grow-take fisheries in Kansas. *North American*  
661 *Journal of Fisheries management*, 41(S1), S314-S322.  
662 <https://doi.org/10.1002/nafm.10469>
- 663 Neumann, R.M., Guy, C. S., & Willis, D. W. (2012). Length, weight, and associated  
664 indices. In A. V. Zale, D. L. Parrish, and T. M. Sutton (Eds.), *Fisheries techniques*  
665 (3rd ed., pp 637–669). American Fisheries Society.  
666 <https://doi.org/10.47886/9781934874295>

- 667 Pollock, K. H., Jones, C. M., & Brown, T. L. (1994). Angler survey methods and their  
668 applications in fisheries management. American Fisheries Society, Special  
669 Publication 25, Bethesda, Maryland. <https://doi.org/10.1007/BF00043009>
- 670 R Core Team. 2023. R: A language and environment for statistical computing. R  
671 Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
- 672 Ranney, S. H., Fincel, M. J., Wuellner, M. R., VanDeHey, J. A., & Brown, M. L. (2008).  
673 Assessing length-related bias and the need for data standardization in the  
674 development of standard weight equations. *North American Journal of Fisheries*  
675 *Management*, 30, 655-664. <https://doi.org/10.1577/M08-097.1>
- 676 Ranney, S. H. (2018a). AnglerCreelSurveySimulation: Simulate an angler creel survey.  
677 Version 1.0.0. <https://CRAN.R-project.org/package=AnglerCreelSurveySimulation>
- 678 Ranney, S. H. (2018b). Quantile regression estimates of body weight at length in  
679 Walleye. *North American Journal of Fisheries Management*, 38, 769-776.  
680 <https://doi.org/10.1002/nafm.10070>
- 681 Réalis-Doyelle, E., Gisbert, E., Alcaraz, C., Teletchea, F., & Pasquet, A. (2018).  
682 Temperature affects growth allometry and development patterns in brown trout  
683 (*Salmo trutta*) fry: a multitrait approach. *Canadian Journal of Fisheries and Aquatic*  
684 *Sciences*, 75, 714-722. <https://doi.org/10.1139/cjfas-2017-0037>
- 685 Roulson, L. H. (2002). *An examination of 1995 and 1996 creel survey data for the*  
686 *Madison River, Montana and evidence of possible effects due to whirling disease,*  
687 *Helena*. Montana Fish, Wildlife & Parks.
- 688 Schaeffer, J. S., & Hoffman, E. M. (2002). Performance of barbed and barbless hooks in  
689 a marine recreational fishery. *North American Journal of Fisheries Management*, 22,



229-235. <https://doi.org/10.1577/1548->

8675(2002)022%3C0229:POBABH%3E2.0.CO;2

Schill, D. J., Griffith, J. S., & Gresswell, R. E. (1986). Hooking mortality of cutthroat trout in a catch-and-release segment of the Yellowstone River, Yellowstone National Park. *North American Journal of Fisheries Management*, 6, 226-232.

[https://doi.org/10.1577/1548-8659\(1986\)6%3C226:HMOCTI%3E2.0.CO;2](https://doi.org/10.1577/1548-8659(1986)6%3C226:HMOCTI%3E2.0.CO;2)

Schreck, C. B., Contreras-Sanchz, W., & Fitzpatrick, M. S. (2001). Effects of stress on fish reproduction, gamete quality, and progeny. *Aquaculture*, 197, 3-24.

[https://doi.org/10.1016/S0044-8486\(01\)00580-4](https://doi.org/10.1016/S0044-8486(01)00580-4)

Simpkins, D. G., & Hubert, W. A. (1996). Proposed revision of the standard weight equation for rainbow trout. *Journal of Freshwater Ecology*, 11, 319-326.

<https://doi.org/10.1080/02705060.1996.9664454>

Skaar, D. (2022). *Summary of the 2021 recreational use survey for the Madison River, Helena*. Montana Fish, Wildlife & Parks.

Thompson, F. (1946). Experiment proves small fish are worth saving. New Mexico Department of Game and Fish, Technical report 11-F.

Titus, R. G., & Vanicek, C. D. (1988) Comparative hooking mortality of lure-caught Lahontan Cutthroat Trout at Heenan Lake, California. *California Fish and Game*, 74, 218-255.

Turek, S. M., & Brett, M. T. (1997). Comment: Trout mortality from baited barbed and barbless hooks. *North American Journal of Fisheries Management*, 17, 807.

<https://doi.org/10.1577/1548-8675-17.3.807>

Vincent, E. R. (1978). *Southwestern Montana Fisheries Investigations: Madison River temperature study, Helena*. Montana Department of Fish and Game.

Wootton, R. J. (1996). *Ecology of Teleost Fishes*. Chapman & Hall, London.

For Peer Review Only

## &lt;A&gt;Tables

Table 1. Estimates of intercept and slope from bootstrapped replicates of quantile regression with  $q = 0.75$  for 20 years' worth of data—every third year—from long-term standardized sampling of Brown and Rainbow Trout from the upper Madison River. Bootstrapped replicates were estimated by resampling with replacement 5,000  $\log_{10}W$  and  $\log_{10}TL$  pairs 1,000 times within each year:species combination.

Species	Year	Intercept ( $\beta_0$ )				Slope ( $\beta_1$ )		
		2.50%	Estimate	97.50%		2.50%	Estimate	97.50%
Brown Trout	Reference	-4.6859	-4.6404	-4.595		2.8634	2.8818	2.9002
	2004	-4.5496	-4.4818	-4.4139		2.8038	2.8305	2.8572
	2007	-4.9495	-4.9021	-4.8547		2.9661	2.9851	3.0041
	2010	-5.0628	-5.0185	-4.9742		3.0076	3.0254	3.0431
	2013	-5.2533	-5.212	-5.1708		3.0790	3.0955	3.1119
	2016	-4.8858	-4.7794	-4.6729		2.8905	2.9319	2.9732
	2019	-4.9808	-4.9355	-4.8902		2.9765	2.9946	3.0127
	2022	-5.1674	-5.1231	-5.0787		3.0459	3.0638	3.0817
Rainbow Trout	Reference	-4.489	-4.43	-4.371		2.7717	2.7955	2.8193
	2004	-4.2363	-4.1444	-4.0525		2.6568	2.6933	2.7299
	2007	-4.7254	-4.6689	-4.6124		2.8662	2.8893	2.9124
	2010	-4.7478	-4.701	-4.6543		2.8714	2.8908	2.9102
	2013	-5.1614	-5.1083	-5.0552		3.0329	3.0545	3.076
	2016	-4.9109	-4.8093	-4.7076		2.8974	2.9373	2.9771
	2019	-4.7391	-4.688	-4.6368		2.8666	2.8876	2.9085
	2022	-4.8855	-4.8223	-4.7591		2.9173	2.9427	2.9681

Table 2. Differences and 95% CIs around the differences in intercepts and slope for each species:year population from the reference (2003) population. The *P*-value is from a test of the null hypothesis that the intercept and slope from the quantile regression, where  $q = 0.75$  for each species:year population, is equal to the reference population intercept and slope. The intercepts for all years for both species are significantly lower than each species' reference (2003) population intercept, except for 2004 in which the intercept is significantly higher than 2003. Slopes for both species for all years were higher than 2003, except for 2004, in which the slope was significantly lower for 2004.

Species	Year	Intercept ( $\beta_0$ )				Slope ( $\beta_1$ )			
		2.50%	Estimate	97.50%	p-value	2.50%	Estimate	97.50%	p-value
Brown Trout	2004	0.0796	0.1587	0.2378	<0.0001	-0.0828	-0.0513	-0.0199	0.0014
	2007	-0.3265	-0.2617	-0.1969	<0.0001	0.0772	0.1033	0.1295	<0.0001
	2010	-0.4381	-0.3781	-0.3181	<0.0001	0.1193	0.1435	0.1677	<0.0001
	2013	-0.6315	-0.5716	-0.5116	<0.0001	0.1895	0.2136	0.2377	<0.0001
	2016	-0.2518	-0.1389	-0.026	0.0159	0.0058	0.0500	0.0943	0.0267
	2019	-0.3558	-0.2950	-0.2342	<0.0001	0.0883	0.1127	0.1372	<0.0001
	2022	-0.5432	-0.4826	-0.422	<0.0001	0.1574	0.182	0.2065	<0.0001
Rainbow Trout	2004	0.1762	0.2856	0.3951	<0.0001	-0.1458	-0.1022	-0.0585	<0.0001
	2007	-0.3217	-0.2389	-0.156	<0.0001	0.0603	0.0938	0.1274	<0.0001
	2010	-0.3419	-0.2710	-0.2002	<0.0001	0.0664	0.0953	0.1242	<0.0001
	2013	-0.7568	-0.6783	-0.5997	<0.0001	0.2272	0.2590	0.2907	<0.0001
	2016	-0.4968	-0.3792	-0.2616	<0.0001	0.0952	0.1418	0.1883	<0.0001
	2019	-0.3338	-0.2579	-0.182	<0.0001	0.0612	0.0921	0.1229	<0.0001
	2022	-0.4786	-0.3922	-0.3059	<0.0001	0.1124	0.1472	0.182	<0.0001

Table 3. Annual predictions and 95% prediction interval estimates of catch and mortality of Brown and Rainbow Trout from a linear model fit to observed data from 2011, 2013, 2015, and 2016. The model was validated with “hold out” data from 2017, 2019, and 2020.

Species	Year	Predicted catch				Predicted mortality		
		2.50%	Fit	97.50%		2.50%	Fit	97.50%
Brown Trout	2023	166,991	235,195	304,670		3,340	4,704	6,093
	2024	174,470	249,459	320,142		3,489	4,989	6,403
	2025	181,405	260,334	342,976		3,628	5,207	6,860
	2026	189,290	275,611	359,286		3,786	5,512	7,186
	2027	194,709	289,967	380,429		3,894	5,799	7,609
	2028	203,201	300,709	401,463		4,064	6,014	8,029
	2029	208,487	314,453	421,155		4,170	6,289	8,423
	2030	216,089	328,666	441,977		4,322	6,573	8,840
Rainbow Trout	2023	282,098	395,059	510,461		22,568	31,605	40,837
	2024	292,742	415,778	541,699		23,419	33,262	43,336
	2025	304,842	438,442	575,719		24,387	35,075	46,058
	2026	315,674	463,034	609,820		25,254	37,043	48,786
	2027	328,031	485,093	641,658		26,242	38,807	51,333
	2028	342,188	507,683	676,942		27,375	40,615	54,155
	2029	352,244	527,682	709,354		28,179	42,215	56,748
	2030	364,356	552,697	740,324		29,149	44,216	59,226

<A>Figures




Figure 1. The Madison River from the dam at Hebgen Lake downstream through Ennis Lake is considered the upper Madison River. Long-term monitoring locations at Varney and Pine Butte are marked and bounded in red (MTFWP 2023).





Figure 2. Quantile regression ( $\square = 0.05$  up to 0.95 by increments of 0.05) prediction and 95% CIs of weight for the midpoint (mm) of each length category [substock (SS), stock-quality (S-Q), quality-preferred (Q-P), preferred-memorable (P-M), memorable-trophy (M-T); Gabelhouse 1984] for the reference data set (Year = 2003) and seven

subsequent years for Brown Trout (A) and Rainbow Trout (B) populations in the upper  
Madison River, Montana.

For Peer Review Only






Figure 3. Linear regression (solid line) of population estimates of Brown Trout (left column) and Rainbow Trout (right column) and constructed 95% confidence intervals (dashed lines) per river kilometer at two long-term monitoring locations in the upper Madison River, Pine Butte (top row) and Varney (bottom row). The overall model fit for Brown Trout at Pine Butte and Varney is significant ( $p = 0.047$  and  $p = 0.014$ , respectively) suggesting that over time, there have been fewer Brown Trout in the upper Madison River. The overall model fits for Rainbow Trout at both sampling locations are not significant at  $\alpha = 0.05$  (Pine Butte:  $p = 0.15$ ; Varney:  $p = 0.055$ ) and Rainbow Trout populations appear to be increasing at Varney.




Figure 4. The observed (filled circles) number of anglers/year on the upper Madison River, Montana. A linear model and 95% CI (solid line and surrounding shaded area) was fitted to the number of anglers/year for years  $\leq 2016$  which was subsequently used to predict (dashed line) the number of anglers/year for 2017 through 2030. Observed data for years 2017, 2019, and 2020 (filled circles) were used to validate the prediction and 95% prediction interval (shaded area around dashed line). The linear model for the number of anglers was significant (adjusted  $R^2 = 0.96$ ;  $\beta_0 = -33991356$ ;  $\beta_1 = 16949$ ;  $p = 0.0134$ ). Data point for 2016 is an estimate from a correlation analysis between reported outfitted trips and total angling pressure (Horton et al. 2018).