

¹ **A geophysical model for the origin of vent clusters in**
² **Colorado Plateau volcanic fields**

F. Deng,¹C. B. Connor,¹ R. Malservisi¹, L. J. Connor¹, J. W. White², A.

Germa¹

Corresponding author: Fanghui Deng, School of Geosciences, University of South Florida,
Tampa, FL, USA. (fanghuideng@mail.usf.edu)

¹University of South Florida, Tampa,
Florida, USA

²U.S. Geological Survey, Austin, Texas,
USA

Abstract. (Chuck)

Distributed volcanic fields are characterized by tens to hundreds of individual vents scattered over broad areas, with each vent active for months to decades, but cumulative activity in entire volcanic fields persisting for up to millions of years. Volcano vent clusters have been identified in numerous distributed volcanic fields globally, but the origin of these vent clusters remains uncertain. We show with new gravity data and numerical modeling that vent clusters in the Quaternary Springerville volcanic field (SVF), Arizona (USA), correlate with gradients in the gravity field. Inverse modeling 299 new gravity stations in and around the SVF using singular value decomposition with Tikhonov regularization indicates that gravity anomalies are explained by density discontinuities that transect nearly the entire crust. These discontinuities are interpreted to be caused by boundaries in the North American crust accreted during the Proterozoic. Vent density is low in regions of high density Proterozoic crust, and high in areas of relatively low density Proterozoic crust. Vent density is highest within the SVF adjacent to crustal boundaries and long vent alignments parallel boundaries. 2D and 3D numerical models of magma ascent are developed to simulate long term average magma migration leading to the development of vent clusters in the SVF, assuming viscous flow of a fluid in a porous media (Darcys Law) is statistically equivalent to full field scale magma migration averaged over geological time through the crust. The location and flux from the magma source region are boundary conditions of the model. Changes in the model conductivity, associated

26 with changes in the bulk properties of the lithosphere, can simulate prefer-
27 ential magma migration paths and alter the estimated magma flux at the
28 surface. Using this model, we find variation in vent density, and occurrence
29 of vent clusters, are explained by changes in conductivity associated with the
30 Proterozoic crust. The implication is that in some distributed volcanic fields
31 large-scale crustal structures, such as inherited tectonic block boundaries,
32 influence magma ascent and the clustering of volcanic vents. Probabilistic
33 models of volcanic hazard for distributed volcanic fields can be improved by
34 identifying crustal structures and assessing their impact on volcano distri-
35 bution with the use of numerical models.

1. Introduction

Distributed volcanic fields are remarkable features, found in a variety of tectonic settings on Earth and nearby planets, with individual fields comprising tens to hundreds of volcanoes scattered across thousands of square kilometers [Williams, 1950; Nakamura, 1977; Hasenaka and Carmichael, 1985; Addington, 2001; Richardson et al., 2013]. Often volcanoes within these fields are thought to be monogenetic, with each volcano, or alignment of nearby volcanoes, representing a single, relatively short-lived magmatic event, such as intrusion and eruption of a dike swarm [Rittmann, 1962]. On Earth, these volcanic fields are predominantly basaltic in composition, although many are bimodal [Bacon, 1982; Mazzarini et al., 2004]. Most volcanoes within these fields are scoria cones, small shields, or lava domes [Valentine and Connor, 2015; Kereszturi and Németh, 2016].

Distribution of vents within volcanic fields is analyzed to delineate trends in volcanic activity, such as migration of the field with lithospheric plate motion [Tanaka et al., 1986; Condit et al., 1989], to better understand the relationship of volcanoes to prominent tectonic boundaries or faults [Conway et al., 1997; Heming, 1980; van den Hove et al., 2017] and to better assess the likely locations of future eruptions [Connor et al., 2012; Cappello et al., 2012]. Statistical analyses of vent distribution have shown that volcanoes cluster within many distributed volcanic fields, rather than being randomly or regularly distributed [Le Corvec et al., 2013]. For example, vent clusters are found in the subduction-zone boundary Michoacán-Guanajuato volcanic field, Mexico [Connor, 1990], the Springerville and San Francisco volcanic fields on the margin of the Colorado plateau (USA) [Condit and Connor, 1996; Conway et al., 1998], the rift-hosted Eifel volcanic field,

Germany [*Schmincke et al.*, 1983; *Jaquet and Carniel*, 2006], or further from active plate boundaries [*Wei et al.*, 2003; *Cas et al.*, 2016; *van den Hove et al.*, 2017].

The origin of vent clusters within distributed volcanic fields remains uncertain. One model is that magma source regions are heterogenous, with some areas of the mantle more prone to partial melting than others leading to more frequent and voluminous activity in some parts of the field compared to others. Alternatively, the crust may act as a filter. Density discontinuities and rigidity contrasts in one part of the crust may tend to enhance sill formation and arrest dike ascent. Structures such as folds [*Wetmore et al.*, 2009] and faults [*van den Hove et al.*, 2017] may alter magma ascent pathways.

In this paper we explore the role of ancient discontinuities in the crust beneath the Springerville volcanic field (SVF), Arizona (USA), in changing patterns of volcanic activity. We accomplish this by re-examining the distribution of volcanic vents in the SVF and by comparing vent distribution to gravity anomalies that we have mapped across and around the field that are likely caused by lateral discontinuities in crustal density, interpreted to have arisen during the accretion of the North American continent approximately 1.5 Ga [*Gilbert et al.*, 2007].

Since it is practical to model the statistical distribution of vents mapped in the field as a continuous density function using kernel density estimation [*Connor and Hill*, 1995; *Connor and Connor*, 2009; *Germa et al.*, 2013], we compare the vent density distribution with a model of bulk magma transport. Bulk magma transport is approximated using the advection-diffusion equation, with ascent from a uniform magma source region, and with heterogeneity within the crust that alters flow paths and gives rise to variations in

magma flux at the surface. A 3D gravity inversion [White *et al.*, 2015] is used to delineate the most prominent lateral changes in crustal density.

We find that using discontinuities derived from the gravity model, the bulk magma transport model gives rise to the major features of vent distribution observed in the SVF. That is, vent clusters, vent alignments, and overall changes in Quaternary vent distribution in the field are explained by the occurrence of ancient crustal discontinuities.

In the following we briefly review the volcanology of the SVF, develop a statistical model of vent distribution, discuss and model the gravity data, and develop a simple model to illustrate the expected magma budget at the surface, given the model of the crust derived from gravity data. We suggest that the data and models we present show that patterns of Quaternary volcanic activity in Colorado Plateau boundary fields is directly impacted by the structure of the crust developed during the Proterozoic.

2. SVF background

(Chuck, Aurelie)

The SVF sits at the southern margin of the Colorado Plateau, where the plateau transitions to the Basin and Range in an area known as the Arizona Transition Zone. Proterozoic crust is exposed along the Mogollon Rim in this transition zone and indicates broad regional changes in Proterozoic crust from primarily greenstones, an assemblage of metavolcanic rocks and pelagic sediments that represent island arcs accreted onto the proto-continent, and Proterozoic sedimentary rocks (Gilbert 2007). Gravity and magnetic maps and structural mapping of the region show that these broad lithologic transitions in the Proterozoic crust are preserved today, primarily creating ENE–WSW gravity and magnetic anomalies which parallel shear zones (faults) mapped at the surface

which have been intermittently activated, for example in Laramide orogeny [*Shoemaker et al.*, 1978; *Seeley and Keller*, 2003]. These investigations have also identified WNW-ESE trending structures in the basement, interpreted to be associated with Proterozoic extension [*Seeley and Keller*, 2003]. Regional gravity anomalies show that both WNW–ENE and ENE–WSW gravity gradients occur in the area of the SVF, suggesting that these Proterozoic boundaries extend through the field.

The Proterozoic boundaries are largely masked at the surface in the SVF area by Paleozoic sedimentary rocks. *Crumpler et al.* [1994], however, mapped monoclinical flexures and faults in SVF that have predominantly WNW–ENE orientations, suggesting that Proterozoic boundaries localized later deformation in the SVF. Quaternary volcanic activity in the SVF is only the latest manifestation of volcanism. Miocene and Pliocene basalt lavas appear to be much more voluminous in the region than subsequent Quaternary volcanism. Lavas from the Mount Baldy shield volcano, located SW of the SVF have radiometric age determinations of 8.7 ± 0.2 Ma and 9.0 ± 0.2 Ma [*Condit and Shafiqullah*, 1985; ?]. Much of the Quaternary SVF erupted onto an older lava flow surface, consisting of hawaiites ranging in age from 7.6 ± 0.4 Ma to 2.9 ± 0.1 Ma [*Condit and Connor*, 1996], and voluminous tholeiites dated at 5.3 ± 0.1 Ma [*Cooper and Hart*, 1990]. Numerous radiometric age determinations, stratigraphic and paleomagnetic studies suggest that latest volcanic activity in the SVF occurred approximately 2.1 – 0.3 Ma [*Cooper and Hart*, 1990; ?; *Condit and Shafiqullah*, 1985; *Aubele et al.*, 1986], with tholeiites erupting early in this episode, and more alkaline basalts erupting later (alkaline-olivine basalt, hawaiiite, mugearite and benmoreite) [*Condit and Connor*, 1996]. Thus, the SVF region has experienced episodic

volcanism at least from approximately 9 Ma, with large chemical heterogeneity in the
basaltic magmas erupted.

A total of 409 Quaternary vents and associated lava flows have been mapped in the
SVF [*Condit et al.*, 1989], which have been grouped into at least 366 eruptive events, as
some eruptions resulted in construction of multiple vents and vent alignment [*Condit and*
Connor, 1996]. These vents form clusters; eruptive activity waxed and waned within
clusters at a much higher rate than in the field on average.

The overall pattern of vent distribution is characterized by a broad ENE–WSW band
of volcanoes, parallel to mapped flexured and inferred Proterozoic lithologic boundaries.
Vent alignments in the field tend to be oriented parallel to these boundaries as well –
ENE trending in the E part of the field, and WNW trending in the W part of the field
[*Connor et al.*, 1992].

Analysis of clinopyroxene-whole-rock pairs in SVF basalts to derive pressure and tem-
perature of crystallization indicates that magmas originate at a wide range of depths, up
to at least 60 km [*Putirka and Condit*, 2003]. Significantly, some relatively high K₂O and
K/Ti basalts appear to stagnate at depths of 0–12 and 23–30 km, within the Proterozoic
section. Significantly, no evidence of stagnation exists at the Moho [*Putirka and Condit*,
2003], suggesting that rheologic boundaries within the crust impact magma ascent in the
SVF. These features of the geochemistry of the SVF and relationship to spatio-temporal
trends in volcanism led us to consider the role of lateral changes in the Proterozoic
crust in magma ascent.

3. Collection and modeling of gravity data

(Fanghui, Laura, JW)

survey design and processing (Table of gravity data - processed) overlay with volcanic
vents on contour map of gravity data development of PEST++ model overlay of volcanic
vents on PEST++ gravity model

4. Finite Difference model

(Fanghui, Rocco) development of the 2D model development of the 3D model
overlay of expected flux through surface and spatial density map.

5. Discussion

(Fanghui, Chuck, Rocco, Aurelie) Comparison with SFVF, Zuni-Bandera, others?

6. Conclusions

(Fanghui)

References

- Addington, E. A. (2001), A stratigraphic study of small volcano clusters on Venus, *Icarus*,
149(1), 16–36.
- Aubele, J. C., L. S. Crumpler, and M. Shafiqullah (1986), K-Ar ages of late Ceno-
zoic rocks of the central and eastern parts of the Springerville volcanic field, Arizona,
Isochron/West, *46*, 3–5.
- Bacon, C. R. (1982), Time-predictable bimodal volcanism in the Coso Range, California,
Geology, *10*(2), 65–69.
- Cappello, A., M. Neri, V. Acocella, G. Gallo, A. Vicari, and C. Del Negro (2012), Spatial
vent opening probability map of Etna volcano (Sicily, Italy), *Bulletin of Volcanology*,
74(9), 2083–2094.

Cas, R. A. F., J. van Otterloo, T. N. Blaikie, and J. vanden Hove (2016), The dynamics of a very large intra-plate continental basaltic volcanic province, the Newer Volcanics Province, SE Australia, and implications for other provinces, *Geological Society, London, Special Publications*, 446, SP446–8.

Condit, C. D., and C. B. Connor (1996), Recurrence rate of basaltic volcanism in volcanic fields: An example from the Springerville Volcanic Field, AZ, USA, *Geological Society of America, Bulletin*, 108, 1225–1241.

Condit, C. D., and M. Shafiqullah (1985), K-Ar ages of late Cenozoic rocks of the western part of the Springerville volcanic field, east-central Arizona, *Isochron/West*, 44, 3–5.

Condit, C. D., L. S. Crumpler, J. C. Aubele, and W. E. Elston (1989), Patterns of volcanism along the southern margin of the Colorado Plateau: the Springerville field, *Journal of Geophysical Research: Solid Earth*, 94(B6), 7975–7986.

Connor, C. B. (1990), Cinder cone clustering in the TransMexican volcanic belt: implications for structural and petrologic models, *Journal of Geophysical Research*, 95(B12), 19395–19405.

Connor, C. B., and L. J. Connor (2009), Estimating spatial density with kernel methods, in *Volcanic and Tectonic Hazard Assessment for Nuclear Facilities*, edited by C. B. Connor, N. A. Chapman, and L. J. Connor, pp. 331–343, Cambridge University Press.

Connor, C. B., and B. E. Hill (1995), Three nonhomogeneous Poisson models for the probability of basaltic volcanism: Application to the Yucca Mountain region, *Journal of Geophysical Research*, 100(B6), 10107–10125.

Connor, C. B., C. D. Condit, L. S. Crumpler, and J. C. Aubele (1992), Evidence of regional structural controls on vent distribution, Springerville volcanic field, Arizona,

Journal of Geophysical Research, *97*, 12 349–12 259.

Connor, L. J., C. B. Connor, K. Meliksetian, and I. Savov (2012), Probabilistic approach to modeling lava flow inundation: a lava flow hazard assessment for a nuclear facility in Armenia, *Journal of Applied Volcanology*, *1*(3), doi:10.1186/2191-5040-1-3.

Conway, M. F., D. A. Ferrill, C. M. Hall, A. P. Morris, J. A. Stamatakis, C. B. Connor, A. N. Halliday, and C. Condit (1997), Timing of basaltic volcanism along the Mesa Butte fault zone in the San Francisco volcanic field, Arizona, from $^{40}\text{Ar}/^{39}\text{Ar}$ ages: Implications for longevity of cinder cone alignments, *Journal of Geophysical Research*, *102*, 815–824.

Conway, M. F., C. B. Connor, B. E. Hill, C. D. Condit, K. Mullaney, and C. M. Hall (1998), Recurrence rates of basaltic volcanism in the SP cluster, San Francisco volcanic field, Arizona, *Geology*, *26*, 655–658.

Cooper, J. L., and W. K. Hart (1990), Mantle sources in the Arizona transition zone and global mantle heterogeneity, *Geology*, *18*(11), 1146–1149.

Crumpler, L. S., J. C. Aubele, and C. D. Condit (1994), Volcanoes and neotectonic characteristics of the Springerville volcanic field, Arizona, *New Mexico Geological Society Guidebook*, pp. 147–164.

Germa, A., L. J. Connor, E. Canon-Tapia, and N. LeCorvec (2013), Spatial distribution of monogenetic eruptive vents from volcanic fields in central Baja California, Mexico, *Bulletin of Volcanology*, *75*(12), 1–14.

Gilbert, H., A. A. Velasco, and G. Zandt (2007), Preservation of Proterozoic terrane boundaries within the Colorado Plateau and implications for its tectonic evolution, *Earth and Planetary Science Letters*, *258*(1), 237–248.

- Hasenaka, T., and I. S. E. Carmichael (1985), The cinder cones of Michoacán–Guanajuato, central Mexico: their age, volume and distribution, and magma discharge rate, *Journal of Volcanology and Geothermal Research*, 25(1-2), 105–124.
- Heming, R. F. (1980), Patterns of Quaternary basaltic volcanism in the northern North Island, New Zealand, *New Zealand Journal of Geology and Geophysics*, 23(3), 335–344.
- Jaquet, O., and R. Carniel (2006), Estimation of volcanic hazards using geostatistical models, in *Statistics in Volcanology*, edited by H. Mader, S. Coles, C. B. Connor, and L. J. Connor, no. 1 in Special Publications of IAVCEI, pp. 89–103, Geological Society of London, London (UK).
- Kereszturi, G., and K. Németh (2016), Sedimentology, eruptive mechanism and facies architecture of basaltic scoria cones from the Auckland Volcanic Field (New Zealand), *Journal of Volcanology and Geothermal Research*, 324, 41–56.
- LeCorvec, N., K. B. Spörli, J. Rowland, and J. Lindsay (2013), Spatial distribution and alignments of volcanic centers: clues to the formation of monogenetic volcanic fields, *Earth Science Reviews*, 124, 96–114.
- Mazzarini, F., G. Corti, P. Manetti, and F. Innocenti (2004), Strain rate and bimodal volcanism in the continental rift: Debre Zeyt volcanic field, northern MER, Ethiopia, *Journal of African Earth Sciences*, 39(3), 415–420.
- Nakamura, K. (1977), Volcanoes as possible indicators of tectonic stress orientation principle and proposal, *Journal of Volcanology and Geothermal Research*, 2(1), 1–16.
- Putirka, K., and C. D. Condit (2003), Cross section of a magma conduit system at the margin of the Colorado Plateau, *Geology*, 31(8), 701–704.

- Richardson, J. A., J. E. Bleacher, and L. S. Glaze (2013), The volcanic history of syria
planum, mars, *Journal of Volcanology and Geothermal Research*, 252, 1–13.
- Rittmann, A. (1962), *Volcanoes and Their Activity*, 305 pp., John Wiley and Sons, New
York, USA.
- Schmincke, H.-U., V. Lorenz, and H. A. Seck (1983), The quaternary Eifel volcanic fields,
in *Plateau Uplift*, pp. 139–151, Springer, Berlin, Germany.
- Seeley, J. M., and G. R. Keller (2003), Delineation of subsurface Proterozoic Unkar and
Chuar Group sedimentary basins in northern Arizona using gravity and magnetics:
Implications for hydrocarbon source potential, *AAPG Bulletin*, 87(8), 1299–1321.
- Shoemaker, E. M., R. L. Squires, and M. J. Abrams (1978), Bright Angel and Mesa
Butte fault systems of northern Arizona, *Geological Society of America Memoirs*, 152,
341–367.
- Tanaka, K. L., E. M. Shoemaker, G. E. Ulrich, and E. W. Wolfe (1986), Migration of
volcanism in the San Francisco volcanic field, Arizona, *Geological Society of America
Bulletin*, 97(2), 129–141.
- Valentine, G. A., and C. B. Connor (2015), *Encyclopedia of Volcanology*, chap. 23: Basaltic
Volcanic Fields, pp. 423–440, Academic Press.
- van den Hove, J. C., J. van Otterloo, P. G. Betts, L. Ailleres, and R. A. F. Cas (2017),
Controls on volcanism at intraplate basaltic volcanic fields, *Earth and Planetary Science
Letters*, 459, 36–47.
- Wei, H., R. S. J. Sparks, R. Liu, Q. Fan, Y. Wang, H. Hong, H. Zhang, H. Chen, C. Jiang,
J. Dong, et al. (2003), Three active volcanoes in China and their hazards, *Journal of
Asian Earth Sciences*, 21(5), 515–526.

Wetmore, P. H., C. B. Connor, S. E. Kruse, S. Callihan, G. Pignotta, C. Stremtan,
and A. Burke (2009), Geometry of the Trachyte Mesa intrusion, Henry Mountains,
Utah: Implications for the emplacement of small melt volumes into the upper crust,
Geochemistry, Geophysics, Geosystems, 10, Q08,006, doi:10.1029/2009GC002,469.

White, J. T., A. Karakhanian, C. B. Connor, L. J. Connor, J. D. Hughes, R. Malservisi,
and P. H. Wetmore (2015), Coupling geophysical investigation with hydrothermal mod-
eling to constrain the enthalpy classification of a potential geothermal resource, *Journal*
of Volcanology and Geothermal Research, 298, 59–70.

Williams, H. (1950), *Volcanoes of the Parícutin Region, Mexico*, *USGS Professional Paper*,
vol. 965, U.S. Geological Survey, Reston, Va, USA.

Acknowledgments.

7. Figures

Laura and Fanghui work on improving the figures. No more than 10 figures, 2 in color.