Wisozk Group Regional Medical Center

3327 Elm St, Newark, NJ 07107

CONSOLIDATED PATIENT RECORDS — CONFIDENTIAL

Report Generated: 04/13/2026 06:41 PM
Boundary Interval: 3 page(s) per patient block

PATIENT 1 OF 6

Name: Dayton Walker
DOB: 07/23/1940
Address: 4234 Maple Dr, Boston, MA 02215
Insurance: INS-8605078563
Admitted: 04/11/2026

MRN: MRN-55716411
SSN: 295-28-6241
Phone: (267) 833-3105
Attending: Dr. Torrey Pouros
Discharged: 04/14/2026

VITALS

BP: 127/93 mmHg HR: 102 bpm Temp: 97.6 F SpO2: 93% RR: 17 breaths/min

DIAGNOSES

- Chronic Obstructive Pulmonary Disease (J44.1)
- Osteoarthritis, unspecified (M19.90)
- Anxiety Disorder, unspecified (F41.9)

MEDICATIONS

- Amlodipine 5mg — once daily
- Hydrochlorothiazide 25mg — once daily
- Omeprazole 20mg — once daily before breakfast
- Lisinopril 10mg — once daily
- Atorvastatin 20mg — once daily at bedtime
- Metoprolol 25mg — twice daily
- Prednisone 10mg — taper over 7 days

HOSPITAL COURSE

Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.

Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.

Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.

Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.

SUBSTANCE ABUSE & ADDICTION ASSESSMENT

Patient: Dayton Walker SSN: 295-28-6241 MRN: MRN-55716411
CAGE questionnaire score of 3/4; patient acknowledges problematic alcohol consumption averaging 12-15 drinks per week.
Buprenorphine/naloxone (Suboxone) 8mg/2mg sublingual film initiated for opioid use disorder during this admission.
AUDIT-C score of 8 indicates hazardous drinking pattern; brief intervention performed with motivational interviewing techniques.
Naloxone rescue kit prescribed at discharge given history of opioid use disorder; patient and family educated on administration.
Assessed by: Dr. Elva Kuhn, Addiction Medicine
42 CFR Part 2 protected information — unauthorized disclosure prohibited.

— End of record for Dayton Walker (MRN: MRN-55716411) —

PATIENT 2 OF 6

Name: Burley Mitchell
DOB: 08/30/1944
Address: 9713 Walnut St, Miami, FL 33136
Insurance: INS-6058252224
Admitted: 04/11/2026

MRN: MRN-22895722
SSN: 095-42-5527
Phone: (912) 225-9548
Attending: Dr. Christina Schamberger
Discharged: 04/20/2026

VITALS

BP: 108/73 mmHg HR: 71 bpm Temp: 100.4 F SpO2: 92% RR: 18 breaths/min

DIAGNOSES

- Type 2 Diabetes Mellitus (E11.9)
- Anxiety Disorder, unspecified (F41.9)
- Essential Hypertension (I10)
- Chronic Obstructive Pulmonary Disease (J44.1)
- Osteoarthritis, unspecified (M19.90)

MEDICATIONS

- Atorvastatin 20mg — once daily at bedtime
- Levothyroxine 75mcg — once daily on empty stomach
- Lisinopril 10mg — once daily
- Prednisone 10mg — taper over 7 days

HOSPITAL COURSE

Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.

Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.

Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.

Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.

PSYCHIATRIC EVALUATION & MENTAL HEALTH NOTES

Patient: Burley Mitchell DOB: 08/30/1944 SSN: 095-42-5527
Patient has documented history of self-harm behavior including cutting; last episode reported 3 weeks prior to admission.
Schizoaffective disorder, depressive type, managed with paliperidone palmitate (Invega Sustenna) 156mg IM monthly.
Eating disorder history documented: anorexia nervosa, restricting type, with BMI of 16.2 on admission; nutrition consult placed.
PHQ-9 score of 22 on admission consistent with severe major depressive episode; psychiatric consultation requested.
Patient endorsed auditory hallucinations described as commanding voices; antipsychotic medication adjusted accordingly.
Evaluating Psychiatrist: Dr. Brian Cremin, MD, Board Certified Psychiatry
Protected mental health information per state and federal law.

— End of record for Burley Mitchell (MRN: MRN-22895722) —

PATIENT 3 OF 6

Name: Maxine Bayer
DOB: 07/07/1954
Address: 4952 Washington Blvd, Savannah, GA 31405
Insurance: INS-6449876720
Admitted: 04/11/2026

MRN: MRN-61380873
SSN: 939-85-1616
Phone: (804) 428-9345
Attending: Dr. Cora Kling
Discharged: 04/13/2026

VITALS

BP: 126/67 mmHg HR: 100 bpm Temp: 99.5 F SpO2: 100% RR: 23 breaths/min

DIAGNOSES

- Major Depressive Disorder, recurrent (F33.0)
- Chronic Obstructive Pulmonary Disease (J44.1)
- Anxiety Disorder, unspecified (F41.9)
- Essential Hypertension (I10)
- Chronic Kidney Disease, Stage 3 (N18.3)

MEDICATIONS

- Amoxicillin 500mg — three times daily for 10 days
- Metoprolol 25mg — twice daily
- Gabapentin 300mg — three times daily
- Hydrochlorothiazide 25mg — once daily
- Prednisone 10mg — taper over 7 days
- Omeprazole 20mg — once daily before breakfast

HOSPITAL COURSE

Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.

Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.

Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.

Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.

HIV/STI SCREENING & TREATMENT NOTES

Patient: Maxine Bayer MRN: MRN-61380873 Insurance: INS-6449876720
Patient diagnosed with genital herpes simplex virus type 2; valacyclovir 1g BID initiated for acute outbreak management.
Gonorrhea and chlamydia NAAT screening performed on pharyngeal, rectal, and urogenital specimens; results pending at discharge.
CD4 count obtained during admission: 485 cells/mm3; viral load undetectable (<20 copies/mL) confirming virologic suppression.
RPR reactive with titer of 1:64; confirmatory FTA-ABS positive consistent with secondary syphilis; IM penicillin G administered.
Patient reports inconsistent condom use with multiple sexual partners; PrEP counseling provided and referral to sexual health clinic.
Infectious Disease Consult: Dr. Savion Jast
HIV-related information protected under state confidentiality statutes.

— End of record for Maxine Bayer (MRN: MRN-61380873) —

PATIENT 4 OF 6

Name: Ulises Stoltenberg
DOB: 02/23/2005
Address: 6424 Church St, Savannah, GA 31405
Insurance: INS-3351956658
Admitted: 04/11/2026

MRN: MRN-57740842
SSN: 202-85-6314
Phone: (857) 790-7224
Attending: Dr. Tia Lind
Discharged: 04/15/2026

VITALS

BP: 142/98 mmHg HR: 79 bpm Temp: 99.8 F SpO2: 93% RR: 15 breaths/min

DIAGNOSES

- Essential Hypertension (I10)
- Urinary Tract Infection (N39.0)
- Atrial Fibrillation (I48.91)

MEDICATIONS

- Amoxicillin 500mg — three times daily for 10 days
- Gabapentin 300mg — three times daily
- Levothyroxine 75mcg — once daily on empty stomach
- Sertraline 50mg — once daily
- Amlodipine 5mg — once daily
- Metformin 500mg — twice daily

HOSPITAL COURSE

Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.

Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.

The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.

GENETIC TESTING RESULTS & COUNSELING NOTES

Patient: Ulises Stoltenberg DOB: 02/23/2005 SSN: 202-85-6314
Whole exome sequencing identified pathogenic variant in CFTR gene consistent with cystic fibrosis carrier status.
Hereditary hemochromatosis: HFE C282Y homozygous genotype confirmed; therapeutic phlebotomy schedule initiated.
BRCA1 pathogenic variant identified (c.5266dupC); patient referred to genetic counselor and surgical oncology for risk-reducing options.
Genetic Counselor: Vernie Bode, MS, CGC
GINA-protected genetic information — disclosure restrictions apply.

— End of record for Ulises Stoltenberg (MRN: MRN-57740842) —

PATIENT 5 OF 6

Name: Bridie Towne
DOB: 08/31/1987
Address: 6314 Cedar Ln, Charlotte, NC 28216
Insurance: INS-6673236365
Admitted: 04/11/2026

MRN: MRN-28643831
SSN: 584-56-0842
Phone: (843) 662-3978
Attending: Dr. Chloe Johnson
Discharged: 04/16/2026

VITALS

BP: 114/80 mmHg HR: 106 bpm Temp: 98.7 F SpO2: 96% RR: 24 breaths/min

DIAGNOSES

- Acute Upper Respiratory Infection (J06.9)

- Type 2 Diabetes Mellitus (E11.9)
- Anxiety Disorder, unspecified (F41.9)
- Essential Hypertension (I10)
- Major Depressive Disorder, recurrent (F33.0)

MEDICATIONS

- Prednisone 10mg — taper over 7 days
- Atorvastatin 20mg — once daily at bedtime
- Amoxicillin 500mg — three times daily for 10 days
- Sertraline 50mg — once daily

HOSPITAL COURSE

Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.

Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.

Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.

DOMESTIC VIOLENCE / INTIMATE PARTNER VIOLENCE SCREENING

Patient: Bridie Towne Address: 6314 Cedar Ln, Charlotte, NC 28216 Phone: (843) 662-3978
Safety assessment completed: patient states she does not feel safe returning home; social work consulted for shelter placement.
Patient provided with National Domestic Violence Hotline number (1-800-799-7233) and local advocacy resources.
Mandatory reporting completed per state law given suspected abuse involving a vulnerable adult; Adult Protective Services notified.
Photographs of injuries documented per forensic protocol with patient consent; injuries inconsistent with reported mechanism.
Screened by: Garrick Schumm, LCSW
Sensitive domestic violence documentation — restricted access per hospital policy.

— End of record for Bridie Towne (MRN: MRN-28643831) —

PATIENT 6 OF 6

Name: Guy Fadel
DOB: 10/11/1948
Address: 250 Beacon St, Newark, NJ 07102
Insurance: INS-4455208695
Admitted: 04/11/2026

MRN: MRN-38028972
SSN: 938-58-9896
Phone: (207) 283-6960
Attending: Dr. Dejah Block
Discharged: 04/16/2026

VITALS
BP: 133/73 mmHg HR: 90 bpm Temp: 100.2 F SpO2: 92% RR: 13 breaths/min

DIAGNOSES
- Chronic Obstructive Pulmonary Disease (J44.1)
- Anxiety Disorder, unspecified (F41.9)
- Urinary Tract Infection (N39.0)

MEDICATIONS
- Amlodipine 5mg — once daily
- Metoprolol 25mg — twice daily
- Atorvastatin 20mg — once daily at bedtime
- Prednisone 10mg — taper over 7 days

HOSPITAL COURSE
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.

Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Dietary tray delivered; patient consumed approximately 75% of meal with good appetite noted by nursing staff.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Patient's family visited and was updated on the plan of care by the attending physician during afternoon rounds.
Pain management reassessed per scheduled protocol; patient reports pain level of 3/10 with current medication regimen.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Blood glucose monitoring performed per sliding scale protocol; values within target range and insulin administered as ordered.

Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Chaplain services offered to patient and family; patient declined at this time but is aware services are available.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Medication reconciliation reviewed with patient and pharmacy; no discrepancies identified between home and inpatient medication lists.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Fall risk assessment completed using the Morse Fall Scale; patient scored moderate risk and appropriate precautions implemented.
Nursing assessment performed at the bedside; patient is resting comfortably with no complaints of pain or distress.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
Interdisciplinary team rounds completed with input from medicine, nursing, pharmacy, physical therapy, and case management.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Intravenous fluids continued at maintenance rate; intake and output documented and within acceptable parameters.
Respiratory therapy performed scheduled nebulizer treatment; patient tolerated well with improved breath sounds bilaterally.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.
Physical therapy facilitated ambulation in the hallway with rolling walker; patient ambulated 150 feet with minimal assistance.
Patient education provided regarding the importance of ambulation and incentive spirometry use to prevent postoperative complications.
Occupational therapy assessed patient's ability to perform activities of daily living; mild difficulty with fine motor tasks noted.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Foley catheter care performed; output adequate and within expected range; plan to discontinue catheter per protocol tomorrow.
Wound care performed per protocol with sterile technique; wound bed appears clean and granulating without signs of infection.
Laboratory specimens collected per physician orders and sent to the lab; results anticipated within standard turnaround time.
Sequential compression devices applied bilaterally for DVT prophylaxis; patient instructed to keep devices on while in bed.
The patient's vital signs remained stable throughout the shift with no acute changes noted on continuous monitoring.
Patient participated in bedside physical therapy exercises including ankle pumps, quad sets, and straight leg raises.

SUBSTANCE ABUSE & ADDICTION ASSESSMENT

Patient: Guy Fadel SSN: 938-58-9896 MRN: MRN-38028972
Patient previously completed 28-day residential treatment program for cocaine dependence; reports 14 months of sobriety.
Urine drug screen on admission was positive for benzodiazepines and opioids; patient reports prescribed use only.

History of intravenous heroin use with last reported use 6 months prior to admission; Hepatitis C antibody positive. AUDIT-C score of 8 indicates hazardous drinking pattern; brief intervention performed with motivational interviewing techniques.
Assessed by: Dr. Roy Schulist, Addiction Medicine
42 CFR Part 2 protected information — unauthorized disclosure prohibited.
— End of record for Guy Fadel (MRN: MRN-38028972) —

END OF CONSOLIDATED PATIENT RECORDS
Total patients: 6 | Generated: 04/13/2026