



Cambridge Assessment International Education

Cambridge International Advanced Subsidiary Level

CANDIDATE NAME			
CENTRE NUMBER		CANDIDATE NUMBER	
MATHEMATICS			9709/23
Paper 2 Pure Mathen	natics 2 (P2)	Oct	ober/November 2019
			1 hour 15 minutes
Candidates answer or	n the Question Paper.		
Additional Materials:	List of Formulae (MF9)		

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 50.





This document consists of 12 printed pages.

© UCLES 2019

		•••••
		•••••
		•••••
		•••••
		•••••
		•••••
		•••••
(ii)	Hence find the largest integer n satisfying the inequality $ 2 \ln n - 7 < 2 \ln n - 9 $.	
(ii)	Hence find the largest integer n satisfying the inequality $ 2 \ln n - 7 < 2 \ln n - 9 $.	
(ii)		

Find the exact value of $\int_{1}^{2} (2e^{2x} - 1)^{2} dx$. Show all necessary working.	
	•••••
	••••••
	•••••
	•••••
	•••••
	••••••
	•••••
	•••••

-	A curve has equation $y = \frac{3 + 2 \ln x}{1 + \ln x}$. Find the exact gradient of the curve at the point for which y
•	
•	
•	
•	

4	The po	olynomia	1 p(x)	is	defined	by

$$p(x) = ax^3 + ax^2 - 15x - 18,$$

where a is a constant. It is given that (x-2) is a factor of p(x).

(i)	Find the value of a .	[2]
(ii)	Using this value of a , factorise $p(x)$ completely.	[3]
		•••••
		••••••
		•••••
		•••••
(iii)	Hence solve the equation $p(e^{\sqrt{y}}) = 0$, giving the answer correct to 2 significant figures.	[2]
		•••••
		•••••

Show that <i>a</i>	$= \sqrt[3]{(3-2\sin 2a)}$	$a-\cos a$).				
•••••			•••••	•••••		• • • • • • • • •
•••••			 		•••••	•••••
•••••			 •••••	•••••		
	••••		 			
•••••	•••••••••••		 ••••••	••••••	•••••	•••••
			 ••••••	••••••		•••••
•••••			 			
			•			
			 	••••••		• • • • • • • • • • • • • • • • • • • •
			 	•••••		•••••
•••••	•••••		 			
•••••			 •••••••	••••••••••	••••••	•
			 	••••••	•••••	• • • • • • • • • • • • • • • • • • • •
			 ••••••			•••••
•••••	•••••		 •	•••••		•••••

-	on in part (i), show			
	•••••	•••••		•••••
				•••••
•••••				
				•••••
				•••••
Use an iterative 3 significant figure	formula, based on res. Give the result	the equation in part of each iteration to 5	t (i), to find the values significant figures.	ue of a corre
Use an iterative 3 significant figure	formula, based on res. Give the result	the equation in part of each iteration to 5	t (i), to find the val	ue of a corre
Use an iterative 3 significant figure	formula, based on res. Give the result	the equation in part of each iteration to 5	t (i), to find the val	ue of a corre
Use an iterative 3 significant figure	formula, based on res. Give the result	the equation in part of each iteration to 5	t (i), to find the val	ue of a corre
Use an iterative 3 significant figure	formula, based on res. Give the result	the equation in part of each iteration to 5	t (i), to find the val	ue of a corre
Use an iterative 3 significant figure	formula, based on res. Give the result	the equation in part of each iteration to 5	t (i), to find the val	ue of a corre
Use an iterative 3 significant figure	formula, based on res. Give the result	the equation in part of each iteration to 5	t (i), to find the val	ue of <i>a</i> corre
Use an iterative 3 significant figure	formula, based on res. Give the result	the equation in part of each iteration to 5	t (i), to find the val	ue of <i>a</i> corre
Use an iterative 3 significant figure	formula, based on res. Give the result	the equation in part of each iteration to 5	t (i), to find the val	ue of a corre
Use an iterative 3 significant figure	formula, based on res. Give the result	the equation in part of each iteration to 5	t (i), to find the val	ue of a corre
Use an iterative 3 significant figure	formula, based on res. Give the result	the equation in part of each iteration to 5	t (i), to find the val	ue of <i>a</i> corre
Use an iterative 3 significant figure	formula, based on res. Give the result	the equation in part of each iteration to 5	t (i), to find the val	ue of <i>a</i> corre
Use an iterative 3 significant figure	formula, based on res. Give the result	the equation in part of each iteration to 5	t (i), to find the val	ue of <i>a</i> corre
Use an iterative 3 significant figure	formula, based on res. Give the result	the equation in part of each iteration to 5	t (i), to find the val	ue of <i>a</i> corre
Use an iterative 3 significant figure	formula, based on res. Give the result	the equation in part of each iteration to 5	t (i), to find the val	ue of <i>a</i> corre
Use an iterative 3 significant figure	formula, based on res. Give the result	the equation in part of each iteration to 5	t (i), to find the val	ue of a corre
Use an iterative 3 significant figure	formula, based on res. Give the result	the equation in part of each iteration to 5	t (i), to find the val	ue of a corre
Use an iterative 3 significant figure	formula, based on res. Give the result	the equation in part of each iteration to 5	t (i), to find the val	ue of a corre
Use an iterative 3 significant figure	formula, based on res. Give the result	the equation in part of each iteration to 5	t (i), to find the val	ue of a corre

6	(a)	Showing all	necessary	working,	solve th	ne equation
---	-----	-------------	-----------	----------	----------	-------------

								_
sec	\sim	0	\sim c	0	~ 1	ν	_	-

for $0^{\circ} < \alpha < 90^{\circ}$.	[5]
	••••••
	••••••
	••••••
	•••••
	•••••
	•••••
	•••••
	••••••

(b)	Showing all necessary working, solve the equation
	$\sin(\beta + 20^\circ) + \sin(\beta - 20^\circ) = 6\cos\beta$
	for $0^{\circ} < \beta < 90^{\circ}$. [4]

)	Find an expression for $\frac{dy}{dx}$ and show that the gradient of the curve at the point $(-1, 2)$ is $-\frac{5}{2}$.

Show that the curve has no stationary points.
Find the <i>x</i> -coordinate of each of the points on the curve at which the tangent is parallel to t <i>y</i> -axis.

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.			

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.