

Cambridge International AS & A Level

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

MATHEMATICS 9709/62

Paper 6 Probability & Statistics 2

February/March 2020

1 hour 15 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid.
- Do not write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

Use a suit 2 errors.	able app	roximatir	ig distrib	ution to	find the	probabil	lity that t	his bookl	et contair	is at
			•••••		•••••					•••••
				•••••	•••••					•••••
••••••	••••••	••••••	••••••	••••••	•••••	•••••	•••••		••••••	•••••
				••••••	•••••					•••••
				•••••	•••••	•••••				•••••
			•••••	•••••	•••••	•••••			••••••	•••••
			•••••	•••••	•••••	•••••			••••••	•••••
				•••••	•••••	•••••			•••••	•••••
•••••	••••••	•••••	••••••	•••••	•••••	•••••	•••••		••••••	
				•••••						•••••
				•••••						
				•••••						•••••
										•••••
										•••••
				•••••						•••••
				•••••	•••••					••••••
			•••••	••••••	•••••				•••••	
				•••••	•••••					
					•••••					
					•••••					

1.25 cm.	
Find α .	[:

3

In the past, the mean time taken by Freda for a particular daily journey was 39.2 minutes. Following

	1	i = 40	$\Sigma t = 1504$	$\Sigma t^2 = 57760$	
(a)	Calculate unbiased estir	nates of t	he population	mean and variance of the new journey time.	. [3
					•••••
b)	Test, at the 5% significa	nce level	, whether the p	opulation mean time has decreased.	[5
		••••••			
					•••••
		••••••			
					•••••
					•••••
		••••••			•••••
			•••••		•••••
		••••••	••••••		•••••
		••••••			•••••
					••••

© UCLES 2020 9709/62/F/M/20

The probability that there will be no accidents during a period of <i>n</i> days is greater than 0.95.
The probability that there will be no accidents during a period of <i>n</i> days is greater than 0.95.
The probability that there will be no accidents during a period of n days is greater than 0.95.
The probability that there will be no accidents during a period of n days is greater than 0.95.
The probability that there will be no accidents during a period of n days is greater than 0.95.
The probability that there will be no accidents during a period of n days is greater than 0.95.
The probability that there will be no accidents during a period of n days is greater than 0.95.
The probability that there will be no accidents during a period of n days is greater than 0.95.
The probability that there will be no accidents during a period of n days is greater than 0.95.

5	Bottles of Lanta contain approximately 300 ml of juice. The volume of juice, in millilitres, in a bottle
	is $300 + X$, where X is a random variable with probability density function given by

$$f(x) = \begin{cases} \frac{3}{4000} (100 - x^2) & -10 \le x \le 10, \\ 0 & \text{otherwise.} \end{cases}$$

(a)	Find the probability that a randomly chosen bottle of Lanta contains more than 305 ml of juice. [3]
(b)	Given that 25% of bottles of Lanta contain more than $(300 + p)$ ml of juice, show that
	$p^3 - 300p + 1000 = 0. [4]$
	p = 300p + 1000 = 0. [4]

© UCLES 2020 9709/62/F/M/20

• • • • • • • • • • • • • • • • • • • •		• • • • • • • • • • • • • • • • • • • •		•••••	• • • • • • • • • • • • • • • • • • • •	•••••
• • • • • • • • • • • • • • • • • • • •				• • • • • • • • • • • • • • • • • • • •		
• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •	•••••	• • • • • • • • • • • • • • • • • • • •	•••••
	• • • • • • • • • • • • • • • • • • • •			• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •
• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •	•••••	• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •
	• • • • • • • • • • • • • • • • • • • •					
• • • • • • • • • • • • • • • • • • • •	•••••	• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •	•••••	• • • • • • • • • • • • • • • • • • • •	•••••
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 - q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 - q) and	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 - q) as	nd (300 +
Given that $p = 3$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) as	nd (300 +
Given that $p = 3$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) as	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) as	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 3$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 3$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 3$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 3$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +
Given that $p = 0$ of juice, find q .	3.47, and that 5	50% of bottles	of Lanta con	tain between	(300 – q) an	nd (300 +

6

The volumes, in millilitres, of large and small cups of tea are modelled by the distributions N(200, 30)

Find the probability that the total volume of a randomly chosen large cup of tea and a random chosen small cup of tea is less than 300 ml.

volume of a randomly chosen small cup of tea.	

7

perc He c	ational survey shows that 95% of year 12 students use social media. Arvin suspects that the centage of year 12 students at his college who use social media is less than the national percentage chooses a random sample of 20 students at his college and notes the number who use social media then carries out a test at the 2% significance level.
(a)	Find the rejection region for the test.

© UCLES 2020 9709/62/F/M/20

•	
	Jimmy believes that the true percentage at Arvin's college is 70%. Assuming that Jimmy
C	correct, find the probability of a Type II error.
•	
•	
•	
•	
•	
•	
•	
•	
•	
•	
•	
•	
•	
•	
•	
•	
•	

Additional Page

must be clearly shown.	estion number(s)
	••••••
	••••••

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.