

A
GO enrichment (704 DEGs)
BP / CC / MF, BH-adjusted; universe = 21,358 genes
epithelial cell proliferation
benzene-containing compound metabolic process
fatty acid metabolic process
cell junction assembly
regulation of epithelial cell proliferation
epithelial to mesenchymal transition
monocarboxylic acid biosynthetic process
basement membrane
sulfur compound binding
heparin binding
glycosaminoglycan binding
bmp binding
phospholipase inhibitor activity
glutathione transferase activity
BP
CC
MF
2%
4%
6%
Gene ratio
−log10 Padj
4
3
2
Count
10
20
30
B
KEGG pathways
11 pathways, FDR < 0.05
cornified envelope formation
drug metabolism - cytochrome p450
drug metabolism - other enzymes
focal adhesion
fluid shear stress and atherosclerosis
glutathione metabolism
aldosterone-regulated sodium reabsorption
chemical carcinogenesis - dna adducts
ecm-receptor interaction
integrin signaling
metabolism of xenobiotics by cytochrome p450
Count
8
12
16
20
0
1
2
−log10 Padj
C
Hallmark gene sets (GSEA)
36 of 50 sets at FDR < 0.05
Myc targets v2
Myc targets v1
E2f targets
G2m checkpoint
Oxidative phosphorylation
Dna repair
Mitotic spindle
P53 pathway
Myogenesis
Estrogen response late
Estrogen response early
Epithelial mesenchymal transition
Interferon gamma response
Interferon alpha response
Uv response dn
Apoptosis
Tnfa signaling via nfkb
Hypoxia
-2
-1
0
1
2
Normalised enrichment score
D
Upstream TFs (ChEA3)
lower rank = stronger evidence
NKX3-1
68
GRHL3
81
SOX15
95
BNC1
100
FOXA1
213
TP63
131
TFCP2L1
99
EHF
104
ELF3
115
FOXQ1
98
ZNF750
99
KLF5
19
0
10
20
30
40
50
ChEA3 mean rank score