

School of Computing and Information Systems
COMP30026 Models of Computation
Week 2: Propositional Logic

If you find yourself getting stuck on a particular question in the tutorial, try to move onto other questions until you have a chance to ask your tutor for help.

Homework problems

P2.1 Which of the following pairs of formulas are equivalent?

- | | |
|---|---|
| (a) $\neg P \rightarrow Q$ and $P \rightarrow \neg Q$ | (e) $P \rightarrow (Q \rightarrow R)$ and $Q \rightarrow (P \rightarrow R)$ |
| (b) $\neg P \rightarrow Q$ and $Q \rightarrow \neg P$ | (f) $P \rightarrow (Q \rightarrow R)$ and $(P \rightarrow Q) \rightarrow R$ |
| (c) $\neg P \rightarrow Q$ and $\neg Q \rightarrow P$ | (g) $(P \wedge Q) \rightarrow R$ and $P \rightarrow (Q \rightarrow R)$ |
| (d) $(P \rightarrow Q) \rightarrow P$ and P | (h) $(P \vee Q) \rightarrow R$ and $(P \rightarrow R) \wedge (Q \rightarrow R)$ |

Solution:

(a) Not equivalent:

P	Q	$\neg P$	\rightarrow	Q	P	\rightarrow	$\neg Q$
0	0	1	0	0	0	1	1
0	1	1	1	1	0	1	0
1	0	0	1	0	1	1	1
1	1	0	1	1	1	0	0
				X			
				X			

We see that the columns for the two implications are different.

(b) Not equivalent:

P	Q	$\neg P$	\rightarrow	Q	Q	\rightarrow	$\neg P$
0	0	1	0	0	0	1	1
0	1	1	1	1	1	1	1
1	0	0	1	0	0	1	0
1	1	0	1	1	1	0	0
				X			
				X			

(c) Equivalent:

P	Q	$\neg P$	\rightarrow	Q	$\neg Q$	\rightarrow	P
0	0	1	0	0	1	0	0
0	1	1	1	1	0	1	0
1	0	0	1	0	1	1	1
1	1	0	1	1	0	1	1
				\checkmark			
				\checkmark			

(d) Equivalent:

P	Q	$(P \rightarrow Q) \rightarrow P$			
0	0	0	1	0	0
0	1	0	1	1	0
1	0	1	0	0	1
1	1	1	1	1	1
\checkmark		\checkmark			

(e), (f) : the pair in (e) equivalent; but the pair in (f) are not equivalent:

P	Q	R	$P \rightarrow (Q \rightarrow R)$	$Q \rightarrow (P \rightarrow R)$	$(P \rightarrow Q) \rightarrow R$
0	0	0	0 1 0 1 0	0 1 0 1 0	0 1 0 0 0
0	0	1	0 1 0 1 1	0 1 0 1 1	0 1 0 1 1
0	1	0	0 1 1 0 0	1 1 0 1 0	0 1 1 0 0
0	1	1	0 1 1 1 1	1 1 0 1 1	0 1 1 1 1
1	0	0	1 1 0 1 0	0 1 1 0 0	1 0 0 1 0
1	0	1	1 1 0 1 1	0 1 1 1 1	1 0 0 1 1
1	1	0	1 0 1 0 0	1 0 1 0 0	1 1 1 0 0
1	1	1	1 1 1 1 1	1 1 1 1 1	1 1 1 1 1
			✓	✓	✗

Note that the formulas in (e) are both equivalent to $(P \wedge Q) \rightarrow R$, which explains why the order of P and Q does not matter here.

(g) Equivalent:

P	Q	R	$(P \wedge Q) \rightarrow R$	$P \rightarrow (Q \rightarrow R)$
0	0	0	0 0 0 1 0	0 1 0 1 0
0	0	1	0 0 0 1 1	0 1 0 1 1
0	1	0	0 0 1 1 0	0 1 1 0 0
0	1	1	0 0 1 1 1	0 1 1 1 1
1	0	0	1 0 0 1 0	1 1 0 1 0
1	0	1	1 0 0 1 1	1 1 0 1 1
1	1	0	1 1 1 0 0	1 0 1 0 0
1	1	1	1 1 1 1 1	1 1 1 1 1
			✓	✓

(h) Equivalent:

P	Q	R	$(P \vee Q) \rightarrow R$	$(P \rightarrow R) \wedge (Q \rightarrow R)$
0	0	0	0 0 0 1 0	0 1 0 1 0 1 0
0	0	1	0 0 0 1 1	0 1 1 1 0 1 1
0	1	0	0 1 1 0 0	0 1 0 0 1 0 0
0	1	1	0 1 1 1 1	0 1 1 1 1 1 1
1	0	0	1 1 0 0 0	1 0 0 0 0 1 0
1	0	1	1 1 0 1 1	1 1 1 1 0 1 1
1	1	0	1 1 1 0 0	1 0 0 0 1 0 0
1	1	1	1 1 1 1 1	1 1 1 1 1 1 1
			✓	✓

P2.2 Define your own binary connective \square by writing out a truth table for $P \square Q$ (fill in the middle column however you like). Can you write a formula which has the the same truth table as $P \square Q$ using only the symbols P, Q, \neg, \wedge, \vee , and \rightarrow ? Repeat the exercise once.

Solution: Here's an example

P	Q	$P \square Q$
0	0	0 1 0
0	1	0 1 1
1	0	1 1 0
1	1	1 0 1

With this truth table for \square , $P \square Q$ is logically equivalent to $\neg(P \wedge Q)$.

P2.3 How many distinct truth tables are there involving two fixed propositional letters? In other words, how many meaningfully distinct connectives could we have defined in the previous question?

Solution: There are four rows in the truth table, so that's 2 choices for each row, which amounts to $2 \times 2 \times 2 \times 2 = 16$ possibilities.

P2.4 Find a formula that is equivalent to $P \leftrightarrow (P \wedge Q)$ but shorter.

Solution: $(P \wedge Q) \leftrightarrow P$ is logically equivalent to $P \rightarrow Q$. This is easily checked with a truth table, but how can we simplify $(P \wedge Q) \leftrightarrow P$ when we don't know what it is supposed to be equivalent to? Well, we can just try. Let us expand the biimplication and obtain $(P \rightarrow (P \wedge Q)) \wedge ((P \wedge Q) \rightarrow P)$. Intuitively, the conjunct on the right is just true, and we can check that with a truth table. So we have found that the original formula is equivalent to $P \rightarrow (P \wedge Q)$, which isn't any shorter, but still. We can rewrite the result as $(P \rightarrow P) \wedge (P \rightarrow Q)$, and now it becomes clear that all we need is $P \rightarrow Q$.

P2.5 Find a formula that is equivalent to $(\neg P \vee Q) \wedge R$ using only \rightarrow and \neg as logical connectives.

Solution: $(\neg P \vee Q) \wedge R$ is logically equivalent to $\neg((P \rightarrow Q) \rightarrow \neg R)$

P2.6 Consider the formula $P \rightarrow \neg P$. Is that a contradiction (is it *unsatisfiable*)? Can a proposition imply its own negation?

Solution: There is no contradiction at all. The formula is true if (and only if) P is false. The point of a conditional formula is to make a claim about the scenario where the premise (P) is true. If the premise of \rightarrow is false, the formula is satisfied. For the same reason, $\neg P \rightarrow P$ is satisfiable; it is not a contradiction. But $P \leftrightarrow \neg P$ is clearly a contradiction. (If you disagree with any of these statements, draw truth tables.)

P2.7 By negating a satisfiable proposition, can you get a tautology? A satisfiable proposition? A contradiction? Illustrate your affirmative answers.

Solution: If you negate a satisfiable proposition, you can never get a tautology, since at least one truth table row will yield false.

You will get another satisfiable proposition iff the original proposition is not valid. For example, P is satisfiable (but not valid), and indeed $\neg P$ is satisfiable.

Finally, if we have a satisfiable formula which is also valid, its negation will be a contradiction. Example: $P \vee \neg P$.

P2.8 For each of the following propositional formulas, determine whether it is satisfiable, and if it is, whether it is a tautology:

- (a) $P \leftrightarrow ((P \rightarrow Q) \rightarrow P)$
- (b) $(P \rightarrow \neg Q) \wedge ((P \vee Q) \rightarrow P)$

Solution: Let us draw the truth tables.

(a)

P	Q	P	\leftrightarrow	$((P \rightarrow Q) \rightarrow P)$
0	0	0	1	0
0	1	0	1	0
1	0	1	1	0
1	1	1	1	1

↑↑

Hence satisfiable, and in fact valid (all 1).

(b)

P	Q	$(P \rightarrow \neg Q) \wedge ((P \vee Q) \rightarrow P)$
0	0	0
0	1	0
1	0	1
1	1	1

\uparrow

Hence satisfiable (at least one 1), but not valid (not all 1). The truth table shows the formula is equivalent to $\neg Q$.

P2.9 Complete the following sentences, using the words “satisfiable, valid, non-valid, unsatisfiable”.

- | | |
|---|--|
| (a) F is satisfiable iff F is not _____ | (e) F is satisfiable iff $\neg F$ is _____ |
| (b) F is valid iff F is not _____ | (f) F is valid iff $\neg F$ is _____ |
| (c) F is non-valid iff F is not _____ | (g) F is non-valid iff $\neg F$ is _____ |
| (d) F is unsatisfiable iff F is not _____ | (h) F is unsatisfiable iff $\neg F$ is _____ |

Solution:

- | | |
|---|--|
| (a) F is satisfiable iff F is not unsatisfiable | (e) F is satisfiable iff $\neg F$ is non-valid |
| (b) F is valid iff F is not non-valid | (f) F is valid iff $\neg F$ is unsatisfiable |
| (c) F is non-valid iff F is not valid | (g) F is non-valid iff $\neg F$ is satisfiable |
| (d) F is unsatisfiable iff F is not satisfiable | (h) F is unsatisfiable iff $\neg F$ is valid |

P2.10 Show that $P \leftrightarrow (Q \leftrightarrow R) \equiv (P \leftrightarrow Q) \leftrightarrow R$. This tells us that we could instead write

$$P \leftrightarrow Q \leftrightarrow R \quad (1)$$

without introducing any ambiguity. Mind you, that may not be such a good idea, because many people (incorrectly) tend to read “ $P \leftrightarrow Q \leftrightarrow R$ ” as

$$P, Q, \text{ and } R \text{ all have the same truth value} \quad (2)$$

Show that (1) and (2) are incomparable, that is, neither is a logical consequence of the other.

Solution: Even with three variables the truth table is manageable, so let us construct it.

(1A)			(1B)					(2)		
P	Q	R	$P \leftrightarrow (Q \leftrightarrow R)$	$(P \leftrightarrow Q) \leftrightarrow R$	$P \wedge Q \wedge R$	$\neg P \wedge \neg Q \wedge \neg R$	\vee	$\neg P \wedge \neg Q \wedge \neg R$	\vee	$\neg P \wedge \neg Q \wedge \neg R$
0	0	0	0	0	0	1	0	1	1	1
0	0	1	0	0	0	0	0	0	0	0
0	1	0	0	0	0	0	0	0	0	0
0	1	1	0	0	0	0	0	0	0	0
1	0	0	1	1	0	0	0	0	0	0
1	0	1	1	0	0	0	0	0	0	0
1	1	0	1	0	0	0	0	0	0	0
1	1	1	1	1	1	1	1	1	1	0
			✓	✓	✗					

B is a logical consequence of A (we write $A \models B$) iff B is true for any assignment of variables which makes A true. So we see that (1) $\not\models$ (2), because cases like 1 0 0 that make (1) true but (2) false. Similarly, (2) $\not\models$ (1), because the case 0 0 0 makes (2) true but (1) false.

P2.11 Let F and G be propositional formulas. What is the difference between “ $F \equiv G$ ” and “ $F \leftrightarrow G$ ”? Prove that “ $F \leftrightarrow G$ ” is valid iff $F \equiv G$.

Solution: The connective \leftrightarrow is part of the language that we study, namely the language of propositional logic. So $A \leftrightarrow B$ is just a propositional formula.

The symbol \equiv belongs to a *meta-language*. The meta-language is a language which we use when we reason *about* some language. In this case we use \equiv to express whether a certain relation holds between formulas in propositional logic.

More specifically, $F \equiv G$ means that we have both $F \models G$ and $G \models F$. In other words, F and G have the same value for every possible assignment of truth values to their variables. The two formulas are logically equivalent.

On the other hand $F \leftrightarrow G$ is just a propositional formula (assuming F and G are propositional formulas). For some values of the variables involved, $F \leftrightarrow G$ may be false, for other values it may be true. By the definition of validity, $F \leftrightarrow G$ is *valid* iff it is true for *every* assignment of propositional variables in F and G .

We want to show that $F \equiv G$ iff $F \leftrightarrow G$ is valid.

- (a) Suppose $F \equiv G$. Then F and G have the same values for each truth assignment to their variables¹. But that means that, when we construct the truth table for $F \leftrightarrow G$, it will have a t in every row, that is, $F \leftrightarrow G$ is valid.
- (b) Suppose $F \leftrightarrow G$ is valid. That means we find a t in each row of the truth table for $F \leftrightarrow G$. But we get a t for $F \leftrightarrow G$ iff the values for F and G agree, that is, either both are f , or both are t . In other words, F and G agree for every truth assignment. Hence $F \equiv G$.

You may think that this relation between validity and biimplication is obvious and should always be expected, and indeed we will see that it carries over to first-order predicate logic. But there are (still useful) logics in which it does not hold.

P2.12 Is $(P \wedge Q) \leftrightarrow P$ logically equivalent to $(P \vee Q) \leftrightarrow Q$?

Solution: Yes, $(P \wedge Q) \leftrightarrow P \equiv (P \vee Q) \leftrightarrow Q$, and both of these formulas are logically equivalent to $P \rightarrow Q$

P	Q	$(P \wedge Q) \leftrightarrow P$					$(P \vee Q) \leftrightarrow Q$				
0	0	0	0	0	1	0	0	0	0	1	0
0	1	0	0	1	1	0	0	1	1	1	1
1	0	1	0	0	0	1	1	1	0	0	0
1	1	1	1	1	1	1	1	1	1	1	1

✓

✓

¹We should perhaps be more careful here, because F and G can be logically equivalent without F having the exact same set of variables as G —can you see how? So we should say that we consider both of F and G to be functions of the *union* of their sets of variables.