School of Computing and Information Systems COMP30026 Models of Computation Week 2: Propositional Logic

If you find yourself getting stuck on a particular question in the tutorial, try to move onto other questions until you have a chance to ask your tutor for help.

Exercises

T2.1 For each pair of formulas, write down and compare their truth tables. In which rows (if any) are they different?

(i)
$$\neg P \rightarrow Q$$
 and $P \rightarrow \neg Q$

(iii)
$$(P \wedge Q) \to R$$
 and $P \to (Q \to R)$

(ii)
$$\neg (P \land \neg Q)$$
 and $P \to Q$

(iv)
$$P \to (Q \to R)$$
 and $(P \to Q) \to R$

T2.2 Find a formula that is equivalent to $(P \land \neg Q) \lor P$ but shorter.

T2.3 Write down the definitions of *satisfiable*, *valid*, *unsatisfiable* and *non-valid*. Explain why any given propositional formula must have exactly two of these properties, and describe the following propositional formulas using these terms.

(i)
$$P \vee Q$$

(iii)
$$((P \to Q) \to P) \to P$$

(ii)
$$P \wedge \neg P$$

(iv)
$$(P \land \neg P) \rightarrow P$$

T2.4 Let's try a more challenging puzzle from Smullyan's Island of Knights and Knaves. Recall that knights always tell the truth and knaves always lie. We meet three people from the island, and we are reliably informed that one of the three is also a magician. They make these statements:

A: B is not both a knave and a magician.

B: Either A is a knave or I am not a magician.

C: The magician is a knave.

Who is the magician? Give a proof, and check that your solution indeed solves the problem.

Hint: If you have a disjunction like $P \vee Q$ or a negated conjunction like $\neg (P \wedge Q)$, negating it (e.g. for the sake of contradiction) produces a conjunction, which is usually much easier to work with.

Homework problems

P2.1 Which of the following pairs of formulas are equivalent?

(a) $\neg P \rightarrow Q$ and $P \rightarrow \neg Q$

(e) $P \to (Q \to R)$ and $Q \to (P \to R)$

(b) $\neg P \rightarrow Q$ and $Q \rightarrow \neg P$

(f) $P \to (Q \to R)$ and $(P \to Q) \to R$

(c) $\neg P \rightarrow Q$ and $\neg Q \rightarrow P$

(g) $(P \wedge Q) \to R$ and $P \to (Q \to R)$

(d) $(P \rightarrow Q) \rightarrow P$ and P

(h) $(P \lor Q) \to R$ and $(P \to R) \land (Q \to R)$

- P2.2 Define your own binary connective \square by writing out a truth table for $P\square Q$ (fill in the middle column however you like). Can you write a formula which has the the same truth table as $P \square Q$ using only the symbols P, Q, \neg, \wedge, \vee , and \rightarrow ? Repeat the exercise once.
- P2.3 How many distinct truth tables are there involving two fixed propositional letters? In other words, how many meaningfully distinct connectives could we have defined in the previous question?
- P2.4 Find a formula that is equivalent to $P \leftrightarrow (P \land Q)$ but shorter.
- P2.5 Find a formula that is equivalent to $(\neg P \lor Q) \land R$ using only \rightarrow and \neg as logical connectives.
- P2.6 Consider the formula $P \rightarrow \neg P$. Is that a contradiction (is it unsatisfiable)? Can a proposition imply its own negation?
- P2.7 By negating a satisfiable proposition, can you get a tautology? A satisfiable proposition? A contradiction? Illustrate your affirmative answers.
- P2.8 For each of the following propositional formulas, determine whether it is satisfiable, and if it is, whether it is a tautology:

(a)
$$P \leftrightarrow ((P \to Q) \to P)$$

(b)
$$(P \to \neg Q) \land ((P \lor Q) \to P)$$

- P2.9 Complete the following sentences, using the words "satisfiable, valid, non-valid, unsatisfiable".
 - (a) F is satisfiable iff F is not _____
- (e) F is satisfiable iff $\neg F$ is _____
- (b) F is valid iff F is not __
- (f) F is valid iff $\neg F$ is _____
- (g) F is non-valid iff $\neg F$ is _____
- (c) F is non-valid iff F is not _____ (d) F is unsatisfiable iff F is not _____
- (h) F is unsatisfiable iff $\neg F$ is _____
- P2.10 Show that $P \leftrightarrow (Q \leftrightarrow R) \equiv (P \leftrightarrow Q) \leftrightarrow R$. This tells us that we could instead write

$$P \leftrightarrow Q \leftrightarrow R$$
 (1)

without introducing any ambiguity. Mind you, that may not be such a good idea, because many people (incorrectly) tend to read " $P \leftrightarrow Q \leftrightarrow R$ " as

$$P, Q, \text{ and } R \text{ all have the same truth value}$$
 (2)

Show that (1) and (2) are incomparable, that is, neither is a logical consequence of the other.

- P2.11 Let F and G be propositional formulas. What is the difference between " $F \equiv G$ " and " $F \leftrightarrow G$ "? Prove that " $F \leftrightarrow G$ " is valid iff $F \equiv G$.
- P2.12 Is $(P \land Q) \leftrightarrow P$ logically equivalent to $(P \lor Q) \leftrightarrow Q$?