

TFPT — Red Team / Stress Test layer

Adversarial targets A–E, the QFT round (F) and the seam round (G): try to *break* the reductions, not confirm them

Stefan Hamann

Alessandro Rizzo

August 14, 2026 · v5.4 (rev 600)

The TFPT document set — what is treated where

Plain language: TFPT (Topological Fixed-Point Theory) is a small discrete compiler. Two inputs — the seam constant $c_3 = \frac{1}{8\pi}$ and the carrier rank $g_{\text{car}} = 5$ — build $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ and read off the Standard-Model skeleton, the constants and the scale grammar. The series is **the introduction plus five numbered papers and five companions** (plus two standalone working notes), best read in order:

0. **introduction** — reading guide, compiler closure, predictions, the dependency DAG and the single proof ledger (`verification/status_ledger.csv`). (*entry point*)
1. **tfpt_1_architecture_e8** — the two axioms, the derivation map, the EM fixed point α^{-1} , the $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ construction, the QBL theorem chain.
2. **tfpt_2_standard_model** — the SM in one φ_0 -ladder formula, flavor from parabolic transport (sheet diamond, dual anchor, Q_+ cohomology), the worked closures.
3. **tfpt_3_e8_audit_bootstrap** — the seven E_8 slices as an audit raster, the cascade bridge, the Möbius bootstrap.
4. **tfpt_4_frontier** — honest status of η_B , m_p/m_e , Koide, dark matter, full QG.
5. **tfpt_5_redteam** — the adversarial audit: declared attacks, what survives, what each reduces to, and the kill tests.

Companions: **R.** **tfpt_research_contracts** — the open interfaces (v_{geo} , G_{net} , F_{transfer}) as numbered contracts; **H.** **tfpt_horizon_readouts** — Appendix H (unit-system reframe): one seam constant as the universal horizon thermal code, the Nariai/anchor surface, the resummed seam clock; **O.** **origin_theory** — why no free fundamental number remains (exact core + one honestly-typed cyclic interpretation); **S.** **tfpt_safeguards** — the verification discipline: every mechanism that defends a claim against coincidence and numerology (status calculus, no-free-pattern, the over-determination map, the fire-wall, frozen predictions + null model, the independent Wolfram and Lean paths, the red team); **P.** **tfpt_prime_front** — research documentation of the prime / zeta line: the seven load-bearing modules on the E_8 glue census (Hecke from geometry through the transport ledger), the sandbox localization program that follows them, and the kill list. Documentation, not a result: it makes no claim about the Riemann Hypothesis.

Working notes: **N1.** **note_e8_gaussian_code** — the Gaussian code bridge: E_8 over $\mathbb{Z}[i]$ via Construction A over the extended Hamming code $[8, 4, 4]$, the canonical four-bit quotient $L/(1+i)L \cong \mathbb{F}_2^4$, and the G_{31} quartic companion (exact algebra, machine-certified); **N2.** **note_hilbert_polya_truncations** — a computable, zeta-free truncation family for the Weil measure: measurements on a Hilbert–Pólya candidate (prediction-freeze methodology; no RH claim).

You are reading document 5 (Red Team).

TFPT status at a glance — read this card the same way in every document

Two inputs, one compiler. From the boundary constant $c_3 = \frac{1}{8\pi}$ (axiom P1) and the five-slot carrier $g_{\text{car}} = 5$ (axiom P2) — jointly the elementary symmetric data of one anchor $a = (1, 1, 2)$ — the discrete compiler $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ is *built*, and the Standard-Model skeleton (gauge group, three families, hypercharges, the flavor matrix), the dimensionless constants ($\alpha^{-1} = 137.0359992$) and the scale grammar are read off as fixed points. The algebraic core is machine-checked; physical readouts run through *named bridges* whose status is typed below.


What is closed, and what genuinely remains. The discrete/algebraic layer is closed ([E]); the everything-open residual reduces to *three named interface problems*:

Interface	Question	Status
v_{geo}	the one metrology unit ($= 1/\sqrt{G} = m/\mu$); No-Unit Thm: no compiler scale	primitive [O]
G_{net}	simple-current extension $(D_5)_1 \otimes (A_3)_1 \rtimes \langle (1, 1) \rangle \cong (E_8)_1$	[C]
F_{transfer}	one functor, four typed interfaces (Koide, η_B , axion, m_p/m_e)	[C]

Gravity is parameter-free. The classical metric-sector field equation is no longer only an $R+R^2$ readout: the fixed-volume entanglement equilibrium (Jacobson; Faulkner et al.), run with TFPT's atoms, gives the *full* covariant Einstein equation $G_{ab} + \Lambda g_{ab} = c_3^{-1} T_{ab}$ with *both* coefficients fixed — $c_3^{-1} = 8\pi$ (no free dimensionless Newton dial; the thermodynamic origin $2\pi/\eta$ *coincides* with the geometric one $|\mathbb{Z}_2|2\pi\chi$ via $|\mu_4| = |\mathbb{Z}_2|\chi(S^2) = 4$, so c_3 is triply over-determined — anchor, geometry, thermodynamics, v358) and Λ from α ($\rho_\Lambda = (3/4\pi^2)e^{-2\alpha^{-1}}$, v60); the Einstein tensor (not Ricci) is forced by Lovelock, so matter conservation is an output (v359). The residual here is only the equation-of-state status (an external candidate action — Bianconi's entropic action, arXiv:2408.14391 — is quantified in v473 without changing the typing) and the one unit v_{geo} ; the global measure (C7 = QG.AMB.01) is now a [C] redundancy (v369), a certification object rather than missing dynamics, conditional on SEAM.EQUIV.01 and Bisognano–Wichmann.

The seam-net structural theorem G_{net} (= SEAM.EQUIV.01: the raw seam *is* the holomorphic $(E_8)_1$ net) now carries two named routes (route split 2026-07-22, no status change in substance): the MMST route SEAM.EQUIV.MMST.01 is *closed modulo cited theorems*; the twistor route SEAM.EQUIV.TWISTOR.01 (Costello–Li, prepared by CELEST.SEAM.01) and the unconditional parent stay [O]. On the MMST route: the explicit lattice model (v367/v368) and the S_3 closure stack pin the target at every computable level — central charge $c = 8$ (v376), the $(E_8)_1$ character with 248 currents and one primary (v377), genus-one torus $\text{GSD} = 1$ (v378) and reflection positivity (v379) — and it is Lean-formalised as FORM.SEAM.MMST.01: the collar's MMST hypotheses are kernel-proved, the MMST scaling-limit and Adamo–Moriwaki–Tanimoto OS-reconstruction theorems enter as named cited axioms, and the #print axioms check is clean. The *only* residual of that route that stays [O] is the abstract continuum existence of the scaling limit (v336) — its 128-spinor extension leg since certified at net level by the peer-reviewed crossed-product package ($h_s=16/16=1 \in \mathbb{Z}$, the Longo–Rehren locality criterion; Böckenhauer; KLM $\mu=4/2^2=1$; AGT/AMT the independent second witness) with the realisation input reduced to invariant level and the parallel Lean route seamResidualClosed' (v469); so the MMST route is *closed modulo cited theorems*, not solved. QGEO.SYM.01 (the μ_4 deck acting geometrically) is its *corollary*: a conformal net's vacuum is rotation-invariant by axiom, so the deck follows with no extra premise (v335, Lean FORM.QGEO.BW.01). The nonperturbative quantum-gravity measure QG.AMB.01 is no longer carried as an open item: it is a [C] redundancy (v369), a certification object rather than missing dynamics, conditional on SEAM.EQUIV.01 and Bisognano–Wichmann (and TFPT is moreover rigorously gap-decoupled from the general Euclidean-QG conformal-factor problem, margin $1.648 > 0$, v76/v330). So the honest residual is the one unit v_{geo} plus the typed F_{transfer} interfaces, above the cited-theorem ceiling on the seam and with QG.AMB.01 a [C] redundancy. Everything carries a claim ID resolving to a row of verification/status_ledger.csv (the single source of truth); **if the text and the ledger ever disagree, the ledger wins.** The historical labels ($U_{\text{wall}}, G_{\text{metric}}, F_{\text{frontier}}$) are retained only for ledger continuity — the live residual is $v_{\text{geo}} \oplus F_{\text{transfer}}$, with the seam G_{net} closed modulo cited theorems on its MMST route (parent [O]) and QG.AMB.01 a [C] redundancy.

Marker key: [E] exact / machine-proven · [C] conditional (holds under named hypotheses) · [O] open / axiom · [X] falsifiable kill test. Per-claim fine type in verification/status_ledger.csv.

[E] exact / machine-proven**[C]** conditional**[O]** open / axiom**[X]** kill test**What this note is**

This is the deliberately *adversarial* layer requested in the v78 review. Its purpose is **not** to confirm TFPT but to attack the five load-bearing reductions at their weakest logical transitions. Each target is run through one fixed protocol — minimal statement, assumptions, logical chain, validity conditions, counterexample search, limiting/degenerate cases, alternative structures, provisional verdict — and produces a deliberately narrow output: where the statement works, where it could fail, and the residual risk. The layer is machine-backed by [verification/redteam/rt_A_e8net.py] ... [verification/redteam/rt_E_vgeo.py], [verification/redteam/rt_F_qft4d.py] and [verification/redteam/run_redteam.py]; a red-team check asserts an *adversarial* fact (a counterexample really exists, a hidden assumption is really needed, a firewall really holds), and the honest outcome lives in the **status** of each target, never in a green pass. The layer has since grown a seam round (Target G): the ten-test side-blind scoreboard on the alignment bit and the first *firing* kill test of the programme — the interacting straddle law (v521/v525/v528/v529), fenced at toy level — since executed as a *selector* with exactly one survivor: reflection positivity dynamically selects the bit $\delta = \pi/2$ (v534).

Target	Minimal claim	Status	Residual risk
A	seam–Calderón measure = $(E8)_1$ net	[C] closed mod. cited thm	one statement: boundary-net holomorphy + $c=8$ ($\Leftrightarrow$ the index-4 inclusion); E_8 and bulk uniqueness then automatic (v83/v87/v89; Lie-level realisation v143). Now closed modulo cited theorems on its MMST route SEAM.EQUIV.MMST.01 (the parent SEAM.EQUIV.01 stays [O]): the lattice model (v367/v368) and the S_3 stack ($c=8$, 248 currents/1 primary, torus GSD=1, RP; v376–v379) pin it at every computable level, Lean-formalised as FORM.SEAM.MMST.01; only the cited MMST/OS continuum existence stays [O] (v336) — its 128-spinor extension leg since re-founded on 1995–2001 peer-reviewed crossed-product theorems ($h_s=1\in\mathbb{Z}$, Longo–Rehren/Böckenhauer/KLM; AGT/AMT second witness) with the realisation input reduced to invariant level (v469). The free-bulk premise is a fixed-point theorem, the cone is eliminated (cone gap = one-particle gap), and the residual factors into the A_2 net assembly + GATE.QGEO (v160–v165).
B	$g_{\text{car}} = 5$ forced by the Pascal rule	survives (narrowed)	boundary derivation of half-spinor exhaustion (degree-2 truncation)
C	$k = c_3/2 = \frac{1}{16\pi}$, $S/A = \frac{1}{4}$	survives	absolute $1/G$ (UV-sensitive) is the one anchor
D	ratios+products $\Rightarrow$ one scale v_{geo}	survives (narrowed)	CP phases ($\delta_{\text{CKM/PMNS}}$) not covered by v_{geo}
E	one scale v_{geo} carries the theory	survives (narrowed)	EW/reheating/leptogenesis scales + seam=Planck identification

Method

The red team treats each reduction as a hostile witness. A confirmatory script that always passes is worthless here: the layer is built so that it *may* downgrade a claim. Four outcomes are allowed — **survives** (stands as worded), **survives (narrowed)** (stands only after a silent assumption is made explicit), **reduced, not closed** (the conservative wording is the correct one), and — reached by Target A after the lattice/ S_3 round — **closed modulo a cited theorem** (pinned at every computable level, only an external published existence theorem left). A fifth, **broken**, is reserved for an actual failure; none occurred on the load-bearing surface. The closest call is deliberately

kept visible rather than filed away: the first genuinely *interacting* seam toy fires Kill-Test 2 of the OS twistor contract *at toy level* — reflection positivity breaks after the *straddle law* (Target G below) — a fenced toy result, not a target downgrade, and simultaneously the first hard selection principle the adversarial layer has produced; run as such, it keeps exactly *one* interacting member alive (v534, Target G).

Target A — the $(E8)_1$ boundary-net identification

Minimal claim. The ambient metric/projective measure is reduced to identifying the seam–Calderón boundary measure with the $(E8)_1$ lattice net, carrier subnet $(D5)_1 \times (A3)_1$. [verification/redteam/rt_A_e8net.py]

Logical chain (established). The Sugawara level-1 central charges are $c(E8)_1 = \frac{248}{31} = 8$, $c(D5)_1 = 5 = g_{\text{car}}$, $c(A3)_1 = 3 = N_{\text{fam}}$, so the conformal-embedding criterion $c_{\text{coset}} = c(E8) - c(D5) - c(A3) = 0$ holds [E]. This is *compatibility*.

Where it breaks: central charge underdetermines the net

Counterexample: $(D8)_1 = SO(16)_1$ also has $c = \frac{120}{15} = 8$, but is *not* holomorphic (four primaries $1, v, s, c$) — a distinct $c = 8$ net. Hence equal central charge is *necessary, not sufficient*: $(E8)_1$ is the unique *holomorphic* $c = 8$ net (even unimodular rank-8 lattice = $E8$), so **holomorphy (single vacuum sector, μ -index 1) is the load-bearing extra assumption**. Dropping chirality is worse: the $c = 8$ Narain family is positive-dimensional. The constructive map seam–Calderón kernel $\rightarrow (E8)_1$ and the uniqueness of bulk reconstruction are not established; the gap $\Delta_{\text{eff}} > 0$ *supports* tightness (clustering) but does not prove it.

Verdict. Keep “reduced to a rigorous boundary-net identification problem”; do *not* write “metric sector closed”. The red team confirms the conservative wording. Status [C]. Residual: holomorphy proof + constructive boundary map + bulk-reconstruction uniqueness.

Target B — carrier rank / the Pascal condition

Minimal formal claim. $2^g = g^2 + g + 2$ has the unique solution $g = 5$ (Lean-verified [E]). This arithmetic is *not* attacked. The attack is upstream: why must the carrier obey the Pascal half-spinor exhaustion $2^{g-1} = \binom{g}{0} + \binom{g}{1} + \binom{g}{2}$? [verification/redteam/rt_B_pascal.py]

Where it narrows: the rule, not the arithmetic, selects 5

Perturbing the constant breaks it: $2^g = g^2 + g + c$ gives $\{5\}$ only at $c = 2$; neighbours give other or empty solution sets. The truncation *degree* is the real choice: with $2^{g-1} = \sum_{k \leq K} \binom{g}{k}$ one finds $K=0 \rightarrow g=1$, $K=1 \rightarrow 3$, $K=2 \rightarrow 5$, $K=3 \rightarrow 7$, i.e. $g = 2K + 1$. So choosing $K = 2$ (to reach $g_{\text{car}} = 5$) is a physical postulate. It is corroborated — the $D5$ half-spinor $\dim S^+ = 2^{g-1} = 16 \Leftrightarrow g = 5$ and the family closure $N_{\text{fam}} = (2^{g-1} - 1)/g$, $g + N_{\text{fam}} = 8$ both pick 5 — but each is itself a postulate, not a theorem.

Verdict. Theorem A’s arithmetic core stands; the Pascal *selection* must be typed [O]/[C], not [E]. Residual: a boundary/seam derivation of half-spinor exhaustion.

Target C — $k = c_3/2$ and the seam-area coefficient

Minimal claim. In reduced units $k = c_3/2 = \frac{1}{16\pi}$ and Fursaev–Solodukhin gives $S/A = \frac{1}{4}$. [verification/redteam/rt_C_kc3.py]

The dimensional firewall holds

The only legal chain is

$$k_{\text{red}} = \frac{c_3}{2} = \frac{1}{16\pi} \text{ (dimensionless),} \quad k_{\text{phys}} = \frac{k_{\text{red}}}{G}, \quad S = 4\pi k_{\text{phys}} A_{\text{phys}} = \frac{A_{\text{phys}}}{4G}.$$

The forbidden form $k_{\text{phys}} = c_3/2$ is dimensionally false (legal only at $G = 1$). A scan of **v67/v68/v73** finds every $c_3/2$ used as the *reduced* coefficient and **zero** naked dimensionful identities; the firewall's teeth are verified against a synthetic violation. Internally consistent with $1/G$ being UV-sensitive (Sakharov/Connes, **v68**).

Verdict. Safe as worded (reduced coefficient). Status **[E]** (structure) + **[O]** for the residual. Residual: the absolute $4D$ Newton scale $1/G$ ($\propto \Lambda^2 f_2$) stays the one dimensionful anchor; nothing *dimensionless* is open.

Target D — $U_{\text{point}} \rightarrow v_{\text{geo}}$

Minimal claim. Ratios plus sector products determine the dimensionless amplitudes, leaving one common scale v_{geo} . [[verification/redteam/rt_D_upoint.py](#)]

Where it narrows: the bijection needs explicit hypotheses

For positive reals, (ratios, product) $\Leftrightarrow$ (individuals) is a bijection, and the TFPT sectors satisfy it (positive rational c , odd size $3 = N_{\text{fam}}$). But it *fails* off-assumption: even sector size leaves a sign ambiguity ($\{2, 3\}$ vs $\{-2, -3\}$ share ratio $3/2$ and product 6); complex amplitudes leave an n -th-root branch ambiguity ($\{1, 1, 1\}$, $\{\omega, \omega, \omega\}$, $\{\omega^2, \omega^2, \omega^2\}$ share all ratios and product 1); a zero amplitude is degenerate. **Genuine residual:** the bijection fixes amplitude *magnitudes* only — CP phases $\delta_{\text{CKM}}, \delta_{\text{PMNS}}$ are complex and are *not* folded into v_{geo} . **Narrowed (T177 mirror, [[verification/v561_cp_channel_identities.py](#)]):** the magnitude data carry *exactly zero* phase on the TFPT sectors ($\arg \prod_i c_i = 0$ exactly for both the lepton and the up-quark triple), and the μ_3 branch group is blind precisely because its order divides $N_{\text{fam}} = 3$ (a μ_4 branch would already shift the product phase) — so the residual is a *fiber* datum (the hexagonal CM unit), not a magnitude datum. **Decided (T178 frame door, [[verification/v564_cp_fiber_pin.py](#)]):** the fiber location is now a *theorem*, not a search result — the fiber acts on the frame by a scalar on one row, so every homogeneous frame block factorises exactly (Euler), and the metric-free invariants of $(\mathbf{1}, d, n)$ have *one* generator (det; the bracket enumeration is exact): no frame-coupled scale-free $\pi/3$ -carrier exists without importing a metric, and every metric import is non-canonical (a continuum of normaliser-relative values). The CP residual is *relocated with a proof*: it is the fiber datum $\text{Im } \rho = \sin(\pi/3)$, and it provably cannot be made frame data.

Verdict. Holds for TFPT once four hypotheses (real, positive, fixed branch, no zeros / odd sector size) are made explicit; they are silent in **v75** and must be named. Status **[E]** (reduction) + **[O]** (v_{geo}). Residual: the CP-phase sector.

Target E — v_{geo} as the unique dimensional floor

Minimal claim. No pure-number theory can derive an absolute dimensionful scale; all scales are ratios to one unit v_{geo} . [[verification/redteam/rt_E_vgeo.py](#)]

Single-scale audit: exact for the certified tiers, conditional beyond

The closed compiler + protected IR readouts all reduce to (pure number) $\times v_{\text{geo}}$: $M_{\text{scal}} = c_3^{7/2} \bar{M}_{\text{Pl}}$, $f_a = (c_3/|\mu_4|)^{7/2} \bar{M}_{\text{Pl}}$, masses $= (\pi/\sqrt{2}) c_f \varphi_0^k v_{\text{geo}}$, and $1/G$ shares the same anchor (**v68/v75**). The audit *surfaces* candidates that are not pure reductions: the seam cutoff Λ (only an *identification* seam=Planck collapses it to v_{geo}), the electroweak matching scale, the reheating temperature, and the leptogenesis scale — all frontier **[C]**/**[O]** inputs that must not be silently absorbed into v_{geo} .

Verdict. v_{geo} is the unique dimensional floor of the certified tiers (true by dimensional analysis). Full single-scale-ness is conditional on the flagged identifications staying honestly typed. Status **[E]** (ratios) + **[O]** (one scale). Residual: explicit reduction (or honest frontier typing) of the EW / reheating / leptogenesis scales + the seam=Planck cutoff.

Target F — the perturbative 4D-QFT + scale round (v269–v275)

Minimal claim. The round published in release 5.2 — the Epstein–Glaser S_{pert} layer (v269/v271/v273), the PMNS Jarlskog CP strength (v270), the absolute ν -scale typing (v272), the scale over-determination (v274) and the QG.AMB.01 roadmap (v275). [verification/redteam/rt_F_qft4d.py]

Two attacks land; the round survives once they are named

(1) **The gravity S_{pert} ghost is a truncation artefact.** Power-counting renormalizability is *not* unitarity: the local R^2/Weyl^2 truncation gives a four-derivative propagator $1/(p^2(p^2 + M^2))$ with a negative-residue massive spin-2 mode (the Stelle ghost). But that is precisely the Seeley–DeWitt truncation of the seam KMS form factor: the entire factor $a(p^2)=e^{p^2/M^2}$ keeps its only pole at $p^2=0$ (v304), the spin-2 sector is ghost-free via the Barnes–Rivers decomposition (v370), and the nearest truncation-zero modulus runs to infinity (v380), so *perturbative* graviton unitarity is established and the ghost is exactly the truncation. The matter+gauge S_{pert} is unitary outright, and now closes *to all orders* as a typed Epstein–Glaser/BRST contract (v381, QFT4D.EG.ALLORDER.01): dimension-4 power-counting $\Rightarrow$ a finite counterterm space, BRST nilpotency $s^2=0$ for $su(3)\times su(2)$ (Jacobi), and the seam gap $\Rightarrow$ the adiabatic limit, with the imported all-order T_n existence (causal factorisation) and Slavnov–Taylor identity; the *nonperturbative* QG.AMB.01 is then a [C] *redundancy* (v369; gap-decoupled from every TFPT readout, not a TFPT structural item, v335), a certification object rather than a new gap. (2) **The over-determination is conditional.** v274’s Route-2 inverts the $\rho_\Lambda/\bar{M}_{\text{Pl}}^4 = (3/4\pi^2)e^{-2\alpha^{-1}}$ prediction, which is itself LAMBDA.BRANCH.01 [C]; so the 0.11% gravity-vs-cosmology agreement is conditional on the Λ -branch (the gravity route is unconditional). **Already-honest:** J_{PMNS} inherits δ ’s [C] provenance (assembled μ_6 phase, not an M_ν output) and the ν -scale is [O] with the NO floor $\Sigma m_\nu = 0.0586$ eV as the one kill test (under the carrier normalisation $y_\nu=y_t$ it is a one-parameter, PS-window-bracketed [O] candidate with the integer c_3 -rung declined, v481 — and meanwhile *excluded* unstructured by the bracketed decision computation, v482).

Verdict. SURVIVES (NARROWED): the round stands as typed; the two narrowings above are now folded into v269 and v274. Residual: the perturbative graviton sector is now established *unitary* — the Stelle ghost is a truncation artefact (v304/v370/v380) — and the nonperturbative QG.AMB.01 is a [C] *redundancy* (gap-decoupled from every TFPT readout, not a TFPT structural item, v335/v369); and the anchor over-determination is conditional on the Λ -branch. On the all-order footing (v381) the SM-non-unification is robust to all orders, and the *optional* carrier-Pati–Salam UV branch is *proton-safe* ([verification/v385_uvbranch_killtest.py]): its minimal $SU(4)$ gauge bosons mediate rare LFV ($K_L \rightarrow \mu e$), *not* $p \rightarrow e^+\pi^0$, so $M_{PS} \sim 3 \times 10^{13}$ GeV clears the binding bound by $\sim 10^7$ — a frozen kill-test surface (κ /forbidden-rep/new-state/LFV) with *no* fake τ_p window (v265).

Under examination: an infinite-derivative narrowing ([C], does *not* close the gate).

The Stelle ghost is a feature of the *local* $R+R^2/\text{Weyl}^2$ truncation: the four-derivative propagator $1/(p^2(p^2+M^2)) = (1/M^2)[1/p^2 - 1/(p^2+M^2)]$ carries the negative-residue spin-2 mode. But that truncation is the a_4 Seeley–DeWitt order of the spectral action $\text{Tr } f(D^2/\Lambda^2)$, whose cutoff is *not* a free scheme — it is the $\beta=1$ seam KMS weight $f(u) = e^{-u}$ (v259, no new parameter). Keeping that exponential cutoff *un-truncated* points to a nonlocal, infinite-derivative graviton form factor $p^2 a(p^2)$; for an *entire* factor $a(p^2) = e^{p^2/M^2}$ (nowhere zero) the dressed propagator $1/(p^2 a)$ keeps its *only* pole at $p^2=0$ — no ghost (Tomboulis 1997; Biswas–Mazumdar–Siegel 2006) — whereas any *polynomial* (finite-derivative) truncation has a real zero = the ghost pole. So *if* the resummed form factor is entire, the perturbative ghost is a truncation artefact: a conditional [C] narrowing of this attack. It does *not* close QG.AMB.01 — the trace regulator f is not yet shown to resum to an entire kinetic form factor $a(\square)$, and a ghost-free *perturbative* graviton is still not the nonperturbative ambient measure. (QGAMB.IDG.01, [C]/[O]; consistent with the SEAM.EQUIV.01 firewall.) [verification/v304_idg_nonlocal_ghost.py]

Sharpened by spin sector ([C]). The Barnes–Rivers spin decomposition makes this precise: of the $5+3+1+1=10$ symmetric-tensor components the Stelle ghost is a *spin-2* state, so the entire form factor $a(p^2)=e^{p^2/M^2}$ cures exactly the spin-2 sector (its only pole stays at $p^2=0$), while the wrong-sign *spin-0* conformal mode ($c_{\text{conf}}(4) = -3/2$) is handled separately by the GHP contour

(v334). So the full *perturbative* graviton is ghost-free sector-by-sector under the TFPT-fixed package — still [C] (it inherits the entire-analyticity assumption) and still not the nonperturbative measure. (GRAV.SPIN2.UNITARITY.01.) [verification/v370_grav_spin2_unitarity.py]

And the “entire” assumption is itself discharged at the truncation-tower level ([E]; [verification/v380_grav_kms_hessian.py]). The seam KMS cutoff $f(u)=e^{-u}$ (v259) gives the kinetic form factor $a(u)=e^u$ ($u=p^2/M^2$), which is entire and nowhere zero — so the spin-2 propagator $e^{-p^2/M^2}/p^2$ has its only pole at $p^2=0$. The Seeley–DeWitt expansion is the *Taylor truncation* of a : the $R+R^2$ order $1+u$ carries the Stelle zero $u=-1$, the Weyl² order $1+u+u^2/2$ the pair $-1\pm i$, and the modulus of the nearest truncation-zero grows monotonically with the order (1, 1.41, ..., 3.07 at order $1\rightarrow 8$) and runs to infinity. So the ghost is *exactly* the truncation and resummation removes it — “entire” means “do not truncate the heat kernel”, upgrading the assumption above to a derived statement. (GRAV.KMS.HESSIAN.01; the exact off-shell curved-space Hessian identification stays the cited heat-kernel step, and it is perturbative, not the nonperturbative measure.)

Target G — the alignment bit and the interacting seam (the first shot that lands)

Minimal claim. The compiler’s residual discrete carrier input is *one symmetry-lift bit* — flag transitivity of the four marks $\iff \tau = i$ (v512) — and the free-seam OS/RP structure that the Woit bridge (WOIT.OS.TWISTOR.01) builds on survives adversarial attack. The red-team questions are two: *can the bit be derived* (in which case calling it an input would be a hidden redundancy), and *can the OS/RP structure be broken?*

Attack 1 — derive the bit (ten failures, honestly banked). Every attack class that could compute the bit from side-blind data has been run and *killed*: the v512 equivalence web itself (no local jet sees the side bit), free RP/ Θ existence (the *eighth* side-blind test, [verification/v521_seam_bit_rp_blind.py]), every Wick-computable mark-decorated state — twist/defect insertions — (the *ninth*; the free-plus-twist class is *exhausted*, [verification/v525_seam_bit_twist_blind.py]), and now the first genuinely *interacting*, non-Wick-computable dynamics (the *tenth*: the interaction sees δ massively, spectral spread 1.32, yet the mark-axis mirror maps $H(m) \rightarrow H(8-m)$ exactly — all side data mirror-equal; [verification/v529_seam_interacting_toy_fk.py]). Ten side-blind tests, ten kills: the bit is *not* derivable from any tested class and remains genuine discrete input. The flip side is a sharper ontology, not a promotion: the bit is now physically *defined* as the twist-class choice — facets #14/#15 of the (now 15-fold) web are exact equivalences both directions, the flip-axis fraction is a gauge-robust order parameter ($O = \frac{1}{2}$ at $m = 4$, 0 elsewhere), and its η holonomy is in principle interferometrically readable ([verification/v528_seam_bit_twist_class_definition.py]; [C] measurement sketch — a definition replaces no derivation, no marker moves). Since the straddle round there is also the first *positive* selection datum — not an eleventh side-blind test but a different question answered: *if* reflection positivity is demanded of the interacting seam dynamics, the surviving member is unique and it *is* the bit ($\delta = \pi/2$, positive coupling; v534 — toy-level evidence for a dynamical origin, not a derivation; the bit stays formal input).

Where it fires: the straddle law (toy level) — the first live ammunition

The minimal interacting Fidkowski–Kitaev quartic on the 16-Majorana NS seam circle (256-dim, exact) is the first adversarial probe with genuinely non-Wick-computable dynamics — and Kill-Test 2 of WOIT.OS.TWISTOR.01 *fires at toy level*. Θ exists exactly on the interacting model ($\Theta^2 = +1$, $[\Theta, H_g] = 0$, $\Theta\Omega_g = \Omega_g$, clock-tower inversion exact), so Kill-Test 1 does *not* fire; but reflection positivity breaks in the interacting ground state for *every* $g > 0$ — inertia ladder $(37, 0, 0) \rightarrow (33, 4, 0) \rightarrow (31, 6, 0) \rightarrow (29, 8, 0)$, Gibbs state at $\beta = 2$ the same ladder, mechanism *interference* (neither crossing nor half-local terms alone break) — and the failure pattern is a law, not noise: on all 24 tested entries **RP fails exactly on quartet-straddled cuts and stays positive definite on quartet-avoiding ones** (the *straddle law*). This is the honest reading of “first live ammunition”: every previous red-team round ended in

survives/narrowed; here an interacting mechanism actually breaks a load-bearing structure — at toy level, under the mandatory fence (one toy, one interaction class, [C] flat-band parent). The same law is the first *hard selection principle* for the interacting algebra: every candidate $\mathcal{A}_{\text{hol}}$ must protect RP against exactly this mechanism on quartet-straddled cuts — a concrete, checkable hurdle for the β_3/γ stages. Threat and filter are one statement. [verification/v529_seam_interacting_toy_fk.py]

The filter, executed (v534). The follow-up runs the straddle filter as a *selector* on the full interacting mark family (independent equivariant couplings on the four mark quartets, both signs, all admissible cuts — including the straddled clock axes 7/15 of the $\pi/2$ member that the 24-entry law had not covered). Result: reflection positivity survives for *exactly one* member $\times$ sign — $\delta = \pi/2$ with positive coupling, PD on all four cuts over the whole grid $g \in \{1/32..8\}$ with the minimum eigenvalue *lifted away* from the RP boundary; every asymmetric member dies ($s=+$ at $g = 1/32$, $s=-$ from $1/8$). The law refines: asymmetric straddling kills RP, the *symmetric* straddling of the self-mirror member protects it — and the literal leading-order cone formalization is dead (the protection is nonperturbative). The filter is not merely a threat: it is a working selection principle whose unique survivor is the alignment bit — the same member that carries the twist-class order parameter $O = \frac{1}{2}$. Same fence (one toy, one interaction class, [C] flat-band parent). [verification/v534_seam_straddle_cone.py]

Verdict. SURVIVES (NARROWED) for the bit claim: ten side-blind kills leave the bit underivable from every tested class, and the twist-class definition makes its negation measurable without moving any marker (the bit stays formal input; [C]/[O] as ledgered). For the OS/RP structure the honest outcome is FIRES AT TOY LEVEL (FENCED): the free and gauge-fixed layers survive (β_1/β_2 , v522/v524, including the grading, the nonsplit $\mathbb{Z}_8$ tower and the OS quotient surviving $g > 0$ constructively), while the interacting straddle threat is the named residual risk of the whole Woit route — WOIT.OS.TWISTOR.01 stays [O], and any future $\mathcal{A}_{\text{hol}}$ that cannot pass the straddle filter is dead on arrival — while the filter’s own execution (v534) shows a passing class *exists* and is unique: the $\delta = \pi/2$ member. Residual: the β_3/γ stages under the straddle constraint; the bit’s derivation (if any) must come from outside the exhausted side-blind classes — the RP-selection route (demand reflection positivity, obtain the bit) is now the named candidate, toy-fenced.

Outcome

What the red team changed. No target broke on the load-bearing surface. The hardest gate (A) is now *closed modulo cited theorems* on its MMST route SEAM.EQUIV.MMST.01 (the lattice/ S_3 round pins it at every computable level, Lean-formalised, only an external published existence theorem left, v367/v368/v376–v379); four targets (B, D, E, F) *survive once a silent assumption is named*; one (C) survives as worded; and the seam round (G) banks ten side-blind kills on the alignment bit while producing the layer’s first toy-level firing — the straddle law (v529), a named threat that doubles as the first hard selection principle for $\mathcal{A}_{\text{hol}}$, since executed as a selector with a unique survivor: the bit itself (v534). The net effect is not new physics but sharper typing: the package now states *where* each reduction would fail and *which* assumptions are truly necessary, exactly at the weakest logical transitions. Reproduce with `cd verification/redteam && python run_redteam.py`.

The falsifiability surface is machine-checked ([verification/v375_observatory_registry.py]). The frozen prediction registry (freeze_file.csv) is verified by a status-typed CI: every prediction carries a kill criterion (no orphan claims), the headline values re-derive from $\{\varphi_0\}$, and a live scorecard records each one’s distance to the current data — θ_{12} vs the JUNO first run (0.3092 ± 0.0087 , 0.28σ , compatible), θ_{13} vs NuFIT 6.0 NO (2.0σ , the openly-flagged most-tensioned core prediction), β_{rad} vs ACT DR6 (0.37σ , a hint with systematics pending), and $r < 0.036$ (BK18, compatible). The observatory states its own weakest point rather than hiding it.

Follow-up — the verdicts were then used as a worklist [E]
 ([verification/v83_e8net_holomorphic_uniqueness.py])

Two residuals were attacked directly after the red-team pass; the result is recorded in ledger rows GATE.METRIC.04 and CAR.PASCAL.01.

Target A, residual 1 — closed [E]. At level 1 the number of primaries equals $\det(\text{Cartan}) = |\text{fundamental group}|$: $A_3=4$, $D_5=4$, $D_8=4$, $\mathbf{E}_8=1$. So holomorphy (single vacuum sector, μ -index 1) is *necessary* — it excludes the same- c rival $(D_8)_1 = SO(16)_1$ ($c = 120/15 = 8$ but four primaries $1, v, s, c$) — and *sufficient*: a holomorphic $c = 8$ chiral CFT is the lattice theory of an even unimodular rank-8 lattice, of which there is exactly one, E_8 , by the Minkowski–Siegel mass formula ($\text{mass} = 1/|W(E_8)| = 1/696729600$, and $|\text{Aut}(E_8)|$ saturates it). The $(D_5)_1 \times (A_3)_1$ embedding $c = 5 + 3 = 8$ stays *compatibility* (16 primaries vs 1). **Consequence:** the constructive map need only show the seam–Calderón boundary net is holomorphic with $c = 8$ (then E_8 is automatic), so Target A drops from *three* residuals to *two*: (i) boundary-net holomorphy + $c = 8$ ($\Delta_{\text{eff}} > 0$ supports, not proves), and (ii) bulk-reconstruction uniqueness — both still [C]/[O].

Target B — reduced [E]. The “ $K = 2$ Pascal truncation” is not free: $K = (g - 1)/2$ is the Pascal-row midpoint $\sum_{k \leq (g-1)/2} \binom{g}{k} = 2^{g-1}$ (odd g), i.e. the half-spinor split; the carrier is the even Clifford half-spinor $\Lambda^{\text{even}}(\mathbb{C}^5) = 1 + 10 + 5 = 16 = \dim S^+$. So B’s residual reduces to the single standard input “carrier = half-spinor of $\text{Spin}(10)$ ” (the Lean arithmetic core stays [E]).

Target B — a backward certificate ([verification/v219_icosahedral_mckay.py]). The choice $(|\mathbb{Z}_2|, N_{\text{fam}}, g_{\text{car}}) = (2, 3, 5)$ that B attacks is, *given* the E_8 closure, the icosahedron: by the McKay correspondence E_8 is the exceptional top of the finite- $SU(2)$ tower ($2T \rightarrow \hat{E}_6$, $2O \rightarrow \hat{E}_7$, $2I \rightarrow \hat{E}_8$), the binary icosahedral group $2I$ (order $120 = |R^+(E_8)|$) has irrep degrees $\{1, 2, 2, 3, 3, 4, 4, 5, 6\}$ equal to the affine- E_8 Kac marks ($\sum = 30 = h(E_8)$), and it contains the 2, 3, 5 rotation-axis orders. This does *not* break B’s verdict — it is a *backward* certificate (it presupposes E_8), so $g_{\text{car}} = 5$ stays the P2 interface [O]; what it removes is the impression that $(2, 3, 5)$ is an arbitrary triple, since no spherical McKay node sits above the icosahedron.

Typed honestly at this stage: A (i)+(ii) (since merged into the single keystone SEAM.EQUIV.01, whose MMST route SEAM.EQUIV.MMST.01 is now *closed modulo cited theorems*, see Target A’s final form below); Target D’s CP-phase residual (at $\theta_{13} = 0$ the Dirac phase is undefined, so this needs a texture correction); Target E’s reheating/leptogenesis scales and the seam=Planck identification. Target C is a dimensional *floor*, not a gap. **No target was promoted.**

Second follow-up round [E] ([verification/v87_bulk_uniqueness_reduction.py], [verification/v88_cp_phase_audit.py], [verification/v86_nstar_reheating.py])

Target A, residual (ii) — conditionally closed [E]: Target A = ONE residual. Bulk-reconstruction uniqueness is *not* independent of residual (i). By the algebraic-QFT classification of full 2D CFTs over a chiral net (Longo–Rehren; Kawahigashi–Longo–Müger; Bischoff–Kawahigashi–Longo–Rehren), the possible bulks are the haploid commutative Q-systems $\simeq$ physical modular invariants of $\text{Rep}(A)$. For a *holomorphic* net $\text{Rep}(A) = \text{Vect}$, so the bulk pairing is *unique*. The contrast is machine-checked: the same- c rival $SO(16)_1$ (discriminant category $\mathbb{Z}_2 \times \mathbb{Z}_2$, μ -index 4) admits *six* modular invariants — diagonal, $s \leftrightarrow c$ swap, both E_8 -extension invariants $|\chi_0 + \chi_{s/c}|^2$ (the lattice glue $E_8 = D_8 \cup (D_8 + s)$, $h_s=1$ bosonic), and two twisted pairings — so without holomorphy the $c = 8$ bulk is *multiply* ambiguous; with it, trivially unique. Hence the entire Target A reduces to the single theorem “seam–Calderón boundary net is holomorphic with $c = 8$ ” ([C]/[O]; ledger GATE.METRIC.05). The theorem also has an equivalent, more tractable *index* form ([verification/v89_carrier_index_lemma.py], ledger GATE.METRIC.06): by Kawahigashi–Longo–Müger, $\mu_A = [B:A]^2 \mu_B$, the carrier inclusion has Jones index $[(E_8)_1 : (D_5)_1 \times (A_3)_1] = \sqrt{16} = 4 = |\mu_4|$ — the glue-group order *is* the inclusion index — and the extension is the μ_4 simple-current extension (all three glue sectors are $h = 1$ currents; $248 = 45 + 15 + 64 + 64 + 60$). Since an index-4 simple-current extension of the carrier has $\mu = 16/4^2 = 1$, **holomorphy follows from the index statement:** Target A $\Leftrightarrow$ “the seam net contains the carrier net with Jones index 4 as its μ_4 simple-current extension” — both ends constructed objects, the index controllable by Longo theory, the gap supporting finiteness. And *which* extension carries no freedom ([verification/v92_glue_uniqueness.py], ledger GATE.METRIC.07): exhaustive classification of the carrier discriminant form $(\mathbb{Z}_4 \times \mathbb{Z}_4, q = (5x^2 + 3y^2)/8)$ shows the extension tower is *completely rigid* — exactly two Lagrangian glues (the two chiralities, identified by the sheet $\mathbb{Z}_2$; the same two objects as the E_8 -extension invariants above), and exactly one halfway extension, whose induced form $q = \{0, 0, 0, \frac{1}{2}\}$

is the D_8 discriminant form: the $SO(16)_1$ rival is the unique intermediate of the tower, not an arbitrary counter-model. Tower: carrier ($\mu=16$) $\rightarrow SO(16)_1$ ($\mu=4$) $\rightarrow (E_8)_1$ ($\mu=1$, sheet pair) — nothing else exists. Target A is thereby the *bare* index statement.

Target A, final form — the Simple-Current Extension Theorem [E]/[C] ([verification/v154_simple_current_theorem.py]). Assembled into one statement: the carrier net $A = (D_5)_1 \otimes (A_3)_1$ extended by the isotropic glue $L = \langle (1,1) \rangle$ has $[B : A] = |L| = 4$, $c=5+3=8$, $\mu(B)=\mu(A)/|L|^2=16/16=1 \Rightarrow$ holomorphic $\Rightarrow B \cong (E_8)_1$. The *entire* adversarial residual of Target A is now the single seam-side identification “the RP seam–Calderón net *is* A ” — the algebra $(B \cong (E_8)_1)$ is exact and machine-checked (v89/v92/v125/v143/v148). That residual is now sharply located: the analytic core (“the kernel is quasi-free”) *follows* from a free RP bulk — the boundary marginal of a Gaussian bulk is the compression of a contraction (v155), the *same* premise the area law and the EH mechanism use — and the net is *explicitly constructed and verified*: DtN = $|k|$, 16 free Majoranas, currents $248=120+128$, holomorphic character $E_4/\eta^8 = j^{1/3}$ with 248 at level 1 (v156). *Final reduction* (v160–v165): the free-bulk premise is recast as a fixed-point theorem (quasi-free $\Rightarrow$ all higher cumulants $\kappa_{2n}=0$, v160); the infinite Schwinger cone is *eliminated* — by Bogoliubov second quantisation the cone gap equals the one-particle gap $(2/3)^6$ (v161), whose value is forced by μ_4 -equivariance (v162); and the remaining 16-dim identification factors into the two *already-open* items — A_2 net existence (an assembly of established net constructions, [C]) and $R_1 = \text{GATE.QGEO}$ (“seam boundary = $\mathbb{P}^1 \setminus \mu_4$ ”) — with the irreducible core $\{\pi, v_{\text{geo}}\}$ a theorem (v165). So Target A carries *no new independent open content*: it is a unification, not an unconditional proof. The two structural residuals are then discharged to a theorem (v175): the CAR second-quantisation functor $\Gamma(t) = \bigoplus_m \Lambda^m(t)$ makes full-cone reflection positivity reduce *for every* m to the one-particle contraction (verified on the complete 2^{16} -dim Fock space), and A_2 is an assembled, verified $(E_8)_1$ certificate; net existence and full-cone RP become [E]. Target A’s single residual is therefore the seam realisation, the named keystone **SEAM.EQUIV.01** — *the raw RP seam is the holomorphic $(E_8)_1$ net at $\tau=i$* — whose MMST route **SEAM.EQUIV.MMST.01** is now *closed modulo cited theorems* (the parent stays [O]): the explicit lattice model (v367/v368) and the S_3 stack ($c=8$, 248 currents/1 primary, torus GSD=1, RP; v376–v379) pin it at every computable level and it is Lean-formalised as **FORM.SEAM.MMST.01**, so the *only* residual that stays [O] is the abstract continuum existence of the scaling limit (the cited MMST/OS theorems, v336); its conformal-deck premise **QGEO.SYM.01** is its *downstream* definitional face, Lean-pinned (**FORM.SEAMEQUIV.01**, v308) to named cited steps plus the derived Recovery gap. The sprint-by-sprint reduction (v176–v302) is recorded in the changelog, not replayed here.

Target A — the keystone residual narrowed from ten further directions [E]/[C] (not promoted). A later convergent round drove the *same* single residual (the cited continuum existence, v336) down further from independent sides, while leaving **SEAM.EQUIV.01** [O]. (i) The seam edge already carries the conformal data of a chiral $c=\frac{1}{2}$ Majorana CFT and reproduces the $c_-=8$ free-fermion scaling limit on the explicit collar ([verification/v444_seam_edge_scaling.py]). (ii) The rigidity step is upgraded from *permit* to *force*: commuting with the order-4 μ_4 clock is *exactly* μ_4 -block-diagonality (entrywise iff, uniform in N ; exact commutant dimension; the order is the discriminator), kernel-hardened in Lean axiom-free as **FORM.SEAM.RIGIDITY.FORCING.01** ([verification/v445_seam_rigidity_forcing.py]). (iii) That clock-invariance is then *derived*, not assumed: the OS canonical transfer of μ_4 -symmetric RP data inherits the clock symmetry and its modular Hamiltonian commutes with it ([verification/v446_seam_clock_invariance.py]), so the residual drops from an operator condition to the structural μ_4 -symmetry of the seam RP data (**QGEO.SYM.01**). (iv) The number $c_-=8$ is re-derived from a logically independent (topological) observable — the $M=1$ collar has integer Chern $C=+1$ (with $C=0/C=-1$ controls) and the bulk–edge correspondence gives $c_-=16 \cdot \frac{1}{2} \cdot 1=8$ ([verification/v447_seam_edge_chern.py]). (v) The one *external* MMST fact is pinned a further order: the edge state is quasi-free ($G^2=G$) and the four-point cross-ratio Cauchy-converges to the conformal value $\sin(5\pi/12)/\sin(\pi/12)=2+\sqrt{3}$ (against an off-scale gapped bulk control) ([verification/v448_seam_edge_fourpoint.py]). (vi) The MMST fact is pinned at the level of *uniformity*: two *independent* edge four-point cross-ratios converge to distinct conformal values ($2+\sqrt{3}$ and 2) at one N -independent $1/N$ rate ($|Q(N)-Q_\infty| \cdot N \leq 4$ for both) — the empirical hallmark the abstract theorem asserts ([verification/v449_seam_mmst_uniform.py]). (vii) The number $c_-=8$ is read a *third* independent way — the Calabrese–Cardy entanglement law gives c per Majorana = $\frac{1}{2}$ ($c_{\text{Dirac}}=1.000$ vs a saturating gapped control), so $c_-=16 \cdot \frac{1}{2}=8$ ([verification/v450_seam_edge_entanglement.py]). (viii) The edge CFT is named by its *operator content*: the Cardy finite-size spectrum reproduces the exact Ising primaries $h_\sigma=\frac{1}{16}$ and $h_\epsilon=\frac{1}{2}$ (L -independent), so the edge is the chiral Ising/Majorana minimal model $M(3,4)$ ([verification/v451_seam_edge_cardy_tower.py]). (ix) The $(E_8)_1$ identity is pinned on

the *torus*: the character $\chi=E_4/\eta^8$ is S -invariant (one primary), T -multiplies by $e^{-2\pi i/3}=e^{-2\pi ic/24}$ ($c=8$) and is holomorphic ($c \equiv 0 \pmod 8$) ([[verification/v452_seam_e8_modular.py](#)]). (x) The last structural premise is *derived*: the four Gauss–Bonnet marks are μ_4 (roots of z^4-1), the form basis is μ_4 -graded and the clock $z \rightarrow iz$ fixes their cross-ratio, so the residual QGEO.SYM.01 of (iii) follows from marks= μ_4 plus the existing QGEO.REALIZE.01 — no new premise ([[verification/v453_seam_mu4_from_marks.py](#)]); the integer backbone of (iv)–(ix) ($c_-=8$, the Chern integers, the order-3 T -phase) is kernel-hardened in Lean axiom-free as FORM.SEAM.EDGE.CHERN.01. (xi) The certification package itself is re-founded ([[verification/v469_seam_crossedproduct_route.py](#)]): the 128-spinor extension is the *local* $\mathbb{Z}_2$ simple-current crossed product — the locality integer $h_s(SO(16)_1)=16/16=1 \in \mathbb{Z}$ is exactly the Longo–Rehren criterion (LR 1995; Böckenhauer 1996; Böckenhauer–Evans 1998; KLM $\mu=4/2^2=1 \Rightarrow$ holomorphic) — so the extension leg rests on 1995–2001 *peer-reviewed* subfactor theory with the AGT/AMT preprint route demoted to an independent second witness; and the realisation input is reduced from model fiat to invariants (R1': quasi-free, gapped, class D, $c_-=8$ from P_1 ; computed $|C|=1$, $\nu=16$, the Kitaev sixteen-fold-way class), Lean-carried as the parallel derivation `seamResidualClosed'`. **None of these promotes the keystone**: the *only* thing that stays [O] is still the abstract continuum-existence theorem (v336); everything else is now corroborated, derived or kernel-checked.

Target D — the CP-phase residual quantified [E] (not closed). With the frozen leading reading $\delta = \pi/3 + 3\lambda_C^2 = 68.65^\circ$: pull vs $\gamma_{\text{PDG}} = 65.7^\circ \pm 3.0^\circ$ is $+0.98\sigma$ (survives). Audit, *not* promoted: the data central value sits 0.07° from the alternative coefficient reading $\pi/3 + 2\lambda_C^2$ ($|\mathbb{Z}_2|$ instead of N_{fam}) — a textbook look-elsewhere trap; the registry keeps coefficient 3 (REG.FREEZE.01). Decision threshold: the readings differ by $\lambda_C^2 = 2.88^\circ$, i.e. 3σ -distinguishable once $\sigma_\gamma \leq 0.96^\circ$. The J -inversion is flagged magnitude-contaminated (source-scale s_{23} vs M_Z). Ledger FLAV.CP.01.

Target E — one flagged scale computed [C]. The reheating temperature is no longer a silent input: $M_{\text{scal}} = c_3^{7/2} \bar{M}_{\text{Pl}}$ [E] + standard physics give $T_{\text{reh}} = 9.6 \times 10^9$ GeV and $N_*(0.05/\text{Mpc}) = 51.4$ [C] (ledger COSMO.NSTAR.01); honestly recorded: this slow Higgs-channel point is A_s -disfavoured (-11.4σ ; the measured A_s requires near-instantaneous reheating), so the point is conditional on the decay channel and the frozen band [50, 60] stays the surface of record; the leptogenesis scale and the seam=Planck identification remain typed inputs. **Again: no target was promoted; one residual was merged (A), one was quantified (D), one input was computed (E).**

Target D — the CP residual geometrically located (not closed) [E]/[C]

The follow-up round only *quantified* the CP residual; this round *locates* it geometrically, in three consistent readings, without promoting it. (i) **A channel typing** ([[verification/v227_degree_exponent_channel_split.py](#)]). The E_8 split $248 = 120 + 128$ is a magnitude/phase split: the exponents sum to $120 = |R^+(E_8)|$ (the *magnitude* channel, where the (ratios, product) $\Rightarrow v_{\text{geo}}$ bijection lives), the fundamental degrees sum to $128 = \text{rank } E_8 \cdot \dim S^+$ (the *phase/glue* channel). The un-covered CP phases of Target D are exactly the 128 phase channel — a clean home, not a new gate; whether 128 also hosts a dark sector is Frontier, *not* asserted (this replaces the over-strong “ S^- is dark matter” reading, now sharpened by v430: the other side S^- is matched $\{2, 128\}$ structure, forced-disjoint from the unmapped $\{12, 14, 18, 20, 24\}$). Strengthened on the fiber by the T177 mirror ([[verification/v561_cp_channel_identities.py](#)]): the typing has an *exact* hexagonal realisation — $\sum_k \rho^{d_k} = 4\rho$ ($\arg = \pi/3$ exactly) against $\sum_k \rho^{m_k} = 4$ ($\arg = 0$ exactly) — with a parity typing forced by integer parity (all degrees even $\Rightarrow$ the phase channel is *sheet-even*; all exponents odd $\Rightarrow$ the magnitude channel is *sheet-odd*), so the phase-channel angle is *sheet-blind*: it carries $\pi/3$ but cannot distinguish quark from lepton (the channel typing stays [C]). (ii) **A hexagonal modulus** ([[verification/v220_cp_hexagonal_modulus.py](#)]). The seam deck is the *square* modulus ($j=1728$, $\mathbb{Z}/4$ clock; the kill-test selects it), which is real and carries no phase; its exceptional partner is the *hexagonal* modulus $j=0$ ($\tau=\rho=e^{i\pi/3}$, $\mathbb{Z}/6$, CM by $\mathbb{Z}[\omega]$). The frozen leading reading $\delta = \pi/3 + 3\lambda_C^2 = 68.65^\circ$ has leading term $\pi/3 = \arg \rho$, the hexagonal CM phase; CP lives in the $\mathbb{Z}/6$ phase *fiber* over the $\mathbb{Z}/4$ seam, not in the deck. The T177 mirror *scopes* the reading ([[verification/v561_cp_channel_identities.py](#)]): only the leading term $\arg \rho = \pi/3$ is character-covariant under the parity/sector involution; the full $\delta = \pi/3 + 3\lambda_C^2$ has *no* character (a CP-even correction added to a CP-odd angle) and is a branch/frame-relative quantity — the frozen reading itself is untouched (REG.FREEZE.01). (iii) **An orientation** ([[verification/v225_dual_normal_frame.py](#)]). The dual normal frame $d = a^\top R^{-1} = -\frac{1}{2}(1, 1, -2)$ (Narai/horizon normal) and $n = (5, -9, 6)$ (first-generation selector, $n^\top R = (8, 0, 0)$) has oriented volume $\det(\mathbf{1}, d, n) = 21 = N_{\text{fam}} \cdot 7$, and rotating by the hexagonal unit gives $\text{Im } \det(\mathbf{1}, d, \rho n) = 21 \sin(\pi/3) \neq 0$ — the CP orientation *reading*, sitting exactly where the magnitude bijection fails. Narrowed sharply by the

T177 mirror ([[verification/v561_cp_channel_identities.py](#)]): that reading is homogeneous of degree +1 in the frame normalisation, so the *number* $21 \sin(\pi/3)$ is frame data, not an invariant; dividing by the oriented volume removes the frame identically ($\text{Im}(\rho \det)/\det = \text{Im} \rho$ exactly), so the invariant core is the fiber datum $\text{Im} \rho = \sin(\pi/3)$ and the *sign* is a declared orientation. *Decided by the T178 frame door* ([[verification/v564_cp_fiber_pin.py](#)]): the removal is not an accident of the functional — in the metric-free (natural) calculus the frame *cannot* couple at all (one bracket generator, every scale-free ratio constant: a theorem), and every metric-declared coupling is non-canonical (the normalisers $\langle d, n \rangle = 8$, $\langle 1, n \rangle = 2$, $\langle n, n \rangle = 142$ give the pairwise-distinct frame factors $21/8$, $21/2$, $441/142$, and a whole continuum q_t — normaliser-relative frame data, none of them claimed); moreover T177's two-class degeneracy is *explained* (the single-sine lemma: on crystallographic fibers $n \in \{1, 2, 3, 4, 6\}$ — exactly the two compiler clocks μ_4 , μ_6 — every carrier emits the same sine, so value-based pinning is impossible by theorem) and *resolved conditional on the classical anchor*: the Jarlskog invariant carries the full character tuple $(-1, -1, -1)$, matching the sheet-odd orientation class only — the unique anchor-matching canonical carrier is the fiber-pure $\text{Im} \rho = \sin(\pi/3)$ (selected, not derived). All three are [E] geometry; the physical CP *derivation* stays the Target-D [C] residual (the seam deck is not re-selected, the registry δ is untouched). (iv) **One hexagonal unit, two sheets** ([[verification/v231_cp_mu6_phases.py](#)]). The reduction is sharper still: *both* CP phases are μ_6 powers of the single hexagonal CM unit $\rho = e^{i\pi/3}$ — $\delta_{\text{CKM}}^{\text{lead}} = \arg(\rho^1) = \pi/3$ (quark) and $\delta_{\text{PMNS}} = \arg(\rho^4) = 4\pi/3$ (lepton, the neutrino phase lattice) — with $\rho^4 = -\rho$, so $\delta_{\text{PMNS}} = \delta_{\text{CKM}}^{\text{lead}} + \pi$ is the quark phase times the $\mathbb{Z}_2$ sheet ($\rho^3 = -1$); the frame orientations of (iii) are correspondingly sheet-flipped ($\pm 21 \sin(\pi/3)$). So Target D's CP sector is *not* two independent phase inputs but *one* hexagonal unit read on the two sheets — the magnitude residual $3\lambda_C^2$ (quark, v88) and the physical derivation stay [C], but one of the two phase inputs is removed. (v) **The universal triality phase** ([[verification/v233_cp_triality_phase.py](#)]). In the canonical $\mu_6 = \mu_3(\text{family/triality}) \times \mu_2(\text{sheet})$ factorisation, $\rho = \omega^2(-1)$ with $\omega = e^{2\pi i/3}$ the $\mathbb{Z}_3$ triality centre (the $2/3$ cusp weight); since $\rho^4 = \rho^1 \rho^3$ with $\rho^3 \in \mu_2$, the quark (ρ^1) and lepton (ρ^4) phases have the *same* family/triality class and differ *only* by the sheet. So the CP phase *is* the universal triality phase, sheet-split — not even a free choice of μ_6 power: the power choice of (iv) is removed, and ω is the μ_3 factor of the order-30 (2, 3, 5) Coxeter/Milnor monodromy that builds the hull (v232). The physical CP derivation still stays [C]. (vi) **The lock as a pre-registered kill test** ([[verification/v320_galois_cp_relation.py](#)]). Because $\delta_{\text{PMNS}} = \delta_{\text{CKM}}^{\text{lead}} + \pi$ is now *forced* (one unit $\rho = \zeta_6$, the μ_4 -power ρ^4 , the $\mathbb{Z}_2$ sheet $\rho^3 = -1$), it is a falsifiable cross-prediction, not a free input: a measured δ_{PMNS} robustly away from 240° at DUNE/Hyper-K (beyond the sub-leading budget, $> 3\sigma$) kills the Galois-CP organisation. The relation is [E], the physical phase reading stays [C], the kill test is [X].

Global look-elsewhere closure [E] ([[verification/v100_numerology_null_mc.py](#)])

The per-target look-elsewhere traps recorded above (the $\pi/3 + 2\lambda_C^2$ alternative in Target D, the $\ln(46080)$ scale reading destroyed in [[verification/v99_koide_flow_time.py](#)]) are local instances of the global adversarial question: *could the whole scorecard be formula-fishing?* The numerology null test answers it with an explicit, declared null model run in adversarial mode: the *entire* complexity-matched formula grammar (which provably contains every scored TFPT formula — the null space includes the theory) is enumerated against conservative data windows. Joint formula-fishing probability $\prod_i p_i = 10^{-25.8}$ over the 13 scored observables; all 94 500 $F_{U(1)}$ variants solved, exactly one hits CODATA ($p_\alpha \leq 1.1 \cdot 10^{-5}$); total $\leq 10^{-30.7}$. The test is falsifiable in both directions: the negative controls (seed perturbation 1%/10% collapses TFPT's own hit count $13 \rightarrow 9/6$; data shuffling $\rightarrow 1.2$) prove it has power, and the census is stable under budget/window variation (< 2 orders). Honest limit, stated as such everywhere: this is a null-model rejection *conditional on the declared grammar* — no statistical test certifies a theory; the tension observables keep their large windows and contribute nothing to the number. Details and the grammar definition: the audit note (tfpt_3) and the module docstring.

Structural	firewall	+	out-of-sample	seed	cross-validation	[E]
([verification/v305_witness_independence.py],			[verification/v306_seed_crossval.py],			
[verification/v465_seed_crosssector_joint.py],			[verification/v466_seed_leptonmass_channel.py])			

The global null test answers “is the *scorecard* fishing?”; two companions answer the deeper adversarial worries “is the same integer counted many times as if independent?” and “is one shared seed dressed up as many predictions?”. **Witness economy** ([[verification/v305_witness_independence.py](#)]): every headline integer ($|\mu_4|=4$, $g_{\text{car}}=5$, $|\mathbb{Z}_2|=2$, $N_{\text{fam}}=3$, $\dim S^+=16$, $\text{rank } E_8=8$, 240 , 248 , $|2I|=120$, $h=30$, $|W(D_5)|=1920$, $\Omega_{\text{adm}}=48$, the gap exponent 6) is a documented image of the *single* anchor $a=(1, 1, 2)$

with π the only transcendental primitive, so the recurring $2/3=|\mathbb{Z}_2|/N_{\text{fam}}$ and $(2/3)^6$ are *one* ratio, not many — 13 atoms from 2 free primitives ($6.5\times$ compression made explicit). The E_8 spine is genuinely overdetermined (≥ 3 structurally distinct derivations sharing only leaf inputs), while the pure-seed readouts are single-witness, so their joint surprise is scored jointly by **v100**, never multiplied; the negative control shows α^{-1} 's “137” is *foreign* to the anchor family (an independent witness via the $F_{U(1)}$ cubic, **v3**). **Seed cross-validation** ([[verification/v306_seed_crossval.py](#)]): treating the axiom seed φ_0 adversarially as a free dial and back-solving it from each of the five seed-grammar observables gives an error-weighted mean reproducing $\varphi_0 = 1/(6\pi) + 48c_3^4$ to 0.04%; leave-one-out predicts every held-out observable within 3σ *out of sample* (the most-tensioned hold-out is $\sin^2 \theta_{13}$, $\sim 2\sigma$), and a data $\leftrightarrow$ formula shuffle explodes the cross-validation χ^2 by $> 100\times$ (the test has power). **Cross-sector slice** ([[verification/v465_seed_crosssector_joint.py](#)]): the minimal externally-runnable form isolates the sharpest shared-origin signature — a correlation no non-shared-origin model predicts — the CMB EB/TB birefringence angle β and the quark Cabibbo angle $\lambda_C=|V_{us}|$ back-solve to the *same* seed within $0.01\sigma_{\text{comb}}$, the three-observable one-parameter fit has $\chi^2/\text{dof}<1$, and the θ_{13} channel is *declared* excluded (adding it raises χ^2 by ~ 4 , the documented $\sim 2\sigma$ pressure of **v328**), so this is the honest θ_{13} -independent statement, not a hidden dilution; a data $\leftrightarrow$ formula shuffle explodes the cluster χ^2 by $> 5\times 10^4$. It upgrades no prediction (a subset of **v306**). **Mass-sector channel** ([[verification/v466_seed_leptonmass_channel.py](#)]): the same seed is read a fourth way, from a *new* experimental sector — the charged-lepton mass ratio $m_e/m_\mu=\frac{12}{17}\varphi_0^2$ back-solves the axiom seed to -0.11% , so the empirical over-determination now spans *five* unrelated measurement sectors (neutrino mixing, CMB, quark mixing, and now charged-lepton *mass* plus the corroborating heavy-quark m_t/m_b , -0.23%); the six-channel cluster keeps $\chi^2/\text{dof}<1$. This is an *empirical* strengthening only: the formula is a source-scale ratio whose source $\rightarrow$ pole transport is the seam-gapped correction class (**v393**) — for m_e/m_μ it largely cancels, so the channel is added as a *consistent new sector*, not a sharper pin — and φ_0 stays fixed by the axioms (a subset of the single seed, **v305** multiplicity 1). All three stay **[E]** as structural/statistical checks, conditional on the same declared grammar. **No-golden-dynamics rule** ([[verification/v305_witness_independence.py](#)]): the golden ratio $\phi = 2\cos(\pi/5)$ lives in the *static* carrier field $\mathbb{Q}(\sqrt{5})$ (the $\mathbb{Z}/5$ hand; minimal polynomial $x^2 - x - 1$, discriminant 5 non-square), while every *dynamic* rate is rational ($\mathbb{Q}$, the $\mathbb{Z}/6 = \text{sheet}\times\text{family}$ hand, e.g. the recovery factor $(2/3)^6$) — so a dynamic rate “explained” by ϕ is a cross-field false friend, and CP phases and recovery rates must run on the order-6 hand (the field firewall of **v314/v315**).

The reverse numerology audit, and what the golden ratio means **[E]/[O]** ([[verification/v354_e8_reverse_audit.py](#)])

The witness-economy box answers “every readout maps onto E_8 — is the *forward* map fishing?”. The deeper adversarial question runs the *other* way: E_8 is a rich object, so how much of its structure maps onto *nothing*, and could the theory be quietly mining that richness? **The reverse audit**. Of E_8 's eight Casimir invariant degrees $\{2, 8, 12, 14, 18, 20, 24, 30\}$ (exponents $+1$), exactly *three* feed a *primary* physical-constant readout — degree 2 (the quadratic/metric), degree 8 (the rank $\rightarrow c_3 = 1/(8\pi)$), degree 30 (the Coxeter number $\rightarrow g_{\text{car}} = 5$ and the order-30 cycle) — and the other *five* carry no primary readout: $\{12, 14, 18, 20, 24\}$. (They do *coincide* with internal structural atoms — A_3 roots, $\dim G_2$, $\det L$, $|W(A_3)|$, the anchor power sum — as **v66** records; but those per-degree coincidences are mostly *unforced*, the numerology temptation that the bandwidth search **v355** declines, see the next box.) So $3/8$ are primary readouts, $5/8$ hull overhead. **Economy, not promiscuity**. A *small fixed* set of E_8 data (rank 8, $h=30$, $\det K=1$, the $D_5\oplus A_3$ sub, the 16 half-spinor) generates the *entire* readout set (gauge group, three families, hypercharges, α^{-1} , flavour, scales); the higher Casimirs are never reached into to fit. A numerological use of E_8 would be *promiscuous* (mine the rich structure for matches); TFPT is *economical* (few invariants, many readouts) and never *reaches into* the higher Casimirs to fit; where those non-readout degrees do carry content it is *forced* internal structure (the spine and the flavour det-ladder, **v431**), not a mined coincidence — the anti-numerology signature is precisely this economy with a forced, *non-readout* remainder. **Why the golden ratio is not numerology**. $\phi = 2\cos(\pi/5)$ sits in exactly that unmapped part: it appears only in *internal* structure (the affine- E_8 attractor eigenvalue **v312**, the icosian lattice **v348**) and in no *frozen physical-constant* readout — the sole, *conditional* exception being the axion misalignment spine angle, where $\theta_i = 3\pi/5$ is the regular-pentagon interior angle and $\cos \theta_i = -1/(2\phi)$ ([[verification/v429_axion_pentagon_phi.py](#)]; **[C]**, branch-level, not a frozen prediction). Numerology means a number *fitted to an observation*; ϕ is fitted to nothing — it is the algebraic signature of the 5-fold ($h=30=2\cdot 3\cdot 5$), the carrier *announcing* that it is icosahedral, and even its one physical-ish appearance

is forced geometry ($\theta_i =$ the $g_{\text{car-gon}}$ angle), not a match to a measured constant. **Honest residual.** The 5/8 hull overhead is exactly that: E_8 is the minimal even-unimodular *hull* (consistency container, $\det K=1$), not the world group, so the higher Casimirs are by-products of unimodular closure — where a future higher-invariant readout could live, or genuine over-capacity (now sharpened: the five non-readout degrees are *not* diffuse slack but the forced two-family ladder $6 \cdot \text{spine} \sqcup \text{det-ladder}$, v431). Stated plainly, not hidden.

Bandwidth search of the unmapped region: the disciplined decline [E]/[C]/[O]
([verification/v355_e8_unmapped_bandwidth.py])

The obvious next move — “then *search* those five degrees for hits we overlooked” — is itself the trap: a structure as rich as E_8 returns a numerical match for almost anything. So the scan is run with a *strict discriminator*: a candidate counts only if it is a *forced* identity (a theorem about E_8), never a coincidence. The honest yield is mostly a *decline*, and that decline is the result. **Forced, set-level (the genuinely overlooked structure — collective, not per-number).** $\sum \deg = 128 = \dim S^+$, the spinor half of $248 = 120 + 128$ (a theorem: $\sum \deg = \# \text{positive roots} + \text{rank} = 120 + 8$) — the invariant “budget” of E_8 *equals the fermion content the carrier reads*; $\sum \exp = 120 = \# \text{positive roots}$; $\prod \deg = |W(E_8)| = 696\,729\,600$; and the exponents $\{1, 7, 11, 13, 17, 19, 23, 29\}$ are exactly the $\varphi(30) = 8$ totatives of 30 (v223), so the unmapped degrees’ exponents $\{11, 13, 17, 19, 23\}$ are part of that forced set, not free. **Suggestive but not a theorem (the discriminator at work).** $12 = h(E_6)$, $18 = h(E_7)$, $30 = h(E_8)$ are all degrees of E_8 (the $2T/2O/2I$ McKay ladder, v219); but “ $\deg(E_8)$ contains its subalgebras’ Coxeter numbers” is not a general theorem, so this appealing pattern is flagged [C], not claimed. **Declined — the numerology temptation.** The per-degree atom matches $14 = \dim G_2$, $20 = \det L$, $24 = |W(A_3)| = 4!$, $12 = |R(A_3)|$, $18 = p_4(a)$ (which v66 already flagged as fittable for 12, 18) each admit ≥ 2 readings from the frozen atoms with no forced selector, and the lone extra prime in $|W(E_8)| = 2^{14}3^55^27$ “=” the scalaron exponent is one of many readings of 7 — all declined. **Reconciliation.** v66 (the eight degrees coincide with the load-bearing integers, expected since E_8 is the hull) and the reverse audit (v354: only 2, 8, 30 are primary readouts) are *both* right once forced set/subchain structure is separated from unforced per-degree coincidence. The search finds *no new forced physical hit*; the disciplined decline *is* the anti-numerology outcome.

The “other side” of the seam vs the unmapped region [E]/[O]
([verification/v430_other_side_reverse_audit.py])

A sharp follow-up runs the audit through the *double cover*: the seam has an *other side* — the one-sided $\mathbb{Z}_2$ collar and the conjugate half-spinor S^- — so *is the leftover where the second sheet lives?* The answer is a disciplined **negative**, the sheet/deck complement of v354/v355. **The deck is the degree-2 invariant (matched).** The one-sided cover contributes $|\mathbb{Z}_2| = 2$ (the $\frac{1}{2}$ in $c_3 = 1/(|\mathbb{Z}_2| 4\pi) = 1/(8\pi)$, v58/v216), and $2 = \min \deg(E_8)$ is the quadratic/metric — one of the *three matched* primary readouts $\{2, 8, 30\}$, never one of the unmapped $\{12, 14, 18, 20, 24\}$. **The two sheets are the 128-spinor (matched, collective).** $\dim S^+ = \dim S^- = 16$ and the spinor block is $128 = \text{rank } E_8 \cdot \dim S^+ = 2^{\text{rank}-1} = \sum \deg$, the spinor half of $248 = 120 + 128$; the other side S^- enters 248 as the conjugate $(\overline{16}, 4)$ ($128 = 2 \cdot 64$), *not* a spare singlet. **Forced-disjoint.** The sheet/deck integer set $\{|\mathbb{Z}_2|, \dim S^+, \dim(S^+ \oplus S^-), 128\} = \{2, 16, 32, 128\}$ has *empty* intersection with the unmapped degrees $\{12, 14, 18, 20, 24\}$; its only contact with the degree alphabet is the matched 2 (and the collective budget $128 = \sum \deg$). **Disciplined decline [O].** Each unmapped degree, “explained” from the sheet atoms, needs an *unforced* coefficient and admits ≥ 2 readings — exactly the mining the v355 discriminator declines, so there is *no* forced sheet $\rightarrow$ unmapped-degree identity. **Consequence.** This consolidates the S^- -dark-matter downgrade (v227) and the no-spare-singlet WIMP no-go of the frontier E_8 scan: the other side is matched conjugate matter, the five unmapped degrees are the hull overhead, and the two do not meet. The only live “other side” question — whether the 128 phase/glue channel hosts a dark sector (v227) — concerns *matched* structure, not the unmapped five.

The unmapped five are forced structure, not diffuse overhead: the degree ladder [E]/[C]
([verification/v431_e8_degree_ladder.py])

The reverse audit (v354/v355) and its sheet/deck complement (v430) leave the five degrees $\{12, 14, 18, 20, 24\}$ with no *primary* readout. The structural follow-up asks the sharper question — *are they diffuse hull overhead, or forced internal structure?* — and the answer is the latter, an *exact*

two-family decomposition $\deg(E_8) = 6 \cdot \text{spine}\{2, 3, 4, 5\} \sqcup (\{2\} \cup \text{det-ladder}\{8, 14, 20\})$. **The split is forced by 30.** The E_8 exponents are exactly the $\varphi(30) = 8$ totatives of $h(E_8) = 30 = 2 \cdot 3 \cdot 5$ (v223), hence coprime to 6, hence $\equiv \pm 1 \pmod 6$ — so the degrees ($= \text{exp} + 1$) occupy *only* the residue classes $\{0, 2\} \pmod 6$: two arithmetic progressions, forced by the prime content of $30 = 2N_{\text{fam}}g_{\text{car}}$. **The $6k$ family is $6 \times$ the spine.** Degrees $\equiv 0 \pmod 6 = \{12, 18, 24, 30\}$, divided by 6, give $\{2, 3, 4, 5\}$ — the v91 spine $\{e_3, p_0, e_1, e_2\}$ of the anchor $a = (1, 1, 2)$; $30 = 6g_{\text{car}}$ is the matched Coxeter readout, 12, 18, 24 were “unmapped.” **The $6k+2$ family is the quadratic Casimir plus the flavour det-ladder.** Degrees $\equiv 2 \pmod 6 = \{2, 8, 14, 20\}$; here $\{8, 14, 20\} = (\det R, \det C, \det L)$ is the winding line $\det M(s, 0) = 6s + 8$ of the flavour diamond (v135), with $C = R + Q \text{diag}(1, 0, 0)$, and 2 is the matched metric Casimir ($8 = \det R$ is over-determined: also rank E_8). **18 is not a holdout.** $18 = 6 \cdot 3 = 6N_{\text{fam}}$ is the spine-family member $6p_0$, not on the det-ladder ($6s + 8 = 18$ has no integer s): it changes *family*, not status, so *no* genuinely orphan degree remains. **Special to E_8 .** The clean $\{0, 2\}$ -mod-6 split needs both $6 \mid h$ and $\text{exponents} = \text{totatives}(h)$; among $\{A_4, D_5, E_6, E_7, E_8, F_4, G_2\}$ it holds for E_8 but *fails* for E_6, E_7, D_5 , so the decomposition is not a generic mod-6 triviality. **The honest v355 line.** The *arithmetic* decomposition is exact [E]; but the *functorial* claim “the $6k+2$ Casimir stratum *is* the flavour sheet operators” needs a representation-theoretic map and stays [C] (degrees 12, 24 still admit two canonical readings each: 6-spine vs $|R(A_3)|/|W(A_3)|$). So this is a *structure* theorem — it reclassifies the 5/8 overhead of v354 to *zero* unfamilied degrees — not a forced functorial flavour map, and it closes no gate.

The two structural integers and all of E_8 are fixed by the degree multiset [E]
 ([verification/v437_e8_degree_joint.py])

A final consolidation of the degree audit (of v6/v66/v355/v431), staying strictly inside the v354/v355 discipline — it adds *no* new per-degree coincidence, only one joint lock. **All of E_8 is fixed by its degrees.** From $\deg(E_8) = \{2, 8, 12, 14, 18, 20, 24, 30\}$ alone: rank = #deg = 8, $h = \max \deg = 30$, #positive roots = $\sum \text{exp} = 120$, #roots = 240, $\dim E_8 = 240 + 8 = 248$, $|W(E_8)| = \prod \deg = 696\,729\,600$, $\sum \deg = 128 = \dim S^+$ — every E_8 integer the compiler uses is a function of the degrees. **The joint quadratic.** The two structural integers ($g_{\text{car}}, N_{\text{fam}}$) are the *unique* root pair of $x^2 - (\text{rank } E_8)x + (h/2) = x^2 - 8x + 15 = (x-3)(x-5)$: the sum of roots is the rank-fill rank $E_8 = 8$ (v6) and the product is $h/2 = 15 = g_{\text{car}}N_{\text{fam}}$ ($2g_{\text{car}}N_{\text{fam}} = h$, v431), and $\{3, 5\}$ is the unique positive-integer factor pair of 15 summing to 8 — so $g_{\text{car}} = 5$ and $N_{\text{fam}} = 3$ are forced *together* by two E_8 degree-invariants, not one at a time. The three primary-readout degrees $\{2, 8, 30\}$ are exactly $\{\min \deg, \text{rank}, \max \deg\}$, the invariants every simple Lie algebra distinguishes. Mirrored in Wolfram; closes no gate.

Table 1: The eight E_8 Casimir invariant degrees, audited ([verification/v66_e8_casimir_degrees.py], [verification/v354_e8_reverse_audit.py], [verification/v355_e8_unmapped_bandwidth.py]; the sheet/deck complement [verification/v430_other_side_reverse_audit.py]; the forced two-family structure [verification/v431_e8_degree_ladder.py]). Only three feed a *primary* physical-constant readout; the other five carry no primary readout but are *not* diffuse overhead — they are the forced two families $6 \cdot \text{spine}\{2, 3, 4, 5\}$ and the det-ladder $\{8, 14, 20\}$, the residue classes $\{0, 2\} \pmod 6$ forced by $h = 30$ (v431). The genuinely forced collective content: $\sum \deg = 128 = \dim S^+$, $\prod \deg = |W(E_8)|$, exponents = totatives of 30.

deg	exp	TFPT reading	forced?	primary readout?
2	1	$ \mathbb{Z}_2 $	forced	yes: metric
8	7	rank E_8	forced	yes: c_3
12	11	$h(E_6) = R(A_3) $	suggestive	no; 6·2 spine
14	13	$\dim G_2$	declined	no; det C ladder
18	17	$h(E_7) = p_4(a)$	suggestive	no; 6·3 spine
20	19	$\det L$	declined	no; det L ladder
24	23	$ W(A_3) = 4!$	declined	no; 6·4 spine
30	29	$h(E_8) = 2 \cdot 3 \cdot 5$	forced	yes: g_{car}
<i>collective (forced):</i> $\sum \deg = 128 = \dim S^+$, $\prod \deg = W(E_8) $, exp = totatives of 30				
<i>two families (forced, v431):</i> $\{0, 2\} \pmod 6$; $6k/6 = \text{spine } \{2, 3, 4, 5\}$, $6k+2 = \{2\} \cup \text{det-ladder } \{8, 14, 20\}$				




Figure 1: The seed-hyperplane test ([`verification/v306_seed_crossval.py`]): each scorecard observable is *independently* inverted to the latent seed φ_0 (with its measurement error). All five project onto the one axiom value $\varphi_0 = \frac{1}{6\pi} + 48c_3^4 = 0.053172$ (green), and their error-weighted mean reproduces it to 0.04% — one real seed, not five fits. The honest pressure point is $\sin^2 \theta_{13}$ ($\sim 2\sigma$ low, a *separate* carrier-trace channel, **v268**; across the *mixing* channels it joins θ_{23} and $|V_{cb}|$ in the named post-hoc **[O]** candidate pattern **v467**, record unchanged); a data $\leftrightarrow$ formula shuffle explodes the cross-validation χ^2 by $> 100\times$. **[E]**.

External-review residual map **[E]/[C]** ([`verification/v182_reviewer_residual_map.py`])

A careful external review (2026-06-14) raised four concerns at the **[C]/[O]** seams: **(A)** the mapping $2D \rightarrow 4D$ (“why does the dynamics read the masses as the Plücker norm 11?”), **(B)** the abstractness of UV gravity, **(C)** the meta-look-elsewhere of the grammar choice, and **(D)** the Koide Möbius flow (non-integer flow time $t \sim 2.84$, missing QFT generator). None adds an *independent* open gap. **A [E]/[C]**: the readout integer is exact ($11 = \sum_{k \leq 2} \binom{4}{k} = \dim S^+ - g_{\text{car}}$, the QBL boundary count, **v174**) and the $2D \rightarrow 4D$ step is the holographic reduction, so $A = \text{QGEO.SYM.01} + F_{\text{transfer}}$. **B [E]**: “gravity is the geometric readout of the boundary” is **QGEO.SYM.01**. **D [E]/[C]**: $\text{Aut}(\mathbb{P}) = \text{PSL}(2, \mathbb{C})$ is the Möbius group and the cusp flow lives on the deck $\mathbb{P} \setminus \mu_4$, so the Möbius *nature* is forced by the geometry; the non-integer $t \sim 2.84$ only locates the physical point and the missing generator is F_{transfer} , so $D = \text{QGEO.SYM.01} + F_{\text{transfer}}$. So A, B, D collapse onto the *two already-named* residuals; only **C** is genuinely *experiment-only*: the grammar is frozen (**v84**), minimal and over-determined (the anchor $a = (1, 1, 2)$ lands in ≥ 6 mutually-independent, not-selected-for structures), a look-elsewhere-resistant signature whose residual — correctly — falls only to out-of-sample data (JUNO, CMB-S4), never to argument. The four concerns are a unification of the residual *structure*, not a closure of the underlying premises, which stay **[O]/[C]**.