

TFPT — Research Contracts for the Remaining Interfaces:

$$v_{\text{geo}}, G_{\text{net}}, F_{\text{transfer}}$$

(U_{wall}): the parabolic flavor wall-selection and (G_{metric}): the full quantum-gravity measure

plus the two continuum contracts CELEST.SEAM.01 and WOIT.OS.TWISTOR.01 (M1–M3 and the δ -chains executed and decided; $\alpha/\beta_1/\beta_2/\beta_3$ executed, γ next)

Stefan Hamann

Alessandro Rizzo

August 7, 2026 · v5.4 (rev 543)

The TFPT document set — what is treated where

Plain language: TFPT (Topological Fixed-Point Theory) is a small discrete compiler. Two inputs — the seam constant $c_3 = \frac{1}{8\pi}$ and the carrier rank $g_{\text{car}} = 5$ — build $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ and read off the Standard-Model skeleton, the constants and the scale grammar. The series is **the introduction plus five numbered papers and five companions** (plus two standalone working notes), best read in order:

0. **introduction** — reading guide, compiler closure, predictions, the dependency DAG and the single proof ledger (`verification/status_ledger.csv`). (*entry point*)
1. **tfpt_1_architecture_e8** — the two axioms, the derivation map, the EM fixed point α^{-1} , the $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ construction, the QBL theorem chain.
2. **tfpt_2_standard_model** — the SM in one φ_0 -ladder formula, flavor from parabolic transport (sheet diamond, dual anchor, Q_+ cohomology), the worked closures.
3. **tfpt_3_e8_audit_bootstrap** — the seven E_8 slices as an audit raster, the cascade bridge, the Möbius bootstrap.
4. **tfpt_4_frontier** — honest status of η_B , m_p/m_e , Koide, dark matter, full QG.
5. **tfpt_5_redteam** — the adversarial audit: declared attacks, what survives, what each reduces to, and the kill tests.

Companions: **R. tfpt_research_contracts** — the open interfaces ($v_{\text{geo}}, G_{\text{net}}, F_{\text{transfer}}$) as numbered contracts; **H. tfpt_horizon_readouts** — Appendix H (unit-system reframe): one seam constant as the universal horizon thermal code, the Nariai/anchor surface, the resummed seam clock; **O. origin_theory** — why no free fundamental number remains (exact core + one honestly-typed cyclic interpretation); **S. tfpt_safeguards** — the verification discipline: every mechanism that defends a claim against coincidence and numerology (status calculus, no-free-pattern, the over-determination map, the fire-wall, frozen predictions + null model, the independent Wolfram and Lean paths, the red team); **P. tfpt_prime_front** — research documentation of the prime / zeta line: the seven load-bearing modules on the E_8 glue census (Hecke from geometry through the transport ledger), the sandbox localization program that follows them, and the kill list. Documentation, not a result: it makes no claim about the Riemann Hypothesis.

Working notes: **N1. note_e8_gaussian_code** — the Gaussian code bridge: E_8 over $\mathbb{Z}[i]$ via Construction A over the extended Hamming code $[8, 4, 4]$, the canonical four-bit quotient $L/(1+i)L \cong \mathbb{F}_2^4$, and the G_{31} quartic companion (exact algebra, machine-certified); **N2. note_hilbert_polya_truncations** — a computable, zeta-free truncation family for the Weil measure: measurements on a Hilbert–Pólya candidate (prediction-freeze methodology; no RH claim).

You are reading companion R (Research Contracts).

TFPT status at a glance — read this card the same way in every document

Two inputs, one compiler. From the boundary constant $c_3 = \frac{1}{8\pi}$ (axiom P1) and the five-slot carrier $g_{\text{car}} = 5$ (axiom P2) — jointly the elementary symmetric data of one anchor $a = (1, 1, 2)$ — the discrete compiler $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ is *built*, and the Standard-Model skeleton (gauge group, three families, hypercharges, the flavor matrix), the dimensionless constants ($\alpha^{-1} = 137.0359992$) and the scale grammar are read off as fixed points. The algebraic core is machine-checked; physical readouts run through *named bridges* whose status is typed below.

What is closed, and what genuinely remains. The discrete/algebraic layer is closed ([E]); the everything-open residual reduces to *three named interface problems*:

Interface	Question	Status
v_{geo}	the one metrology unit ($= 1/\sqrt{G} = m/\mu$); No-Unit Thm: no compiler scale	primitive [O]
G_{net}	simple-current extension $(D_5)_1 \otimes (A_3)_1 \rtimes \langle (1, 1) \rangle \cong (E_8)_1$	[C]
F_{transfer}	one functor, four typed interfaces (Koide, η_B , axion, m_p/m_e)	[C]

Gravity is parameter-free. The classical metric-sector field equation is no longer only an $R+R^2$ readout: the fixed-volume entanglement equilibrium (Jacobson; Faulkner et al.), run with TFPT's atoms, gives the *full* covariant Einstein equation $G_{ab} + \Lambda g_{ab} = c_3^{-1} T_{ab}$ with *both* coefficients fixed — $c_3^{-1} = 8\pi$ (no free dimensionless Newton dial; the thermodynamic origin $2\pi/\eta$ *coincides* with the geometric one $|\mathbb{Z}_2|2\pi\chi$ via $|\mu_4| = |\mathbb{Z}_2|\chi(S^2) = 4$, so c_3 is triply over-determined — anchor, geometry, thermodynamics, v358) and Λ from α ($\rho_\Lambda = (3/4\pi^2)e^{-2\alpha^{-1}}$, v60); the Einstein tensor (not Ricci) is forced by Lovelock, so matter conservation is an output (v359). The residual here is only the equation-of-state status (an external candidate action — Bianconi's entropic action, arXiv:2408.14391 — is quantified in v473 without changing the typing) and the one unit v_{geo} ; the global measure (C7 = QG.AMB.01) is now a [C] redundancy (v369), a certification object rather than missing dynamics, conditional on SEAM.EQUIV.01 and Bisognano–Wichmann.

The seam-net structural theorem G_{net} (= SEAM.EQUIV.01: the raw seam *is* the holomorphic $(E_8)_1$ net) now carries two named routes (route split 2026-07-22, no status change in substance): the MMST route SEAM.EQUIV.MMST.01 is *closed modulo cited theorems*; the twistor route SEAM.EQUIV.TWISTOR.01 (Costello–Li, prepared by CELEST.SEAM.01) and the unconditional parent stay [O]. On the MMST route: the explicit lattice model (v367/v368) and the S_3 closure stack pin the target at every computable level — central charge $c = 8$ (v376), the $(E_8)_1$ character with 248 currents and one primary (v377), genus-one torus GSD = 1 (v378) and reflection positivity (v379) — and it is Lean-formalised as FORM.SEAM.MMST.01: the collar's MMST hypotheses are kernel-proved, the MMST scaling-limit and Adamo–Moriwaki–Tanimoto OS-reconstruction theorems enter as named cited axioms, and the #print axioms check is clean. The *only* residual of that route that stays [O] is the abstract continuum existence of the scaling limit (v336) — its 128-spinor extension leg since certified at net level by the peer-reviewed crossed-product package ($h_s=16/16=1 \in \mathbb{Z}$, the Longo–Rehren locality criterion; Böckenhauer; KLM $\mu=4/2^2=1$; AGT/AMT the independent second witness) with the realisation input reduced to invariant level and the parallel Lean route *seamResidualClosed'* (v469); so the MMST route is *closed modulo cited theorems*, not solved. QGEO.SYM.01 (the μ_4 deck acting geometrically) is its *corollary*: a conformal net's vacuum is rotation-invariant by axiom, so the deck follows with no extra premise (v335, Lean FORM.QGEO.BW.01). The nonperturbative quantum-gravity measure QG.AMB.01 is no longer carried as an open item: it is a [C] redundancy (v369), a certification object rather than missing dynamics, conditional on SEAM.EQUIV.01 and Bisognano–Wichmann (and TFPT is moreover rigorously gap-decoupled from the general Euclidean-QG conformal-factor problem, margin $1.648 > 0$, v76/v330). So the honest residual is the one unit v_{geo} plus the typed F_{transfer} interfaces, above the cited-theorem ceiling on the seam and with QG.AMB.01 a [C] redundancy. Everything carries a claim ID resolving to a row of *verification/status_ledger.csv* (the single source of truth); **if the text and the ledger ever disagree, the ledger wins.** The historical labels ($U_{\text{wall}}, G_{\text{metric}}, F_{\text{frontier}}$) are retained only for ledger continuity — the live residual is $v_{\text{geo}} \oplus F_{\text{transfer}}$, with the seam G_{net} closed modulo cited theorems on its MMST route (parent [O]) and QG.AMB.01 a [C] redundancy.

Marker key: [E] exact / machine-proven · [C] conditional (holds under named hypotheses) · [O] open / axiom · [X] falsifiable kill test. Per-claim fine type in *verification/status_ledger.csv*.

[E] exact / machine-proven**[C]** conditional**[O]** open / axiom**[X]** kill test

$$\mathbf{P1} \, c_3 = \frac{1}{8\pi} \quad \mathbf{P2} \, g_{\text{car}} = 5 \quad (\text{two inputs, [O] axioms})$$

$$\text{Anchor } a = (1, 1, 2): e_1, e_2, e_3 = 4, 5, 2 \quad ([E] \text{ exact})$$

$$\text{Discrete compiler: } D_5 \oplus A_3 + \mu_4 \Rightarrow E_8, h = 30, \varphi_0 \quad ([E] \text{ exact})$$

$$\text{SM skeleton: three families, hypercharges, flavor matrix } R \quad ([E])$$

$$\text{Physical bridges: } v_{\text{geo}}, G_{\text{net}}, F_{\text{transfer}} \quad ([C] \text{ bridges})$$

$$\text{Frontier readouts: } \Lambda, \text{ inflation, } \eta_B, \text{ dark matter, QG} \quad ([C]/[O])$$

What this document is about

After the compiler closure the entire residual is

$$\text{Rest} = (U_{\text{wall}}) \oplus (G_{\text{metric}}) \oplus (F_{\text{frontier}})$$

This note turns the two genuine research gates into *contracts*: a numbered chain of lemmas, the single theorem that closes each gate, and — for every step — whether it is machine-certifiable today. (F_{frontier}) (Koide, η_B , axion relic scale, m_p/m_e) is *not* a gate: those are deliberately typed QFT/cosmology interfaces (see `tfpt_4_frontier`). **Priority:** (U_{wall}) *first* (finite, algebraic, falsifiable), then (G_{metric}) (deep analytic programme).

Operative aliases (current state of the reductions; the historical gate names above are kept for ledger continuity). The lemma chains below have since *discharged* large parts of both gates, so the operative residual reads sharper:

$$\text{Rest}_{\text{operativ}} = v_{\text{geo}} \oplus G_{\text{net}} \oplus F_{\text{transfer}}$$

— (U_{wall}) $\rightarrow v_{\text{geo}}$: the quark *ratios* are closed combinatorially (Readout Rigidity + Plücker, U2/v49/v71), U_{point} collapses to the *one* dimensionful unit v_{geo} shared with $1/G$ (v75/v68) — what remains is a scale anchor, not a flavor gap (the *full* amplitude matrix beyond the ratios is the only place where U_{point} still appears); (G_{metric}) $\rightarrow G_{\text{net}}$: holographically reduced to the single boundary-net statement “the seam-Calderón inclusion has Jones index $4 = |\mu_4|$ ” — holomorphy, $(E_8)_1$ and the unique 2D bulk then *follow* (v89/v92/GATE.METRIC.05-07); and the carrier-side QBL input now merges into the *same* statement: the tower carrier $\rightarrow SO(16)_1 \rightarrow (E_8)_1$ carries one and the same 16-fermion content, whose single quasi-free kernel (rank = c ; Wick/Pfaffian exact) is the certified Calderón datum — closing G_{net} therefore also closes the carrier choice **[E]** (`[verification/v113_quasifree_kernel.py]`); (F_{frontier}) $\rightarrow F_{\text{transfer}}$: named QFT/cosmology *transfer* interfaces, never compiler powers — and *not* four unrelated interfaces but *one* discrete geodesic on the Sheet Diamond (the $K \rightarrow C \rightarrow F$ sheet axis $M(1, t)$, with the Koide source $\rightarrow$ pole as its 1-D Möbius projection), along which the external continuous solver merely evaluates (`[verification/v224_diamond_ftransfer_path.py]`); the discrete data of that geodesic are now arithmetically framed — $\det M(1, t) = N(\alpha_t)$ with $\alpha_t = (4t+7+\sqrt{-7})/2$ integral, the path one translation $\alpha_{t+1} = \alpha_t + 2$ of constant discriminant -7 (`[verification/v533_ftransfer_disc7_norm.py]`), and the preregistered next test is whether the four external solvers share *one* fractional-linear action on α_t (kill: four incompatible arithmetic actions) — *executed in its literal group form* (2026-08-02, `[verification/v632_ftransfer_pgl2.py]`): one common parabolic PGL_2 action on the discrete half (the declared readings exactly intertwined with the base translation), unifold degeneration on the native continuous half ($g_{\text{Boltzmann}} = g_{\text{pole}}$ identically, g_{QCD} exactly conjugate to the base translation, relic degenerate); the kill criterion is *not* met, and the residual — the external anchor clocks/jets — stays fenced by `CONTRACT.F.01`.

The residual is certification, not construction (`[verification/v384_residual_certification.py]`). A ledger-CI audit classifies every remaining item into exactly three *external* kinds, with *zero* open

internal TFPT mechanisms: [E] (A) an external math proof — SEAM.EQUIV.01 continuum existence (the cited MMST theorem plus the crossed-product extension package, v336/v469, with AGT/AMT as second witness), ALPHA.QUILLEN.EXACT.01 (v382, its level and normalisation sharpened by v470, the determinant-line/moduli bridge lemma exhibited at the finite level by v472 — the continuum ζ -det identification stays the open part) and the constructive QG.AMB.01 measure (a [C] redundancy, v369); [E] (B) theorem-forbidden — v_{geo} , the one unit, by the No-Unit theorem (v153/v364); [E] (C) external physics — F_{transfer} (v371–v374, firewalled v187). The two load-bearing facts are checked (No-Unit dimensionlessness; the gap-decoupling margin $1.648 > 0$), so the only hard things left are theorems and external physics, not missing dynamics (RESIDUAL.CERTIFICATION.01).

The Coxeter-clock facet symmetry, resolved as a structural lemma (RES.COXETER.SYMMETRY.01, [verification/v394_coxeter_atoms.py]/[verification/v409_coxeter_prime2_lemma.py]). The order-30 clock $30 = 2 \cdot 3 \cdot 5 = h(E_8)$ has all three primes facet-typed, and they are exactly the three anchor atoms ($2=|Z_2|$, $3=N_{\text{fam}}$, $5=g_{\text{car}}$) wearing number-field hats ($\mathbb{Q}(i)/\mathbb{Q}(\sqrt{-3})/\mathbb{Q}(\sqrt{5})$, v390). The table is *not* symmetric: the seam (prime-2) appears as the *common factor* of both dynamic rates ($2/3 = |Z_2|/N_{\text{fam}}$ and $(\varphi+2)/4 = (\varphi+2)/|\mu_4|$, $|\mu_4|=2^2$), not as an independent attractor. *The structural lemma (v409) closes the falsifiable corollary:* the sheet involution $\sigma = \text{diag}(1, -1, -1)$ has spectrum $\{+1, -1\}$ and projector $(I+\sigma)/2$ spectrum $\{0, 1\}$ — none in the open interval $(0, 1)$, so prime-2 supplies a *parity/projection*, never a nonzero contraction rate; the genuine rates $2/3$ and $(1/3)^6$ lie in $(0, 1)$ and carry prime-2 only as a *factor*. Hence prime-2 is *necessary* (admissible boundary transport is sheet-paired) but *not autonomous*: **no prime-2-only attractor exists** — dynamics begins only after coupling to family prime-3 or carrier prime-5 ($|\mu_4|=2^2$ the Galois bridge). This closes the corollary [E] and types the necessity [C], sharpening v383’s “every sector is the same gapped operator” towards “the same *seam-driven* operator”; the full “appears in every gap” universality stays the v383 reading. The corollary is **machine-proved in Lean 4** (TfptCarrier/CoxeterPrime2.lean, FORM.COXETER.PRIME2.01: axiom-pure, lake build clean). [E]/[C].

QFT layer (this round). The boundary QFT is now *one relative object* $\text{TFPT}_{\text{QFT}} = (\mathcal{A}_\Sigma, \omega_\Sigma, \Delta_\Sigma, \rho, A_F, H_F, D_F, J, \gamma, S_{\text{rel}})$, closed modulo the *single* keystone SEAM.EQUIV.01 (whose downstream conformal-deck face is QGEO.SYM.01): its finite Dirac is a covariance induction of the seam KMS state (v258), its spectral-action cutoff *is* that KMS weight (v259), its seam, carrier-16 and E_8 live on one Kummer/K3 surface (v260), and the whole assembly is certified as the *Modular Spectral Closure* (v261). The ambient quantum-gravity measure (G_{metric}) is gap-decoupled and kept separate by design — the QFT layer does not wait on it.

The two continuum contracts, and how far they are executed. Beside the two historical gates this note carries the two *continuum-route* research contracts, both still [O] as contracts but with most of their preregistered stages now *executed*. CELEST.SEAM.01 (the celestial and twistor continuum route): WP1–WP4 and both WP5d stages executed, the WP5e stages $\alpha/\beta/\gamma/\delta_2/\varepsilon_2/\varepsilon_1$ executed, all three back-reaction milestones M1–M3 executed with their preregistered kills evaluated (v515–v517), the δ_1 chain *decided* (kill under the derived measure, v518), the declared-vs-derived measure question *decided at probe level* for the declared completion reading (single-valuedness from F-independence + the Quillen pairing, v520), and the w_m normalisation itself *derived constructively* ($1/\det_j$ the Atiyah–Bott/zeta fixed-point factor from three independent sources, $\psi = 64$ reproduced, v523) — nothing in the measure chain is declared any more at that level; the residual [O] is the global BCOV integral beyond the fibre zero-mode factor. WOIT.OS.TWISTOR.01 (the Osterwalder–Schrader twistor bridge — the actual bridge from compiler to physics): the α stage executed (Θ exists at the free level and free RP selects the same family, v519), β_1 executed (the μ_4 clock is the euclidean rotation, the gaugeable datum is the GSO $\mathbb{Z}_2$, gauge-fixed RP holds, v522), β_2 executed (the OS quotient explicit, $\mathcal{H}_{\text{phys}}$ positive definite at both levels, the clock a positive transfer operator with spectral calculus, v524), β_3 executed (the $\mathbb{PT} \leftrightarrow \mathbb{PT}^*$ duality typed against σ_{std} : the OS cut and the $(2, 2)$ signature *forced*, the kill branch empty by algebra, the induced member = the Θ_{phys} carrier, v565) — γ is next, still under the *straddle-law* constraint: the first interacting seam toy fires Kill-Test 2 at toy level (RP breaks on exactly the quartet-straddling cuts, v529) — an honest threat and simultaneously the first hard selection principle for $\mathcal{A}_{\text{hol}}$, since *executed* as such: run as a selector on the interacting mark family, reflection positivity keeps exactly *one* member alive — $\delta = \pi/2$ with positive coupling, the alignment bit itself (v534, toy-fenced; the literal leading-order cone formalization is dead, the protection is nonperturbative). Both contracts stay [O]; no marker moves.

Research Contract 1 — (U_{wall})

Goal. Prove

$$\nabla_F^* = \text{Sel}_{D_4, \text{cusp}, \det R=8, \text{Spec}(Q_+)=\{1,2,3\}}(\mathcal{M}_{\text{par}}(\mathbb{P}^1, \mu_4, E, \alpha)),$$

with $E = \mathcal{O}(-2) \oplus \mathcal{O}(-1) \oplus \mathcal{O}(-1)$ and $\alpha = \{0, \frac{1}{3}, \frac{2}{3}\}$ at the four points of μ_4 . The substrate is standard: degree-zero stable parabolic bundles correspond to unitary representations with prescribed local monodromies (Mehta–Seshadri); tame non-abelian Hodge gives the Higgs / flat-connection side. The TFPT content is to pin the *one* D_4 -symmetric, rank-3, four-point point that realises the selectors.

Lemma 1 (U0 — normalisation). *The gate data are equivalent to the $SU(3)$ character variety $\mathcal{X}_{\text{cusp}} = \{(M_1, \dots, M_4) \in C_{\text{cusp}}^4 : M_1 M_2 M_3 M_4 = \mathbf{1}\} / SU(3)$ with $C_{\text{cusp}} = \text{Ad}_{SU(3)} \text{diag}(1, \omega, \omega^2)$, $\omega = e^{2\pi i/3}$, and $\pi_1(\mathbb{P}^1 \setminus \mu_4) = \langle \gamma_1, \dots, \gamma_4 \mid \prod \gamma_i = 1 \rangle$. To show: $\mathcal{M}_{\text{par}}^{D_4} \simeq \mathcal{X}_{\text{cusp}}^{D_4}$ as polystable Mehta–Seshadri / tame Hodge realisation. Machine: the group presentation and D_4 -action are finite and codable (Sage/Wolfram).*

Finite rigidity — now proven [E]. The finite half of U0 is closed exactly: μ_4 has cross-ratio 2 and Möbius stabiliser $\langle z \mapsto iz, z \mapsto 1/z \rangle$ of order $8 = |D_4|$ (a 4-cycle of the marks + a reflection, faithful); $H^1(\mathbb{P}^1 \setminus \mu_4)$ has rank $4 - 1 = N_{\text{fam}} = 3$; and the forms $\omega_k = z^{k-1} dz / (z^4 - 1)$, $k = 1, 2, 3$, are μ_4 -eigenforms ($z \mapsto iz : \omega_k \mapsto i^k \omega_k$) carrying the three nontrivial characters χ_1, χ_2, χ_3 with weights $(1, 2, 3) = \text{Spec}(Q_+)$. Hence a genus-zero boundary with faithful D_4 , four parabolic marks and this μ_4 -decomposition is Möbius-equivalent to $\mathbb{P}^1 \setminus \mu_4$ with the $(1, 2, 3)$ grading (QGeo.RIGID.01, [E], [verification/v168_qgeo_rigidity.py]). The single residual is the seam-side *realisation* — that these hypotheses follow from the seam collar itself — which stays QGeo.REALIZE.01 ([C]).

Seam Collar Realisation Theorem — the one central proof obligation [E]/[O]
([verification/v176_seam_collar_realisation.py])

Stated as one theorem and certified as a reduction — not a fabricated proof.

Theorem (target). *Given* (1) an oriented one-sided RP seam collar Σ ; (2) a sheet-odd Calderón involution; (3) quasi-free one-particle seam data with gap $(2/3)^6$; (4) a faithful D_4 action on the boundary marks; (5) four parabolic marked points with μ_4 -character decomposition; (6) an admissible $c=8$ boundary net — *then* the conformal boundary of Σ is Möbius-equivalent to $\mathbb{P}^1 \setminus \mu_4$, its H^1 character basis is canonically the generation basis with grading $(1, 2, 3)$, the seam Calderón contraction is the rank-8 K_Σ polarisation, and the index-4 simple-current extension is $(E_8)_1$.

The proof decomposes into six steps, five already [E]:

- 1–2 Geometry / uniformisation [E] (v168):** μ_4 has cross-ratio 2 and a faithful Möbius D_4 stabiliser of order 8; four genus-0 marks with faithful D_4 force the μ_4 square up to Möbius; $H^1(\mathbb{P}^1 \setminus \mu_4)$ has rank $4 - 1 = N_{\text{fam}} = 3$.
- 3 Calderón data [E] (v156):** the Dirichlet-to-Neumann operator of the 2d Laplacian on the half-space has symbol $|k|$ (harmonic extension $e^{ikx - |k|y}$), the free chiral dispersion — bulk-detail-independent.
- 4 H^1 character = generation grading [E] (v137/v168):** $\omega_k = z^{k-1} dz / (z^4 - 1)$ carry μ_4 characters i^k with weights $(1, 2, 3) =$ the A_3 exponents $= \text{Spec}(Q_+)$ — natural, not merely isomorphic.
- 5 One-particle contraction [E] (v162):** a μ_4 -equivariant operator is diagonal in the cusp basis $(\frac{1}{|\mu_4|} \sum_j U^j X U^{-j} = \text{diag } X$ for $U = \text{diag}(1, i, -1, -i)$); its spectrum is the cusp weights $\{0, \frac{1}{3}, \frac{2}{3}\}$, so the transfer eigenvalues $(1 - \alpha)^{p_2}$ ($p_2=6$) are $\{1, (2/3)^6, (1/3)^6\}$, gap $(2/3)^6$ — forced, no free knob.
- 6 AQFT closure [E] (v154/v175):** $(D_5)_1 \otimes (A_3)_1$ with the isotropic μ_4 glue has index $4 = |\mu_4|$, $c = 5+3 = 8$, μ -index $16/16 = 1 \Rightarrow$ holomorphic $\Rightarrow (E_8)_1$ ($248 = 120+128$); full-cone RP holds for all m via the CAR second-quantisation functor.

The proof tree — and the two genuine open doors [O]
([verification/v177_seam_marking_kernel.py]). Taking the marks/ D_4 as a *hypothesis* (as the six-step form above does) is a near-circular trap; the honest decomposition factors the theorem as

$$\text{REALIZE} = \text{MARKS} \cdot \text{UNIFORM} \cdot \text{COHOM} \cdot \text{MODULE} \cdot \text{KERNEL} \cdot \text{NET},$$

of which *four nodes are now closed constructively*: UNIFORM [E] (an order-4 Möbius map conjugates to $z \mapsto iz$, a non-fixed orbit scales to μ_4 , and $\sigma\rho\sigma = \rho^{-1}$ for $\sigma : z \mapsto 1/z$ gives a faithful D_4 — so a genus-0 four-marked faithful- D_4 boundary *is* $(\mathbb{P}^1, \mu_4)$), strengthening v168's cross-ratio to the full uniformisation — and now **Lean-formalised** as FORM.QGEO.02: $\rho^4 = \text{id}$, $\rho^2 \neq \text{id}$, $\sigma^2 = \text{id}$, $\sigma\rho\sigma = \rho^{-1}$, the orbit scaling to μ_4 , the μ_4 permutation, and the multiplier classification “order-4 $\Leftrightarrow \zeta = \pm i$ ” are machine-proved in `TfptCarrier/MobiusUniformisation.lean`, AUDIT: PASS); COHOM [E] ($\rho^* \omega_k = i^k \omega_k$, grading (1, 2, 3) — also **Lean-formalised** as FORM.QGEO.03: the pullback characters i^1, i^2, i^3 and the reflection parity $\sigma^* \omega_1 = \omega_3$, $\sigma^* \omega_2 = \omega_2$, $\sigma^* \omega_3 = \omega_1$ are machine-proved in `TfptCarrier/CohomologyGrading.lean`, AUDIT: PASS); MODULE [E] (a *unique* μ_4 -equivariant iso — multiplicity-free + residue normalisation, reflection $\omega_1 \leftrightarrow \omega_3$, ω_2 fixed; the parity is the Lean FORM.QGEO.03 above, the *uniqueness* stays symbolic); and NET [E] (v154/v175). The residual is therefore *exactly two* named obligations, neither a finite computation: QGEO.MARKS.01 — the *raw* RP seam collar canonically produces the genus-0 four-parabolic-marked D_4 boundary (the four marks from the anchor atom $e_1(a) = 4 = |\mu_4|$, collisions excluded, $\tau\rho\tau = \rho^{-1}$) — and QGEO.KERNEL.01 — the *raw* seam Calderón operator equals the μ_4 -equivariant free $c=8$ gapped one-particle contraction *as an operator* ($C_\Sigma = U^{-1}C_{\mu_4}U$), not merely the same spectral list. **The whole structural residual of the theory is these two doors**; everything between them is [E]. They need a human constructive-geometry/AQFT argument (or a proof assistant), not more arithmetic.

How far each door opens by computation ([`verification/v178_marks_kernel_attempt.py`]). Neither door is closed, but each splits into a finite [E] core + one *sharper* premise. For MARKS: *given* a genus-0 double with an order-4 clock ρ (fixed points $\{0, \infty\}$) and reflection τ , the marks are forced to be a generic ρ -orbit $\{1, i, -1, -i\} = \mu_4$ (not the fixed points), distinct iff $b_1 = 3 = N_{\text{fam}}$ (any collision drops b_1), with $\tau\rho\tau = \rho^{-1}$ giving faithful D_4 — so MARKS reduces to the *single* topological premise “the seam double is genus-0 with an order-4 clock whose marks form one orbit”. For KERNEL: on the 3-dim multiplicity-free H^1 the μ_4 characters (1, 2, 3) are distinct, so by Schur a μ_4 -equivariant operator is forced *diagonal* and is *completely determined by its eigenvalue per character* — two such with the forced transfer spectrum $\{1, (2/3)^6, (1/3)^6\}$ and the unique residue-normalised basis are *equal as operators*, not merely isospectral; the “spectrum vs operator” distinction *vanishes* on the finite block. So KERNEL reduces to the *single* premise “the full DtN on L^2 reduces to this 3-dim gapped transfer block + the fixed rank-8 polarisation” (principal symbol $|k|$, Lee–Uhlmann v156; no drift v158). The two doors are thereby *milder*, but still open.

Marks without a geometric posit: a Lefschetz/character argument ([`verification/v195_marks_lefschetz_character.py`]). The marks need not be *assumed* as D_4 boundary points: they follow from the H^1 character alone. The order-4 clock $\rho: z \mapsto iz$ fixes exactly $\{0, \infty\}$, so a puncture at a fixed point would contribute a *trivial* character to H^1 ; but the carrier datum is $H^1 = \chi_1 \oplus \chi_2 \oplus \chi_3$ (the A_3 exponents (1, 2, 3), with $\text{Tr}(\rho|H^1) = i + i^2 + i^3 = -1$), which has *no* trivial component — so no puncture sits at a fixed point. With $\text{rank } H^1 = n - 1 = 3 \Rightarrow n = 4$ punctures, all off the fixed locus and each in a free order-4 orbit, the only possibility is a *single* free μ_4 orbit (cross-ratio 2), and the closed-surface Lefschetz number $L(\rho) = 1 + 1 = 2 = |\{0, \infty\}|$ is consistent. So MARKS reduces from the geometric posit “ D_4 marks” to the milder *representation* premise “the raw seam H^1 carries the A_3 -exponent rep” — a carrier-algebra statement, strictly less circular, still [O].

And the two doors are in fact ONE ([`verification/v179_conformal_realisation.py`]). MARKS's residual is not even three independent assumptions: (1) genus-0 *is* P1 — the one-sided Gauss–Bonnet $c_3 = 1/(\|\mathbb{Z}_2\| 2\pi \chi(S^2)) = 1/(8\pi)$ uses $\chi(S^2) = 2$, and $\chi = 2 - 2g \Rightarrow g = 0$, so “genus-0” is the *same* $\chi = 2$ that supplies the “8” in c_3 (v58/v73); (2) the order-4 clock *is* the carrier — it is the Coxeter element of $W(A_3) = S_4$ (v117), order $h(A_3) = 4 = |\mu_4|$; (3) marks = one orbit is *forced* — the parabolic marks are not the two elliptic fixed points $\{0, \infty\}$, and a free order-4 orbit has exactly $4 = e_1(a)$ points. So MARKS reduces to “the carrier clock acts *conformally* on the genus-0 double”, and KERNEL then *follows* from that same geometry (Lee–Uhlmann $|k|$ + the $c = 8$ rigidity v158 + the cusped-surface spectral split, residual = $H^1 = \mathbb{C}^3 = N_{\text{fam}}$). **Hence the entire structural residual of the theory is a single geometric premise** QGEO.CONF.01: *the raw RP seam normal double is conformally* $(\mathbb{P}^1, \mu_4)$ *with the carrier clock as the order-4 Möbius automorphism* $z \mapsto iz$. Given it, everything — genus-0, the four D_4 marks, the (1, 2, 3) grading, the DtN block, the rank-8 polarisation and the $(E_8)_1$ net — follows. This single premise is a constructive-geometry statement (the conformal class of the raw seam *is* the cusped sphere), *not* a finite computation, and is left honestly [O].

And even that conformal premise reduces to a milder isometry premise ([`verification/v180_clock_is_mobius.py`]). QGEO.CONF.01 need not be assumed as a conformal-class identity: by *uniformisation* (Poincaré–Koebe) the genus-0 double (P1) with its RP-metric conformal

structure *is* $\mathbb{P}^1$ — the target is *produced*, not assumed — and by Kerékjártó (1919; Constantin–Kolev 2003) a finite-order orientation-preserving homeomorphism of S^2 is conjugate to a rotation, so the order-4 clock ($h(A_3)=4$) is topologically $z \mapsto iz$. An orientation-preserving *isometry* of a Riemannian surface is holomorphic (angle-preserving), hence a Möbius map of $\mathbb{P}^1$, and an order-4 Möbius map is $z \mapsto iz$ (elliptic, multiplier i , free orbit μ_4). So QGEO.CONF.01 is *implied* by the strictly milder, non-circular premise QGEO.ISO.01: “the carrier clock is an order-4 orientation-preserving *isometry* of the RP seam metric” — which does *not* name $\mathbb{P}^1/\mu_4$ and is the natural statement that the seam transport is metric-compatible (a holonomy preserving the RP inner product). **That isometry premise is the single, now milder, structural residual of the whole theory**; its surface realisation from the raw seam stays honestly [O].

And one step further reaches bedrock ([verification/v181_clock_is_conformal_symmetry.py]). By *equivariant uniformisation* (Nielsen realisation; for cyclic groups Kerékjártó + uniformisation) a finite-order orientation-preserving action on a genus-0 surface preserves a complex structure for which it acts by Möbius maps — so a *conformal automorphism* (strictly weaker than an isometry: every isometry is conformal, conformal allows any positive scale) already suffices to put the order-4 clock in the form $z \mapsto iz$. Since the clock is the Coxeter element of $W(A_3) = S_4$ (v117), i.e. the μ_4 deck, and a deck transformation preserves the pulled-back conformal structure *by definition*, the residual collapses to the *definitional* identification QGEO.SYM.01: *the seam’s conformal structure is the μ_4 -deck structure of the carrier clock* (the carrier μ_4 glue *is* the conformal monodromy of the seam). This is a physical/definitional statement tying P2 to the seam’s conformal geometry — *not* a finite computation and *not* reducible further without relabeling. **This is bedrock**: the chain REALIZE $\rightarrow$ theorem $\rightarrow$ MARKS+KERNEL $\rightarrow$ CONF $\rightarrow$ ISO $\rightarrow$ SYM (v175–v181) ends at a single definitional premise, left honestly [O]; everything above it is a theorem or an established citation.

A concrete geometric model for the bedrock premise: the μ_3 -cover program ([verification/v597_mu4_cover_model.py]–[verification/v601_equivariant_dual.py]). The bedrock premise now has its first explicit *candidate realisation*: the cyclic cover $y^3 = x^4 - 1$ of $\mathbb{P}^1$ branched at μ_4 (order-3 cusp monodromies, exactly the contract’s cusp structure). Machine-exact, in five rounds: the cover carries the rank-3 Eisenstein H_1 , the projective D_4 via the deck, the cusp-class torsion $(\mathbb{Z}/3)^4$ of order 81, and the unique invariant hermitian form of Lorentz signature $(1, 2)$ with $\det J = 8$ (v597/v599); the anti-holomorphic reflection (the real structure R , $R^2=1$, unique up to scale) exists and closes the operator dictionary (v599); the compiler pair (U, V) itself is *realised* — $\mathbb{Z}[\omega]$ -integral, GL_3 -conjugate to the compiler pair, integer on the saturated $\text{Fix}(R)$ lattice, with saturation index $81 = 3^4$ in the exact v566 divisor pattern $(1, 1, 1, 3, 3, 3, 3)$ (v600, Lean-certified as FORM.QGEO.04 in TftCarrier/CoverEmbedding.lean: determinant-free Smith certificates, kernel **decide**); and the presentation is canonical — the J-adapted cusp projectors give a manifestly D_4 -equivariant pair family (mark choice = winding choice) realising the *dual* parabolic, with the two parabolic types exchanged by the J-adjoint, and the v590 Klein four-group living integrally on the curve (v601/[verification/v602_duality_forms.py]). Honest status: this is a *model*, not yet the seam — the identification of the raw RP seam with the cover (the Weil-twisted positivity comparison against the v572 Gram data, and the geometric identity of the seam line) stays [O]; QGEO.SYM.01 does not move.

A sharper, operator-checkable restatement: QGEO.ENERGY.01 ([verification/v192_qgeo_energy_clock.py]). The bedrock can be re-expressed as an energy identity rather than a conformal-class statement: *the carrier clock ρ preserves the Calderón / Dirichlet-to-Neumann energy form of the RP seam double*, $\rho^* \Lambda_\Sigma \rho = \Lambda_\Sigma$. In two dimensions Dirichlet-energy invariance *is* conformal invariance, so this feeds the same chain (conformality $\rightarrow$ Möbius $\rightarrow z \mapsto iz \rightarrow \mu_4$). Its appeal is checkability: $\Lambda_\Sigma = |k|$ is the rank-8 Calderón polarisation already in the net construction, so the premise becomes an operator identity rather than an abstract identification. **Honest scope, however**: “ ρ preserves Λ_Σ ” is plausibly *equivalent* to “ ρ is the conformal deck” (QGEO.SYM.01), so QGEO.ENERGY.01 *relocates* the bedrock to a more falsifiable surface but does *not* eliminate it — it becomes a theorem only if ρ is defined *independently* (from the carrier algebra) and the invariance is then *proven*, which is open. The bedrock therefore stays [O]; the gain is a sharper operator-analytic target, not a closure.

The non-circular sharpening: QGEO.ENERGY.02 ([verification/v193_qgeo_energy_commutator.py]). The way to make “ ρ defined independently” precise is the *energy commutator* $[\rho, \Lambda_\Sigma] = 0$ on the rank-8 Calderón polarisation, with three sub-claims: (1) ρ is the Coxeter element of $W(A_3) = S_4$ (order $4 = |\mu_4|$, v117), defined from the *carrier* algebra, not the seam; (2) *the non-circularity hinge* — Λ_Σ must be the DtN of the *raw* RP seam (definable from the RP data alone), not the μ_4 normal-form operator;

(3) $[\rho, \Lambda_\Sigma] = 0$ forces the μ_4 character grading $(1, 2, 3)$ on H^1 , already established (QGE0.COHO01). An exact consistency rider pins the direction: the induced EH coefficient $k = (\ln m)/(12\pi)$ (v150) reproduces $c_3/2$ iff $\ln m = q(A_3) = \frac{3}{4}$ (the *family* glue norm), while $q(D_5) = \frac{5}{4}$ (the carrier norm) misses by $g_{\text{car}}/N_{\text{fam}} = \frac{5}{3}$ — so the seam energy form must reproduce the *family* norm (gravity is family-geometry-induced). If sub-claim (2) holds, $[\rho, \Lambda_\Sigma] = 0$ turns QGE0.SYM.01 from definitional bedrock into an *energy-rigidity theorem*; until then it is a sharp, falsifiable *proof target*, [O].

Subclaim (2) tackled — one notch closer, not closed ([verification/v194_raw_seam_dtn_rp.py]).

The non-circularity hinge is essentially *met* at the canonical level: Osterwalder–Schrader reconstruction yields a canonical transfer operator $T = e^{-H}$ from the RP state and reflection alone — no coordinates (v54; §(4) of tfpt_2) — on a *quasi-free* seam state (v155: the boundary marginal of a Gaussian bulk is the compression $P\Gamma P$, a contraction). So Λ_Σ is RP-definable without ever naming $\mathbb{P}^1 \setminus \mu_4$. The commutator $[\rho, \Lambda_\Sigma] = 0$ then *closes on the finite cohomology* H^1 [E]: $\rho: z \mapsto iz$ acts on $\langle w_1, w_2, w_3 \rangle$ ($w_k = z^{k-1} dz / (z^4 - 1)$) as $\rho^* w_k = i^k w_k$, distinct characters $i, -1, -i$, so any ρ -equivariant operator is diagonal there (v177). What does *not* close is the *full- L^2* commutation beyond H^1 — it is exactly the statement that the quasi-free seam state’s modular flow is *geometric* (Bisognano–Wichmann). So the bedrock *reduces* from a bare definitional postulate to “intrinsic BW geometric modular covariance of the quasi-free seam state” — a named, standard (hard) AQFT property, one notch sharper, but still [O] (and the BW step must be established *intrinsically*, else it re-presupposes the conformal covariance it should produce).

A concrete variational handle on the residual ([verification/v196_seam_energy_functional.py]).

The open BW statement can be made a discretisable extremum problem via the failure functional $E_{\text{fail}}(\Sigma, \rho) = \|[\rho, \Lambda_\Sigma]\|_{HS}^2 + \|\rho^4 - 1\|^2 + \|\Theta\rho\Theta - \rho^{-1}\|^2$, whose zero locus is exactly the QGE0.SYM.01 conditions (the energy commutator, the order-4 clock, and the RP-reflection twist). On the finite H^1 block $E_{\text{fail}} = 0$ at the μ_4 configuration ($\rho = \text{diag}(i, -1, -i)$, Λ μ_4 -equivariant, $\Theta = \text{conjugation}$) and > 0 for any μ_4 -breaking perturbation, so μ_4 is a genuine minimiser [E]; minimising E_{fail} over the *full* raw-seam DtN is precisely the BW residual above. So the bedrock is now a sharply-posed *variational target*, not a bare definition — but still [O].

The decisive crack: principal symbol exact + Tomita–Takesaki ([verification/v198_modular_commutator_reduction.py]). Two observations collapse the residual to a single non-circular state-invariance. *First*, the DtN principal symbol is $|k| = \sqrt{-\partial_\theta^2}$ (Lee–Uhlmann, v156); in the Fourier basis $e^{in\theta}$ it is $\text{diag}(|n|)$ and the clock $\rho: z \mapsto iz$ (rotation by $\pi/2$) is $\text{diag}(i^n)$, both *diagonal*, so $[\rho, |k|] = 0$ *exactly* on all of L^2 — the leading-order commutation is free on the whole boundary, never the open part. *Second*, by Tomita–Takesaki the modular operator Δ_ω is intrinsic to (M, Ω) , so a state-preserving symmetry ($\rho\Omega = \Omega$) commutes with Δ and hence with $\log \Delta \sim \Lambda_\Sigma$: thus $[\rho, \Lambda_\Sigma] = 0$ *follows from* $\omega \circ \rho = \omega$ — using only the state and reflection (which RP supplies) and *no* conformal covariance, which **removes the BW circularity worry above**. Since the finite H^1 block is already μ_4 -invariant (v177), the entire residual is now the single *state-invariance* “the raw quasi-free seam state is μ_4 -invariant” ($\omega \circ \rho = \omega$, equivalently the Gaussian bulk measure is deck-invariant) — the maximally-operational, checkable form of QGE0.SYM.01. A foundational symmetry postulate cannot be derived from nothing (the role $c = \text{const}$ plays in relativity); it is now in its sharpest, non-circular, falsifiable form, [O].

The residual, fully localised and numerically probed ([verification/v199_seam_state_invariance.py], [verification/v200_seam_variational_scan.py]). Attacking $\omega \circ \rho = \omega$ on the *raw* seam (not the μ_4 normal form) confirms it is non-circular — ρ is the Coxeter element of $W(A_3) = S_4$ ($\rho^4 = 1$), an automorphism of the raw CAR algebra, defined entirely carrier-side — and reduces it, via the quasi-free covariance $C_\Sigma = \frac{1}{2}(1 + \text{sgn } H_1)$, to $[\rho, H_1] = 0$, i.e. H_1 is *block-diagonal in the four μ_4 -character classes* $\{n \equiv r \pmod{4}\}$. The principal symbol $|k| = \text{diag}(|n|)$ passes automatically, so the entire residual is the single bounded statement “the sub-principal symbol of the raw DtN has no off-character matrix elements”. The companion variational scan minimises $E_{\text{fail}}(\theta) = \|\rho(\theta)^4 - 1\|^2 + (\text{faithfulness})$ over a free clock angle and finds its *unique* minimum at $\theta = \pi/2$ — the μ_4 clock $z \mapsto iz$ (the only faithful order-4 grading $i, -1, -i$) — a numerical sign that the variational principle selects μ_4 . This is the “next executable theorem” a working mathematician can take up; it does not close the postulate, [O].

The bounded sub-principal residual is the image of the μ_4 marks, not a free postulate ([verification/v201_seam_subprincipal_marks.py]). One genuine step further: the seam DtN is $\Lambda = |k| + M_f$ with M_f the bounded sub-principal piece = multiplication by a curvature function $f(\theta)$ on the seam circle (the standard Steklov/DtN structure), so $\langle n | M_f | n' \rangle = f_{n-n'}$. Then M_f is μ_4 -character-block-

diagonal *iff* f has Fourier support only on modes $\equiv 0 \pmod{4}$, i.e. *iff* f is $\mathbb{Z}_4$ -invariant (the principal $|k|$ already commutes, v198). But a curvature sourced by the four marks, $f(\theta) = \sum_{j=0}^3 g(\theta - 2\pi j/4)$ for *any* profile g , has $f_m = 4g_m$ [$m \equiv 0 \pmod{4}$] — automatically $\mathbb{Z}_4$ -invariant. So the μ_4 -mark orbit (forced by v195) *forces* the sub-principal symbol block-diagonal; 3 equally-spaced marks ($\mathbb{Z}_3$) or 4 generic marks both break it (negative controls). The residual of $\omega \circ \rho = \omega$ thus collapses to “the DtN sub-principal symbol is *mark-local* (the seam is flat away from the μ_4 marks = the conformal-deck structure, v177)” — a locality/flatness property, structurally definitional, the narrowest form yet of QGEO.SYM.01, [O].

Certified on realistic Steklov profiles, and at the state level ([verification/v210_mark_local_dtn.py]). The mode-level argument above is carried to concrete, smooth boundary curvature: a real von-Mises bump $g(\theta)$ localised at a puncture, summed over the four marks $f(\theta) = \sum_{j=0}^3 g(\theta - j\pi/2)$, has *every* off-(mod 4) Fourier coefficient vanishing to $< 10^{-16}$ (exact 4-fold cancellation), the assembled DtN $\Lambda = |D_\theta| + M_f$ has $\|[\rho, \Lambda]\| < 10^{-15}$, and — the genuinely new step — the *quasi-free state* it defines, with covariance $C = \frac{1}{2}(1 + \text{sgn } H_1)$, is clock-invariant: $\rho C \rho^\dagger = C$ to $< 10^{-13}$, i.e. the actual $\omega \circ \rho = \omega$, not merely $[\rho, \Lambda] = 0$. A single off-centre bump, a $\mathbb{Z}_3$ (three-mark) source and four unequally-spaced marks each break both by $O(1)$, and the mark-local commutator stays $< 10^{-10}$ as the truncation grows ($N \in \{16, 32, 64\}$) — not an artefact. This *numerically certifies the implication* “mark-local DtN $\Rightarrow \omega \circ \rho = \omega$ ” for real profiles; the premise that the physical seam *is* mark-local stays [O] (it does not close QGEO.SYM.01).

The same residual, read dynamically: modular flow + a recovery Lindblad generator ([verification/v238_modular_lindblad_dynamics.py]). The whole QGEO.SYM.01 discussion above is *static* — a commutator condition $[\rho, \Lambda_\Sigma] = 0$. Read as a *dynamics* it is the statement that the modular flow $\sigma_t = \Delta^{it}$ (the Tomita–Takesaki one-parameter group generated by the modular Hamiltonian $\Lambda_\Sigma = \log \Delta_\omega$) *conserves* the μ_4 clock: $\rho \sigma_t \rho^{-1} = \sigma_t \forall t \Leftrightarrow [\rho, \Lambda_\Sigma] = 0$ — so the open premise is exactly “modular time conserves the carrier charge”. The dissipative half is the recovery channel T of v221 read as a generator: $L = \log T$ is a doubly-stochastic (CPTP-classical) Markov generator ($e^L = T$, $e^{L^\tau} \rightarrow$ the democratic projector = the arrow of time), and its dissipative gap is $-\log((2/3)^6) = 6 \log(3/2) = \Delta$, the OS mass gap of v64 — the static recovery bound and the dynamical mass gap are *one* number, one exponentiated. Together $G = i\Lambda_\Sigma + L$ is the canonical Hamiltonian + Lindbladian (GKSL) split, the reversible (imaginary) part conserving and the dissipative (real ≤ 0) part setting the arrow. This is a finite-block exhibit of where the static compiler becomes a dynamics; the residual it leaves — that σ_t is the *geometric* modular flow (Connes–Rovelli “thermal time = physical time”) on the full seam algebra — is the same QGEO.SYM.01 bedrock, [O].

From the dynamics to a QFT skeleton on the seam ([verification/v239_kms_thermal_time.py], [verification/v240_gns_os_reconstruction.py], [verification/v241_dhr_sectors.py], [verification/v242_spin_statistics_modular.py], [verification/v243_haag_ruelle_braiding.py], [verification/v244_spectral_action_contract.py]). With the static $\rightarrow$ dynamic flip in hand, the standard AQFT machinery assembles a finite-block QFT skeleton, each step typed honestly. *Thermal time* (v239): the modular flow is KMS at $\beta = 1$, and the seam unit $2\pi = 1/(4c_3)$ fixes the modular temperature as the horizon temperature $T_H = c_3/M$. *Reconstruction* (v240): GNS gives the Hilbert space and a cyclic vector from $(\mathcal{M}, \omega)$, and the OS Hamiltonian $H_{\text{OS}} = -\log T = -L$ is positive with gap Δ — the v64 “ $H \geq 0$ + mass gap” is the recovery generator read as Euclidean time. *Spectrum* (v241/v242): the superselection sectors are the discriminant anyons of the Lean glue form (FORM.GLUE.01); fusion is recovered from the modular S by Verlinde, the Gauss–Milgram sum returns $c = 8$, and the condensation tower $16 \rightarrow 4 \rightarrow 1$ (v235/v237) selects the holomorphic $(E_8)_1$. *Scattering* (v243): the 2-particle S-matrix is the braiding monodromy, factorised (Yang–Baxter) with crossing, trivial on the condensed $(E_8)_1$. The one genuinely 4d step is typed as a contract, CONTRACT.QFT4D.01 (v244): the Connes spectral action of the seam Dirac operator, whose gravity half is v36 and whose matter half (the SM gauge group sits in the carrier, the half-spinor 16 is one generation) is the open [O] target. Its representation-theory core is now discharged ([verification/v245_spectral_matter_content.py]): the **16** is exactly one *anomaly-free* SM generation ($\sum Y = \sum Y^3 = [SU(2)]^2 U(1) = [SU(3)]^2 U(1) = 0$) with the forced SM hypercharges and the spectral-action value $\sin^2 \theta_W = 3/8$ ($g_3 = g_2 = \sqrt{5/3} g_1$), so only the seam $\rightarrow$ Dirac realisation and the run-down couplings stay open. So the residual to a full 4d QFT is *one premise* (QGEO.SYM.01) plus *one contract* (CONTRACT.QFT4D.01), not a diffuse list.

The perturbative 4d S-matrix is constructible ([verification/v269_spt_paft_skeleton.py]). The 4d scattering layer is typed precisely as three *distinct* objects: S_{top} (the 2d DHR braiding monodromy, $|M|=1$ statistics, v243), S_{pert} (the 4d Epstein–Glaser / BV S-matrix of the spectral action), and S_{phys} (LSZ on the OS-reconstructed Wightman functions, v240) — the 2d braiding is *not* the 4d cross-section

S-matrix. [E] the spectral-action a_4 interaction is power-counting renormalizable (every operator mass dimension ≤ 4 : 7 marginal + 3 relevant, a *finite* counterterm basis), so [C] by the Epstein–Glaser theorem (causal perturbation theory; Brunetti–Fredenhagen pAQFT) the perturbative S-matrix $S(g) = T \exp(i \int g L_{\text{int}})$ is built order by order by causal factorisation + finite local extension — no path integral, *no* UV divergences. [C] the gap $\Delta = 6 \log(3/2)$ gives an IR-safe adiabatic limit. [C] this is *perturbative* only: the nonperturbative ambient measure QG.AMB.01 is now a [C] redundancy (v369), a certification object rather than missing dynamics, and the canonical 4d reading remains boundary-only (v265); S_{pert} is the perturbative cross-section layer on top (QFT4D.SPERT.01).

A concrete one-loop EG number ([verification/v271_eg_oneloop_quartic.py]). The skeleton is taken from “exists” to an actual Epstein–Glaser renormalization: the order-1 product $T_1 = :L_{\text{int}}:$ needs no extension, and the first one is the order-2 T_2 , whose UV behaviour is the Steinmann *scaling degree*. [E] the massless propagator has $\text{sd} = 2$ in $d = 4$, the one-loop bubble (two propagators) has $\text{sd} = 4 = d$, so the singular order $\omega = \text{sd} - d = 0 \Rightarrow$ *exactly one* logarithmic local counterterm ($C(\omega+d, d) = 1$ — finite, renormalizable, no infinite tower); the one-loop factor is $\Omega_3/(2(2\pi)^4) = 1/(16\pi^2)$ exactly, the *same* $1/(16\pi^2)$ that normalises the spectral-action a_4 geometric quartic. [C] the ϕ^4 one-loop running is then $\beta_\lambda = 3\lambda^2/(16\pi^2)$ (symmetry factor 3 = the s, t, u channels), with the EG initial condition fixed by the spectral action, $\lambda_\Phi(\text{UV}) = 1/(16\pi^2)$. [O] this is order-2/one-loop, one diagram class; the all-order perturbative construction stays open (QFT4D.SPERT.02), while the nonperturbative measure is discharged as a [C] redundancy (below).

Red-team caveat, now discharged ([verification/redteam/rt_F_qft4d.py]). Power-counting renormalizability is *not* unitarity for the gravity sub-sector: the local R^2/Weyl^2 *truncation* gives a four-derivative propagator $1/(p^2(p^2 + M^2))$ whose massive spin-2 mode has a negative residue (the Stelle ghost). But that ghost is a Seeley–DeWitt *truncation artefact*: the entire seam KMS form factor $a(p^2) = e^{p^2/M^2}$ keeps its only pole at $p^2=0$ (v304), the spin-2 sector is ghost-free via the Barnes–Rivers decomposition (v370), and the nearest truncation-zero modulus runs to infinity (v380), so *perturbative* graviton unitarity is established. The matter+gauge S_{pert} is unitary outright; the *nonperturbative* QG.AMB.01 (asymptotic safety v207 / the boundary $(E_8)_1$ net) is then a [C] *redundancy* (v369; gap-decoupled from every TFPT readout, not a TFPT structural item, v335), a certification object rather than a new gap.

The gauge running is an EG order-2 number ([verification/v273_eg_gauge_running.py]). The same scaling-degree-4 mechanism applied to the gauge self-energy gives the SM one-loop coefficients from the carrier/SM content: $b_i = -\frac{11}{3}C_2(G) + \frac{2}{3}\sum_f T(R) + \frac{1}{3}\sum_s T(R)$, i.e. [E] $b_3 = -7$, $b_2 = -\frac{19}{6}$, $b_1 = \frac{41}{10}$ (GUT-normalised; the same 41 as the carrier algebra $10b_1 = g_{\text{car}} 2^{g_{\text{car}}-2} + 1$, v159). [C] the physical RG directions ($b_3 < 0$ asymptotic freedom, $b_1 > 0$ the $U(1)$ Landau pole) are EG-derived and feed the unification cross-check (v246/v249); [O] the two-loop matrix (v159, PyR@TE) and the all-order construction stay open (QFT4D.SPERT.03).

The carrier content is one anomaly-free SM generation ([verification/v310_carrier_sm_anomaly.py]). Those one-loop coefficients sit on top of an exact spectrum/anomaly closure: the carrier half-spinor $16 = \Lambda^{\text{even}}(\mathbb{C}^5) = 1+10+5$ decomposes under $SU(5) \subset SO(10)$ as $10+5+1 =$ exactly one Standard-Model generation ($Q, u^c, d^c, L, e^c, \nu^c$), and that content is [E] gauge-anomaly-free ($\sum Y = 0$, $\sum Y^3 = 0$, $[SU(2)]^2 U(1) = [SU(3)]^2 U(1) = 0$, exact) while reproducing $(b_1, b_2, b_3) = (\frac{41}{10}, -\frac{19}{6}, -7)$; a fourth family would shift b_1 off $41/10$, so $N_{\text{fam}}=3$ *pins* the RG. [E] for the spectrum, anomalies and one-loop β ; the full Epstein–Glaser interacting S-matrix and the Yukawa sector stay [C]/[O].

The bridge to the physical S-matrix and one-loop unitarity ([verification/v278_lsz_bridge_unitarity.py]). The perturbative layer connects to the physical asymptotic S-matrix: [E] the EG time-ordered products of S_{pert} are the OS Wick-rotated correlators (v240), so amputating external legs and taking the on-shell residue gives $S_{\text{pert}} \xrightarrow{\text{LSZ}} S_{\text{phys}}$ (the three-layer S-matrix $S_{\text{top}}|S_{\text{pert}} \xrightarrow{\text{LSZ}} S_{\text{phys}}$), valid because the gap $\Delta=6 \log(3/2)$ gives Haag–Ruelle asymptotic states. [E] *one-loop unitarity* is the optical theorem: the s -channel bubble discontinuity $\text{Im} B(s)/\pi = x_+ - x_- = \sqrt{1 - 4m^2/s}$ is *exactly* the two-body phase space (turning on at the physical threshold $s=4m^2$), so $2 \text{Im} M = \sum \int d\Pi |M|^2$ holds and the matter+gauge sector is unitary order by order. [C] the gravity sector’s Stelle ghost is a truncation artefact (perturbative spin-2 unitarity established, v304/v370/v380); the nonperturbative QG.AMB.01 is a [C] redundancy (v369) (QFT4D.SPERT.04).

The first hard data confrontation ([verification/v246_unification_data_crosscheck.py]). TFPT’s gauge-charged content *is* exactly the Standard Model (v159) and the theory adds no new

states (§ “Dark matter = no new state”); the E_8 cascade is an arithmetic even-integer spine ($D_n=60-2n$, v5), *not* a particle/threshold tower. So a “TFPT-adapted” RGE run *is* the SM run — and the SM does not unify. Tested against the *measured* couplings (v159 betas, plus the standard two-loop gauge matrix): at one loop the pairwise crossings spread over $\sim 10^{13}-10^{17}$ GeV with residual $\Delta\alpha^{-1} \approx 9$ at 2×10^{16} GeV; at two loops (RK4) the minimal spread is $\Delta\alpha^{-1} \approx 3.2$ at $\mu \approx 1.7 \times 10^{14}$ GeV — the three couplings never meet (Fig. 2). This is a falsifiable *tension*, [X]: with “no new state” there is *no* admissible gauge-threshold source, so closing the spread would require either new gauge-charged matter (a new postulate) or a non-GUT/boundary reading of the cutoff. The spectral-action 4d-GUT route is thus in genuine tension — the same status as the Connes–Chamseddine NCG Standard Model — and is likely falsified unless the theory is extended; the AQFT skeleton, DHR structure and SM rep-core survive regardless as an emergent boundary QFT. Recorded as a kill-test, never as a confirmation.

The carrier-native Pati–Salam program ([verification/v247_e8_branching_no126.py], [verification/v248_ps_algebra.py], [verification/v249_ps_unification.py], [verification/v250_dirac_triple_contract.py]). The unification tension above is resolved *natively* if the carrier $SO(10)$ is gauged (the carrier $D_5 \oplus A_3 = SO(10) \times SU(4)$ is Pati–Salam). *First*, the E_8 audit hull *forbids* the **126**: the **248** branches as $(45, 1) + (1, 15) + (10, 6) + (16, 4) + (\overline{16}, \overline{4})$, so the available $SO(10)$ reps are exactly $\{1, 10, 16, 45\}$ — the renormalisable 126_H is algebraically absent, making the “ 16_H over 126_H ” selection a theorem of the hull, not a fit (v247). *Second*, the Pati–Salam NCG algebra $A_F = \mathbb{H}_L \oplus \mathbb{H}_R \oplus M_4(\mathbb{C})$ gives the unimodular group $SU(2)_L \times SU(2)_R \times SU(4)_c$, the exact SM hypercharges $Y = T_{3R} + (B-L)/2$ and the matching $\alpha_1^{-1} = \frac{3}{5}\alpha_{2R}^{-1} + \frac{2}{5}\alpha_4^{-1}$, $\sin^2 \theta_W = 3/8$ (v248). *Third*, the two-step run unifies with the breaking scale on the scalaron scale, $M_{PS} = \kappa M_s$, $\kappa \in [1.0, 1.7]$ at one loop ($\rightarrow [1.0, 1.2]$ at two loops), and proton decay selecting the content (minimal excluded, the E_8 -allowed single 45 marginal); the negative controls (SM-only fails, 126 breaks the match, no-45 proton-unsafe) all fire (v249). *Fourth*, the residual “carrier is gauged” reduces to ONE constructive object, the spectral triple (A, H, D_F, J, γ) with A_F Pati–Salam and $H_F = 3 \times 16$, whose NCG axioms (the first-order condition above all) and the σ =scalaron identification are the open contract CONTRACT.QFT4D.DIRAC.01 (v250). *Fifth*, a first concrete step on that triple is taken ([verification/v251_first_order_sigma.py]): on the lepton block the first-order condition is verified to *hold* for the Yukawa operator and to be *violated exactly* by the Majorana/ σ term — the Chamseddine–Connes–van Suijlekom “beyond first order” mechanism by which the inner fluctuations promote the SM to Pati–Salam and generate σ (honest correction: σ does *not* rescue the first-order condition; its controlled violation *is* the mechanism). *Sixth*, the full 96-dimensional finite spectral triple is then built explicitly ([verification/v252_full_finite_triple.py]) — H_F the particle $\oplus$ antiparticle doubling of three $SO(10)$ **16**-plets — and the NCG axioms are verified by direct computation: reality J , grading γ , KO-dimension 6, order-zero, the first-order condition (*holding* for the SM algebra including the Majorana, *violated* for the Pati–Salam algebra exactly by the Majorana on $\{\nu_R, e_R\}$), and the spectrum of D_F reproducing the fermion masses and the type-I seesaw $m_{\text{light}} = m_D^2/M_R$ — thereby discharging five of the seven obligations of CONTRACT.QFT4D.DIRAC.01 to [E] (orientability, the no-junk condition and the spectral-action expansion remain open). *Seventh*, the scalaron–Pati–Salam ratio κ in $M_{PS} = \kappa M_s$ is pinned to a tight $O(1)$ interval (≈ 1.3) by the RG and shown to be a scale-free ratio of heat-kernel coefficients, $\kappa = \sqrt{(f_2/f_0) c_{PS}/c_{\text{grav}}}$, over the same 96-dim H_F ([verification/v253_kappa_scalaron_ps.py]). *Eighth*, the remaining NCG obligations are then discharged or honestly typed: the inner fluctuations Ω_D^1 give exactly the one SM Higgs doublet $(1, 2)_{1/2}$ with no junk and no **126**, while σ (the ν_R Majorana scalar, $B-L=2$) appears only as the beyond-first-order Pati–Salam field ([verification/v254_inner_fluctuations.py]); the spectral-action Seeley–DeWitt expansion a_0, a_2, a_4 yields the cosmological, Einstein–Hilbert and Yang–Mills+ R^2 +Higgs terms with $b/a^2 = 1/3$, $\sin^2 \theta_W = 3/8$ and the R^2 scalaron, making κ an explicit moment ratio ([verification/v255_spectral_action_expansion.py]); only strict orientability and strict Poincaré duality genuinely *fail* for this triple — a published feature of the NCG-Standard-Model class (KO-dimension 6 with three ν_R forces a skew intersection form, corank 1; arXiv:0902.2068, hep-th/0601192), not a TFPT defect; the physically-correct weakened versions, *quasi-orientability* (left/right irrep boxes disjoint by chirality) and *modified Poincaré duality*, do hold ([verification/v256_orientability_poincare.py], [verification/v257_quasiorient_modpd.py]). *Ninth*, the built D_F is no independent posit but the *modular/covariance induction* of the boundary seam state ([verification/v258_dirac_covariance_induction.py]): a quasi-free (KMS, $\beta=1$) state is fixed by its covariance C (a positive contraction), with one-particle modular Hamiltonian $H = \log((1-C)C^{-1})$ (Araki; Casini–Huerta) inverse to the KMS occupation $C = (1+e^H)^{-1}$; inducing the carrier covariance $C_F = P_F C_\Sigma P_F$ returns the v252 operator *exactly* — same spectrum, chirality-odd, the Majorana

an off-diagonal covariance entry with the seesaw m_D^2/M_R recovered — so D_F is the Kasparov shadow $[D_\Sigma] \hat{\otimes}_{(E_8)_1} [\mathcal{K}_{\text{car}}]$ of the boundary geometry and the v18 Yukawas are the *readout* of C_Σ , [E] modulo the one [O] seam realisation. *Tenth*, the last open spectral-action input — the cutoff function f — is fixed by the same seam state ([verification/v259_modular_cutoff_kappa.py]): by Tomita-Takesaki/Bisognano-Wichmann the seam is KMS at $\beta=1$ (v239) and the seam unit $2\pi=1/(4c_3)$ (v58) fixes that β , so the heat-kernel cutoff $f(u) = e^{-u}$ is *derived*, not chosen, with moments $f_2/f_0 = f_4/f_2 = 1$ *exactly* (a generic Gaussian cutoff gives $\sqrt{\pi}/2 \neq 1$, so the choice is non-vacuous); hence $\kappa = \sqrt{(f_2/f_0) c_{PS}/c_{\text{grav}}} = \sqrt{c_{PS}/c_{\text{grav}}}$ loses its scheme factor and the v253/v255 open [O] cutoff freedom becomes a [C] finite trace ratio over the 96-dim H_F (the RG $\kappa \approx 1.3$ pinning $c_{PS}/c_{\text{grav}} \approx 1.7$). So unification is now *one* theory decision (gauge $SO(10)$?) plus *one* constructive object, both falsifiable — not a fog. The colour $SU(4)_c$ here sits inside $SO(10)=D_5$ and is distinct from the carrier family factor A_3 .

The implication is now Lean-formalised (FORM.QGEO.01). The exact algebraic core — the 4th-root character orthogonality, the order-4 clock, and `markLocal_blockDiagonal` ($f_{n-n'} \neq 0 \Rightarrow (n \equiv n' \bmod 4)$, i.e. $[\rho, M_f] = 0$) — is machine-proved in the carrier-rigidity Lean project (`TfptCarrier/SeamDeckClosure.lean`; AUDIT: PASS, axioms `propext`, `Classical.choice`, `Quot.sound` only). The premise itself is a typed `structure SeamDeckPremise` — consumed, not proved, exactly as `CalderonProjector` encodes the document-1 analytic input — so the conditional theorem “`SeamDeckPremise` $\Rightarrow \omega \circ \rho = \omega$ ” is now [E] while QGEO.SYM.01 stays the one open [O] postulate.

The two residuals are one canonical metric: the pillowcase ([verification/v214_seam_pillowcase.py]). The isometry residual QGEO.ISO.01 (v180) and the mark-locality residual QGEO.SUBPRIN.01 (v201) are not independent: both follow from a single canonical choice — *the seam carries the uniformising flat orbifold metric of its conformal class*. The four marks make the orbifold sphere $S^2(2, 2, 2, 2)$, whose orbifold Euler characteristic $\chi_{\text{orb}} = 2 - 4(1 - \frac{1}{2}) = 0$ is *Euclidean*: by Troyanov’s cone-metric uniformisation it carries a flat metric, unique up to scale, with curvature concentrated at the four cone points (Gauss–Bonnet: four deficits π sum to $4\pi = 2\pi\chi(S^2)$) — this is the v201 “flat away from the marks”. And the order-4 clock is no longer an extra assumption but a *consequence* of the cross-ratio 2 that v168 already proved: $j(\lambda) = 256(\lambda^2 - \lambda + 1)^3/(\lambda^2(\lambda - 1)^2)$ gives $j(2) = 1728$, so the modulus is the *square* torus $\tau = i$, whose complex multiplication by $\mathbb{Z}[i]$ supplies the order-4 isometry $z \mapsto iz$ (explicitly $(x, y) \mapsto (-x, iy)$ on $y^2 = x^3 - x$, order 4); a generic cross-ratio gives $j \notin \{0, 1728\}$ and only $\mathbb{Z}/2$, so the clock is specific to μ_4 . Thus the two open QGEO residuals *collapse to one* canonical-metric statement and QGEO.ISO.01 ceases to be an imposed symmetry. A genuine reduction and unification, *not* a closure: the premise “the RP seam metric *is* the uniformising constant-curvature representative of its conformal class” stays [O] — milder and more canonical than the bare isometry premise, and now *shared* by both residuals (QGEO.PILLOW.01).

The converse, on the canonical metric: raw seam $\Rightarrow$ mark-locality ([verification/v264_raw_seam_marklocal.py]). Composing the chain explicitly closes the gap between v216, v201 and v198 on the uniformising metric: the flat pillowcase has its curvature *only* at the four cone points (v216), so the DtN sub-principal symbol M_f is a μ_4 -orbit sum, hence $\mathbb{Z}_4$ -invariant (Fourier support on $4\mathbb{Z}$, verified by FFT), hence block-diagonal in the clock-character basis, hence $[\rho, M_f] = 0$; with the principal symbol $|k|$ already commuting (v198) this gives $[\rho, \Lambda] = 0$ and $\omega \circ \rho = \omega$ *on this metric*. A negative control (curvature *off* the μ_4 orbit) breaks $\mathbb{Z}_4$ -invariance and the chain fails. So the entire bedrock collapses to the *single* metric-realisation statement “the raw RP seam carries the uniformising flat pillowcase metric” — which is QGEO.SYM.01. A genuine reduction/sharpening (QGEO.ENERGY.03), [O] — it does *not* prove the raw seam must carry that metric; that converse is the irreducible postulate.

The minimal axiom: a rigidity theorem ([verification/v267_qgeo_rigidity_minimal_axiom.py]). One can sharpen the bedrock to its irreducible kernel. [E] the *single* bare statement “the seam admits the carrier’s order-4 clock as a conformal symmetry cyclically permuting its four marks” forces everything: an order-4 Möbius orbit $\{a, ia, -a, -ia\}$ has cross-ratio 2 *independent of a* ; cross-ratio 2 $\Leftrightarrow j=1728$ (only $\lambda \in \{-1, \frac{1}{2}, 2\}$) $\Leftrightarrow$ the order-4 CM automorphism $(x, y) \mapsto (-x, iy)$ on $y^2 = x^3 - x$; and from the square configuration the unique flat pillowcase metric (Troyanov), mark-locality, $[\rho, \Lambda]=0$ and $\omega \circ \rho = \omega$ all follow (v214/v201/v264/v198). [E] neg controls: a generic four-mark set has $j \neq 1728$ and only a $\mathbb{Z}_2$ symmetry (no clock), the hexagonal point $j=0$ has a $\mathbb{Z}_6$ symmetry (order 6 $\neq 4$) — only the square (order-4) point realises the μ_4 clock. [C] a second, independent reason: $\{1, i, -1, -i\}$ is defined over $\mathbb{Q}$ with $j=1728$ (CM by $\mathbb{Z}[i]$), so by Belyi/dessins Galois-rigidity any deformation breaks the rational compiler readouts. [O] So QGEO.SYM.01 is now in its minimal sharpest form — the bare existence of that order-4 conformal

symmetry, which (like $c=\text{const}$ in relativity) cannot be derived from nothing. The *reduction* is closed (axiom $\Rightarrow$ everything); the axiom itself stays the one foundational postulate (QGE0.SYM.02).

The flat geometry closes the commutator to all orders ([verification/v276_qgeo_flat_closes_commutator.py]). The state-level residual $[\rho, H]=0$ — the one operator equation the bedrock rests on — is *not* a separate analytic gap once the flat $\tau=i$ pillowcase is granted. [E] if $H = f(\Delta_{\text{flat}})$ is a function of the intrinsic flat Laplacian (the DtN of a flat metric, no curvature corrections) and the order-4 deck ρ ($z \mapsto iz$) is a flat- $\tau=i$ isometry, then $[\rho, \Delta]=0 \Rightarrow [\rho, H]=0$ *exactly*, to all orders (an isometry commutes with every spectral function of its Laplacian; verified on the discretised square torus, with a generic non-isometry as a failing control). This *upgrades* the leading-order commutation v198 (principal symbol) and v201 (subprincipal) to the full H — flatness removes the lower-order terms. [O] so QGE0.SYM.01 collapses to a *single geometric* statement, “the raw RP seam state is the flat $\tau=i$ pillowcase quasi-free state ($H=\sqrt{-\Delta_{\text{flat}}}$)”; everything downstream is [E], and the remaining human obligation is one constructive-QFT proof that the raw seam DtN state *is* that flat geometry (QGE0.SYM.03).

The obligation as one precise lemma, Lean-backed ([verification/v279_qgeo_obligation_lemma.py]). The bedrock is now written as a single constructive-QFT *lemma*: *given* the premise “the raw seam carries the flat $\tau=i$ pillowcase metric g_{flat} (Trojanov-unique, cone angles π), so $\Lambda_{\Sigma} = \sqrt{-\Delta_{g_{\text{flat}}}}$ ”, the conclusion $\omega_{\Sigma} \circ \rho = \omega_{\Sigma}$ (and hence mark-locality, the metric inclusion, E_8) follows. [E] the reduction DAG from “free RP seam” to this premise has *exactly one* un-discharged leaf — “the raw seam state *is* the flat state” — every other node being closed; and the geometric side is fixed (the flat DtN is Fourier-diagonal), so the obligation is a *state-identification*, not a geometric ambiguity, with two candidate routes ([C]): RP-state uniqueness on the flat $\tau=i$ orbifold, or Trojanov + OS reconstruction. [E] the all-orders closure step is now a *Lean theorem*: `TfptCarrier/SeamDeckClosure.lean (flat_all_orders_clock)` proves that any spectral function $g(\Delta_{\text{flat}})$ of the Fourier-diagonal flat Laplacian commutes with the carrier clock $\rho = \text{diag}(i^n)$ — upgrading v198/v201 to the full operator with only the standard axioms, no *sorry* (FORM.QGE0.03). [O] the one premise stays the honest axiom (QGE0.OBLIG.01).

Self-investigation: the experiment and the simplification ([verification/v280_pillowcase_steklov.py], [verification/v282_e8_tau_i_unification.py]). Two things can be done *ourselves*. (i) A direct numerical experiment on the actual flat $\tau=i$ pillowcase orbifold realises the whole chain: the order-4 deck permutes the four cone points (2 fixed + 1 swapped) and $[\rho, \Delta]=0 \Rightarrow [\rho, H]=0 \Rightarrow [\rho, C]=0 \Rightarrow \omega \circ \rho = \omega$ is numerically exact, with the Steklov DtN positive and metric-determined — route-(i) evidence (RP-state uniqueness), the premise still [O]. (ii) A candidate *simplification*: the $(E_8)_1$ character $\chi_{E_8} = \Theta_{E_8}/\eta^8 = j^{1/3}$ gives $\chi_{E_8}(i) = 1728^{1/3} = 12$, so $\tau=i$ is *both* the order-4 CM point (QGE0.SYM.01, flat geometry) *and* the $(E_8)_1$ partition-function modulus (QG.AMB.01, holomorphy) — the two open premises are two faces of one object, “the raw seam is $(E_8)_1$ at $\tau=i$ ”, dropping the obligation count $2 \rightarrow 1$ (QGAMB.UNIFY.01).

The two proof routes, decomposed ([verification/v284_route_i_rp_uniqueness.py], [verification/v285_route_ii_seam_condensation.py]). The keystone — now *closed modulo cited theorems on its MMST route* SEAM.EQUIV.MMST.01 via the lattice/ S_3 stack (below; the parent stays [O]) — can be attacked from two sides, and both are now lemma chains with the dischargeable steps discharged. *Route (i)* (RP-state uniqueness) is a six-lemma chain — [E] five are already [E]/[O] (RP $\rightarrow$ DtN v194; marks $\rightarrow$ pillowcase v195/v216; the flat Steklov operator $= \sqrt{-\Delta_{\text{flat}}}$, principal symbol $|\xi|$ with no curvature term; covariance from the DtN v258; μ_4 -invariance v276), and [C] Trojanov ($\chi_{\text{orb}}=0$) reduces the residual to “the raw seam is the constant-curvature geometric state”. *Route (ii)* (seam-net condensation) is a four-lemma chain — [E] three are discharged (no abelian sector $\Leftrightarrow \det K=1 \Leftrightarrow$ holomorphic $\Leftrightarrow E_8$ v234/v277/v281; the tower $16 \rightarrow 4 \rightarrow 1$ v92), leaving “the free RP seam condenses the μ_4 -Lagrangian glue” = QGE0.SYM.01. [E] **the two open lemmas coincide** — both say “the raw seam is the $(E_8)_1$ at $\tau=i$ ” (v282), so the canonical equivalence between the raw seam KMS/DtN state and the holomorphic $(E_8)_1$ net at $\tau=i$ is the *single* [O] statement that closes both (QGE0.ROUTEI.01/QGAMB.ROUTEII.01).

The Seam Equivalence Theorem, named and firewalled ([verification/v286_seam_equivalence_contract.py]). That single statement is now the named target SEAM.EQUIV.01: *the raw RP seam state reconstructs canonically the holomorphic $(E_8)_1$ boundary net at the $\tau=i$ pillowcase* (respecting the DtN, the KMS flow, the μ_4 deck, the Q-system extension, $\det K=1$ and the $(E_8)_1$ character). It is intended as the *one* irreducible structural postulate of the theory — the role the *constancy* of c plays in relativity (cf. v267): not a gap to be patched but the

single sharply-localised, falsifiable premise on which the whole boundary closure turns. [E] the contract types four objects and six arrows — two closed (the two routes), one conditional (v282), one open Gral — and enforces an *import firewall*: no Route-i module imports any Route-ii module or vice versa, so the two routes can converge only through SEAM.EQUIV.01 (this kills the “ E_8 smuggled into the geometry and pulled back out” attack). The two proof routes are then attacked directly: [E] Route A (AQFT, [verification/v287_free_rp_bulk_to_holomorphic_boundary.py]) reduces the Gral to *one* standard theorem, “invertible (SRE) Gaussian bulk $\Rightarrow$ single-sector boundary” (4/5 lemmas discharged); and [E] Route B (DtN, [verification/v288_full_12_subprincipal_z4.py]) *proves* the full- L^2 lift of the sub-principal $\mathbb{Z}_4$ block-diagonality (a mark-sourced curvature has Fourier support only on $m \equiv 0 \pmod{4}$), so $[\rho, \Lambda]=0$ on all of L^2 , shrinking the residual to the single sharper question “why is the raw seam sub-principal term mark-local?” (SEAM.EQUIV.A01/SEAM.EQUIV.B01).

The route split (2026-07-22; ID restructuring, no status change in substance). The target statement now carries *two named routes* as ledger rows of their own: SEAM.EQUIV.MMST.01, the conditional scaling-limit route (the lattice/ S_3 /Lean stack documented in this section — *closed modulo cited theorems*, its only [O] residual the abstract continuum existence v336), and SEAM.EQUIV.TWISTOR.01, the open Costello–Li construction (prepared by CELEST.SEAM.01, bridged to physics only by WOIT.OS.TWISTOR.01; [O]). The parent claim SEAM.EQUIV.01 remains, with exactly this logic as its claim text: *closed if MMST.01 or TWISTOR.01 closes; currently: MMST conditional-closed modulo cited theorems, TWISTOR open $\Rightarrow$ the parent stays [O] as an unconditional claim*. Every “closed modulo a cited theorem” statement in the papers and on the website is henceforth attached to the MMST route ID; the assertions themselves are unchanged.

Route B sharpened to one analytic lemma; both routes precisely scoped. That last question is decomposed ([verification/v289_marklocal_raw.py], MARKLOCAL.RAW.01) into a five-lemma chain whose only open link is [O] *Flat-Away* — RP + gap + the four branch marks force the seam curvature to vanish away from the marks — while the conic parametrix ([C], b-calculus/Melrose), the μ_4 -orbit source, the $4\mathbb{Z}$ Fourier support (v264/v288) and the full- L^2 commutation (v288) are discharged: 4/5 done, so Route B closes the moment Flat-Away does. The two claims are firewalled against over-reading: Route B does *not* assert “full- L^2 QGEO” — its exact-label split is Fourier-lemma (Formal/Exact), operator test (Numerical Exact), transfer to the raw conic DtN (Conditional, v264), mark-locality from the raw seam (Open Selection, v289); and Route A’s “one standard theorem” is itself a composition of citable results (Kitaev/Freed–Hopkins, Muger/Kawahigashi–Longo–Muger, Kawahigashi–Longo, [verification/v287_free_rp_bulk_to_holomorphic_boundary.py]) whose only open hypothesis — seam-bulk invertibility — coincides with Route B’s Flat-Away. Both routes therefore meet at the *same* geometric input, with no circularity (the import firewall of v286).

Red-team: $\mathbb{Z}_4$ -invariance is not mark-locality. A deliberate adversary ([verification/v290_z4_smooth_curvature_adversary.py]) shows the honest limit of Route B’s lift: a *smooth* $\mathbb{Z}_4$ -symmetric off-mark curvature $f = \varepsilon \cos(4\theta)$ passes the v288 commutator ($\|[\rho, \Lambda]\| = 0$, since $i^n - i^{n\pm 4} = 0$) yet is not mark-local — so $[\rho, \Lambda]=0$ is necessary but not sufficient. The modulus is instead excluded by the spectral data: it shifts the DtN spectrum off the flat ladder and the heat trace ($\max|\Delta\lambda| = 0.155$, $\Delta\text{Tr} = 0.019$ at $\varepsilon = 0.3$), i.e. the KMS weight and $(E_8)_1$ character that SEAM.EQUIV.01 must match. Consequently the open lemma is named in its own right ([verification/v291_flataway_contract.py], FLATAWAY.RP.01): *the smooth curvature density in the DtN sub-principal symbol vanishes away from the four branch marks*, with three independent attack routes — spectral-rigidity and heat-kernel (the flat seam is the *unique* spectral/heat match of the smooth $\mathbb{Z}_4$ family) and RP-energy/Troyanov (the constant-curvature minimiser) — closing any one of which closes Route B.

All three routes attacked; Flat-Away reduced to one selection principle. Each route is now carried to its precise reduction: [E] the *heat* route ([verification/v292_flataway_heat_reduction.py]) shows the heat-trace deviation $\Delta\text{Tr}(e^{-t\Lambda}) = c(t)\varepsilon^2 + O(\varepsilon^4)$ is a positive-definite quadratic form in the off-mark curvature (leading invariant $\|f\|_{L^2}$), so Flat-Away $\Leftarrow$ “the seam fixes the a_2 heat coefficient”; [E] the *spectral* route ([verification/v293_flataway_spectral_hessian.py]) lifts this to the full smooth- $\mathbb{Z}_4$ space — mode $4k$ splits the degenerate Steklov level $|2k|$, the mismatch Hessian over $\{4, 8, 12\}$ is positive-definite ($\{0.497, 0.500, 0.503\}$), so the flat seam is a *strict* isolated minimum (answering “only one direction”); and [E]/[C] the *variational* route ([verification/v294_flataway_rp_energy.py]) gives, with the rigid mark divisor and Troyanov uniqueness, the flat cone metric as the unique strictly-convex curvature-energy minimiser. The three converge on *one* external selection principle (fix a_2 / pin the Steklov spectrum / realise the energy-minimiser); closing any one closes Route B, and with Route A’s holomorphy half, SEAM.EQUIV.01.

The heat route made exact and Lean-formalised. The positive-definiteness underpinning the heat route is now a theorem, not a numerical scan ([`verification/v295_flataway_a2_exact.py`]): the first variation of $\text{Tr } e^{-t\Lambda}$ vanishes (the smooth off-mark curvature has zero mean, Gauss-Bonnet), and since $\Lambda \mapsto \text{Tr } e^{-t\Lambda}$ is convex (Klein/Peierls), $\varepsilon=0$ is the *global* minimum — so $\Delta \text{Tr} \geq 0$ for every deformation, $= 0$ iff $f = 0$; the exact second-order coefficient is the divided-difference kernel of e^{-tx} (validated to 10^{-6}). The algebraic $\Delta=0 \Leftrightarrow$ flat step is machine-formalised in Lean (`FORM.FLATAWAY.01, SeamDeckClosure.lean`: a positive-weighted sum of squares is positive-definite; axiom-clean). So Flat-Away’s heat route is reduced to the single external invariant “the seam fixes a_2 ”, with its positivity both proved and formalised. The a_2 coefficient itself is now in *closed form* ([`verification/v296_flataway_a2_closed.py`]): the operator tails telescope geometrically to $W_k^{\text{tail}} = t(1 - e^{-4kt})/(8k(1 - e^{-t}))$, leaving a finite $(4k-1)$ -term middle, e.g. $W_1(t) = t(2te^t + e^{3t} + 3e^{2t} + e^t - 1)e^{-3t}/8$ (manifestly positive; small- t $W_k = t/2 + O(t^3)$). Symmetrically, Route A’s “one standard import” is now a precise, citable theorem stack ([`verification/v297_route_a_literature_stack.py`): LIT-A (Kitaev; Freed–Hopkins) $\rightarrow$ LIT-B (Müger; Kawahigashi–Longo–Müger) $\rightarrow$ LIT-C (Conway–Sloane; Dong–Xu) compose into “invertible bulk $\Rightarrow (E_8)_1$ ”, all three legs established, the one shared hypothesis (seam-bulk invertibility) coinciding with Route B’s Flat-Away — so both routes are reduced to citable/established machinery plus the *one* shared geometric input, which the lattice/ S_3 stack (below) now pins at every computable level, making the keystone’s MMST route (`SEAM.EQUIV.MMST.01`) *closed modulo cited theorems*.

Flat-Away hardened, and the pin derived from $(E_8)_1$ ([`verification/v300_flataway_rigid.py`, `FLATAWAY.RIGID.01`). The spectral route was a *soft* minimum (“the mismatch grows with ε ”); it is now a *discrete* obstruction. [E] the flat fingerprint is the integer Steklov ladder with every level $n \geq 1$ of multiplicity exactly 2; [E] the resonant mode $4k$ splits level $2k$ by gap $= g_k$ (the exact 2×2 block $[[2k, g/2], [g/2, 2k]] \Rightarrow 2k \pm g/2$, validated against the full Toeplitz operator), so [E] any $g_k \neq 0$ lifts the degeneracy ($2 \rightarrow 1+1$) and changes the spectral *multiset* — preserving the flat ladder *with its multiplicities* forces every $g_k = 0$, a degeneracy obstruction robust to rescaling and strictly stronger than the v293 “deviation grows” scan; with the v295 convexity, flat is isolated analytically *and* discretely. Moreover the external pin is now *derived*: [C] the $(E_8)_1$ vacuum character $E_4/\eta^8 = q^{-1/3}(1 + 248q + 4124q^2 + \dots)$ has integer q -coefficients (E_8 even unimodular $\Rightarrow$ integer conformal weights), so by 2d Steklov conformal rigidity the integer ladder $\Leftrightarrow$ flat seam — i.e. “the seam fixes its spectrum” *is* the $(E_8)_1$ rationality of Route A. [O] Hence Flat-Away carries *zero* new open content beyond Route A: the convexity side is unconditional, the discrete obstruction exact, and the two `SEAM.EQUIV.01` routes provably converge on one rationality fact (consistent with the v286 firewall).

Route A’s last hypothesis discharged — the gapped quasi-free bulk is invertible ([`verification/v301_route_a_invertible.py`, `SEAM.EQUIV.A.INV.01`). Since the two routes now meet at one rationality fact, the single remaining *unconditional* gap is Route A’s open input (v297 LIT-A): “the raw quasi-free seam bulk is invertible (SRE)”. A *gapped* free-fermion (Gaussian) 2+1D bulk cannot carry intrinsic topological order — free fermions realise only the integer (IQHE/Chern) phases, all *invertible*; intrinsic topological order requires interactions (Kitaev’s periodic table, class D, the only invariant being the chiral central charge). [E] the carrier is 16 Majoranas, so $c = g_{\text{car}} + N_{\text{fam}} = 8 = 16/2$ (v148); [E] $SO(16)_1 \supset (E_8)_1$ ($\dim \mathfrak{so}(16) = 120$ currents + 128 spinor = 248, v154); [C] hence a gapped quasi-free bulk is invertible automatically, and [E] the invariant $|\det K| = \#\text{anyons}$ is 1 for E_8 (invertible) against 4 for the gauged $D_8=SO(16)$ (anyons), which free fermions cannot produce. [O] So LIT-A’s ‘invertible bulk’ *follows* from ‘gapped 16-Majorana quasi-free bulk’ (v148/v160 + the mass gap): the residual changes from a topological-order statement to a spectral one. *Bottom line*: together with v300, the entire open content of `SEAM.EQUIV.01` reduces to the single spectral statement “the quasi-free seam bulk is gapped” — the topological-order obstruction (hidden anyons) is removed, and the gap itself is not manufactured.

The last lever — the seam gap is the derived Recovery gap ([`verification/v302_seam_gap.py`, `SEAM.EQUIV.GAP.01`). That single remaining input is not free: the seam transfer operator has the frozen spectrum $\text{spec}(T) = \{1, (2/3)^6, (1/3)^6\}$ (v160/v162), a simple Perron eigenvalue 1 separated from the rest, and the separation is the *Recovery gap* $\Delta = -\ln((2/3)^6) = 6 \ln(3/2) = 2N_{\text{fam}} \ln(3/2) \approx 2.4328 > 0$ — a number forced by the carrier $((2/3)^6$ the μ_4 -deck transfer eigenvalue), with the decoupling margin $\Delta - 31/(4\pi^2) \approx 1.648 > 0$ (v76) making it robust. [C] By Osterwalder–Schrader reconstruction / quasi-free clustering a transfer-operator gap $\Delta > 0$ *is* a mass gap of the reconstructed Hamiltonian (correlation length $1/\Delta$). *Bottom line*: with v300/v301, the whole residual of `SEAM.EQUIV.01` is now a composition of standard cited theorems (OS/quasi-free clustering + Kitaev free-fermion invertibility + the v297 AQFT stack) over established TFPT facts (16 quasi-free Majoranas; transfer gap $6 \ln(3/2)$): *no undischarged TFPT-internal assumption remains*. It stays [O] (not machine-proved end-to-end), but

the open front is reduced to citable machinery over a spectral gap that is provably positive.

The continuum chain assembled, and the residual tied to the seam one-sidedness ([[verification/v356_continuum_mmst_applicability.py](#)], [SEAM.EQUIV.CONTINUUM.03](#)). Writing the chain out link-by-link: **S1** the gapped (v302) quasi-free 16-Majorana collar $\rightarrow$ **S2** invertible (Kitaev free fermions, v301) $\rightarrow$ **S3** the *nontrivial* invertible phase ($c_- = 8 \neq 0$) $\rightarrow$ **S4** a non-gappable anomalous chiral edge (anomaly inflow / bulk-edge) $\rightarrow$ **S5** its chiral-CFT scaling limit (MMST, v336; range $\text{rank} \leq c \leq D$, i.e. $8 \leq 8 \leq 16$) $\rightarrow$ **S6** the order-4 μ_4 clock + the AQFT stack pin $(E_8)_1$ (v351/v297). Every link is a theorem or an established TFPT fact *except* S3, and S3 is precisely the seam being *one-sided* (chiral, $c_- \neq 0$) — the *same* one-sidedness that makes $c_3 = 1/(8\pi)$ a one-sided Gauss–Bonnet ($|\mathbb{Z}_2| \nmid K$, v58). So the lone realization input is tied to the constant that *defines* the theory, not a free dial; the residual is now “the abstract collar is realized as a genuine lattice chiral free-fermion invertible phase”, [\[O\]](#) but theorem-bounded on both sides.

MMST in-class verified for the collar, so the scaling limit is $(E_8)_1$ ([[verification/v366_mmst_seam_collar.py](#)], [SEAM.MMST.INCLASS.01](#)). The continuum leg is now checked *hypothesis-by-hypothesis* for the explicit seam collar: it is a $D = \dim S^+ = 16$ Majorana quasi-free (CAR) state (v155/v160), gapped with $\Delta = 6 \log \frac{3}{2} > 0$ *derived* (v302), and chiral with $c_- = D/2 = 8 = g_{\text{car}} + N_{\text{fam}}$, so it sits squarely in the MMST free-lattice-fermion class with $\text{rank} \leq c \leq D$ reading $8 \leq 8 \leq 16$ (*in* range, saturating $\text{rank} = c = 8$). The MMST massless-scaling-limit theorem then applies as a cited result, and the order-4 μ_4 clock pins the target to $(E_8)_1$ ($\det K = 1$, one primary) over the same- c rival $SO(16)_1$ ($\det K = 4$, v351). [\[E\]](#) the class arithmetic, the derived gap, the chirality and the $(E_8)_1$ pinning; [\[C\]](#) the CAR-class typing and the cited MMST theorem; [\[O\]](#) the lone residual is exactly S3 above (the lattice realization). The continuum side of [SEAM.EQUIV.01](#) is thereby reduced to a single, named, theorem-bounded input. The lattice $\rightarrow$ conformal-net step itself is now a *rigorous AQFT construction*: Adamo–Giorgetti–Tanimoto (arXiv:2506.01008, 2025) build the two-dimensional conformal net $\mathcal{A}_Q$ of an even lattice Q and *classify* every two-dimensional extension of the Heisenberg net (under a discreteness assumption) as such a lattice net — so “the lattice is a conformal net” is a theorem of the same operator-algebraic lineage the continuum leg invokes, not a hope; and since (xxxii) ([[verification/v469_seam_crossedproduct_route.py](#)]) the extension step no longer even *needs* the preprint lineage — the local $\mathbb{Z}_2$ simple-current crossed product ($h_s = 1 \in \mathbb{Z}$, Longo–Rehren 1995; Böckenhauer 1996; Böckenhauer–Evans 1998; KLM 2001) certifies it on peer-reviewed ground, with AGT/AMT as the independent second witness.

S3 made concrete: an explicit gapped chiral-Majorana lattice model ([[verification/v367_seam_s3_lattice.py](#)], [[verification/v368_seam_s3_inflow.py](#)]). The S3 input is no longer abstract. The $p+ip$ Bloch model $h(k) = \sin k_x \sigma_x + \sin k_y \sigma_y + (M - \cos k_x - \cos k_y) \sigma_z$ is, at $M=1$, a gapped free-fermion phase with numerically-computed Chern number $|C|=1$ ($C=0$ trivially at $M=3$), so $N_{\text{Maj}} = \dim S^+ = 16$ copies give a chiral edge of $c_- = 16/2 = 8 = g_{\text{car}} + N_{\text{fam}}$, with the edge K-matrix $SO(16)_1$ ($\det 4$) condensed to $(E_8)_1$ ($\det 1$) by the order-4 μ_4 clock (v351/v369). A finite STRIP of the same model carries an edge-localised in-gap chiral branch (absent in the trivial phase) — so by anomaly inflow the chiral edge *exists* as a consequence of the invertible bulk (the S4 leg), discharging the “does the edge exist?” question on the lattice. [\[E\]](#) the gapped Chern model, the $c_- = 8$ counting and the edge existence; [\[C\]](#) the μ_4 condensation to $(E_8)_1$; [\[O\]](#) only the genuine *continuum scaling limit* (MMST, v336) stays open — S3 is a runnable lattice model that pins the target at every computable level, so the keystone’s MMST route ([SEAM.EQUIV.MMST.01](#)) is *closed modulo cited theorems*.

The S3 closure stack: $c = 8$, the $(E_8)_1$ character, the torus count and reflection positivity — all computed on the lattice ([[verification/v376_seam_s3_centralcharge.py](#)], [[verification/v377_seam_s3_e8character.py](#)], [[verification/v378_seam_s3_modular.py](#)], [[verification/v379_seam_s3_rp.py](#)]). On that explicit model the target of the scaling limit is now pinned at every level a computation can reach. *Central charge* (v376): the Calabrese–Cardy finite-size entanglement scaling gives $c = 1.0000$ per critical complex mode, so the 16-Majorana (= 8 complex) collar has $c = 8 = g_{\text{car}} + N_{\text{fam}}$, and it is chiral ($c_- = 8$). *Convergence* ([[verification/v392_seam_s3_scalinglimit.py](#)]): across five sizes $L = 66 \dots 514$ the slope $c_1(L)$ is a monotone Cauchy-like sequence approaching 1 and $1/L$ -Richardson-extrapolates to $c_1(\infty) \approx 1.000$ ($c \approx 8.00$ in the limit) — the existence-relevant *convergence* the two-size value did not test (evidence, not a proof of existence; the residual stays [\[O\]](#)). *The net* (v377): the exact affine character $\chi = E_4/\eta^8 = q^{-1/3}(1 + 248q + 4124q^2 + \dots)$ gives $c = 8$ and 248 level-1 currents (one primary), and the μ_4 /GSO promotion $248 = 240 + 8 = 120 + 128$ ($240 = 16 \cdot 5 \cdot 3$) pins $(E_8)_1$ over the same- c rival $SO(16)_1$ (120 currents, four primaries). *Genus-1* (v378): the KLM condensation $\mu(\text{carrier}) = 16 \rightarrow \mu(E_8) = 16/4^2 = 1$ and the single character’s modular T -eigenvalue $e^{-2\pi i/3}$ give torus ground-state degeneracy 1 — the

holomorphy / $\det K=1$ statement that the genus-0 arguments could not reach (v344). *Reflection positivity* (v379): the gapped collar’s Euclidean two-point function has a positive spectral measure, so the OS reflection Gram matrix is a *positive-mixture Gram* $M = V^\top \text{diag}(1/K)V$, $V_{ki} = e^{-\varepsilon_k \tau_i}$ — hence $x^\top M x = \sum_k w_k (v_k \cdot x)^2$ is a sum of squares and positive semidefiniteness holds *exactly*, as an identity rather than to a tolerance (hardened 2026-07-28 after T133 certified that the *assembled double-precision* matrix has a negative direction at -7.75×10^{-18} , inside the old 10^{-10} tolerance; the min-eigenvalue check is retained as a diagnostic measurement, and a ghost mode still breaks the RP test). [E] all four (numerical c , exact character, genus-1 count, RP); [O] the one thing a computation cannot supply is the abstract *continuum existence* of the limit (the cited MMST theorem, v336). With RP in hand the ambient measure C7 is then discharged by the redundancy theorem (v369), so S3 is the master key: the target is pinned exactly, only the existence proof is external.

A ground-state witness for premise (i): the modular commutator measures $c_- = 8$ from the bulk state alone ([verification/v489_seam_modular_commutator.py], SEAM.CCC.01). The chiral central charge entered so far through band topology (the Chern number, v367) and through counting (the KLM/det-Cartan arithmetic, v378). The Kim–Shehab–Kim modular commutator $J(A, B, C) = i \text{Tr}(\rho_{ABC}[K_{AB}, K_{BC}]) = (\pi/3) c_-$ (PRL 128 (2022)) reads c_- off the *entanglement data of the exact BdG ground state* of the same collar — no band bundle, no representation theory enters — via the Peschel covariance ($W_X = -2 \text{artanh } \Gamma_X$) on a disk cut into three 120° wedges: per layer $c_- = 0.499998$ at $L=32$ (converged 0.500000 at $L=48$, deviation -1.8×10^{-7}), the trivial $M=3$ control gives 0.000000, the orientation flip $M=-1$ gives -0.499986 (sign flip), layer additivity is exact (10^{-14}), and the 16-layer carrier lands on $c_- = 8 = g_{\text{car}} + N_{\text{fam}}$ (direct run 7.9967 at $L=16$, additivity route 8.0000). [E] the third, mutually independent lattice witness of premise (i) of SEAM.EQUIV.01 — and the first read from ground-state entanglement alone; [O] unchanged: the continuum scaling limit (MMST, v336) and the cited KLM condensation step stay exactly where v367/v378 left them — SEAM.EQUIV.01 stays [O].

The spin-structure parity census: $\mu_{\text{pre}}=4$ **witnessed on the lattice** ([verification/v490_seam_parity_census.py], SEAM.MU.01). Step 1 of the KLM chain (v378) — “the carrier sits in the $SO(16)_1$ class, $\mu_{\text{pre}}=4$, *before* the μ_4 condensation” — was arithmetic; it is now a lattice ground-state measurement. The T^2 fermion parity per spin structure (AA, AP, PA, PP) is determined *twice* (Read–Green TRIM product and real-space Majorana Pfaffian, spectra matching to 9×10^{-15}): per layer $(+, +, +, -)$ — the $\nu=1$ Ising signature — and for the 16-layer carrier parity¹⁶ = + in *all four* sectors, so the gauged torus GSD is $4 = \mu(SO(16)_1) = |\det \text{Cartan}(D_8)|$. The 16-fold-way bridge then runs on *measured* numbers: $\theta_v = e^{2\pi i \nu / 16} = 1$ exactly at $\nu = 2c_- = 16$ (v489) — the gauged vortex is a boson, condensable, while the rivals $\nu = 1, 2, 8$ are not holomorphically condensable. Honest fences: the fermionic Kitaev–Preskill TEE is identically zero for *both* $M=1$ and $M=3$ ($\gamma_f = 0.000001/0.000000$) — an invertible fermionic phase has no TEE, so γ_f does *not* discriminate $(E_8)_1$ vs $SO(16)_1$; and the KLM step $\mu : 4 \rightarrow 4/2^2 = 1$ stays the cited theorem ([C], the condensed model is interacting). The numerics pin exactly the premise package $\{c_-=8, \mu_{\text{pre}}=4, \theta_v=1\}$ of the condensation; SEAM.EQUIV.01 stays [O].

And that external step is now Lean-formalised as “closed modulo a cited theorem” (FORM.SEAM.MMST.01, TftptCarrier/SeamScalingLimit.lean). The continuum leg is machine-formalised in the audit-contract style of SeamEquivChain.lean: the collar’s MMST applicability hypotheses are *provable* arithmetic facts by Lean kernel **decide** (no axioms) — $D=16$ Majoranas, rank= $c=8$, the range $8 \leq 8 \leq 16$, the chirality $c_-=8 \neq 0$, and the holomorphy discriminator $\det K=1$ vs $SO(16)$ ’s 4; the MMST scaling-limit theorem and the Adamo–Moriwaki–Tanimoto OS reconstruction are declared as *named cited axioms*; and the conclusion $(E_8)_1$ follows by a well-typed composition whose **#print axioms** pins the residual to exactly those two published theorems (plus the carrier gap), with no hidden assumption and no **sorry**. Every TFPT-internal hypothesis is thus kernel-checked; only the two cited theorems are assumed — the honest closure ceiling for SEAM.EQUIV.01. *Two of the seven convergent routes below are now kernel-hardened the same way*: FORM.SEAM.BW.HSMI.01 (TftptCarrier/SeamStandardPair.lean) proves the four Borchers/Wiesbrock *standard-pair relations* exactly over $\mathbb{Z}$ on the centred ladder ($[K, P]=P$, $J^2=1$, $JKJ=-K$, $JPJ=P_+$, all by kernel **decide**, no domain axiom), with the cited Borchers step and the one continuum input as named axioms; and FORM.SEAM.APPLICABILITY.01 (TftptCarrier/SeamApplicabilityLedger.lean) kernel-checks the audit *count* itself — of the 12 cited-theorem hypotheses exactly 11 are internal and 1 external (and $8 \leq 8 \leq 16$, $\det K=1 \neq 4$), all axiom-free — so the very bookkeeping the keystone’s honesty rests on is machine-verified, leaving only the one named continuum fact open.

TOE-roadmap follow-ups — the keystone as a typed certificate, the bedrock mod-

ular form, and the gap-decoupled measure ([verification/v308_seam_equiv_chain.py], [verification/v309_modular_bedrock.py], [verification/v311_gap_decoupled_measure.py]). Three machine-checked steps push the keystone arc further. (i) *Composition certificate* ([verification/v308_seam_equiv_chain.py]): the closing implication is one well-typed logical chain $\text{gap} > 0 \rightarrow \text{invertible} \rightarrow \text{holomorphic } c=8 \rightarrow (E_8)_1$ whose arithmetic discriminators are exact — $|\det \text{Cartan}(E_8)|=1$ against $|\det \text{Cartan}(D_8)|=4$, the carrier tower $4 \cdot 4=16 \rightarrow 4 \rightarrow 1$, $c=g_{\text{car}}+N_{\text{fam}}=8$ — so the only undischarged premise is the cited OS step; [E] the discriminators, [O] the keystone. (ii) *Modular bedrock* ([verification/v309_modular_bedrock.py]): with the four marks derived from Gauss–Bonnet (v216) and the order-4 clock from cross-ratio 2 (v214), the quasi-free modular Hamiltonian $K = \log((1-C)/C)$ is built explicitly and $[\rho, K]=0 \Leftrightarrow \omega \circ \rho = \omega$ (a $\mathbb{Z}_2$ profile breaks it), so the bedrock reduces *non-circularly* to the Bisognano–Wichmann intrinsicity of the raw collar state; [E] the model, [O] the intrinsicity. (iii) *Gap-decoupled measure* ([verification/v311_gap_decoupled_measure.py]): the admissible sector clusters exponentially at rate $(2/3)^6$ (correlation length $1/\Delta$, finite susceptibility $729/665$) and the v76 margin $\Delta - 31/(4\pi^2) > 0$ decouples it from the ambient, so the physical sector is a convergent object while QG.AMB.01 is a [C] redundancy (v369). Each pins the residual to a single named step; with the lattice/ S_3 stack the keystone’s MMST route (SEAM.EQUIV.MMST.01) is now *closed modulo cited theorems*. (iv) *The intrinsic-BW residual as a cited reduction* ([verification/v424_seam_lto_rp_reduction.py]): the open bedrock lemma of (ii) — the raw collar’s intrinsic Bisognano–Wichmann property — is reduced to the recent commuting-projector theorem of Naaijken–Penneys–Wallick (arXiv:2605.10693), whose reflection-positivity axiom (LTO-RP) is exactly $u_\Theta = J$ (the reflection *is* the modular conjugation), i.e. the BW condition $\theta K \theta = -K$ — *not* $\omega \circ \rho = \omega$. [E] the μ_4 clock is a modular *symmetry* ($[\rho, K]=0$), not the flow ($\sigma^\psi = \rho$ is type-wrong, a one-parameter group versus a finite order-4 element); a positive character-block-diagonal K has $[\rho, K]=0$ yet, being positive definite, never $\theta K \theta = -K$, so clock-invariance does *not* imply (LTO-RP). [O] make-or-break: NPW26 prove (LTO-RP) for *topologically ordered* commuting-projector models (Toric Code via a trace, Levin–Wen anyonic), whereas the seam is the opposite corner — invertible ($|\det \text{Cartan}(E_8)|=1$, no anyons) and a $\beta=1$ KMS state (not a trace) — so two sub-steps stay open: realising the raw seam as a $\mathbb{Z}_2$ -reflection commuting-projector LTO, and extending (LTO-RP) to the invertible KMS case. An MMST-style reduction (cf. v308/v336), *not* a closure. *Sub-step (ii) is now constructively exhibited* ([verification/v426_seam_lto_rp_kms.py]): a gapped reflection-symmetric μ_4 -even collar at $\beta=1$ KMS satisfies $\Theta K \Theta^{-1} = -K$ ($u_\Theta = J$) and $[\rho, K]=0$, so (LTO-RP) holds there by direct Tomita–Takesaki — the corner NPW26 leave open — with $|\det \text{Cartan}(E_8)|=1$ (invertible, no anyons) and the state a genuine KMS, not a trace; only the continuum sub-step (i) remains [O]. On the covariance side, the recent theorem of Adamo–Giorgetti–Tanimoto (arXiv:2508.07109, 2025) — every DHR representation of a conformal net extending a loop-group/Virasoro net is *automatically* positive-energy and diffeomorphism-covariant — defuses the old “Bisognano–Wichmann presupposes the very covariance it should produce” circularity (v215): once the seam net is in hand, geometric covariance is a *consequence*, not an input.

Twenty-six convergent attacks on the one continuum residual — intrinsic BW, a bulk continuum motor and its chiral-edge counterpart, the β -homotopy, the applicability audit, uniform rigidity with its forcing converse and the derivation of clock-invariance, three independent reads of the edge central charge, the four-point and uniform-in- N pinning of the external fact, the Ising operator tower, the $(E_8)_1$ torus modular data, the μ_4 -from-marks derivation, the kill tests, the lattice Sugawara current algebra, a uniform-in- N Tomita–Takesaki tower, the chirality-from- P_1 forcing, an $(E_8)_1$ character kill test, the exact MMST citation audit, the complementary lattice-VOA route, the explicit flat-band commuting-projector LTO realisation, the strict-locality (Kapustin–Fidkowski) obstruction, the character-level 128-spinor extension, the $c=8$ holomorphic-uniqueness pin and the one-particle realisation rigidity ([verification/v438_seam_hsmi_borchers.py], [verification/v439_seam_lieb_robinson.py], [verification/v440_seam_lto_rp_beta.py], [verification/v441_seam_applicability_ledger.py], [verification/v442_seam_rigidity_uniform.py], [verification/v443_seam_e8_discriminator.py], [verification/v444_seam_edge_scaling.py], [verification/v445_seam_rigidity_forcing.py], [verification/v446_seam_clock_invariance.py], [verification/v447_seam_edge_chern.py], [verification/v448_seam_edge_fourpoint.py], [verification/v449_seam_mmst_uniform.py], [verification/v450_seam_edge_entanglement.py], [verification/v451_seam_edge_cardy_tower.py], [verification/v452_seam_e8_modular.py], [verification/v453_seam_mu4_from_marks.py], [verification/v454_seam_edge_virasoro.py], [verification/v455_seam_bw_uniform.py], [verification/v456_seam_chirality_from_c3.py], [verification/v457_seam_character_killtest.py],

[verification/v458_seam_mmst_citation_audit.py], [verification/v459_seam_lattice_voa_route.py],
 [verification/v460_seam_s3_lto.py], [verification/v461_seam_strict_locality.py],
 [verification/v462_seam_spinor_continuum.py], [verification/v463_seam_c8_holomorphic_uniqueness.py],
 [verification/v464_seam_oneparticle_rigidity.py], [verification/v469_seam_crossedproduct_route.py]].

None closes SEAM.EQUIV.01; together they ground the one continuum input on both its halves, reduce it to a single analytic fact and make the claim falsifiable. (v) *The intrinsic Bisognano–Wichmann route* ([verification/v438_seam_hsmi_borchers.py]): the seam supplies a Longo *standard pair* / +half-sided modular inclusion — a positive lightlike translation $P \geq 0$, the boost K with $[K, P] = P$ (the Borchers scaling $\Delta^{it} P \Delta^{-it} = e^{-t} P$), and a reflection J with $JKJ = -K$, $JPJ = P_-$ — so by Borchers’ theorem the modular flow is geometric *intrinsically*, *without* presupposing the rotation-covariance of the vacuum; [E] the ladder algebra and the $sl(2, \mathbb{R})$ positivity (min eig > 0 , with the indefinite boost as control), [C] the synthesis, [O] assembling the finite algebra and the positive-energy rep into one continuum standard pair (the MMST/NPW26 wall). (vi) *The continuum-existence motor* ([verification/v439_seam_lieb_robinson.py]): the explicit $p+ip$ collar (v367) has a uniform-in- N gap, exponential clustering with N -independent ξ , and a finite Lieb–Robinson light cone, so by Nachtergaele–Sims its *bulk* thermodynamic limit exists — grounding the bulk half of the cited MMST existence in our model and narrowing the open step to the chiral massless *edge* scaling limit; [E] gap/clustering/light-cone, [C] the lift, [O] the edge CFT. (vii) *The β -homotopy* ([verification/v440_seam_lto_rp_beta.py]): the (LTO-RP)/BW data ($\Theta K \Theta = -K$, $[\rho, K] = 0$) are β -independent, and along the whole path $C_\beta = 1/(1+e^{\beta K})$ from the NPW26-proven *trace* state ($\beta=0$) to the seam KMS state ($\beta=1$) every covariance is reflection-related, μ_4 -invariant and valid — a norm-continuous, obstruction-free homotopy, upgrading the single point of v426 to a path argument for sub-step (ii). (viii) *The applicability audit* ([verification/v441_seam_applicability_ledger.py]): a hypothesis-by-hypothesis ledger of MMST/NPW26/Adamo finds 11 of 12 hypotheses are TFPT-internal computed facts ([E]) and exactly one is external — the continuum existence of the chiral scaling limit — covariance being an Adamo consequence, not an input; the keystone’s external dependence is pinned to that one analytic fact. (ix) *Uniform rigidity* ([verification/v442_seam_rigidity_uniform.py]): the band-limited μ_4 -character rigidity of v398 is uniform in N — exact 4-sector commutation to $N=256$, a uniform gap, and an N -independent clock-breaking lower bound $\|[\rho, V]\| = 2$ — so it is not a small- N artifact (the full- L^2 forcing stays [O]). (x) *The kill tests* ([verification/v443_seam_e8_discriminator.py]): mode D of the closure-mode classification (v347) as concrete *falsifiable* discriminators against the nearest $c=8$ rival $SO(16)_1$ — torus GSD 1 vs 4, level-1 currents 248 vs 120, lattice theta 240 vs 112 norm-2 roots — with $c=8$ alone unable to decide (v277) and only $\det K=1$ (the genus-1 statement, v90) selecting E_8 ; a seam with GSD >1 would falsify $(E_8)_1$. [X] the tests, [O] the realisation. (xi) *The chiral-edge scaling motor* ([verification/v444_seam_edge_scaling.py]): the counterpart of (vi) on the *edge*, attacking exactly the chiral massless scaling limit that the bulk motor left open. On the same $p+ip$ collar (v367/v368) the lattice edge already carries the conformal data of a chiral $c=\frac{1}{2}$ Majorana CFT — a linear, chiral in-gap dispersion (velocity $v \approx 0.98$, $R^2 = 0.9999$, with *opposite* per-edge velocities), an *algebraic* edge two-point $\sim r^{-\eta}$ with $\eta \approx 1$ (versus the *exponential* bulk row and the exponential trivial-phase control), and a Cauchy-convergent scaling-collapse under refinement ($\eta = 1.001 \pm 0.007$, successive spread $0.028 \rightarrow 0.013$) — so the 16-copy edge is a chiral $c_- = 8$ $(E_8)_1$ CFT; [E] the kinematics, correlator and collapse, [C] the central-charge reading, [O] the rigorous uniform-convergence theorem (cited MMST). With (vi) for the bulk and (xi) for the edge, *both halves* of the one continuum residual are numerically grounded and pinned to a single cited theorem. (xii) *The rigidity forcing converse* ([verification/v445_seam_rigidity_forcing.py]): the converse of (ix), upgrading rigidity from “block-diagonal *permits* commuting” to “commuting with the order-4 clock *forces* block-diagonal”. For $\rho = \text{diag}(i^n)$, $[\rho, K] = 0$ iff $K_{ij} = 0$ whenever $i \not\equiv j \pmod 4$ (verified entrywise, swept $N=4..128$), so the commutant is *exactly* the 4-sector block algebra: $\dim\{K : [\rho, K] = 0\} = \sum_s n_s^2$, a proper subspace of the full N^2 algebra ($N=16$: $64 < 256$); every off-block unit has $\|[\rho, E_{ij}]\| \geq \sqrt{2}$ (zero slack), and the order-2 clock leaves a strictly larger commutant (64 vs 128) — the *four* Gauss–Bonnet marks (order $4 = |\mu_4| = h(A_3)$) force strictly more structure than two, and only the index-4 extension is $(E_8)_1$ ($\det = 1$, $v281/v154/v351$). [E] the entrywise iff, exact commutant dimension, off-block sharpness and order discriminator; [C] the synthesis (RP-definability + clock-invariance *force* the block-diagonal Λ_Σ , v442’s “permit” upgraded to “force”); [O] the single residual is now precisely that the physical transfer lies in the commutant (clock-invariance = QGEO.SYM.01, v323/v335). (xiii) *Clock-invariance derived* ([verification/v446_seam_clock_invariance.py]): closing (xii)’s residual one notch further — the operator condition $[\rho, \Lambda_\Sigma] = 0$ is itself a *consequence* of a structural symmetry, not an extra assumption. On explicit finite OS data, the canonical transfer $\Lambda = C(1)C(0)^{-1} = e^{-h}$ (the transfer fixed by RP+reflection,

v54) of a μ_4 -invariant covariance satisfies $[\rho, \Lambda]=0$ (swept $N=4.32$; iff-sharp μ_4 -breaking neg control), Λ is a positive contraction, and the free modular Hamiltonian $K=\log((1-C)C^{-1})$ (v258) of the invariant state has $[\rho, K]=0$ (Tomita–Takesaki). [E] the inheritance, contraction and modular commutation; [C] so with (xii), RP-definability *plus* the four marks force the block-diagonal Λ_Σ ; [O] the residual reduces from an operator commutation to the structural “the seam RP data is μ_4 -symmetric” (QGEO.SYM.01, v216/v323). (xiv) *The independent topological edge reading* ([verification/v447_seam_edge_chern.py]): a second, fit-free route to the edge central charge of (xi). On the same $p+ip$ collar (v367/v368) the bulk Chern number is the integer $C=+1$ (Fukui–Hatsugai–Suzuki link variables; $C=0$ in the trivial phase, $C=-1$ at the opposite mass, matching (xi)’s opposite per-edge velocities), the open strip carries exactly $|C|=1$ chiral edge branch per boundary, and the Li–Haldane entanglement spectrum has in-gap modes at $\frac{1}{2}$ only in the topological phase; [E] the Chern integer, channel count and entanglement signature, [C] the bulk–edge reading $c_- = N_{\text{Maj}} \cdot \frac{1}{2} \cdot |C| = 16 \cdot \frac{1}{2} = 8 = g_{\text{car}} + N_{\text{fam}}$ — the same $c_- = 8$ as (xi) from a logically independent (topological) observable. (xv) *The four-point pinning of the external fact* ([verification/v448_seam_edge_fourpoint.py]): extending (xi)’s two-point scaling limit to the next order to support exactly the one external fact (viii) isolates. The edge state is quasi-free (the ground-state correlation matrix is an exact projector $G^2=G \sim 10^{-13}$, the MMST CAR hypothesis), and the edge four-point cross-ratio $Q=[G_{01}G_{23}]/[G_{02}G_{13}]$ at commensurate positions is Cauchy-convergent under refinement to the free-fermion conformal value $\sin(5\pi/12)/\sin(\pi/12)=2+\sqrt{3} \approx 3.732$ (the bulk control off-scale, exponential); [E] the quasi-free property, four-point convergence and conformal form, [C] so the one external MMST fact is empirically supported at *both* two- and four-point order. (xvi) *The uniform-in- N pinning* ([verification/v449_seam_mmst_uniform.py]): strengthening (xv) from convergence at a single cross-ratio to the *uniformity* the cited theorem asserts — two *independent* edge four-point cross-ratios converge to *distinct* conformal values ($Q(1, 2, 4, 7)/12 \rightarrow 2+\sqrt{3}$ and $Q(1, 3, 5, 9)/12 \rightarrow 2$) at one N -independent $1/N$ rate ($|Q(N)-Q_\infty| \cdot N \leq 4$ for both, $N_x \in \{36, 48, 60\}$), the bulk control off-scale; [E] the two limits and the uniform rate, [C] so the MMST uniform-convergence *hypothesis* is empirically met. (xvii) *The entanglement reading of c_-* ([verification/v450_seam_edge_entanglement.py]): a *third*, correlator-independent route to the edge central charge — the Calabrese–Cardy block-entropy law gives $c_{\text{Dirac}}=1.000$ for the critical free-fermion chain (vs a saturating, ℓ -independent gapped control), so c per Majorana $= \frac{1}{2}$ and $c_- = 16 \cdot \frac{1}{2} = 8 = g_{\text{car}} + N_{\text{fam}}$; [E] the law and the control, [C] the assembly. (xviii) *The Ising operator tower* ([verification/v451_seam_edge_cardy_tower.py]): naming the edge CFT by its *operator content* — the Cardy finite-size spectrum reproduces the exact Ising chiral primaries $h_\sigma = \frac{1}{16}$ (the periodic–antiperiodic ground split) and $h_\epsilon = \frac{1}{2}$ (the lowest NS pair gap), both L -independent, so $\{c, h_\sigma, h_\epsilon\} = \{\frac{1}{2}, \frac{1}{16}, \frac{1}{2}\}$ is the free-Majorana minimal model $M(3, 4)$; [E] the dimensions and conformal scaling, [C] the identification. (xix) *The $(E_8)_1$ torus modular data* ([verification/v452_seam_e8_modular.py]): pinning the $(E_8)_1$ identity on the *torus*, complementing the edge reads (xi)/(xiv)/(xvii)/(xviii) — the character $\chi = E_4/\eta^8$ is S -invariant ($\chi(-1/\tau) = \chi(\tau)$, one primary), T -multiplies by $e^{-2\pi i/3} = e^{-2\pi i c/24}$ ($c=8$) and is holomorphic ($c \equiv 0 \pmod{8}$, T -phase order 3); [E] the full modular signature. (xx) *The μ_4 -from-marks derivation* ([verification/v453_seam_mu4_from_marks.py]): folding the last structural premise into the existing one — the four Gauss–Bonnet marks are μ_4 (roots of z^4-1), the form basis ω_k is μ_4 -graded ($\rightarrow i^k \omega_k$, weights $(1, 2, 3)=A_3$ exponents) and the order-4 clock $z \rightarrow iz$ fixes their cross-ratio $\lambda=2$ (the D_4 stabiliser, v168), so the residual QGEO.SYM.01 of (xiii) follows from marks $= \mu_4$ plus the existing QGEO.REALIZE.01 — [E] the three mark facts, [C] no new premise: the chain (xx) $\rightarrow$ (xiii) $\rightarrow$ (xii) closes modulo realisation. The integer backbone of (xiv)–(xix) ($c_- = 8$, the Chern integers, the order-3 T -phase) is kernel-hardened in Lean axiom-free as FORM.SEAM.EDGE.CHERN.01. (xxi) *The lattice Sugawara current algebra* ([verification/v454_seam_edge_virasoro.py]): testing the cited MMST “Koo–Saleur generators $\rightarrow$ continuum Virasoro with the correct c ” fact on the edge *algebra*, not just its central charge. The critical free-fermion ring has the Affleck–Cardy Casimir energy $E_0(L) = e_\infty L - \pi v c / (6L)$ fitting $c_{\text{Dirac}} = 1.000$ ($\frac{1}{2}$ per Majorana, $L=64.1024$), the lattice $U(1)$ current closes its affine *level* exactly ($\langle 0 | J_n J_{-n} | 0 \rangle = n$, $n=1..8$; vanishing for a gapped control), and the level-1 Sugawara central charge $c = \dim G / (1+h^\vee) = 8$ for *both* $SO(16)_1$ (120/15) and $(E_8)_1$ (248/31); [E] Casimir, Kac–Moody level and Sugawara, [C] $16 \cdot \frac{1}{2} = 8$ is the level-1 current-algebra c_- . (xxii) *The uniform-in- N Tomita–Takesaki tower* ([verification/v455_seam_bw_uniform.py]): the inductive-limit companion of v426 for sub-step (i) of (iv) — a family of reflection-odd, μ_4 -even, gapped collars ($N=8..128$) on which the intrinsic Bisognano–Wichmann condition $u_\Theta = J$ ($\|\Theta K \Theta + K\| = 0$) holds *exactly at every N* , with an N -independent gap ($6 \log \frac{3}{2}$) and an N -independent neg-control margin ($\|\Theta K \Theta + K\| = 2\|K\|$ for a side-even reflection); [E] the uniform BW, gap, clock symmetry and teeth, [C][O] so the inductive limit inherits $u_\Theta = J$ (the abstract continuum theorem stays the cited

backbone). (xxiii) *The chirality-from- P_1 forcing* ([[verification/v456_seam_chirality_from_c3.py](#)]): the S3 input of the continuum chain (v356) is removed as a free choice — c_3 's “8” is the one-sided count $8=|\mathbb{Z}_2| \cdot (\int K/\pi)=2 \cdot 4$ ($8\pi=|\mathbb{Z}_2| \int K=2 \cdot 2\pi\chi(S^2)$, v73), and a reflection sends the edge Chern integer $C \rightarrow -C$ (verified $C=+1 \rightarrow -1$ on the explicit $p+ip$ model), so any *two-sided* (reflection-symmetric) gapped boundary forces $C=-C=0$ while the *one-sided* seam — the same $|\mathbb{Z}_2|$ quotient that gives c_3 its 8π — has no such reflection and forces $C \neq 0$; [E] the one-sided count and the $C \rightarrow -C$ flip, [C] chirality (S3) is a *consequence* of P_1 , magnitude $c_-=8$. (xxiv) *The $(E_8)_1$ character kill test* ([[verification/v457_seam_character_killtest.py](#)]): a *falsifiable* sharpening of the (x)/(x”) discriminators — the $(E_8)_1$ tower $\chi=E_4/\eta^8=q^{-1/3}(1+248q+4124q^2+34752q^3+\dots)$, the weight-1 count $248=120+128$ (vs the free 120) and the sector count $|\det \text{Cartan } E_8|=1$ (vs $|\det \text{Cartan } D_8|=4$); a measured 120, four sectors, or a wrong tower coefficient would *falsify* $(E_8)_1$, and the data give $248/1/\{1, 248, 4124\}$ (passed). [X] the kill criterion, [E] the exact discriminators. (xxv) *The exact MMST citation audit* ([[verification/v458_seam_mmst_citation_audit.py](#)]): a hypothesis-by-hypothesis reading of the cited MMST theorem (arXiv:2107.13834) against the collar — H1 CAR class, H2 massless limit, H3 the per-copy $c=\frac{1}{2}$ Thm 4.11, H4 decoupled additivity to $c=8$, and H5 the WZW range $8 \leq 8 \leq 16$ with its 120 bilinear currents all *match*, isolating the single residual H6 to the 128-spinor extension $SO(16)_1 \rightarrow (E_8)_1$ (the $\det K \rightarrow 1$ holomorphy step), *outside* MMST's bilinear method; [E] the per-copy theorem, [C] the additivity and range, [O] the one open spinor extension — since (xxxii) certified at net level by the peer-reviewed crossed product. (xxvi) *The complementary lattice-VOA route* ([[verification/v459_seam_lattice_voa_route.py](#)]): the second, independent construction that supplies exactly H6 — for the even unimodular E_8 the lattice net A_{E_8} (Adamo–Giorgetti–Tanimoto, arXiv:2506.01008; OS arXiv:2407.18222) has weight-1 space $248=8+240$ with the 240 roots splitting 112 integer + 128 *half-integer* (the spinor, $248=120+128$), so the lattice route builds the 128-spinor currents MMST (xxv) leaves open; [E] the even unimodularity and root split, [C] AGT constructs the holomorphic extension — since (xxxii) an *independent second witness*, the primary net-level certification being the peer-reviewed crossed product — [O] the routes leave one shared realisation input (the collar's scaling limit = A_{E_8}), tied to P_1 by (xxiii). The G-block arithmetic (the one-sided $8=|\mathbb{Z}_2| \int K/\pi$, the reflection $C \rightarrow -C$, $c_-=8$, the range $8 \leq 8 \leq 16$, $248=8+240=120+128$, $240=112+128$, $\det K \rightarrow 1$ vs 4) is kernel-hardened in Lean (FORM.SEAM.RESIDUAL.01, TfptCarrier/SeamResidualAxiom.lean), reducing the residual to ONE named TFPT-internal realisation axiom (CollarRealizedAsLatticePhase) plus the two cited existence theorems. (xxvii) *The explicit flat-band commuting-projector (LTO) realisation* ([[verification/v460_seam_s3_lto.py](#)]): NPW26 prove the intrinsic Bisognano–Wichmann lemma of (iv) for *commuting-projector* (local-topological-order) models, so sub-step (i) asks for the seam realised as such a $\mathbb{Z}_2$ -reflection LTO. On the same v367 $p+ip$ collar the flattened Bloch $Q=h/|h|$ is a frustration-free flat-band parent — $Q^2=I$ and the occupied $P=(I-Q)/2$ is an exact projector EQUAL to the ground-state projector — whose real-space kernel is gap-protected quasi-local ($\xi \approx 1.14$ at $M=1$, $|P(20,0)| < 10^{-6}$; a clean power law $|R|^{-2.1}$ at the gapless point, so locality needs the gap), and whose flattened modular Hamiltonian $K_{\text{flat}} = \text{sign}(K)$ inherits the $\mathbb{Z}_2$ reflection $\Theta K_{\text{flat}} \Theta = -K_{\text{flat}}$ and the μ_4 clock $[\rho, K_{\text{flat}}]=0$ EXACTLY; [E] the flat-band/projector and reflection algebra, [C] the quasi-locality, [O] the chiral $c_-=8 \neq 0$ obstructs *strictly* finite-range commuting-projector locality (Kapustin–Fidkowski, arXiv:1810.07756), so the realisation is exactly the quasi-local LTO net NPW26 use — advancing sub-step (i) of (iv) from abstract existence to an explicit gap-protected net, strict locality the cited residual. (xxviii) *The strict-locality obstruction made explicit* ([[verification/v461_seam_strict_locality.py](#)]): upgrading (xxvii)'s “strict locality is the cited residual” from a citation to an EXHIBITED topological obstruction. On the same v367 collar the FHS Chern is $C=+1$ (chiral) / 0 (trivial) / -1 (opposite mass), and the hybrid Wannier-centre (Wilson-loop) phase $\theta(k_y)$ winds by $|C|=1$ (chiral) and 0 (trivial) over the BZ — exponentially-localised Wannier functions exist *iff* that winding is 0 (Brouder–Panati et al.; Thouless), so the chiral collar has none; both phases are gapped, yet only the trivial one (winding 0) admits a strict finite-range commuting projector, so it is the *chirality* $c_-=8 \neq 0$, not the gap, that obstructs. [E] the Chern integer, the winding = $|C|$ identity and the gapped neg-control, [C][O] Kapustin–Fidkowski (arXiv:1810.07756): a strictly finite-range commuting projector would force winding 0 hence $C=0$, so for $C=1 \neq 0$ none exists — sub-step (i)'s realisation is *provably* the quasi-local NPW26 LTO net, strict locality topologically forbidden rather than a missing premise. (xxix) *The character-level 128-spinor extension* ([[verification/v462_seam_spinor_continuum.py](#)]): the constructive companion of (xxv)/(xxvi), exhibiting the 128-spinor extension three ways — the exact Jacobi/ E_8 identity $\theta_2^8 + \theta_3^8 + \theta_4^8 = 2E_4$ makes the holomorphic character the sum of the $SO(16)_1$ vacuum and spinor sectors, $\chi_{(E_8)_1} = E_4/\eta^8 = \chi_o^{SO16} + \chi_s^{SO16}$ (weight-1 counts $120+128=248$); the finite 16-Majorana Fock space already carries them ($C(16,2)=120$ currents and $2^{16/2}=256=128+128$

Ramond states); and a critical free-fermion ring converges to the chiral CFT (Casimir $c_- \rightarrow 8$, lowest scaled gap $x_1 \rightarrow 1$ with $|x_1 - 1| \sim 1/N^2$). [E] the θ identity, the 120+128=248 split and the finite- L convergence, [C][O] the rigorous holomorphic extension is certified at net level by the crossed product (xxxii), with AGT/AMT (xxvi) as the independent second witness. (xxx) *The $c=8$ holomorphic-uniqueness pin* ([verification/v463_seam_c8_holomorphic_uniqueness.py]): the positive lever that makes the *identification* half classification-forced rather than assumed. At level 1 the central charge $c = \dim \mathfrak{g}/(1+h^\vee) = 8$ has *three* simple solutions — A_8 (dim 80), $D_8 = \mathfrak{so}(16)$ (dim 120) and E_8 (dim 248) — so $c=8$ is necessary but *not* sufficient, and the $\det K=4$ $SO(16)$ rival of (xxiii) is a genuine same- c competitor, not a strawman. But a holomorphic (single-sector) $c=8$ theory has an $SL(2, \mathbb{Z})$ -covariant character, and the only weight-0 form with leading $q^{-1/3}$ and integer coefficients is $j^{1/3} = E_4/\eta^8$, whose q^1 coefficient (the one-dimensional candidate space, vacuum $a_0=1$) is 248 — so $\dim V_1=248$ is *forced*, selecting E_8 uniquely and excluding A_8/D_8 . [E] the three candidates, the forced 248, the tower $\{1, 248, 4124, 34752, 213126\} = E_4/\eta^8 = j^{1/3}$; [C] Dong–Mason/ Schellekens give the holomorphic $c=8$ VOA unique $= V_{E_8}$ (the even unimodular rank-8 lattice unique $= E_8$, Mordell–Witt), so “ $c=8$ holomorphic limit $= (E_8)_1$ ” is classification-forced. (xxxi) *The one-particle realisation rigidity* ([verification/v464_seam_oneparticle_rigidity.py]): attacking the *realisation* input R_1 itself. Since the seam is quasi-free (v113/v160/v161), by Araki’s self-dual CAR the whole net is the second quantisation of its one-particle symbol P , so R_1 collapses to a finite-per-mode statement about P : the gapped ground state is the *unique* quasi-free state with $P =$ the spectral projection onto $K < 0$ ($P^2 = P$ to $< 10^{-12}$, K fixed by gap + chirality + reflection v426); its real-space kernel is Cauchy on every fixed window ($\max |P_{2N} - P_N| \sim 1/N \rightarrow 0$, so P_∞ exists), and the limit carries $c=1$ per Dirac (entanglement $S(L) = (c/3) \ln L$, 16 copies $\Rightarrow c_- = 8$, tying v450). [E] idempotency, Cauchy convergence and the c -fit; [C] Araki/Shale–Stinespring give the state $\leftrightarrow$ symbol bijection and implementable Bogoliubov map, so R_1 is upgraded from a bald axiom to “the unique quasi-free realisation, its one-particle limit exhibited, modulo the named CAR functor”. (xxxii) *The net-level crossed-product route + the invariant-level realisation* ([verification/v469_seam_crossedproduct_route.py]): both halves of the residual re-founded on harder-to-reject ground. The 128-spinor extension needs no preprint: the $SO(16)_1$ spinor has $h_s = N/16 = 16/16 = 1 \in \mathbb{Z}$, so its statistics phase is $+1$ — *exactly* the Longo–Rehren locality criterion — and the $\mathbb{Z}_2$ simple-current crossed product $SO(16)_1 \rtimes \{1, s\}$ is a *local* index-2 extension by 1995–2001 *peer-reviewed* subfactor theory (Longo–Rehren, Rev. Math. Phys. 7 (1995); the level-1 $so(N)$ net with its DHR sectors from even-CAR: Böckenhauer, Rev. Math. Phys. 8 (1996); simple-current extensions of nets: Böckenhauer–Evans, Comm. Math. Phys. 197 (1998)); KLM (Comm. Math. Phys. 219 (2001)) gives $\mu(B) = 4/2^2 = 1 \Rightarrow$ holomorphic $\Rightarrow (E_8)_1$ by the (xxx) pin — so the AGT/AMT route (xxvi) is *demoted to an independent second witness*. The index-4 μ_4 glue runs on the same integer, $h(J^k) = \{1, 1, 1\}$ for $k=1, 2, 3$ (the v125 isotropy $q(k(1, 1)) = k^2$ at net level, $\mu = 16/4^2 = 1$). And the realisation input R_1 is reduced from model fiat to *invariants* (R_1'): quasi-free [C] (v155/v160) + gap [E] (v302) + class D + $c_- = 8$ [E] ((xxiii), from P_1) — computed on the collar as FHS Chern $|C|=1$ (control 0), $\nu = 16 \times |C| = 16$, $c_- = \nu/2 = 8$; and $\nu \equiv 0 \pmod{16}$ is exactly Kitaev’s class-D phase whose edge admits a purely bosonic description = the $(E_8)_1$ level-1 state (the sixteen-fold way; c_- a *phase* invariant, Kapustin–Spodyneiko; Kim et al.), so any realisation with the R_1' invariants is phase-equivalent to the v367 stack. [E] the locality integers, the KLM μ -arithmetic, the computed invariants and the composed chain; [C] the re-founded citation package; [O] the cited theorems stay cited. The full G-block arithmetic — now including the (xxviii) winding obstruction, the (xxix) finite-Fock spinor counts, the (xxx) holomorphy selector and the (xxxii) locality/ μ /sixteen-fold joints — is kernel-hardened in Lean (FORM.SEAM.RESIDUAL.01), the two formerly separate cited theorems are *merged* into one combined external axiom `seam_realisation_theorem`, so the residual reduces to ONE realisation axiom plus ONE cited theorem (`#print axioms` drops from six to four); and the (xxxii) route is Lean-carried as the *parallel* derivation `seamResidualClosed'` (the invariant-level axiom `collar_invariants` plus the single combined package `crossedproduct_route_theorem`, its joints — `lr_locality_integer`, `glue_locality_integers`, `klm_holomorphic`, `sixteenfold_pin` — kernel facts with no axioms). Across (v)–(xxxii), SEAM.EQUIV.01 stays [O]: with (xxx) pinning the identification, (xxxi) reducing the realisation to its one-particle data and (xxxii) re-founding the extension leg on peer-reviewed ground, the residual is now *entirely certification* — a named, hypothesis-audited package of published theorems (MMST, Longo–Rehren/Böckenhauer/Böckenhauer–Evans/KLM, Dong–Mason, Araki, Shale–Stinespring; AGT/AMT as second witness) with no open internal mechanism — yet still rests on cited continuum-existence theorems we do not re-prove.

The TOE/QFT closure contracts as three citable certificates
 ([verification/v398_seam_state_rigidity.py], [verification/v399_gravity_complete.py],

[[verification/v401_metrology_closure.py](#)]). Three companion contracts package the closure status as named targets without fabricating any closure. (i) *Seam State Rigidity* (SEAM.RIGIDITY.01): the scattered state-invariance reductions (v177/v199/v308/v309/v335) as ONE theorem target — [E] the RP-definability hinge (OS canonical transfer from RP + reflection alone, v54, quasi-free v155), [E] the band-limited μ_4 -character-block-diagonal core swept $N=4.64$ (beyond the 3-dim H^1 , made uniform in N to $N=256$ by v442), [E] holomorphy $\Rightarrow (E_8)_1$; and [E] the forcing converse (v445) that commuting with the order-4 clock is *exactly* μ_4 -block-diagonality, so the residual is no longer “does block-diagonal permit” but only $[\rho, \Lambda_\Sigma]=0$ on the full L^2 — the physical transfer lying in that commutant (intrinsic Bisognano–Wichmann). [E] Moreover (v446) that operator commutation is itself *derived*: the OS canonical transfer of μ_4 -symmetric RP data inherits the clock (with the modular Hamiltonian commuting, v258), so the residual reduces further to the *structural* statement that the seam RP data is μ_4 -symmetric (the four marks, QGEO.SYM.01), leaving [O] only that structural symmetry on the raw continuum collar — for which the four marks now have an *established* continuum frame: the four-interval multilocal modular geometry (v480, keybox below), where the invariance is manifest at the state level in the exactly solvable class. (ii) *Gravity-complete = Boundary-complete + Ambient-redundancy* (GRAVITY.COMPLETE.01): the local equation parameter-free (v358/v359) plus the redundancy discriminators (v369), with $\mathcal{O}_{\text{phys}} \subset \mathcal{A}_\Sigma$ checked over five gravitational readout classes. (iii) *Metrology closure* (METROLOGY.CLOSURE.01): the final input set $\{a, \pi, v_{\text{geo}}\}$ — 0 dimensionless dials + 1 unit, the dimensional bookkeeping that every dimensionful observable is a number $\times v_{\text{geo}}^d$ (No-Unit, v153), and v_{geo} theorem-forbidden ($\text{TOE}_{\text{strict}} = \text{TOE}_{\text{dimensionless}} + [v_{\text{geo}}]$, v384).

One surface for seam, carrier and lattice: the Kummer/K3 ([[verification/v260_k3_kummer_unification.py](#)]). The pillowcase is the elliptic-involution quotient $T_{\tau=i}^2/(z \mapsto -z)$, its four marks the four 2-torsion fixed points; squaring the construction to complex dimension two gives the *Kummer surface* $\text{Km}(T^2 \times T^2)$, a K3, whose three facets reproduce three TFPT objects at once. [E] the 16 nodes (the 2-torsion $|A[2]|=2^4=16$ of the abelian surface) resolve to 16 exceptional $\mathbb{P}^1$'s = $\dim S^+ = \Lambda^{\text{even}}(\mathbb{C}^5) = 16$ (v197), the carrier-16 as the *square* of the seam-4; [E] the cup-product form on $H^2(K3, \mathbb{Z})$ is the even unimodular lattice $U^3 \oplus E_8(-1)^2$ of signature (3, 19), so the same E_8 the carrier glues to (v1) is the surface's intersection form. (Honestly: the 16 node classes span the rank-16 Kummer lattice, a sublattice *distinct* from the $E_8(-1)^2$ summand — both exact, not identified.) Thus the v1 “glue $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ ” is the lattice shadow of a single smooth object that already carries the seam and the carrier; E_8 appears both *locally* as the du Val $\mathbb{C}^2/2I$ singularity (the Poincaré-sphere link, v232/v236; by Kronheimer that local model *is* the hyper-Kähler quotient of the marks quiver, v479) and *globally* as this H^2 summand. A unification (ARCH.K3.01), *not* a closure: the lattice arithmetic is [E], the carrier $\leftrightarrow$ node identification [C].

Synthesis — the Modular Spectral Closure ([[verification/v261_modular_spectral_closure.py](#)]). The ten steps above are not ten independent results: they are facets of one relative spectral datum

$$\text{TFPT}_{\text{QFT}} = (\mathcal{A}_\Sigma, \omega_\Sigma, \Delta_\Sigma, \rho, A_F, H_F, D_F, J, \gamma, S_{\text{rel}})$$

(seam algebra, KMS state, modular generator, μ_4 clock; the carrier finite triple; the relative spectral action), governed by a single *Modular Spectral Universality* statement: *the unique reflection-positive, holomorphic, index-4 KMS seam state ω_Σ minimises the modular clock defect, induces via the Kasparov/covariance map (v258) the 96-dim KO-6 finite triple D_F (v252), whose inner fluctuations (v254) generate the gauged Pati–Salam spectral action with the seam KMS cutoff (v259) — so Dirac, cutoff, gauging, glue and orientability are not separate choices but readouts of one seam state. The certificate v261 pins the cross-consistency: one number $4 = [B:A] = |\mu_4| = 2\chi = |(\mathbb{Z}/2)^2|$ (index = marks = fixed points = glue order), one carrier-16 ($\dim S^+ = \Lambda^{\text{even}}(\mathbb{C}^5) = \text{Kummer nodes}$, $H_F = 96$), and one gap $\Delta = 6 \log(3/2)$ (OS mass gap = recovery rate = Lindblad gap, static and dynamic). The honest account of completeness is therefore:*

- **[E]/[C] QFT-complete (modulo one premise).** The admissible OS sector, the boundary $(E_8)_1$ net, the relative spectral action, D_F (as covariance induction), the cutoff (as the KMS weight) and the gauging (as inner fluctuations) are one relative object — closed once QGEO.SYM.01 (the seam realisation: *the raw quasi-free seam state is μ_4 -invariant*) is granted. That single [O] premise is the only structural input the QFT layer still consumes.
- **[C] Ambient QG — a redundancy, separate by design.** The full covariant metric-sector / ambient quantum-gravity measure (QG.AMB.01) is gap-decoupled ($\Delta_{\text{eff}} = 1.648 > 0$) and is now a [C] *redundancy* (v369), *not* a blocker for the QFT layer: a certification object, not a missing piece of the boundary QFT.

So “solving the QFT” here means exactly this: the boundary QFT is a single relative object reduced to one foundational symmetry postulate, with ambient QG held apart — not a diffuse list of open interfaces. The MSC closes the boundary relative QFT object modulo QGEO.SYM.01; it does *not* assert SM-only 4D gauge unification, and the canonical 4D reading is boundary only.

4D-QFT fork policy — a frozen decision tree, not an open ambiguity ([verification/v265_qft4d_fork_freeze.py])

The 4D reading is a *branch policy*, machine-frozen in `qft4d_fork_freeze.json`:

- **A (canonical) — boundary only.** [C] *Boundary-only is the canonical 4D reading*: TFPT delivers the boundary QFT, the relative spectral action and the SM readout; a 4D-GUT is *not claimed by default* (E_8 is the audit hull, never an unbroken 4D gauge group).
- **SM-only 4D-GUT unification is killed.** [X] the plain SM couplings do not meet (at the $\alpha_1=\alpha_2$ scale α_3^{-1} is off by ~ 5.6 ; spread ~ 8.8 at 2×10^{16} GeV, v246). This is the canonical *expectation*, not a failure.
- **B (conditional) — carrier-native Pati–Salam UV branch.** [C]/[O] permitted *only* with the E_8 -allowed content $\{1, 10, 16, 45\}$ (no **126**, v247), $\sin^2 \theta_W = 3/8$ (v248), $\kappa = M_{PS}/M_s$ in the frozen $O(1)$ band (v249/v253), and a threshold model with a proton-decay kill test; carrier gauging itself stays the contract CONTRACT.QFT4D.DIRAC.01.
- **Proton-decay rule, now applied** ([verification/v266_ps_threshold_proton.py]). [X] the explicit two-step thresholds give $d=6$ $\tau_p(p \rightarrow e^+ \pi^0)$: the *minimal* 16-Higgs content ($M_{GUT} \sim 2.4 \times 10^{15}$) gives $\tau_p \sim 4.4 \times 10^{33}$ yr $<$ Super-K (2.4×10^{34}) and is *killed*; the E_8 -allowed (15, 1, 1) [the single 45] raises $M_{GUT} \sim 5.9 \times 10^{15} \Rightarrow \tau_p \sim 1.6 \times 10^{35}$ yr, proton-safe *now* but sitting at the Hyper-K reach ($\sim 1.4 \times 10^{35}$ yr) — so branch B is *content-selected* (must be $+(15, 1, 1)$) and *Hyper-K-decisive within the decade*, not an open-ended hope (only one 45 in the hull, so it stays structurally marginal; τ_p carries an $O(3)$ hadronic band).

The fork is thus a frozen decision tree with explicit promotion/demotion rules, *not* an open ambiguity.

And the bedrock is now a falsifiable kill-test, not a proof-promise ([verification/v215_seam_deck_killtest.py]). A foundational symmetry cannot be derived from nothing (Bisognano–Wichmann would presuppose the very covariance it should produce); the honest move is to recast QGEO.SYM.01 as four independently-falsifiable predictions about the raw seam DtN. The non-circular reduction (v198/v199/v201) gives $\omega \circ \rho = \omega \Leftrightarrow [\rho, C] = 0$ for the quasi-free covariance, with the principal symbol $|k|$ commuting exactly, so the whole residual sits in the bounded sub-principal M_f (equivalently: the raw Gaussian bulk form A is μ_4 -deck-invariant). The four levers are **K1** exactly four marks ($b_1 = \# \text{marks} - 1 = 3 = N_{\text{fam}} \Rightarrow \# \text{marks} = 4 = |\mu_4|$); **K2** the *square* configuration (cross-ratio $2 \Rightarrow j = 1728$, $\text{Aut} = \mathbb{Z}/4$; the clock order $h(A_3) = 4$ selects $\mathbb{Z}/4$, not the $j=0$ $\mathbb{Z}/6$, and a generic mark set has only $\mathbb{Z}/2$); **K3** the mod-4 sub-principal support ($[\rho, M_f] = 0 \Leftrightarrow f$ is $\mathbb{Z}_4$ -invariant, automatic for a μ_4 -mark sum); and **K4** the transfer gap $(2/3)^6 = 64/729$. These are frozen in `freeze_file.csv` (`seam_deck_symmetry`) and carrier-side green, so any future drift fails the suite. The *decisive* kill is deferred to QGEO.REALIZE.01: a first-principles raw-seam-collar DtN must *produce* four square marks without inputting them — an *emergence* test, the non-circular alternative to assuming the symmetry. The *light* version of that deferred test is now *executed* ([verification/v503_qgeo_emergence_light.py], QGEO.EMERGE.LIGHT.01), with the result *rigidity, not dynamical selection*: the free field does *not* select the square — the global det’ winner at fixed area is the *hexagonal* torus ($\log F(\rho) = -1.03352 > \log F(i) = -1.05469$, Osgood–Phillips–Sarnak), $\tau = i$ is a saddle of the full moduli space, the only critical points are the CM points ($E_2^*(i) = E_2^*(\rho) = 0$), and criticality at the square is Palais symmetric criticality — while the square *is* pinned sharply by the clock: the raw free-field DtN is mod-4 block-diagonal *iff* $\tau = i$ (an isolated zero of the K3 indicator as a modulus function, 8×10^{-15} at $\tau = i$, strictly monotone away), and $\text{Aut}(E_\rho) = \mathbb{Z}_6$ carries no order-4 element. The clock input itself is now *reduced to one bit* ([verification/v506_seam_clock_rigidity.py], SEAM.CLOCK.RIGIDITY.01): the deck has exactly two Möbius square roots, both automatically of order 4, and a marks-preserving root exists *iff* the deck is the central involution of the mark-stabiliser D_4 — given that alignment bit, order, uniqueness, the free 4-cycle and $\tau = i$ are all theorems (harmonicity alone is insufficient on bare mark data, exact counterexample — on free-deck circles it becomes sufficient, v510); on the Fock side the NS deck implementation is *forced* to order 4 ($U^2 = (-1)^F$ exactly, nonsplit $\mathbb{Z}_4$; the R sector splits), so the fermionic $\mathbb{Z}_4$ explains the *order* and only the spatial alignment remains the

carrier input. That bit has now survived its *tautology attack* ([[verification/v507_seam_bit_origin.py](#)], SEAM.BIT.ORIGIN.01): the marks are the four half-periods of the seam double and *every* mark-free collar deck is a half-period translation — the deck’s *class* is derived from the common torus origin — yet the origin does not force the alignment (the marks-preserving root exists iff the deck pairs separate harmonically, $\lambda \in \{-1, 2, \frac{1}{2}\}$, and the silver counterexample sits in-family as “ $\tau = i$ with a non-CM-fixed half-period”); the bit is an equivalence tetrad (clock $\iff \tau=i$ with the CM-fixed half-period $\iff$ deck central in the mark- D_4 $\iff$ harmonic deck pairs $\iff$ nonsplit NS Fock lift) whose physical face is the Fidkowski–Kitaev-type extension class (central $U^2 = (-1)^F$ vs edge $U_e^2 = +1$, split) — distinguished, but not derived: only the *position* (which half-period) remains the carrier input. That position half is now *topology* ([[verification/v510_seam_bit_freedom.py](#)], SEAM.BIT.FREEDOM.01): the collar deck is a *covering* deck, hence free on the seam circle; the unique free involution is the antipode ($\text{Aut}(C_{16}) = D_{16}$, complete census), nonsplit $\iff$ free over all 17 involutions in exact $\text{Cl}(16)$, and the edge/silver class — whose mark circle passes through the deck poles, the restriction of the *branched* pillowcase involution — is excluded; harmonic + free solves exactly to $b = \pm i$, so the bit reduces to the square-modulus datum $\tau = i$ alone. That datum now carries an equivalent *discrete* form ([[verification/v512_seam_tau_flag.py](#)], SEAM.TAU.FLAG.01): on the free seam circle bare mark-transitivity is automatic for every deck-invariant configuration $M(\delta) = \{\pm 1, \pm e^{i\delta}\}$ (the pair-exchanging V_4 ; no local jet sees the side bit), and what selects $\delta = \pi/2 \iff \tau = i$ is exactly *flag* transitivity ($V_4 \rightarrow D_4$ symmetry lift), welded into a 13-fold exact equivalence web (since 15-fold by v528 facets #14/#15) — the remaining input is *one discrete symmetry-lift bit*, its negation concretely measurable (odd-doublet split $(2/\pi) \ln \cot(\delta/2) \neq 0$). The candidate [E]-closure route through free OS positivity is since *decided negative* ([[verification/v521_seam_bit_rp_blind.py](#)], SEAM.BIT.RPBLIND.01): free reflection positivity and anti-linear Θ existence (the v519 battery) are *side-blind* along the whole $M(\delta)$ family — bond-cut Gram spectra identical to 40 digits for every tested δ , the $N = 16$ odd- m failure a lattice-placement artifact resolved at $N = 32$, the continuum kernel exactly $1/\sin((s+t)/2)$ with the axis position dropping out — so the free RP/ Θ battery is the *eighth* side-blind test on the v512 scoreboard ($7 \rightarrow 8$); the only δ -sensitive clauses presuppose the clock (the group generated by the admissible OS reflections closes to D_4 exactly at $\delta = \pi/2$: face #13 of the web, a typed [C] reformulation), and the honest residual [O] gap is the mark-decorated *state* class (defect/twist insertions, the interacting $\mathcal{A}_{\text{hol}}$) — the mark-decorated half of that gap is since decided side-blind too ([[verification/v525_seam_bit_twist_blind.py](#)], SEAM.BIT.TWISTBLIND.01: every well-defined mark-decorated state functional in the free-plus-twist class keeps RP positive along the whole $M(\delta)$ ladder — the *ninth* side-blind test, $8 \rightarrow 9$; the free-plus-twist class is exhausted) and the interacting half is since decided too ([[verification/v529_seam_interacting_toy_fk.py](#)], SEAM.INT.FKTOY.01: the *tenth* side-blind test, $9 \rightarrow 10$, on genuinely non-Wick dynamics — Θ_{15} mirrors $H(m) \rightarrow H(8-m)$ exactly despite δ -seeing spectra) — while the bit itself is now *physically defined* as the twist-class choice ([[verification/v528_seam_bit_twist_class_definition.py](#)], SEAM.BIT.TWISTCLASS.01: facets #14/#15 both directions, order parameter O , unification with flag-transitivity, web $13 \rightarrow 15$; [C] interferometric measurement sketch; the bit remains formal input) — and since carries the first *positive* selection datum: the straddle filter, run as a selector on the interacting mark family, keeps exactly *one* member alive — $\delta = \pi/2$ with positive coupling ([[verification/v534_seam_straddle_cone.py](#)], SEAM.STRADDLE.CONE.01; toy-level evidence for a dynamical origin, not a derivation). The full QGEO.REALIZE.01 stays [O]; no marker moves. [E] carrier-side (the four levers), [O] seam-side (the emergence): a kill-test, *not* a closure (QGEO.KILL.01).

And the mark count is now *derived*, not assumed: four marks from Gauss–Bonnet ([[verification/v216_marks_gauss_bonnet.py](#)], [[verification/v217_marks_emergence_scan.py](#)]). The K1 lever (“exactly four marks”) is the one that can be *executed* today. If the marks are $\mathbb{Z}_2$ branch points (cone angle π , deficit π) on the flat seam sphere ($\chi=2$ — the same χ that supplies the 8 in c_3), Gauss–Bonnet *forces* their number: $n\pi = 2\pi\chi = 4\pi \Rightarrow n = 2\chi = 4 = |\mu_4| = N_{\text{fam}} + 1$. The closed Euclidean sphere 2-orbifolds are exactly $(2, 3, 6)$, $(2, 4, 4)$, $(3, 3, 3)$, $(2, 2, 2, 2)$, and *three independent* criteria — all cone orders = 2 (the $|\mathbb{Z}_2|$ sheet branch), rank $H^1 = \#\text{marks} - 1 = 3 = N_{\text{fam}}$ (the three generations pick the 4-mark square over the 3-mark hexagonal $(3, 3, 3)$), and a free order-4 deck — converge on the pillowcase $(2, 2, 2, 2)$. The numerical sibling v217 confirms it on the actual DtN/state: scanning the number n and the positions freely, the 4-equally-spaced (square) configuration is the unique joint minimiser of {clock-invariance, 3-family cohomology, transfer gap $(2/3)^6$ }. So the mark *count* and *equality* move from input to derived; only the square *modulus* (cross-ratio 2, the order-4 clock) stays the one carrier input, and the from-zero raw-seam emergence QGEO.REALIZE.01 stays [O]. The light emergence test ([[verification/v503_qgeo_emergence_light.py](#)]) sharpens this residual to a *convergence*:

given the clock, the modulus freezes automatically ($P(iZ) = P(Z)$ forces $a_3 = a_2 = a_1 = 0$ with $j = 1728$ identically, [v493](#)), and the order-4 element on H^1 is the Coxeter element of $W(A_3)$ — exactly the P2 cusp-class carrier datum of [v499](#): the QGEO modulus rest and the P2 typing rest R1 are *one* common residual (“why does the raw seam carry the order-4 clock”), not two. And the clock-rigidity theorems ([v506](#)) reduce that common residual to *one alignment bit* — “the collar deck is the central involution of the mark- D_4 ” — with the clock’s *order* fermionically forced ($U^2 = (-1)^F$ exact); the bit-origin theorems ([v507](#)) then derive the deck’s *class* (every mark-free collar deck is a half-period translation of the seam double) and characterise the bit as an equivalence tetrad with a physical extension-class face, so only its *position* — which half-period — stays the input; the freedom theorems ([v510](#)) finally type that position half as *topology* (the collar deck is a covering deck, hence free; the edge class is excluded), so the residual is the square-modulus datum $\tau = i$ alone — restated by the flag-transitivity web ([v512](#)) as one discrete symmetry-lift bit ($V_4 \rightarrow D_4$: flag transitivity $\iff \tau = i$), and since [v528](#) equivalently as the *twist-class choice*, in principle interferometrically readable (SEAM.BIT.TWISTCLASS.01; the bit remains formal input). [\[E\]](#) (count + equality), [\[O\]](#) (the square modulus and the raw-seam realisation) — QGEO.MARKS.03 / QGEO.EMERGE.01.

And the four marks are a *four-interval* modular geometry ([\[verification/v480_multilocal_four_interval.py\]](#)). The seam circle minus its four marks is the $n=4$ disjoint-interval region of a chiral free fermion — the one continuum class whose *multi-interval* modular theory is exactly known (Casini–Huerta mixing term; Longo–Xu multi-interval subfactors; Rehren–Tedesco multilocal fermions; and Kawahigashi–Longo–Müger: *holomorphic* $\iff$ trivial multi-interval inclusion, $\mu=1$ — the same one bit as $\det K=1$). Until now the suite used only the single-ball CHM boost ([v358](#)); [v480](#) exhibits the multilocal mechanism on the exactly solvable antiperiodic critical ring: [\[E\]](#) the quarter-turn clock is a signed permutation with $\rho^4=-1$ *exactly* (the fermionic μ_4 clock is the order-8 double-cover lift — the same $\mathbb{Z}_2$ -lift pattern as $2I$ over I); [\[E\]](#) its primitive-8th-root character projectors decouple the 4-interval covariance at 10^{-15} (the diagonal 4-fold-cover isomorphism), each sector a *single-interval* quarter-ring kernel with boundary twist (10^{-14} spectra); [\[E\]](#) $\rho C \rho^{-1} = C$ at 10^{-14} , so $\omega \rho = \omega$ holds *manifestly* at the 4-interval state level — and a one-site displacement of a single interval re-couples the sectors at 10^{-2} (the invariance is the μ_4 -symmetric configuration, not an artifact); and the cross-interval modular coupling peaks on the conjugate-point diagonal, the lattice shadow of the Casini–Huerta mixing term (numerical). [\[C\]](#) the continuum multi-interval theory is cited, not re-derived; [\[O\]](#) the raw-seam premise (QGEO.SYM.01) is untouched — this puts the bedrock and the AP2 continuum leg ([v476/v478](#)) in one established, exactly solvable frame; it does not close them.

QGEO.SYM.01 is *not* a theorem — it is the final seam-identification premise

The single load-bearing bolt of the whole construction, on one page.

- What is identified.** That the conformal structure of the raw RP seam double *is* the μ_4 -deck structure of the carrier clock — i.e. the carrier (P2) and the seam’s conformal geometry are the same object. Equivalently, in Riemann–Roch *index* language: the seam’s degree-4 mark divisor on the genus-0 double has $h^0 = \deg + 1 = 5 = g_{\text{car}}$ and $\text{rank } H_1 = \deg - 1 = 3 = N_{\text{fam}}$ (with $\Lambda^{\text{even}}(\mathbb{C}^5) = 1+10+5 = 16 = \dim S^+$), so P2 is the genus-0 mode count of the four-mark seam divisor ([\[verification/v228_rr_index_gate.py\]](#)) — the same definitional identification, in index form, conditional on the seam producing that degree-4 divisor.
- Why it is weaker than QGEO.CONF.01/ISO.01.** It does *not* name $\mathbb{P}^1$ or μ_4 (uniformisation *produces* them) and asks only for a conformal *symmetry*, not a metric isometry or a pre-named conformal class — strictly the mildest of the three.
- What follows automatically (the conditional theorem).** *Given* QGEO.SYM.01, equivariant uniformisation puts the order-4 clock in the form $z \mapsto iz$ on $\mathbb{P}^1$, the four parabolic marks are its free orbit μ_4 , H^1 carries the $(1, 2, 3)$ grading, the seam Calderón contraction is the rank-8 K_Σ polarisation, and the index-4 simple-current extension is $(E_8)_1$ — the *entire* AQFT closure of [v175–v181](#).
- What would falsify or unseat it.** A seam double of genus > 0 (uniformisation to $\mathbb{P}^1$ fails); a carrier clock that is *not* a conformal symmetry of the seam geometry (a non-deck transport); or a fourth chiral generation ($N_{\text{fam}} \neq 3$ collapses $b_1 = N_{\text{fam}}$ and the four-mark μ_4 structure). The first two are constructive-geometry questions about the raw seam; the third is the standard family-count kill test.

Honest framing. The correct external statement is *conditional*: “Given QGEO.SYM.01, the RP seam




Figure 1: The emergent-QFT skeleton on the seam (v238–v246): time as the modular flow, the arrow as the recovery generator, particles as DHR sectors, the S-matrix as the braiding, and the 4d matter content as the carrier’s $SO(10)$ **16**. Each step is machine-checked; the one residual — SEAM.EQUIV.01, whose MMST route SEAM.EQUIV.MMST.01 is closed modulo cited theorems (v367–v379) with the cited continuum scaling-limit existence (v336) as its [O] leg (parent [O]) — is in red, the frozen 4D fork and the QG certification layer in gold. The whole skeleton is *conditional* on that keystone and is neither solved nor experimentally confirmed.

boundary realises the μ_4 deck structure, and the AQFT closure to $(E_8)_1$ follows through the verified chain v175–v181” — *not* “the theory *derives* the seam geometry”. The premise stays [O]; everything downstream of it is [E]. Per v323 (Bisognano–Wichmann) QGEO.SYM.01 is itself the *downstream* conformal-deck face of the single named keystone SEAM.EQUIV.01 (the raw RP seam = the holomorphic $(E_8)_1$ net at $\tau=i$), which is now *closed modulo cited theorems on its MMST route* SEAM.EQUIV.MMST.01 (the lattice/ S_3 stack pins it at every computable level, Lean FORM.SEAM.MMST.01; only the cited continuum existence stays [O]; the twistor route SEAM.EQUIV.TWISTOR.01 and the unconditional parent stay [O]).

This is not a gap to apologise for — it is TFPT’s one fundamental postulate. Every fundamental theory rests on an irreducible premise (the constancy of c in relativity, the equivalence principle, the canonical commutator). QGEO.SYM.01 is that premise for TFPT: a single geometric identification (seam conformal structure = carrier μ_4 deck) from which the entire algebraic and topological structure unfolds *by theorem*. The achievement is not that there is no postulate — it is that the whole theory has been compressed onto *exactly one*, sharply located and explicitly falsifiable.




Figure 2: Honest data cross-check (v246): the spectral-action prediction $g_1=g_2=g_3$, $\sin^2 \theta_W=3/8$ versus the measured couplings run with the SM RGEs (one-loop solid, two-loop dashed). The three lines do *not* meet at any scale (1-loop residual $\Delta\alpha^{-1} \approx 9$ at 2×10^{16} GeV; 2-loop minimal spread ≈ 3.2 at 1.7×10^{14} GeV) — a falsifiable tension. With SM content and no new states there is no admissible threshold source: not a fit, and not rescued by a cascade.

The Seam-Holomorphy Theorem — the one named closing target [E]/[O]
([verification/v234_seam_holomorphy_selection.py])

The bedrock above can be stated as *one condition with three provably-equivalent faces*, so that “closing the theory” becomes a single, sharply-bounded analytic theorem. **The condition:** the seam carries *no nontrivial abelian sector*. Its three faces all force E_8 :

1. **AQFT** (v83/v154): the boundary chiral net is *holomorphic* (μ -index 1, single vacuum sector). Holomorphic + $c = 8 \Rightarrow$ the unique even unimodular rank-8 lattice net $(E_8)_1$, and $c = 8 = g_{\text{car}} + N_{\text{fam}}$ is itself structural.
2. **Geometry** (v232): the seam link is a *homology 3-sphere* $\Leftrightarrow$ its binary group Γ is *perfect* $\Leftrightarrow \Gamma = 2I$ (the Poincaré sphere $S^3/2I$).
3. **Representation theory** (v219): exactly *one* 1-dim irrep (the trivial one).

These are the *same* number: $\#(1\text{-dim irreps}) = |\Gamma^{\text{ab}}| = |H_1(S^3/\Gamma)| = \#(\text{mark-1 affine-Dynkin nodes})$, which equals 1 *only* for E_8 ($A_n \rightarrow n+1$, $D_n \rightarrow 4$, $E_6 \rightarrow 3$, $E_7 \rightarrow 2$, $E_8 \rightarrow 1$). So $2I$ is the unique nontrivial perfect finite $SU(2)$ subgroup and $S^3/2I$ the unique homology-sphere space form. **Consequence:** Research Contract 2 (G_{net} /Target A), the carrier P2, and the Kleinian seam are *not* separate gates — they are this *one* condition.

The closing chain (what a complete closure would be), [O] at one step:

$$\begin{aligned} &\text{RP} + \text{gap} + \text{chirality} \xrightarrow{\text{v160}} \text{quasi-free bulk} \xrightarrow{?} \text{holomorphic boundary net} \\ &\xrightarrow{[\text{E}]} (E_8)_1 \xrightarrow{\text{v219/v228}} 2I, (2,3,5), \text{P2} \xrightarrow{\text{v179-v181+Kleinian genus-0}} \text{QGEO.SYM.01.} \end{aligned}$$

The free-bulk premise is already a fixed-point theorem (v160–v165); the Kleinian model supplies the genus-0 of the uniformisation chain for free. **The single remaining analytic step is “free Gaussian RP bulk $\Rightarrow$ holomorphic (single-sector) boundary net”** — the bulk-boundary statement that an invertible (Gaussian) bulk has a holomorphic chiral boundary. That step is now *closed modulo cited theorems* (the MMST route SEAM.EQUIV.MMST.01): the lattice/ S_3 stack pins it at every computable level (v367/v368, v376–v379, Lean FORM.SEAM.MMST.01), only the cited continuum existence staying [O]

(v336); v_{geo} stays the No-Unit metrology primitive and F_{transfer} stays external physics. This box certifies that the three faces coincide and force E_8 ([E]); the analytic step is closed modulo that cited theorem.

A physical realisation of the step ([verification/v235_seam_chern_simons.py]). The step lands in standard physics. A free gapped *bosonic* 2+1d bulk is an abelian Chern–Simons / K -matrix theory (even integer K), where $\#\text{anyons} = |\det K|$, the edge chiral central charge is $c = \text{signature}(K)$, and the edge net is *holomorphic* $\Leftrightarrow |\det K| = 1$ — so “no nontrivial abelian sector” is precisely $\det K = 1$. The extension tower of v92 is the anyon-condensation tower read by determinant: $D_5 \oplus A_3$ (det 16, 16 anyons) $\rightarrow SO(16)_1 = D_8$ (det 4) $\rightarrow (E_8)_1$ (det 1, holomorphic), all rank 8 with $c = \text{signature} = 8 = g_{\text{car}} + N_{\text{fam}}$, so the holomorphic end is the *Kitaev E_8 quantum-Hall state*. Holomorphy $\Leftrightarrow$ condensing the order- $|\mu_4| = 4$ *Lagrangian* glue (det $16 \rightarrow 16/4^2 = 1$); a partial order-2 isotropic only reaches det 4 (D_8). The residual is thereby a single integer condition — “the free RP seam condenses the order- $|\mu_4|$ Lagrangian glue” — the same sheet/ μ_4 selection as GATE.QGEO, but now with holomorphic $\Leftrightarrow$ det 1 a theorem (GATE.HOLO.02, [E]/[O]).

The residual as a physical statement ([verification/v237_seam_sre_closure.py]). The same integer condition is the physics of topological order: a K -matrix phase has ground-state degeneracy $|\det K|^g$ on a genus- g surface, so $\det K = 1 \Leftrightarrow$ no topological ground-state degeneracy on any closed surface $\Leftrightarrow$ the bulk is *short-range-entangled* (invertible). Among rank-8 even positive-definite K ($c = 8 = g_{\text{car}} + N_{\text{fam}}$) only E_8 is SRE — the unique nontrivial bosonic SRE 2+1d phase, the *Kitaev E_8 state*, whose edge is the holomorphic $(E_8)_1$. TFPT already has a unique vacuum on the plane (RP + gap + cluster + OS); the plane cannot see $\det K$ (no non-contractible cycles), the torus can. So the one analytic step is the sharp, falsifiable physical statement “the free RP gapped seam bulk is *short-range-entangled* (no topological order)” ($= \det K = 1 =$ the *Kitaev E_8 phase*, GATE.HOLO.03, [E]/[C]) — and the lattice/ S_3 stack now answers it at every computable level (torus $\det K = 1$, v378), so the keystone’s MMST route (SEAM.EQUIV.MMST.01) is *closed modulo cited theorems*, only the cited continuum existence (v336) staying [O] (the parent stays [O]).

Lemma 2 (U1 — wall landscape, corrected). *Parabolic degree zero forces the line weight-sums onto the wall (w_1, w_2, w_3) = (2, 1, 1). An explicit balanced representative (weights in $\frac{1}{3}$) is*

$$W_{\text{wall}} = \frac{1}{3} \begin{pmatrix} 2 & 1 & 2 & 1 \\ 1 & 0 & 0 & 2 \\ 0 & 2 & 1 & 0 \end{pmatrix},$$

each column a permutation of (0, 1, 2), row sums (2, 1, 1). **Honest result:** the $6^4 = 1296$ column-permutation matrices give 144 walls; under the full group $D_4 \times S_3(\text{lines}) \times S_3(\text{values})$ they form five orbits, not one. So W_{wall} is not singled out by symmetry — the uniqueness must come from the selector (Lemma U_4), not from a symmetry quotient. The wall landscape is nonetheless a finite, explicitly enumerated set; W_{wall} is one of the five. Machine: fully certified $C_U^{(1)}$ ([verification/v29_research_contract_certs.py]).

The splitting type is now algebraic (RH route, [verification/v32_rh_splitting.py]). Passing from the monodromy to the Fuchsian residues A_k (eigenvalues = weights $\{0, \frac{1}{3}, \frac{2}{3}\}$, $A_k = U^{k-1} A_0 U^{-(k-1)}$, $U = \text{diag}(1, i, -i)$) the twisted $\mathbb{Z}_4$ -average collapses exactly: $\sum_k U^k A_0 U^{-k} = 4 \text{diag}(A_0)$, so the exponents at infinity are $4 \text{diag}(A_0)$. A flat bundle with regular ∞ needs integer exponents summing to $4 = -\deg E$; with the cusp weights the only options are perms of (2, 1, 1) and (2, 2, 0), i.e.

$$\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2 \iff \text{diag}(A_0) = \left(\frac{1}{2}, \frac{1}{4}, \frac{1}{4}\right)$$

(the sibling $(2, 2, 0) \rightarrow \mathcal{O}(0) \oplus \mathcal{O}(-2)^2$). Such a Hermitian A_0 (spectrum $\{0, \frac{1}{3}, \frac{2}{3}\}$, that diagonal) exists by Schur–Horn. So the splitting is an algebraic condition on $\text{diag}(A_0)$. Honest wall: the global $\prod M_k = \mathbf{1}$ is the path-ordered monodromy (not $\exp(2\pi i \sum A_k)$), constraining off-diag(A_0) via the genuine RH solve; and R-extraction still needs the $C_6 \leftrightarrow \mathbb{P}^1 \setminus \mu_4$ bridge. The RH route fixes the splitting algebraically but not c_u/c_d .

Lemma 3 (U2 — the kill switch, run and hardened [E]). *The selected polystable wall representative must not collapse to the diagonal abelian $U(1)^3$ point if it is to give non-trivial $SU(3)_F$ transport*

mixing: (A) ∇_F^ polystable but not simultaneously diagonal (the desired breakthrough), or (B) diagonal collapse, in which case c_u/c_d stays honestly [C].* **Result: the cheap test is run robustly and lands on A.** The kill-switch fires iff the D_4 -symmetric cusp representation is reducible (commutant non-scalar). Across 400 random cusp-class generators of the $\mathbb{Z}_4$ family $M_k = U^{k-1}MU^{-(k-1)}$ the commutant is scalar at every point (irreducible 400/400), the reducible locus is hit with probability 0 (measure-zero, codimension ≥ 1), and the mixing strength stays bounded away from zero (min off-diag = 0.35 > 0.05). So case A is the generic behaviour, not an accident of the one explicit point of [verification/v33_explicit_flat_bundle.py]: (U_{wall}) survives its own cheapest falsification test. [verification/v38_uwall_killswitch.py]

Lemma 4 (U3 — D_4 fixed locus as an explicit variety, positive-dimensional). With $M_k = U^{k-1}AU^{-(k-1)}$ ($U = \text{diag}(1, i, -i)$, the $\mathbb{Z}_4$ rotation) and the reflection $M_1 = VM_1^{-1}V^{-1}$, $M_3 = VM_3^{-1}V^{-1}$, $M_2 = VM_4^{-1}V^{-1}$ (with $V = -[[1, 0, 0], [0, 0, 1], [0, 1, 0]]$, $V^2=1$, $VUV^{-1}=U^{-1}$), the relations become polynomials in the entries of $A \in C_{\text{cusp}}$; $\mathcal{X}_{\text{cusp}}^{D_4}$ is cut out as $\bigcup_k C_k$, parametrised by trace coordinates $x = \text{tr}(M_1M_2)$ (note $\text{tr} A = 1+\omega+\omega^2 = 0$). **Result** ($C_U^{(2)}$, [verification/v30_d4_character_variety.py]): adding the reflection to the $\mathbb{Z}_4$ locus of v19 does not isolate the point — the full D_4 -fixed product locus is still positive-dimensional ($|\text{tr}(M_1M_2)|$ varies continuously over $\approx [0, 3]$). So even the full D_4 symmetry cannot select ∇_F^* . Machine: dimension confirmed numerically; Gröbner/homotopy for the exact components (medium).

Lemma 5 (U4 — selectors as equations). The readouts $\rho \mapsto \det R(\rho)$, $\rho \mapsto \text{Spec}(Q_+(\rho))$, $\rho \mapsto \Lambda_{f,j}(\rho)$ are well-defined (cyclotomic/algebraic) on $\mathcal{X}_{\text{cusp}}^{D_4}$. **Acceptance criterion:**

$$\#\{\rho \in \mathcal{X}_{\text{cusp}}^{D_4} : \det R = 8, \text{Spec}(Q_+) = \{1, 2, 3\}\} = 1$$

in the polystable quotient. If more than one survives, the theory needs a further input and the gate is not closed. **The bottleneck (made precise by U3):** since the D_4 -fixed variety is positive-dimensional, the selector must do all the cutting — and evaluating $\det R(\rho)$, $\text{Spec}(Q_+(\rho))$ as functions on $\mathcal{X}_{\text{cusp}}^{D_4}$ requires the parabolic-degree $\leftrightarrow$ residue dictionary $R(\rho)$, i.e. the H2 equivalence. That dictionary was the single remaining analytic input of the (U_{wall}) gate. Status update: since v118–v122 the dictionary itself is exact — hexagon = sign-twisted family spectrum, lepton coefficients = resolvent determinants of the $W(A_3)$ monodromy, addresses pinned, margins theorems (boxes below); what remains is the strictly smaller R_4' , the establishment of the selectors ($n = (5, -9, 6)$ + the diamond determinants; since [verification/v134_dual_anchor.py]: the pair (d, n)).

What $R(\rho)$ is ([verification/v31_R_dictionary.py]). Two facts pin its nature. (i) Case A is alive: every D_4 -fixed cusp monodromy found is irreducible, so the wall point carries non-trivial $SU(3)_F$ mixing — the U2 kill switch does not fire to the diagonal case B. (ii) $R(\rho)$ is not algebraic: R is integer-valued, but the algebraic invariants of ρ (e.g. $\text{tr}(M_1M_2)$) vary continuously on the locus; an integer readout cannot be a continuous algebraic function. Hence $R(\rho)$ is the discrete invariant of the non-abelian Hodge / Mehta–Seshadri map (the parabolic-degree / Hodge-filtration jumps of the harmonic bundle E_ρ): integer output, but its computation is the transcendental harmonic-metric (Hitchin) solve on the generic Higgs locus. **Refinement (U4'')**, [verification/v40_harmonic_metric.py]: at the polystable point TFPT actually selects this pessimism does not apply — see the box below.

Progress: the selector side of U4 is closed on the explicit point ([E]). While $R(\rho)$ as a function on the whole locus needs the Hitchin solve, the two U4 selectors are algebraically accessible on the explicit flat bundle of [verification/v33_explicit_flat_bundle.py] — no PDE. Computed exactly from A_0 : the exponents at infinity $\text{eig}(\sum_k A_k) = \{2, 1, 1\}$ give the splitting type $E = \mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$, i.e. the parabolic anchor $a = (1, 1, 2)$; the parabolic degree is 0 ($\deg E = -4 = -|\mu_4|$ plus four cusp weight-sums of 1), so the point is *polystable* (Mehta–Seshadri substrate). The two selectors then read off the bundle directly:

$$\begin{aligned} \det R &= 8 = n \cdot a \quad (\text{anchor pairing on the splitting type}), \\ \text{Spec}(Q_+) &= \{1, 2, 3\} = 3\alpha + 1 \quad (\alpha = \text{cusp weights } \{0, \tfrac{1}{3}, \tfrac{2}{3}\}) \end{aligned}$$

So $\det R = 8$ is the anchor pairing of the splitting type and $\text{Spec}(Q_+)$ is the affine image ($d=3$ the weight denominator) of the cusp weights — *both selectors are read off the explicit bundle's algebraic data*. **Residual (sharpened below by Readout Rigidity):** the quark ratio c_u/c_d turns out *not* to need the Hitchin datum at all — it is *constant* on this discrete selector stratum (v49), hence an integer Plücker readout. The only genuinely transcendental input left is the *absolute* amplitude normalisation (U_{point}), which is an anchor, not a missing PDE. [verification/v39_uwall_selectors.py]

Breakthrough: the harmonic metric is finite linear algebra, not a PDE ([E]). The U4 “transcendental Hitchin solve” is the worst case for a *generic* Higgs bundle, but the point TFPT selects is *polystable*, and by Mehta–Seshadri a parabolic-degree-0 polystable bundle is *unitary* — a unitary representation has the *constant* invariant Hermitian harmonic metric and zero Higgs field $\Phi = 0$. So at the selected point the harmonic metric is finite linear algebra. On the explicit bundle of [verification/v33_explicit_flat_bundle.py] this is realised: the raw monodromies M_k are non-unitary ($\|M_k^\dagger M_k - \mathbf{1}\| \approx 0.83$), but there is a *unique* common invariant Hermitian form H ($\dim \ker\{M_k^\dagger H M_k = H\} = 1$), it is *positive-definite*, and the unitarised holonomy $\widetilde{M}_k = H^{1/2} M_k H^{-1/2}$ is unitary ($\sim 10^{-8}$), lies in the cusp class $\{1, \omega, \omega^2\}$ with $\prod \widetilde{M}_k = \mathbf{1}$, and has *clean* matrix elements $|\widetilde{M}_k|_{ij} \in \{0, \frac{1}{2}, \frac{1}{\sqrt{2}}\}$. **So $C_U^{(3)}$ reduces from a transcendental PDE to: this finite unitarisation (done) + the combinatorial word-length R (closed: $H\mathbf{1} + \det R = 8$).** *Residual:* the feared PDE is gone; the quark ratio c_u/c_d is then fixed combinatorially by Readout Rigidity (below), and only the *absolute* amplitude scale remains as an anchor. [verification/v40_harmonic_metric.py]

The final leg-assignment test (honest result) ([E]). Does the harmonic-frame holonomy $\{0, \frac{1}{2}, \frac{1}{\sqrt{2}}\}$ feed the $\delta = \frac{1}{2}$ resolvent to give the lepton amplitudes $\Lambda = (0.475, 1.107, 0.917)$? **(i) Ruled out as a literal identity:** $\Lambda_\mu = 1.107 > 1$, while unitary holonomy entries have modulus ≤ 1 , so the amplitudes *cannot* be holonomy matrix elements — they are the non-unitary resolvent Green function (as v19 already noted). **(ii) Suggestive positive:** the harmonic-frame holonomy diagonal modulus is exactly $|\text{diag } \widetilde{M}_0| = (0, \frac{1}{2}, \frac{1}{2})$, and that $\frac{1}{2}$ *equals* the distinguished lepton transport value $\delta = \frac{1}{2}$ that v20 used by hand — so the geometry plausibly *supplies* δ (an observation at the explicit point; genericity across the locus is future work). **(iii) Clean negative:** no natural holonomy construction (e.g. $(\mathbf{1} - \frac{1}{2}\widetilde{M})^{-1}$) maps the holonomy to the amplitude vector. **Verdict:** the amplitudes are the $\delta = \frac{1}{2}$ resolvent (closed combinatorially via the word-length R); the holonomy supplies δ and the leg selection, not the amplitude magnitude. c_u/c_d is *not* obtained from the holonomy alone — a clean, honest negative on the literal reproduction, with one suggestive geometric link ($\delta = \frac{1}{2}$). The negative is now *explained*: the scalar leg is wrongly typed — see the Exterior Leg Lemma. [verification/v41_leg_assignment.py]

The torsion normal form and the $\delta = \frac{1}{2}$ theorem ([E]). The v41 “genericity is future work” item is now executed — and yields a theorem. *(i) Flatness = μ_4 torsion [E]:* for any M (symbolic), $\prod_k U^k M U^{-k} = (MU)^4$ — the $\mathbb{Z}_4$ -family flatness constraint *is* the torsion statement “ $T = MU$ is a fourth root of unity”: the μ_4 atom appears as the *order* of the twisted cusp generator. *(ii) The $\delta = \frac{1}{2}$ theorem [E] (both directions):* on the *involutive* branch ($T^2 = \mathbf{1}$) T is a Hermitian reflection $2vv^\dagger - \mathbf{1}$, and the cusp trace condition $2v^\dagger U^{-1}v = 1$ splits *exactly* into $|v_1|^2 = \frac{1}{2}$ (real part) and $|v_2| = |v_3|$ (imaginary part), forcing

$$\text{diag } M = (0, \frac{i}{2}, -\frac{i}{2}), \quad \text{spec } M = \{1, \omega, \omega^2\} \text{ automatic}$$

($\chi_M = \lambda^3 - 1$ from unitary + trace 0 + $\det 1$). So $\delta = |M_{22}| = \frac{1}{2}$ holds *exactly and constantly* on the whole two-parameter branch — the distinguished lepton transport value that v20 used by hand is a μ_4 -torsion identity, not an observation. *(iii) Reflection lemma [E]:* the D_4 reflection forces $\text{diag } M = (r, z, \bar{z})$ with r real and $\text{Re } z = -r/2$ — the whole D_4 diagonal freedom is *one* real parameter, and $\delta = \frac{1}{2} \iff r = 0$. *(iv) Branch census [E] (seeded):* the full D_4 -fixed flat cusp locus realises only $\text{tr}(MU) \in \{1, -1\}$; the involutive branch has the diagonal *constant* $(0, \frac{1}{2}, \frac{1}{2})$ to machine precision, while on the $\{1, i, -i\}$ branch d_1 varies over $[0, 1]$ — and the explicit v33/v40 point sits exactly in its $d_1 = 0$ slice, the *same* δ value. *Residue [C] (recorded):* whether the anchor splitting

$\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$ (Mehta–Seshadri side) forces the $d_1 = 0$ slice — or selects the involutive branch — is the remaining question; $\delta = \frac{1}{2}$ itself is no longer the open item. [\[verification/v114_torsion_delta.py\]](#)

The anchor pins the residue: an exact normal form for the (U_{wall}) bundle ([E]). The **v114** branch question is answered — and the **v33** *numerical* Riemann–Hilbert point acquires an *exact closed form*. (i) μ_4 -average lemma [E]: $\sum_k U^k X U^{-k} = 4 \text{diag } X$ for any X — μ_4 conjugation-averaging is the diagonal projection, so the exponents at infinity are $4 \text{diag } A_0$ and the anchor splitting holds iff $\text{diag } A_0 = (2, 1, 1)/4 = a/|\mu_4|$: the residue diagonal is the anchor over μ_4 (v33’s diagonal was forced, not chosen). (ii) The $(8, 0, 5)/144$ lemma [E]: the anchor diagonal and the cusp spectrum $\{0, 1, 2\}/N_{\text{fam}}$ pin $\sum |a_{ij}|^2 = \frac{13}{144}$, and with $a_{13} = 0$ the determinant condition solves *uniquely* to

$$\left(|a_{12}|^2, |a_{13}|^2, |a_{23}|^2 \right) = \frac{(8, 0, 5)}{144} \quad \begin{array}{l} 8 = \text{rank } E_8, \ 5 = g_{\text{car}}, \\ 144 = (|\mu_4| N_{\text{fam}})^2, \ 8+5=13=\Delta_Q \end{array}$$

— the off-diagonal weights are compiler atoms, and the total 13 is the *same* Δ_Q that denominates $c_u/c_d = 55/117 = 5 \cdot 11/(9 \cdot 13)$. (iii) *Exact normal form* [E]: $A_0^* = \begin{bmatrix} 1/2 & \sqrt{2}/6 & 0 \\ \sqrt{2}/6 & 1/4 & \sqrt{5}/12 \\ 0 & \sqrt{5}/12 & 1/4 \end{bmatrix}$ has $\chi = \lambda(\lambda - \frac{1}{3})(\lambda - \frac{2}{3})$ *exactly*. (iv) A_0^* is the RH solution [E]: its big-circle monodromy is trivial to 10^{-10} , $\prod M_k = \mathbf{1}$, and the unitarised holonomy has $|\text{diag } \widetilde{M}_0| = (0, \frac{1}{2}, \frac{1}{2})$, $\text{tr}(\widetilde{M}_0 U) = 1$ — the **v33/v40** point is gauge-equivalent to A_0^* (the a_{12}, a_{23} phases are diagonal gauge). (v) *The anchor forces the $d_1 = 0$ slice* [E]: a multi-seed RH scan over the anchor-pinned spectral manifold lands at *every* flat solution on the *same* gauge orbit with $|\text{diag } \widetilde{M}_0| = (0, \frac{1}{2}, \frac{1}{2})$ — combined with **v114**, the whole harmonic diagonal is **anchor + torsion forced**; $\delta = \frac{1}{2}$ needs no selection. *Residue* [C]: $a_{13} = 0$ and the gauge-orbit uniqueness are numerical (one component across all seeds); promoting them to [E] via Gröbner on the RH constraints is the remaining U_{point} step. [\[verification/v115_anchor_residue.py\]](#)

The resonance theorem: $M_\infty = \mathbf{1} \iff a_{13} = 0$ — **one gauge orbit** [E]
([\[verification/v116_resonance_uniqueness.py\]](#))

The anticipated Gröbner promotion turned out to be a *linear* computation. Writing the equivariant system at infinity as $dY/dw = -(1/w)(B_0 + B_1 w + \dots)Y$ with $B_m = \sum_k i^{km} U^k A_0 U^{-k}$, the *twisted* μ_4 averages collapse [E]:

$$B_0 = 4 \text{diag } A_0, \quad B_1 = 4a_{12}E_{12} + 4\overline{a_{13}}E_{31}$$

(only the eigenvalue-ratio $-i$ cells survive). The anchor exponents $\text{diag}(2, 1, 1)$ have resonance gap 1; the level-1 formal-gauge equation $(1 - \text{ad } R_0)H_1 = R_1$ is singular exactly on the cells $\{(2, 1), (3, 1)\}$, where B_1 has entries $(0, 4\overline{a_{13}})$; all higher levels are non-resonant and the formal monodromy is $e^{-2\pi i \text{diag}(2, 1, 1)} = \mathbf{1}$. Hence [E]:

$$M_\infty = \mathbf{1} \iff a_{13} = 0$$

— the transcendental Riemann–Hilbert condition collapses to *one linear equation*. *Uniqueness corollary* [E]: $a_{13} = 0$ + the $(8, 0, 5)/144$ moduli + phases = diagonal gauge $\Rightarrow$ the flat anchor locus is **exactly one gauge orbit**, the exact A_0^* : within the μ_4 -equivariant anchor class the (U_{wall}) bundle datum is *algebraically unique*. *Sharpness* [E]: $\|M_\infty - \mathbf{1}\| \sim 8.5 |a_{13}|$ (0.17 at $a_{13} = 0.02$) vs 10^{-10} at A_0^* . *Honest scope* [C]: the equivariant ansatz is the established D_4 selector input (**v19/v30**); the *residue* side is now [E], while the holonomy values (the harmonic diagonal $(0, \frac{1}{2}, \frac{1}{2})$) remain [E] — transcendental monodromy of an exact system.

The monodromy is the Weyl group of A_3 — the holonomy is exact [E]
([\[verification/v117_monodromy_weyl_a3.py\]](#))

The last [E] item of the chain — the holonomy *values* — closes: the unitarised monodromy of the exact

$A_0^\star$ system is itself *exact*, with entries in $\frac{1}{2}\mathbb{Z}[i]$:

$$\widetilde{M}_0 = \begin{pmatrix} 0 & -\frac{1+i}{2} & \frac{1-i}{2} \\ -\frac{1+i}{2} & -\frac{i}{2} & -\frac{1}{2} \\ \frac{1-i}{2} & -\frac{1}{2} & \frac{i}{2} \end{pmatrix}$$

unitary, $\det = 1$, $\text{tr} = 0$, $\chi = \lambda^3 - 1$ (the cusp class *exactly*), $\widetilde{M}_0^3 = \mathbf{1}$, and $\text{diag } \widetilde{M}_0 = (0, -\frac{i}{2}, +\frac{i}{2})$ — so $d_1 = 0$ and $\delta = \frac{1}{2}$ are now *matrix entries*, not observations; $(\widetilde{M}_0 U)^4 = \mathbf{1}$ with $\text{tr}(\widetilde{M}_0 U) = 1$ (the $\{1, i, -i\}$ branch, $d_1 = 0$ slice — exactly where **v114/v115** located the physical point). The group **[E]**: exact enumeration of $\langle U, \widetilde{M}_0 \rangle$ gives order 24 with order statistics (1, 9, 8, 6) and character values (3, -1, 0, 1) — uniquely S_4 in its standard $\otimes$ sign representation, i.e. the (twisted) reflection representation of the **Weyl group of the family lattice A_3** :

$$\text{monodromy}(U_{\text{wall}}) = W(A_3) = S_4, \quad |W(A_3)| = 4! = 24$$

— the flavor wall realises the A_3 factor of the glue $D_5 \oplus A_3 + \mu_4$ as its Weyl group (U = a 4-cycle, the μ_4 deck; $\widetilde{M}_0$ = a 3-cycle, the family rotation). *Identification [E]*: the ODE monodromy of $A_0^\star$ equals the exact $\widetilde{M}_0$ to 3.5×10^{-10} , and the invariant form H is U -invariant to 10^{-11} . *Status*: holonomy values **[E]**; only the analytic identification stays **[E]**. The δ thread (**v20** $\rightarrow$ **v40/v41** $\rightarrow$ **v114** $\rightarrow$ **v115** $\rightarrow$ **v117**) is closed end to end.

The hexagon is the sign-twisted family spectrum — first exact H2 piece [E]
(**[verification/v118_hexagon_family_dictionary.py]**)

The missing “ $C_6 \leftrightarrow \mathbb{P}^1 \setminus \mu_4$ dictionary” (H2) acquires its first exact entry. *Sign-twist lemma [E]*: $-\widetilde{M}_0$ has order 6 and

$$\text{spec}(\widetilde{M}_0) \cup \text{spec}(-\widetilde{M}_0) = \mu_6$$

— the six-site hypercharge hexagon spectrum *is* the family monodromy spectrum together with its sheet twist ($\mathbb{Z}_6 = \mathbb{Z}_2 \times \mathbb{Z}_3$ realised as $(-1) \times \langle \widetilde{M}_0 \rangle$); the **v20** denominators $\frac{5}{4} - \cos(r\pi/3)$ are exactly the eigen-denominators $|1 - \zeta_6^r/2|^2$ of the two family resolvents $(\mathbf{1} \mp \frac{1}{2}\widetilde{M}_0)^{-1}$. *Cyclotomic determinant [E]*: $\det(\mathbf{1} - t\widetilde{M}_0) = 1 - t^3 = (1 - t)\Phi_3(t)$, so $\det(\mathbf{1} \mp \frac{1}{2}\widetilde{M}_0) = (\frac{7}{8}, \frac{9}{8})$ — and the lepton coefficients become resolvent determinants: $c_e = |\mu_4|/(2\det_-) = \frac{16}{7}$, $c_\mu = N_{\text{fam}}/(2\det_+) = \frac{4}{3}$, $c_\tau = |\mu_4|\det_-/N_{\text{fam}} = \frac{7}{6}$, product rule $= |\mu_4|/\det_+ = \frac{32}{9}$. The lepton 7 and 9 *are* the family-resolvent determinants; $\delta = \frac{1}{2} = |\langle \widetilde{M}_0 \rangle_{22}|$ is supplied by the matrix itself. *Sheet-extended group [E] (audit)*: $\langle U, \widetilde{M}_0, -\mathbf{1} \rangle$ has order $48 = \Omega_{\text{adm}} = N_{\text{fam}} \dim S^+$ — the seed’s quartic coefficient. *Residue [C]*: the dictionary’s *values* are exact; its *address table* (which hexagon site each fermion occupies; the quark composition) remains **v20**’s established input / open — nothing re-fished.

The address table: the lepton words are the compiler atoms [E]/[C]
(**[verification/v120_address_table.py]**)

The address table acquires its structure — on word-length integers *frozen* in **v18/v20**, long before today’s readings (the red-team firewall). (i) *The lepton words are the atoms [E]*:

$$(L_e, L_\mu, L_\tau) = (8, 5, 3) = (\text{rank } E_8, g_{\text{car}}, N_{\text{fam}}) = (p_0 + e_2, e_2, p_0)$$

in mass order (longest word = lightest lepton). (ii) *Addresses = division by the hexagon [E]*: $(r, w) = L \bmod p_2$ with $p_2 = 6 = |R^+(A_3)|$ itself an atom: $e \rightarrow (2, 1)$, $\mu \rightarrow (5, 0)$, $\tau \rightarrow (3, 0)$; $\sum r = 10 = p_3 = \mathcal{A}_\Lambda$, $\sum L = 16 = \dim S^+$. (iii) *Sheet parity [E]*: by the **v118** dictionary, e sits on the *untwisted* sheet ($\omega \in \text{spec } \widetilde{M}_0$), μ, τ on the *twisted* sheet $(-\omega, -1)$; the anchor lepton τ occupies the unique *real* twisted eigenvalue -1 — exactly the site where **v20**’s product rule replaces the resolvent. (iv) *Quark sum rules [E] (frozen v18 words 7, 7, 5, 3, 2, 0)*: up-type sum $= 10 = p_3$; down-type sum $= 14 = p_1 + p_3$; **all quarks** $= 24 = |W(A_3)|$ (the monodromy group order, **v117**); **all nine charged fermions** $= 40 = p_1 p_3 = a^T R a$ (**v119**); quark $\sum r = 2p_2$, $\sum w = |\mathbb{Z}_2|$; the top sits at the vacuum site $(0, 0)$ — zero word, no suppression, the established ladder anchor. *Residue [C]*: five independent closures (16, 10, 14, 24, 40) constrain the address map; the per-fermion *assignment* remains the **v18/v20** input — deriving it is the remaining H2

content.

The address table is pinned: R is the unique hexagon matrix with atom margins [E] ([verification/v121_address_pinning.py])

The per-fermion assignment closes at the *information* level. The established identity $L = R + 2U$ (v95) says the word-length table *is* $L[\text{sector}][\text{generation}]$ — rows (up; down; lepton) = $((7, 3, 0); (7, 5, 2); (8, 5, 3))$, the v18/v20 words verbatim — so the addresses are the entries of the *residue operator* R plus one hexagon turn for the first generation, and R 's first column is the **anchor** $a = (1, 1, 2)$. The margins are atoms [E]: row sums $(4, 8, 10) = (e_1, \text{rank } E_8, p_3)$, column sums $(4, 13, 5) = (e_1, \Delta_Q, g_{\text{car}})$. And the census closes it [E]:

among all 3×3 matrices with entries in the hexagon $\{0..5\}$,
exactly 17 have the atom margins — and exactly ONE has $\det = 8$

— and it is R (also the unique one with $\text{SNF} = (1, 1, 8)$; all 17 determinants are distinct, 8 occurs once — control computed explicitly). **The assignment table carries zero residual information beyond the atom margins and the rank determinant.** Within-sector ordering is the established transport reading ($m \sim \lambda_Y^L$: longer word = lighter fermion). *Residue [C]*: the open H2 item shrinks from “derive 9 per-fermion integers” to “derive the six atom margins + $\det = \text{rank}$ ”. *Firewall*: R , L and all word lengths were frozen in v4/v18/v20/v71 long before this census — no degrees of freedom spent.

The margins are theorems ([verification/v122_margin_theorem.py]). Even the margins are not input. Three selectors, all established and frozen *before* the address question was posed — (S1) the D_4 annihilator $n = (5, -9, 6)$ with $n^T R = (8, 0, 0)$ (it kills the unchanged (c_2, c_3) plane; $n \cdot a = 8$), (S2) $\det R = 8$, (S3) $\det K = 4$ with $K = R + Q\Sigma$ — pin R *uniquely* among hexagon matrices [E]: the (S1) census has $4 \times 4 \times 4 = 64$ candidates, (S2) leaves 12, (S3) leaves **exactly one**, and it is R . Hence the atom margins, the anchor column $Re_1 = a$ and all sum rules above are *consequences* — their input status is revoked. *Bonus lemma [E]*: on the (S1) + (S2) locus $\det(M + 2U) = 20 = \det L$ holds for *all* 12 candidates — the fourth quartet determinant is redundant, the compiler's familiar overdetermination. *Residue [C]*: H2 reduces to the establishment of the selectors themselves (n and the diamond determinants — pre-existing inventoried objects); the address question introduces *no new residual class*.

The dual-normal selector: (d, n) **pins** R **columnwise** ([verification/v136_dual_normal_selector.py]). The selector list shrinks once more. With the dual anchor $d = a^T R^{-1}$ ([verification/v134_dual_anchor.py]) and the torsion normal n , each *column* of R obeys $d \cdot c_j = a_j$ and $n \cdot c_j = 8\delta_{1j}$; the common kernel of (d, n) is the primitive vector $(6, 8, 7)$ (audit: p_2 , rank E_8 , scalaron-7), so within the address box $\{0..5\}^3$ each column is the *unique* lattice point (census: 1 of 216 each) — $(d, n) \Rightarrow R$, and the 6^9 census above collapses to three column problems [E]. *Honesty*: K is *not* pinned directly ($d^T K = (1, \frac{1}{2}, 1)$); $\det K = 4$ comes via the μ_4 wall ([verification/v135_det_surface.py]). *The mechanism is bounded*: the zero mode of A_0^* carries the exact spectral normal $\sigma = (2, -9, 5) = (|\mathbb{Z}_2|, -N_{\text{fam}}^2, g_{\text{car}})$ (signed squares; $\|v\|^2 = \frac{16}{9}$ with $16 = \dim S^+$, $\text{adj } A_0^* = \frac{2}{9} P_v$ with $\frac{2}{9}$ = the SdS entropy curvature), $\sigma \cdot \mathbf{1} = -|\mathbb{Z}_2|$, $\sigma \cdot a = N_{\text{fam}}$, $n \cdot \sigma = 121 = \|\text{Pl}(K)\|_1^2$; and the exact $W(A_3)$ orbit control shows *no* group image of σ is proportional to n — the naive cofactor/orbit lift fails, the step $\sigma \rightarrow n$ is a genuine mechanism beyond the monodromy group. R_4' *restated [C]*: derive the *pair* (d, n) ; then R is forced columnwise.

The selector triangle ([verification/v139_selector_triangle.py]). The pair itself now collapses further. First, d is *pure anchor data*: $d = \frac{3}{2}a - 2 \cdot \mathbf{1}$ exactly — the closed form never mentions R , so the first selector is *derived*, not residual. Second, the frame $(\mathbf{1}, a, \sigma)$ is a basis of $\mathbb{Z}^3$ with $\det(\mathbf{1}|a|\sigma) = 11 = \|\text{Pl}(K)\|_1$ (the quark constant as the frame volume), and n is the *unique* integer covector with the three atom pairings

$$n \cdot \mathbf{1} = 2 = |\mathbb{Z}_2|, \quad n \cdot a = 8 = \text{rank } E_8, \quad n \cdot \sigma = 121 = \|\text{Pl}(K)\|_1^2$$

— the cofactor definition is replaced by frame data plus three atoms. On the $(\mathbf{1}, a)$ -constrained line the σ -pairing is $11(11+t)$: divisible by 11 for *every* t , a perfect square exactly at the true n ($t=0$). The review's predicted lift is exact as an audit identity: $n = \text{reverse}(\sigma) + |\mu_4|e_3$ (generation flip + μ_4 lift at the third slot).

Frame integrality: three pairings $\rightarrow$ two ([verification/v142_frame_integrality.py]). The third

pairing is *not* independent. The pairing triples $(m \cdot \mathbf{1}, m \cdot a, m \cdot \sigma)$ of *integer* covectors form an index-11 sublattice of $\mathbb{Z}^3$, cut out by the single congruence

$$x - 3y + z \equiv 0 \pmod{11} \quad (\text{index } 11 = \|\text{Pl}(K)\|_1 = \det(\mathbf{1}|a|\sigma)),$$

so with the two line pairings at their atoms $(x, y) = (|\mathbb{Z}_2|, \text{rank } E_8) = (2, 8)$, integrality *forces* $z \equiv 0 \pmod{11}$ — the $11(11+t)$ line *is* the integer line. Primitivity forces t *even* (odd t makes every entry of $n(t)$ even), and $t=0$ is the unique even t with a square σ -pairing for $|t| < 88$ (the global pick keeps the square selector — recorded honestly). Both line pairings are Cramer-reducible to inventoried identities ($n \cdot a = \det R$ by Laplace along the anchor column; $n \cdot \mathbf{1} = 8(R^{-1}\mathbf{1})_1$ with $R^{-1}\mathbf{1} = \frac{1}{4}(1, 1, -1)$, **v134**) — restructuring, not establishment.

The pairing values are atom identities (`[verification/v145_pairing_atoms.py]`). Even the two line-pairing *values* carry no independent information. Reading the **v139** lift structurally — $n = w_0(\sigma) + |\mu_4|e_3$ with w_0 the A_3 *exponent duality* $m \leftrightarrow h-m$ (an established Lie involution; lift slot = the top-exponent line, lift size = the glue order) — all three pairings become atom arithmetic:

$$n \cdot \mathbf{1} = |\mu_4| - |\mathbb{Z}_2| = |\mathbb{Z}_2|, \quad n \cdot a = |\mu_4| a_3 = 8 = \text{rank } E_8 \text{ via } |\mu_4| + g_{\text{car}} = N_{\text{fam}}^2 \quad (4 + 5 = 9, \sigma \perp w_0(a)),$$

and $n \cdot \sigma = \|\sigma\|^2 - N_{\text{fam}}^2 + |\mu_4|g_{\text{car}} = 110 - 9 + 20 = 121$ with $\|\sigma\|^2 = |\mathbb{Z}_2|^2 + N_{\text{fam}}^4 + g_{\text{car}}^2 = 110 = |\mathbb{Z}_2|g_{\text{car}}\|\text{Pl}(K)\|_1$. R_4' *residue* **[C]** (*final form*): derive the *one* map $n = w_0(\sigma) + |\mu_4|e_3$ — no independent number remains; everything else in the chain anchor $\rightarrow (d, n) \rightarrow R$ is exact.

Both dual normals are anchor data (`[verification/v149_cusp_normal.py]`). The remaining opacity dissolves in the canonical frame: the pairings of n with the *cusp eigenbasis* (the geometric basis of **v98**) are exactly

$$n \cdot e_1 = 6 = p_2(a), \quad n \cdot e_2 = 3 = p_0(a), \quad n \cdot e_3 = 5 = e_2(a)$$

— one anchor invariant per Q_+ degree, and the cusp matrix is unimodular, so these three pairings determine n uniquely. With $d = \frac{3}{2}a - 2 \cdot \mathbf{1}$ the dual normal *pair* (**v134**) is anchor data through and through — one normal per frame; the cusp data $(6, 3, 5)$, the frame pairings $(2, 8, 121)$ and the w_0 -lift are three presentations of one covector, all atomic (equivalences exact). *Audit*: $\sigma \rightarrow (5, 1, 2)$; alternate reading $(6, 3, 5) = (|R^+(A_3)|, N_{\text{fam}}, g_{\text{car}})$; sum $14 = \dim G_2$. R_4' *residue* **[C]** (*cusp form*): derive the *assignment* — why the Q_+ degrees $(1, 2, 3)$ carry (p_2, p_0, e_2) in that order; one discrete assignment, zero continuous freedom.

Exterior Leg Lemma — the quark u/d leg is a $\Lambda^2 F$ area, not a scalar **[E]/[C]**

The **v41** negative is a *category error*, not a dead end. u and d share the hexagon-radial data $(r, w) = (1, 1)$ ($L = 7$ for both), and the C_6 resolvent $c(r, w) = |\mu_4|^w / (\frac{5}{4} - \cos \frac{r\pi}{3})$ reads *only* (r, w) — so any *scalar* leg gives $c_u/c_d \equiv 1$. A scalar sees radius, not orientation. The u/d separation lives in the smallest *non-scalar* invariant of the $SU(3)_F$ composition: the exterior square $\Lambda^2 F$, i.e. the oriented anchor-plane area $\text{Pl}_{1,a}(K)$. The “11” is then *generated*, not read:

$$(K, \mathbf{1}, a) \rightarrow B_K = \left(\frac{13}{18} \frac{8}{11} \frac{4}{6}\right) \rightarrow \text{Pl}(K) = (-1, 6, 4) = (-N_\Phi, |R^+(A_3)|, |\mu_4|) \rightarrow \|\text{Pl}(K)\|_1 = 11,$$

$$\frac{c_u}{c_d} = \frac{g_{\text{car}} \|\text{Pl}_{1,a}(K)\|_1}{N_{\text{fam}}^2 \Delta_Q} = \frac{5 \cdot 11}{9 \cdot 13} = \frac{55}{117} \quad (\text{anchor microcode} = \frac{e_2(a)(p_3(a)+1)}{p_0(a)^2(2p_2(a)+1)}).$$

Typing: **[E]** for the Plücker / microcode identities ($\text{Pl}(K) = (-1, 6, 4)$, $\|\cdot\|_1 = 11$, $5 \cdot 11 / (9 \cdot 13) = 55/117$); and — by Readout Rigidity (**v49**, below) — **[E]** also for the *physical* ratio $\mathcal{C}_{ud}(\rho^*) = \text{this } \Lambda^2 F \text{ readout}$, which is *constant* on the derived selector stratum and so does *not* need any point-selection on $\Lambda^2 F$. The quark residual is thereby re-typed from “find the right scalar y ” to a closed exterior readout for the *ratio*; only the *absolute* amplitude scale (U_{point}) stays as an anchor, no new scalar, no fishing.

Audit (the 121 lemma, `[verification/v119_review_validation_2.py]`): the “11” has a second, independent appearance **[E]**: with $h = \mathbf{1} + a = (2, 2, 3)$ in the canonical anchor ordering, $h^T R h = 121 = 11^2 = \|\text{Pl}(K)\|_1^2 = (p_3 + 1)^2$ — the quadratic *anchor norm* of the residue operator R reproduces the Plücker integer (sensitivity control: the three orderings of h give $\{105, 121, 135\}$; only the canonical one yields 121). Bonus checksums: $\mathbf{1}^T R \mathbf{1} = 22 = 2 \times 11$ (the entry sum of R) and $a^T R a = 40 = p_1 p_3 = 10 b_1 - 1$. Typed *audit*, not load-bearing — the oriented Plücker mechanism above remains the derivation. `[verification/v42_exterior_leg.py]`

Bridge test ([[verification/v43_exterior_bridge.py](#)]). The typing is confirmed: for the $SU(3)$ holonomy the exterior square is the conjugate representation, $\Lambda^2(\widetilde{M}) = \widetilde{M}$ exactly — the exterior leg *is* the $\bar{3}$. But the *continuous* action $\Lambda^2(\widetilde{M}) \cdot (\mathbf{1} \wedge a)$ does *not* equal the *integer* $\text{Pl}(K) = (-1, 6, 4)$: the integer Plücker is the combinatorial K datum. So the bridge $\rho^* \rightarrow \text{Pl}(K)$ is the *discrete* non-abelian-Hodge invariant (the integer R ; $\det R = 8$ and $\text{Spec}(Q_+) = \{1, 2, 3\}$ already confirmed on the bundle, Lemma U4 box), not a continuous exterior-square computation — exactly the H2 equivalence, now pinned from the $\Lambda^2 F$ side.

Lie-level grounding — no special math ([[verification/v44_carrier_exterior.py](#)]). The discrete invariant has a home already in TFPT: the carrier half-spinor is the *even exterior algebra* of the 5-slot carrier, $16 = \Lambda^{\text{even}}(5) = \binom{5}{0} + \binom{5}{2} + \binom{5}{4} = 1 + 10 + 5$, and under $SU(5) \subset SO(10)$ the up quark sits in $10 = \Lambda^2(5)$ while the down sits in $\bar{5} = \Lambda^4(5)$. So the *exterior degrees* are $\deg(u^c) = 2$, $\deg(d^c) = 4$, with $\deg(u) + \deg(d) = 6 = |R^+(A_3)|$ (the hexagon) and $\deg(d) - \deg(u) = 2 = |\mathbb{Z}_2|$ (the sheet). The “ Λ^2 ” of the exterior leg is therefore not exotic Hodge data — it is the carrier’s own Λ^2 grading (the up quark *lives* in $\Lambda^2(5)$). This re-types the residual once more: from “discrete non-abelian-Hodge invariant” to “the carrier exterior degree” — a standard [E] fact. *Honest scope*: this grounds the *type*; the Plücker *number* 55/117 stays the family $\text{Pl}(K)$ [E], and the carrier- $\Lambda^2 \leftrightarrow$ family- $\Lambda^2 F$ link (via family $\otimes$ carrier = 15 = 16–1) plus the normalisation remain [C].

The “11” is Pascal — a third, simplest origin ([[verification/v45_family_exterior.py](#)]). Following that link, $\text{Pl}(K)$ is the *exterior algebra of the family fundamental* $4 = \mu_4$ (the $A_3 = SU(4)$ fundamental): $\Lambda^k(4) = \binom{4}{k}$ is Pascal row 4 = (1, 4, 6, 4, 1) with full sum $2^4 = 16 = \dim S^+$, and $|\text{Pl}(K)| = (1, 4, 6) = \{\Lambda^0, \Lambda^1, \Lambda^2\}(4)$ are exactly its distinct exterior powers. Hence

$$\|\text{Pl}(K)\|_1 = \sum_{k=0}^2 \binom{4}{k} = 11 = 2^4 - \binom{4}{3} - \binom{4}{4} = 16 - g_{\text{car}}$$

(the cumulative $\Lambda^\bullet(\mu_4)$ up to degree 2 = $\deg(u^c)$), and family $\otimes$ carrier = 15 = $\dim \mathfrak{su}(4) = \Lambda^2(5) \oplus \Lambda^4(5) = 10 + 5 = N_{\text{fam}} g_{\text{car}}$ read three ways. So the “11” is generated a *third* way — the cumulative family exterior algebra up to the up-quark’s degree — pure Pascal/branching, *no special math*. The [C] physical normalisation is unchanged; the algebraic origin of 55/117 is now triply grounded.

Lemma 6 (U5 — the “11” worry, *discharged* by Readout Rigidity). *The original worry was that $c_u/c_d = C_{ud}(\rho^*)$ must be evaluated at the uniquely selected ρ^* without using 11 as input, so that until then 55/117 would be a mere table-level rational shadow. **This is now resolved (v49):** C_{ud} is constant on the discrete selector stratum $S_{8,Q}$, so the “11” is earned as the rigid Plücker readout $\|\text{Pl}(K)\|_1$ (v42) — independently of point-uniqueness. Point-uniqueness of ρ^* enters only the absolute amplitude scale, not the ratio.*

Lemma 7 (U6 — the four certificates). $C_U^{(1)}$ wall enumeration (done); $C_U^{(2)}$ D_4 -fixed character solver; $C_U^{(3)}$ selector uniqueness (one S -equivalence class); $C_U^{(4)}$ amplitude extraction — the ratios $R, Q, c_u/c_d$ are read off the (derived) selector stratum (closed, v49/v71); only the absolute U_f^*, c_q normalisation is the anchor.

Progress: an explicit valid flat bundle is realised (RH solve) ([E]). A numerical Riemann–Hilbert solve produced an *explicit* Hermitian residue A_0 (eigenvalues $\{0, \frac{1}{3}, \frac{2}{3}\}$, $\text{diag} = (\frac{1}{2}, \frac{1}{4}, \frac{1}{4})$) for which the Fuchsian system $A(z) = \sum_k U^{k-1} A_0 U^{-(k-1)} / (z - i^k)$ simultaneously satisfies: the cusp class at each puncture; the splitting $\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$ (exponents $\sum_k A_k = \text{diag}(2, 1, 1)$); the *path-ordered* monodromy at infinity is trivial, $\|M_\infty - \mathbf{1}\| \sim 10^{-9}$, i.e. $M_1 M_2 M_3 M_4 = \mathbf{1}$; and the representation is *irreducible* — **case A** (non-trivial $SU(3)_F$ mixing, not the diagonal case B). So a valid (U_{wall}) flat bundle *exists and is constructed*; the U2 kill switch lands on A. *Honest scope*: this is one realising point (existence + case A), likely one of a residual family; the unique ∇_F^* still needs the $\det R = 8$ selector and c_u/c_d still needs the $C_6 \leftrightarrow \mathbb{P}^1 \setminus \mu_4$ dictionary (H2). [[verification/v33_explicit_flat_bundle.py](#)]

H2 bridge probed ([[verification/v34_h2_bridge_attempt.py](#)]). The explicit per-puncture monodromies M_k (cusp class, $\prod M_k = \mathbf{1}$ to 10^{-5}) were computed; but the natural resolvent-dressed diagonal extraction ($|\text{diag } M_k| = (0, \frac{1}{2}, \frac{1}{2})$ times the C_6 resolvent factor) does *not* reproduce the lepton amplitudes (0.475, 1.107, 0.917). So the precise $\Gamma_{ij}^{\min}$ geodesic-to-word dictionary (how the 6-site

hypercharge hexagon combines with the 3-dim family ρ^*) was the genuine remaining analytic input — it is *not* fixed by a natural guess, and we do *not* fish for 55/117. *Resolved since (v118): the correct extraction is the resolvent determinant of the $W(A_3)$ monodromy, not the diagonal dressing — the probe's failure is thereby explained, and the dictionary is exact.*

Theorem U — the four-way split (sharper than one monster gate) [E]/[O]

D_4 fixes the admissible chamber, not the physical point (the D_4 -fixed locus is positive-dimensional, U3); the discrete selectors $\det R = 8$, $\text{SNF}(R) = (1, 1, 8)$, $\text{Spec}(Q_+) = \{1, 2, 3\}$ do the cutting. The gate splits into four pieces, three of them essentially closed:

- U_{unitary} [E]: parabolic degree 0 + polystable $\Rightarrow$ unitary (Mehta–Seshadri) $\Rightarrow \Phi = 0$; the harmonic metric is the *finite* invariant form H ($M_k^\dagger H M_k = H$), not a Hitchin PDE ([verification/v40_harmonic_metric.py]).
- U_{H_2} [E]: R, Q, K are read off the bundle (splitting type = a , $\det R = 8$, $\text{Spec}(Q_+)$) ([verification/v39_uwall_selectors.py]).
- U_{Λ^2} [E]: **Readout Rigidity** — on the discrete stratum $\mathcal{S}_{8,Q} = \{\det R = 8, \text{SNF}(R) = (1, 1, 8), \text{Spec}(Q_+) = \{1, 2, 3\}\}$ the anchor-plane readout is *constant*, $\mathcal{C}_{ud}(\rho) = g_{\text{car}} \|\text{Pl}_{1,a}(K)\|_1 / (N_{\text{fam}}^2 \Delta_Q) = \frac{55}{117}$, *independent* of the continuous D_4 position — so c_u/c_d needs no point uniqueness ([verification/v49_readout_rigidity.py], [verification/v42_exterior_leg.py]).
- U_{point} [O]: full uniqueness of ρ^* — needed only for the *complete* U_f^* amplitude matrix, *not* for c_u/c_d .

Headline: the remaining flavor bridge is not a PDE gate — it is a finite exterior *readout-rigidity* theorem; the only residual, U_{point} , is not an independent gate — it reduces to the single dimensionful anchor v_{geo} (Gate 1 below).

The simple closure ([verification/v71_simple_r_bridge.py]). The selector stratum $\mathcal{S}_{8,Q}$ is now *fully derived*: $\det R = 8 = \text{rank } E_8$ with $\text{SNF}(R) = (1, 1, 8)$ is the lattice invariant (cf. $\det Q = 3 = N_{\text{fam}}$, v70), and $\text{Spec}(Q_+) = \{1, 2, 3\}$ is the D_4 -equivariant result (v69). The four flavor operators are integer lattice maps with compiler-atom determinants ($\det Q, \det K, \det R, \det L$) = (3, 4, 8, 20), product $1920 = |W(D_5)|$. So by Readout Rigidity the quark *ratios* ($c_u/c_d = g_{\text{car}} \cdot 11 / (N_{\text{fam}}^2 \cdot 13) = \frac{55}{117}$, etc.) are *pure integer Plücker readouts* — *no transcendental monodromy solve*. The earlier monodromy machinery was over-engineering for the ratios; the only transcendental piece, U_{point} , is the *absolute* amplitude normalisation — an anchor of the same nature as the one dimensionful scale, not a missing computation.

Gate 1 complete: $U_{\text{point}} \rightarrow v_{\text{geo}}$ ([verification/v75_upto_vgeo.py]). The absolute amplitudes are not free either. The lepton amplitudes are exact ($c = (\frac{16}{7}, \frac{4}{3}, \frac{7}{6})$, v20), every cross-sector c -ratio is closed (Plücker), and each *sector product* is a clean φ_0^{ret} -power (Grand Mass Volume, v46: $\det M_{\text{sector}} \sim (\varphi_0^{\text{ret}})^{6,9,10}$; the lepton product is $\frac{16}{7} \cdot \frac{4}{3} \cdot \frac{7}{6} = \frac{32}{9} = 2^{g_{\text{car}}} / N_{\text{fam}}^2$). Since (pairwise ratios) + (product) $\Leftrightarrow$ (individuals) is a bijection, the nine charged-fermion amplitudes are *fixed up to one overall scale* v_{geo} . **So U_{point} is not an independent gate: it reduces to v_{geo} , the same one irreducible dimensionful anchor as gravity's $1/G$ (v68) — the two [O] anchors collapse to one.**

Research Contract 2 — (G_{metric})

Goal. Construct the TFPT metric measure

$$Z[J] = \lim_{\chi \rightarrow \infty} \int_{G_\chi / \text{Diff}} \exp(-S_{\text{rel},\chi}[g, A, \psi] + J \cdot O) d\mu_{\text{Cal},\chi},$$

$$S_{\text{rel},\chi} = \text{Tr} f\left(\frac{D_{\text{rel}} + A_\Sigma}{\chi}\right) - \text{Tr} f\left(\frac{D_{\text{ref}}}{\chi}\right) + \frac{i\pi}{2} \Delta\eta_\Sigma,$$

the relative spectral action whose low-curvature shadow is the already-closed $R + R^2$ equation.

Lemma 8 (G0–G1 — configuration space & BRST/Hodge Fredholm). *For $s > 5/2$ a Sobolev space of Θ -symmetric metrics on the doubled seam collar M_Σ admits a gauge slice; the gauge-fixed gravitational complex $\Omega^0(TM) \xrightarrow{K} \Gamma(S^2 T^*M) \xrightarrow{K^*} \Omega^0(TM)$ with $\Delta_{\text{gh}} = K^*K$ and $\Pi_{\text{phys}} =$*

$\Pi_{\text{BRST/Hodge}}\Pi_{\text{TT}}$ is elliptic Fredholm; the conformal mode is removed as BRST-exact / non-propagating in the Calderón polarisation (not rhetorically). Machine: principal symbols in xAct/-Cadabra; linear-algebra certificates in Lean (medium).

Lemma 9 (G2 — finite relative spectral action, *heat-kernel grounded* [E]). $S_{\text{rel},\chi}$ is finite, differentiable, with a local Seeley–DeWitt expansion $\text{Tr } f(D^2/\chi^2) \sim f_4\chi^4 a_0 + f_2\chi^2 a_2 + f_0 a_4 + \dots$. For the Lichnerowicz Dirac operator $D^2 = \nabla^2 + \frac{R}{4}$ (so $E = -\frac{R}{4}$, spinor trace $\text{tr } \mathbf{1} = 4$) the Gilkey coefficients give, per $(4\pi)^{-2}$,

$$a_2 = \frac{1}{6} \text{tr}(6E + R \mathbf{1}) = -\frac{R}{3} \quad (\text{Einstein–Hilbert } \int R), \quad a_4|_{R^2} = \frac{1}{360} \text{tr}(180E^2 + 60RE + 5R^2 \mathbf{1}) = \frac{R^2}{72}$$

(the R^2 /Starobinsky term), so the spectral action structurally produces $S_{\text{eff}} = \int \sqrt{g} \frac{\bar{M}_{\text{Pl}}^2}{2} (R + \frac{R^2}{6\bar{M}_{\text{Pl}}^2}) + \dots$ with the scalaron mass ratio fixed by the cutoff moment, $M^2/\bar{M}_{\text{Pl}}^2 = 6(4\pi)^2/f_0$. The TFPT closure $M^2/\bar{M}_{\text{Pl}}^2 = c_3^7$ fixes $f_0 = 6(4\pi)^2/c_3^7$, i.e. $M = c_3^{7/2} \bar{M}_{\text{Pl}} = 3.06 \times 10^{13} \text{ GeV}$ — the boundary normalisation c_3 is the gravitational scale — and hence the Starobinsky slow-roll forms $n_s \simeq 1 - \frac{2}{N}$, $r \simeq \frac{12}{N^2}$. The $R + R^2$ structure is convention-independent (standard Gilkey); the precise rational $\frac{1}{72}$ and factor 6 are the standard Dirac conventions used here ([verification/v36_spectral_action_g2.py], [verification/v28_gravity_fR.py]).

The simpler resolution: “full QG” for physics is the gap-decoupled admissible sector ([E]). The infinite-dimensional ambient measure (G6) is the summit, but the admissible IR sector does not wait for it. The metric coupling is gap-dominated (G5): $2\|V_{\text{metric}}\| = 0.785 < \Delta = 6 \log \frac{3}{2} = 2.433$, so the admissible sector keeps a *strictly positive* effective gap $\Delta_{\text{eff}} = 1.648 > 0$ under the stated RP and metric-coupling assumptions. A positive gap dynamically *isolates* the admissible sector from the un-built ambient/deep-UV: under those assumptions the admissible (IR) measure is closed, and the ambient completion is decoupled from it (we do *not* assert it is physically irrelevant — that would itself be a physical interpretation). So the honest reading is

admissible IR sector = closed under stated RP; ambient metric measure (G6) = G_{metric} gate
--

G2 supplies the local action ($R + R^2$, heat-kernel); G5 supplies the decoupling of the admissible sector; together they reduce (G_{metric}) to the ambient projective limit (G6). **Precise status (Alessandro’s boundary):** the admissible IR sector is closed under the stated RP and gap-dominance assumptions; the ambient metric projective measure is now discharged as a [C] *redundancy* (v369, the holographic-route keybox below), a certification object rather than a missing piece. Its absence does *not* affect the bounded IR claim. [verification/v36_spectral_action_g2.py]

The glue Q-system: the index-4 object of the G_{net} statement exists explicitly ([E]/[C]). The G_{net} target — “the seam–Calderón inclusion has Jones index $4 = |\mu_4|$ ” — no longer refers to an unknown object. On the glue group algebra $A = \mathbb{C}[\mathbb{Z}_4]$ (multiplication $\delta_a \otimes \delta_b \mapsto \delta_{a+b}$, unit δ_0) all Longo Q-system axioms hold exactly [E]: associativity, unit law, the Frobenius identity $(m \otimes 1)(1 \otimes m^*) = m^* m = (1 \otimes m)(m^* \otimes 1)$, and specialness $mm^* = |\mathbb{Z}_4| \mathbf{1}$ — a special C^* Frobenius algebra with

$\dim \theta = 4 = \mu_4 = [B : A]$

exactly the Carrier Index Lemma value (KLM: $\mu_{\text{carrier}} = 16 = 4^2 \mu_{E_8}$ re-verified). *Locality = isotropy* [E]: on the Lagrangian glue $\langle(1,1)\rangle$ the discriminant form gives $q(k(1,1)) = k^2 \in \mathbb{Z}$ for every k — the braiding obstruction vanishes (v92 in Q-system language); the two Lagrangian glues (the sheet pair) are the only maximal isotropic choices. *Halfway step* [E]: $\{0, 2\} < \mathbb{Z}_4$ closes and gives the unique intermediate Q-system $\mathbb{C}[\mathbb{Z}_2]$ = the $SO(16)_1$ tower step, with index factorisation $4 = 2 \times 2$. *R2 sharpened* [C] (recorded, not claimed): what remains is the operator-algebraic identification “the seam–Calderón inclusion of the 16-Majorana net (v113) realises *this* Q-system”

— matching two constructed objects, not a search; the three loads (metric + carrier + QBL, v123) hang on this one identification. [2026-08-03 update, the index side measured on the ladder: [verification/v726_phys_car_pp_index.py] (57/57, T2-SLICE-G0) measures the inclusion on the CAR ladder $N = 48\text{--}384$: the μ_4 -average conditional expectation has optimal Pimsner–Popa constant *exactly* $1/4$ (explicit rank-one minimizer + pinching theorem) and Watatani index *exactly* 4 via an explicit quasi-basis with *forced* weights (uniform weights break scalarness; the $\mathbb{Z}_2$ -only average gives index 2 — μ_4 distinguished from μ_2), with μ_4 *derived* from the clock $L = S^{N/12}$ (order 4 forced by the NS spin structure) and the Ramond sector healed *state-preservingly* by the Klein zero-mode dressing ($[\rho, U'] = 0$ exactly). The remaining half is unchanged in type — the *identification* of this ladder Q-system with the constructed $\mathbb{C}[\mathbb{Z}_4]$ object — now registered with kill criteria as the open half of GNET.RAMIFIED.01 (Frobenius mismatch; index $\neq 4$ on the state-preserving ladder; dressing incompatible with the σ descent); no marker move.] [2026-08-05 update, the finite precursor of the identification ([verification/v779_gnet_gns_arf.py], 27/27 + 27/27): the coherent commuting system of index-4 expectations along the Haar scaling tower (restriction coherence, $[E_I, E_J] = 0$, joint idempotency, all exact) is the finite half of the Longo Q-system identification, and the exact dim law $d_c(\ell) = 4^{\ell-1} + 2^{\ell-1} \cos(\pi c/2)$ makes the 4-element $\mathbb{C}[\mathbb{Z}_4]$ group-layer quasi-basis obstruction exactly $2^{-\ell}$ at every finite level ($d_0 - d_2 = 2^\ell > 0$), restored only asymptotically — the crossed-product structure is *exactly asymptotic*, and $d_2(1) = 0$ is the $\ell = 1$ index-3 anomaly. Typing unchanged; no marker move.] [2026-08-06 registration — GNET.MARTINGALE.LIMIT.01, [O] (the analytic remainder), finite hypotheses measured in [verification/v802_gnet_martingale.py] (15/15, MARTINGALE-HYPOTHESES-MET): the Haar ladder is an exact filtration with one index-4 Watatani expectation for the whole tower (quasi-basis explicit, 185/3041 elements at $\ell = 2/3$); the local-state defects are summable at measured rate ~ 0.25 per doubling (one power better than the $2^{-\ell}$ dim-law envelope; reported, not forced); the unique limit state follows by elementary telescoping and is verified on every rung; the split/normality precursors are uniform (determinant witness ≥ 0.985 , stabilizing at 0.9928); the decimation control converges to the wrong degenerate limit — the coherent Haar bond is load-bearing. The registered remainder (steps 5–7, typed, not executed): GNS reconstruction of ω_∞ ; local normality on each $A(W)$ (the uniform S_4 witnesses as the finite input); split property and the μ_4 simple-current extension (GATE.METRIC.11) on the limit net. The open QGEO majorant residues (b)/(c) are bypassed for limit existence and relocated into the normality step — a reallocation, not a proof. Kill: non-summable defects at a deeper reachable rung, or a determinant-witness collapse. No marker move.]

The hull is the graded module of the Q-system ([verification/v128_graded_hull.py]). The Q-system is not merely an abstract companion of E_8 — E_8 *realises* it [E]: the 240 roots, built as explicit norm-2 vectors in $D_5 \oplus A_3$ +glue coordinates, decompose as $240 = 52 + 64 + 60 + 64$ over the glue $\mathbb{Z}_4$ (the v89 branching as vectors), and for *all* 6720 root pairs whose sum is again a root, $\text{coset}(r_1+r_2) = \text{coset}(r_1) + \text{coset}(r_2) \bmod 4$:

E_8 is a $\mathbb{Z}_4$ -graded Lie algebra over the carrier; the grading group is the glue = the Q-system

— the checksum hull is the Q-system’s graded module, realised in 248 dimensions. The R2 identification thus matches the Calderón inclusion against a structure that E_8 already *carries*. [verification/v125_glue_qsystem.py]

Graded Frobenius ([verification/v143_graded_frobenius.py]). The identification holds at the finite Lie level: the graded module is in fact a graded *Frobenius* structure, i.e. everything the Q-system demands of its module is already exact inside E_8 [E]: (i) *coset duality* — $\text{coset}(-r) = -\text{coset}(r) \bmod 4$ for all 240 roots ($-C_1 = C_3$ bijectively, C_0, C_2 self-dual), so the Killing form restricts to a *nondegenerate* invariant pairing $g_k \times g_{-k} \rightarrow \mathbb{C}$; (ii) the glue character average $\frac{1}{4} \sum_k i^{k \deg}$ is *exactly* the carrier projector (the conditional-expectation shadow, quasi-basis index $|\mathbb{Z}_4| = 4$); (iii) each glue sector is *one* $W(D_5) \times W(A_3)$ orbit with multiplicity-one dimensions $64 = 16 \cdot 4 = (\mathbf{16}, \bar{\mathbf{4}})$, $60 = 10 \cdot 6 = (\mathbf{10}, \mathbf{6})$ — simple currents, sector ring $= \mathbb{Z}_4$ = the $\mathbb{C}[\mathbb{Z}_4]$ fusion, halfway subgroup $\{0, 2\} = SO(16)_1$. R2 residue [C] (final form): exactly *one* conformal-net statement remains — the seam–Calderón inclusion of the 16-Majorana *net* carries this Q-system — with no remaining

finite freedom.

Where that statement lives: the NS/R sector census (`[verification/v148_fock_census.py]`). The remaining work is scoped exactly. Every *untwisted* (Neveu–Schwarz) state of the 16 Majoranas has *integer* ($D_5, SO(6)$) weights, and integer weights lie in the even discriminant classes — so the NS module supports only $\{(0, 0), (2, 0), (0, 2), (2, 2)\}$: **the odd glue classes $k(1, 1)$ — the actual μ_4 simple currents — have zero NS support; they are twisted (Ramond-type) modules.** The Ramond zero-mode module ($2^8 = 256$ ground states) carries exactly the four odd class pairs, 64 each, and splits $128 + 128$ into the odd sectors of the *two* Lagrangian glues $\langle(1, 1)\rangle$ and $\langle(1, 3)\rangle$ (v92): *choosing the glue is choosing the Ramond projection* ($128 =$ one $SO(16)$ spinor chirality), and E_8 assembles as $248 = 120_{\text{NS}} + 128_{\text{R}}$. *Consequence [C]*: the index-4 extension cannot even be *stated* on the untwisted module alone; R2’s one remaining statement is irreducibly about the twisted-sector structure of the seam–Calderón net, with the sheet choice realised as the R-projection; the finite shadow is exhausted (v113/v125/v143 + this census). [2026-08-03 update, the NS/R grading acquires an arithmetic home: `[verification/v722_phys_ramified_ns_r.py]` (9/9, RAMIFIED-SEES-NSR) shows the NS/R grading *is* the parity character of $E_8(\mathbb{Z}[i])/(1+i) = \mathbb{F}_2^4$: $(1+i)L$ lies in the even/NS sublattice (proved on a $\mathbb{Z}$ -basis), the 240 roots fall into 15×16 classes of *pure* parity (7 NS + 8 R), and parity is a homomorphism $\mathbb{F}_2^4 \rightarrow \mathbb{F}_2$ — the Ramond projection is $(1+i)$ -adic, at the one ramified edge of the $\mathbb{Z}[i]$ - E_8 Hecke tower (both controls fire: the $\mathbb{Z}[i]^4$ census is not 15×16 , scrambled labels break purity). Together with the ladder index measurement (`[verification/v726_phys_car_pp_index.py]`; $\chi = 1 \leftrightarrow$ half-integer labels on both sides, the ramified defect in $\ker \chi$) this gives the groupoid reading of G_{net} its first *exact* witnesses; the gate typing is unchanged.]

Hecke from geometry on the glue census (HECKE.GEOM.01). The μ_4 -refined E_8 class thetas that underlie the glue Q-system now carry an *in-suite* Hecke package (consolidated from discovery T27/T31/T32; 25 checks, ~ 11 s). (A) Kneser p -neighbours realise $\#_{\text{iso_lines}} = \sigma_3(p) \# \mathbb{P}^3(\mathbb{F}_p)$ identically and at $p = 2, 3, 5, 7$ ($135/1120/19656/137600$). (B) The marked neighbour-sum is the affine Hecke element $\nu_p = a \text{Id} + b T_p$ with cusp rule $a_p = b - \sigma_3$, recovering $a_3 = -4$, $a_5 = -2$ ($p = 7$ consistency $\lambda_{\text{odd}} = 19840$). (C) The census form space has $\dim_{\mathbb{Q}} V = 7 = 5+2$ as the 2-adic oldform hull of E_4 and f_8 , with recovery $\pi_{\text{cusp}} = (28 - T_3)/32$ and newform via $(1 - V_2 U_2)$. Classical theorems named classical; the claim is the compiler identification (correspondence $\rightarrow$ operator $\rightarrow a_p$; redundancy $= \mu_4/2$ -adic). Fence: both new systems stay weight 4 — $GL(1)$ /weight-drop transport remains [O]; no RH claim; no marker moves. Machine-checked: `[verification/v535_hecke_from_geometry.py]`. [E] (mechanics) / [O] (weight drop)

Eichler trace layer at the E_8 lattice (HECKE.GEOM.EICHLER.01). Consolidated promotion of discovery T33/T36/T37/T42 (23 checks, ~ 30 s; builds on HECKE.GEOM.01). Witt $\lambda_{\text{Eis}} = \sigma_3(\# \mathbb{P}^3 - \sigma_3)$ closed in p ; Eichler $\lambda_{\text{geom}} = \lambda_{\text{Eis}} + a_p^2$ two-sided at $p = 3$ (live type-(i)/(ii): $352 = 364 - 12$) and $p = 5$ (T36 freeze: $3784 = 3906 - 122$) with assembler $R = \sigma_3 - 1 - c(p^2)/8$; Type-A/B densities $N_A = \min(240(1 + p^3), \#_{\text{iso}} - 1)$ ($p = 7$ live $82560/743040$; injectivity $p = 11, 13$); signed $a_p = -c(p)/8 \Rightarrow (-4, -2, +24)$ and $b \in \{24, 124, 368\}$; $O(1)$ assembler for odd $p \leq 31$. Classical scaffolding named classical; claim = in-suite two-sidedness + closed compiler densities. Fences: $\text{Shell}(p^2)$ for $p \geq 7$ out of scope; fibre refinement scoped to $\{5, 7, 11\}$; weight-4 $\rightarrow GL(1)$ stays [O]; no RH; no marker moves. Machine-checked: `[verification/v536_eichler_trace_layer.py]`. [E] (mechanics) / [O] (weight drop)

Half-integral bridge (HECKE.GEOM.HALFINT.01). Consolidated promotion of discovery T38/T41/T44/T45 (20 checks, ~ 90 s; companion to HECKE.GEOM.01/HECKE.GEOM.EICHLER.01). Unique signed Shimura preimage $\text{Sh}_{t=2, \psi=1}(g) = -8 f_8$ in the 70-element compiler weight-5/2 theta-monoid (witness $g = \vartheta_2(q^2)^2 \vartheta_3(q^2) \vartheta_4(q^2)$; three further monoid elements lift to scales $+8, -16, +16$); exact $T(p^2)$ -eigenform with $\lambda = a_p(f_8)$ for $p = 3, 5, 7, 11, 13$; $U_4(g) = 0$ and level $32 = 4 \cdot 8$ outside Kohnen 1982 (plus-route structurally closed — negative result part of the claim); AFE-based Waldspurger/Baruch–Mao quotient $R(d) = b(|d|)^2 / (|d|^{3/2} L(f_8 \times \chi_d, 2))$ constant on ten fundamental $d \equiv 1 \pmod{8}$ ($R = 23.1873585645\dots$, spread $\leq 10^{-12}$, including $d = 89$), with $d \equiv 5 \pmod{8}$ vanishing via root number -1 . Classical Shimura/Kohnen/Waldspurger/Baruch–Mao

named classical; claim = unique monoid preimage + in-suite chain + *measured* R . Fences: GL(2) centre $s = 2$ (not the ξ -line); no RH; ZETA.HP.CARRIER untouched; no marker moves. Machine-checked: [verification/v537_halfintegral_bridge.py]. [E] (Shimura/Hecke/Kohnen fence) / [C] (measured R)

Compiler relative-trace identity (HECKE.GEOM.RTF.01). Promotion of discovery T53 (18 checks, ~ 14 s): HECKE.GEOM.01/HECKE.GEOM.EICHLER.01/HECKE.GEOM.HALFINT.01 are three projections of one finite relative-trace identity [E_8 /glue orbital counting] = [Newform/Waldspurger spectral sum]. (R1) $\text{Tr}_V(\nu_p \circ \pi)$ bilateral independent at $p \in \{3, 5, 7\}$ across five projectors (anchors e.g. $p=7$: 727680 full / 19840 on f_8). (R2) Eichler $\lambda_{\text{geom}} = \lambda_{\text{Eis}} + a_p^2$ is the RTF orbit split ($\lambda_{\text{Eis}} = 336/3780/19264$, $a_p^2 = 16/4/576$). (R3) Period side = quaternary lattice counting with $R = 23.1873585645$ (rel $\sim 10^{-12}$). Verdict ONE-FORMULA. Classical Jacquet-type RTF / Eichler / Witt / Waldspurger named classical; NEW = compiler-native bilateral independence. Fences: identity *finite*; infinite RTF named open; no RH; not ZETA.HP.CARRIER-relevant; no marker moves. Machine-checked: [verification/v538_relative_trace_identity.py]. [E] (R1/R2) / [C] (R3)

Weil structure of the compiler family (RTF.GNS.WEIL.01). Promotion of discovery T55/T61/T62/T63 (25 checks, ~ 6 s): the compiler family possesses a fully identified Weil structure up to two *explicitly isolated* obstructions (obstructions are load-bearing verified content, not residual footnotes). (A) GNS = direct integral over the Gelfand spectrum of the character subalgebra (fibre-orthogonal; twist-mix per fibre $\leq 5.037 \times 10^{-16}$). (B) Trivial Sato–Tate isotype = GL(1) core $G_0 = (1 + Y)/(1 - Y) = \zeta_p(w - 3)^2/\zeta_p(2w - 6) = \sum 2^{\omega(n)} n^{-u}$ globally. (C) Exact linear relation $Q_{\text{fam}} = 2Q_\zeta(g) - 2Q_\zeta(g_b) + \text{Arch} + \text{Corr}$. **Obstruction 1:** minus sign on the doubling term (family positivity does *not* imply $Q_\zeta \geq 0$). **Obstruction 2:** Corr is non-automorphic (no polar/arch absorption; no finite Euler factor; $e^{-\sum p^{-u}}$ -type). Finite-class $Q_{\text{fam}} \geq 0$ measured [C]; dense-class positivity open / RH-adjacent — *not* claimed. Classical Weil/Gelfand/SU(2)/Sato–Tate/ 2^ω /GNS named classical. Fences: not “almost RH”; ZETA.HP.CARRIER untouched; no marker moves. Machine-checked: [verification/v539_weil_structure_family.py]. [E] (A/B/C/obstructions) / [C] (finite-class positivity)

Amplitude route and positive linear carrier (RTF.GNS.AMP.01). Promotion of discovery T67–T72 (34 checks, ~ 3 s): the route out of the square plane behind the two RTF.GNS.WEIL.01 obstructions. (A) Amplitude Dirac $D = \begin{pmatrix} 0 & V \\ V^\top & 0 \end{pmatrix}$ with $D^2 = \text{diag}(VV^\top, K)$, $V^\top V = K$ (rel 2.7×10^{-16}), exists exactly and is Hecke-equivariant ($p = 3, 5, 7$ intertwining exact; Shimura $b(dp^2) = b(d)(a_p - \chi_d(p)p)$ on 644 pairs). (B) Geometric polarisation $b = N_+ - N_-$ exact; $\Theta = N_+ + N_-$ is a *pure* Siegel–Weil Eisenstein σ_3 -eigenform ($p \leq 13$, cusp component zero) with Cohen seed $\Theta(d) = -48 L(-1, \chi_d)$ exact-rational on 159 live d (Cohen 1975). (C) Every coefficient bilinear form inherits the even- k deletion (Cauchy–Littlewood window lemma, five channels; numerator $1 - p^6 X^2$ pair-independent); the deletion is exactly the square-class double counting (fam + $2b + \delta_{k1} = [2, 2, 2, 2]$; Möbius square sieve exact). (D) Positive linear carrier $\ell^2(d, \mu)$, $\mu = 48 |L(-1, \chi_d)| |d|^{-a}$: full weights $\lambda_k = 1 + p^{3k} - (\chi p)^k$ (layer $[1, 1, 1, 1]$) and plus balance $Q = Q_\zeta(g_-) + Q_\zeta(g_+)$ (rel 2×10^{-15}); the b /doubling kernel enters *with plus* (sign inverted vs the square plane). (E) Completed FE $\Lambda_\Theta(s) = 8^{1-s} \Lambda_{\Theta^\dagger}(5/2 - s)$ exact (rel $\sim 10^{-40}$; Fricke closed); the guaranteed cone is FE-self-dual. **Open boundary as content:** Euler-region positivity only (edge L -values, not the central line); the residual distance to the Weil cone is the FE-covariant gap functional λ^* on $n \equiv 6 \pmod{8}$ (one exact Farkas witness; no finite signed theta library removes it). Classical Cohen 1975 / Siegel–Weil / Jacobi–Fricke / Cauchy–Littlewood / Shintani / Weil 1952 / Farkas named classical. Fences: not “almost RH”; ZETA.HP.CARRIER untouched; no marker moves. Machine-checked: [verification/v540_amplitude_linear_carrier.py]. [E] (A–E exact) / [C] (sampled cone facts; λ^* named open)

Matching lemma and transport ledger package (RTF.GNS.LEDGER.01). Promotion of the discovery proof package T78/T79/T80/T81/T82/T85 (33 checks, ~ 10 s; every core check recomputed, not cited). (A) The matching lemma is *proved exact-integer* on $[4, 10^6]$: $7S(j) < 40A(j)$

at every atom (0 violations over 939 870 clash atoms; proven no-overflow; big-integer rechecks); exact margin $X = 22242575693/270725400000 = 0.082159$ at $j^* = 554400$; $\rho_{\text{crit}} = 1.1440 > 21/20$; maximal-greedy identity $\Rightarrow$ uniform in (M, λ) ; structure laws (8-law, seed-tower, w -table, Cohen seeds $\{-48, -80, -8, -8\}$, bracket $\Theta \leq 2|\psi| \leq 3\Theta$) at 0 tolerance. (B) The transport ledger closes exactly: $Q_{\text{Weil}} = Q_{\text{cert}} + \Delta_{\text{arch}} + \Delta_2$ with $\Delta_{\text{pole}} \equiv \Delta_{\text{conv}} \equiv 0$ *proven* (kernel collapse sympy-exact; residual 4.4×10^{-16} on 24 rows); the odd-prime side of the Weil functional equals the certified plus combination $P_{\zeta}(g_-) + P_{\zeta}(g_+)$ exactly (rel 3.9×10^{-16}). (C) The signed envelope is character-exact: $-\psi = (\chi_{-4} + \frac{1}{4}\chi_8 + \frac{1}{4}\chi_{-8}) \cdot \Theta$ (all odd $n \leq 50000$, both builds); signed certificate 0 violations (margin $8.86\times$); confinement set equality zero-credit $\Leftrightarrow$ coherent on 10^6 . (D) The archimedean term is *internal*: Legendre duplication $(2\pi)^{-s}\Gamma(s) = \frac{1}{2}\Gamma_{\mathbb{R}}(s)\Gamma_{\mathbb{R}}(s+1)$; kernel identity $k_{\zeta} = K_{\text{fam}} - K_{\text{shift}}$ pointwise (7.9×10^{-31}); $\Delta_{\text{arch}} = A_{\text{fam}} - A_{\text{shift}}$ at 9.9×10^{-16} . (E) The coherent class ($= \mathbb{Z}[i]$ -norms, set equality) is closed by the λ -equivariant channel: CM carrier g_{λ} exact in two independent routes; $\mu_1 = c_1/(c_0 d^2) \in [-1, 1]$ exact; λ -window certificate 0.0653 vs 0.2365 unlifted ($3.6\times$), exact Fraction rechecks. (F) Frontier facts [C]: coherent-chain crossing $k^* = 14$ ($\log_{10} N^* = 23.1$, exact Fractions); Robin tail factor 6.16 — the constant route stays open. **Named limits as load-bearing content:** (i) *one* classically-shaped open lemma (correlated cancellation on credit-rich non-coherent tail atoms, uniform form; provably-shaped $\neq$ formal proof); (ii) I5 in one-family form ($Q_{\text{cert}} + \Delta_2 + A_{\text{fam}} - A_{\text{shift}} \geq 0$) as the single remaining TFPT-specific object — by the closed ledger *equivalent* to Weil positivity $\Leftrightarrow$ RH (equivalence typing, no progress claim). Classical Gronwall / Robin 1983 unconditional / Cohen 1975 / Hecke 1918–1920 / Cornacchia / Dirichlet / Mertens-AP / Landau / Legendre duplication / Weil 1952 / Alaoglu–Erdős named classical. Fences: no RH-evidence language; not “almost RH”; ZETA.HP.CARRIER untouched; no marker moves. Machine-checked: [verification/v541_matching_lemma_ledger.py]. [E] (A–E exact) / [C] (frontier facts; two named limits open)

The margin-chain identities of phase 2 (PRIME.MARGIN.IDENT.01). Promotion of the *identity/theorem block* of the discovery parts T128–T131 (44 checks, ~ 0.2 s; every check recomputed on small frame-A windows). Scope fence, declared with the row: identities and *per-instance* theorems only — no fit, no graded floor, no Lanczos value, no uniform-in-zone statement. Battery: 24 seam instances ($n = 3 \dots 97$, $\rho = 1.25 \dots 4.00$; every inverted or diagonalised matrix ≤ 300), 43 graded/fine grid pairs and 400 randomised profiles; every identity is a residual against a *preregistered* tolerance plus a mutation control that must fail by $\geq 10^{-3}$ relative. (1)–(2) $\sum_i (1 - g_i) = t_l + t_r$ exactly (5.5×10^{-13}) $\Rightarrow$ the κ weights are a probability vector and $\kappa_{\text{flat}} = 1$; the (T) four-term identity $\tau_{\text{dn}} = \|y\|^2 - m_{\text{prot}} + m_{\text{fill}} - V_{\text{bord}}$ at 2.2×10^{-16} (two independent routes for τ). (3)–(4) $p_{N-1} = 2 - p_0 + 2E$ at 2.3×10^{-16} $\Rightarrow$ the equivalence $2E > p_0 \Leftrightarrow p_{N-1} > 2$ (0 exceptions, both branches realised); the Abel/Hölder chain $\kappa \leq 2 - p_0 + \frac{N-2}{N} \sum_j |\Delta p_j - \overline{\Delta p}| + (p_{\text{max}} - p_{N-1})$ per instance (0 violations; the same chain *without* the curvature term is violated on 88/400 profiles). (5)–(6) Matrix-free two-scale assembly $= J^T Q J$ (5.2×10^{-16}), graded overlap $= J$; the C ea/Strang defect identity $S_{\text{graded}} - S_{\text{uniform}} = R^T X^{-1} R$ as a *matrix* identity with PSD defect on all 43 grids (2.1×10^{-14}) $\Rightarrow$ the compression error is one-sided by construction. (7)–(9) The secular sandwich $\varepsilon/(\|u\|^2 + \varepsilon/\mu_1) \leq \lambda_{\min}(Q) \leq \varepsilon/\|u\|^2$ on all 48 sections where $A > 0$ and $\varepsilon > 0$ are *checked* (width ≤ 1.4448) with Albert’s sign equivalence in *both* directions; $S^{-1} > 0$ entrywise (measured 48/48) $\Rightarrow$ sign-constant and simple ground vector (Perron–Frobenius), with a control where hypothesis and conclusion fail together; and $n_{\text{run}} \leq n_{\text{cross}} + 1 \leq s_2 + 2$ with $\sum_j |\Delta p_j - \overline{\Delta p}| = \text{TV}(P)$, 0 exceptions on a non-vacuous battery (s_2 up to 31). **Named limits as load-bearing content:** (i) per instance on *small* windows — nothing uniform in the zone index; (ii) the κ law is *not* claimed (T129 found it false in general), only the chain underneath it; (iii) positivity of the pole-free section at depth (the open M25d) is neither assumed nor extended and no floor value is carried; (iv) the Perron step is an *implication* with a measured hypothesis; (v) the defect identity says nothing about the defect’s *size* (the open T130 exponent). Classical Abel / Hölder / Schur–Haynsworth 1968 / Albert 1969 / C ea 1964–Strang / Yserentant 1986 / Perron–Frobenius–Ostrowski 1937 named classical; Weil 1952 cited, never used as a criterion. Fences: zero-firewall AST-checked; no RH-evidence language; not “almost RH”; ZETA.HP.CARRIER untouched; no marker moves. Machine-checked:

[verification/v542_margin_chain_identities.py]. [E] (per-instance identities; nothing uniform)

The lumped M -matrix pair of phase 2 (PRIME.MMATRIX.IDENT.01). Promotion of the *identity/theorem block* of the discovery part T136 (35 checks, ~ 1 s; every check recomputed on small frame-A windows). Scope fence, declared with the row: matrix identities, per-instance theorems with *checked* hypotheses, and one *negative* result — no fit, no D -exponent, no uniform-in-zone statement, no floor value. Battery: 20 seam windows ($n = 4 \dots 59$, $\rho = 1.25 \dots 4.00$, $h = 25 \dots 299$; every factorised or diagonalised matrix ≤ 300) giving 39 matrices with positive off-diagonals, 312 shifted ladder rungs, 8 whitened two-grid rungs; every statement is a residual against a preregistered tolerance plus a mutation control that must fail by $\geq 10^{-3}$ relative. (1)–(2) With $L_\Delta = \text{diag}(\Delta \mathbf{1}) - \Delta$ the Laplacian of the positive off-diagonal part, $A_B = A + L_\Delta$ is Stieltjes, row-sum preserving (1.6×10^{-15}) and $A_B \succeq A$ — an *upper* bound on $\lambda_{\min}(A)$, a floor never; the congruence $A = A_B^{1/2}(I - W)A_B^{1/2}$, $W = A_B^{-1/2}L_\Delta A_B^{-1/2} \succeq 0$, is a matrix identity (4.4×10^{-15}) with $\lambda_{\max}(W) = \rho(L_\Delta A_B^{-1})$, and the *declared* equivalence $\lambda_{\max}(W) < 1 \iff A \succ 0$ is checked in *both* directions — the Loewner sandwich is circular by itself, which is the fence. (3)–(4) With $x = A_B^{-1} \mathbf{1} \geq 0$ and $A_B^{-1} \geq 0$ entrywise (Ostrowski 1937, *checked*), $\lambda_{\min}(A_B) \geq 1/\max_r x_r$ on all 39 rows (min slack 3.0×10^{-3} , sharpest ratio 0.9070) — one solve, one sign check, no eigenvalue; $(A - sI)_B = A_B - sI$ (2.5×10^{-16}) and $(A_B - sI)^{-1} \geq A_B^{-1} \geq 0$ entrywise (Berman–Plemmons 1979) $\Rightarrow$ the shifted ladder is *structurally empty* (0 exceptions of 312). (5)–(6) For the regular splitting $A_B = D_0 - N_0$ (all three hypotheses checked), Varga’s $\rho(J) = \tau/(1 + \tau)$ holds at 4.8×10^{-15} ($\rho(J) = 0.7347\text{--}0.9860$), and the Collatz–Wielandt bracket at the anchor vector gives an *a-priori* $\rho(J) \leq 0.9861 < 1$ with no eigenvalue anywhere, sharp on the measured gap by 1.0062–1.0531. (7) The k -scanned M18 majorant is a *valid* bound at every k (39 (rung, k) pairs, 0 violations) and does *not* reach the preregistered bar 1/2 on any rung (best 3.080) while the measured bad mass is ≤ 0.0120 : a *negative* result, the localisation term alone exceeding the bar (0.542), and the bar is not moved. **Named limits as load-bearing content:** per instance on small windows, nothing uniform in the zone index, in D or in the gap; the T136 D -exponents (anchor $D^{-0.56}$, gap $D^{2.72}$, slack $D^{0.12}$) are fits and stay in the sandbox; (2) carries no floor and M25d stays open; (3) and (5) are implications with hypotheses checked at the instance; (4) kills a route and says nothing about the true $\lambda_{\max}(W)$. Classical Stieltjes / Ostrowski 1937 / Fan 1958 / Berman–Plemmons 1979 / Varga 1962 / Collatz 1942–Wielandt 1950 / Haynsworth 1968 named classical; Weil 1952 cited, never used as a criterion. Fences: zero-firewall AST-checked; not “almost RH”; ZETA.HP.CARRIER untouched; no marker moves. Machine-checked: [verification/v543_lumped_pair_identities.py]. [E] (per-instance identities) + [X] (one closed route)

The long-lag support structure of phase 2 (PRIME.LONGLAG.SUPP.01). Promotion of the *structure block* of the discovery part T137 (24 checks, ~ 0.2 s) on the same 20 windows (3–25 prime-power atoms, 20555 positive off-diagonal edges, 455 loaded stripes, 39 bracketed lumped comparisons). Scope fence: support and amplitude structure, one two-sided bracket, and one *death certificate* for a whole family of bounds — no fit, no decay exponent, and *no* structure bound that closes. (1)–(3) The kernel splits exactly as $c = c_{\text{arch}} - c_{\text{comb}}$ with $c_{\text{comb}} \geq 0$, and the archimedean section alone has *all* off-diagonals ≤ 0 on 20/20 windows $\Rightarrow$ every positive off-diagonal of A is comb-generated (control: the full section carries $+2.0 \times 10^{-1}$); all 20555 positive edges have their Hankel index $M - 1 - r - t$ inside the atom-hat index set (per-window fraction exactly 1.000000) $\Rightarrow \text{supp}(\Delta)$ is a union of anti-diagonal stripes at prime-power atom positions (converse reported, *not* claimed); each stripe is a perfect matching (0 endpoint collisions), so its edge vectors are exactly orthogonal and L_Δ per stripe is a direct sum of 2×2 blocks. (4) $\Delta_{rt} \leq c_{\text{comb}}[M - 1 - r - t]$ on all 20555 edges (excess -1.9×10^{-6} , sharpest ratio 0.99998532) — no eigenvalue, no norm; the Toeplitz lag in place of the Hankel index fails on 20415/20555. The tighter one-atom refinement is *honestly undecided* at $h \leq 300$ (the atom hats never overlap) and is not promoted. (5) For the lumped comparison $B = D_0 - N_0$ the anchor identity $q_J = 1 - \min_r 1/(d_r x_r) < 1$ (8.2×10^{-16}) needs no measurement; the Neumann partial sum to $K = 8$ is then an *entrywise* lower bound for B^{-1} *because the discarded terms are nonnegative*, and partial sum plus anchor-weighted tail an entrywise upper bound, on 39 rows. (6) Since Gershgorin, every row-sum norm and every Collatz–Wielandt quotient

at every *positive* weight are upper bounds for $\rho(|E|)$, $E = B^{-1}L_\Delta$, one rigorous *lower* bound decides the family at once: $\rho(|E|) \geq 1.3141 \geq 1$ on 20/20 rows (cheapest four members ≥ 1.5937), so no member can *ever* certify $\lambda_{\max}(W) < 1$ — while the *signed* $\rho(E) = \lambda_{\max}(W) \leq 0.99995781$ on the same rows: the absolute value alone destroys the cancellation. **Named limits as load-bearing content:** per instance on small windows, no decay profile and no amplitude law (T137’s decay measurements are fits); (2) is one inclusion; (4) bounds the amplitude and *not* $\lambda_{\max}(W)$, and the structure bound assembled from (2)–(4) does *not* beat the margin (T137: 0 of 35 windows) and is not promoted; (5) brackets the *lumped* comparison, not the target, whose entrywise inverse positivity at depth stays open; (6) closes a route, not the question. Classical Gershgorin 1931 / the Neumann series / Haynsworth 1968 named classical; Demko–Moss–Smith 1984 cited as a model and *not* applied (the comparison is dense); Weil 1952 cited, never used as a criterion. Fences: zero-firewall AST-checked; not “almost RH”; ZETA.HP.CARRIER untouched; no marker moves. Machine-checked: [verification/v544_long_lag_support.py]. [E] (per-instance structure) + [X] (one family closed)

PRIME.HARDY.IDENT.01 — the Hardy-core identities of phase 2 (T140, T141). (1) The form lifting $\text{Gram} = CHC^\top$ with $C = \text{diag}(\sqrt{\Delta})M$ (3.0×10^{-15}), hence $\text{rank}(\text{Gram}) \leq h - 1$. (2) The exact finite-core reduction: the closed geometric covering kernel $K_{rs} = W([r \wedge s, r \vee s])$ equals $M^\top \Delta M$ (1.3×10^{-15}) and $\rho(W) = \lambda_{\max}(K^{1/2}HK^{1/2}) = 1 - \lambda_{\min}(A, A_B) = \lambda_{\max}(\text{Gram})$ on three independent routes. (3) $H = \text{diag}(s) + L_N$ (4.7×10^{-16}). (4) The checkerboard split $\text{Band}_b = D + A_{\text{even}} + A_{\text{odd}}$ entrywise exact with a three-step Weyl sequence. (5) The four Hardy identities — the K^+ Rayleigh form, $L_N = D^\top BD$ with the signed crossing kernel, $Q = B_+ \mathbf{1}$ and $DKD^\top = L_\Delta$ on the interior nodes. (6) The closed Muckenhoupt quantity $A_{M0} = \max_k Q_k(\Delta \mathbf{1})_{k+1}$ by two routes. (7) The additive Hardy chain and the joint shape $\Lambda(H, Y) \cdot \Omega(Y)$ as *valid* upper bounds on 11/11 windows — dominance only. (8) As a negative typing, the sign-versus-absolute separation of the measured spreads ($1.36\times$ against $\geq 54.94\times$). **Named limits as load-bearing content:** per instance on small windows, nothing uniform in the zone index or in D ; every exponent is a fit; (7) promotes validity and not size — neither shape beats the target, the additive shape is dead as a *shape* by its own exact Weyl floor and the joint shape is decided at the normalisation Ω ; (2) is an identity and not a bound, so the zone-uniform discrete Hardy inequality stays *open*. Fences: Muckenhoupt 1972 / Bradley 1978 / Opic–Kufner 1990 / Gantmacher–Krein 1950–1960 named classical; Weil 1952 cited, never used as a criterion; zero-firewall AST-checked; no marker moves. Machine-checked: [verification/v545_hardy_core_identities.py]. [E] (per-instance identities) + [X] (one route typed)

PRIME.CAPCHAIN.IDENT.01 — the capacity-chain identities of phase 2 (T142–T145). (1) The capacity decomposition $K^{-1} = D^\top J^{-1}D + xx^\top/\text{cap}$ with $J = DKD^\top$ (entrywise, 3.8×10^{-15}), $x = K^{-1}\mathbf{1}$, $\text{cap} = \mathbf{1}^\top K^{-1}\mathbf{1}$, as a matrix identity (1.4×10^{-13}) whose Dirichlet half is an *orthogonal projection* under the congruence — $\Omega = 1$ exactly (5.1×10^{-12}). (2) The exact capacity–Rayleigh identity for $1 - \rho(W)$: numerator/denominator assembly equals the direct quotient for every v (9.5×10^{-14}), eigenvalue reproduced under the declared 1/gap conditioning, inf property verified, the constant vector carried entirely by the capacity term. (3) Cholesky nestedness: $\text{cap}_E([a, a+j])$ as a prefix sum of squared forward-substitution entries of one triangular solve (3404 interval capacities at 2.0×10^{-14} , exhaustive on the median window; the reversed ordering fails at $\geq 4.6 \times 10^{-1}$). (4) The whitening certificate $\lambda_{\max}(W) \leq \kappa_{\text{up}} = 1.2245\text{--}1.4236 \leq \text{Gershgorin } 2.1213$, direction-correct, refusing at the shrunken shift. (5) The T145 M4 split: the pointwise mass-half theorem ($\sum_k f_k^2 \leq \psi^2$ exact), $\sigma_{\text{tot}} = 0.2197\text{--}0.4425 < 1$ per window, licence 4 with $|R|$ and not R^+ (witness $[[1, -\frac{1}{2}], [-\frac{1}{2}, 1]]$, $3 > 2$, built in), the ordered sign-free chain, and $S1'$ *certified per window on both routes* (better route $c_0 = 2.5154\text{--}4.2265$ against Maz’ya’s Dirichlet value 4), with Charikar’s cited density constant bracketed by exhaustive enumeration. **Named limits as load-bearing content:** per instance on small windows, nothing uniform in the zone index or in D ; the c_0 of (5) is *measured* at the minimiser’s own layer cake and the a-priori level lemma (T145’s L1) stays open, unclaimed; Maz’ya’s strong-type half is not used as a theorem anywhere — it is the thing the cycle transcribes. Fences: Maz’ya 1985 / Muckenhoupt 1972 / Fukushima–Ōshima–Takeda 1994 / Miclo 1999 / Charikar 2000 / Goldberg 1984 / Gershgorin 1931 named classical; Weil

1952 cited, never used as a criterion; zero-firewall AST-checked; no marker moves. Machine-checked: [verification/v546_capacity_chain_identities.py]. [E] (per-instance identities and certified inequalities)

PRIME.LEVEL.LEMMA.01 — the level-lemma identities of phase 2 (T146). (1) The union identity for nested cakes: $\sum_{j,l} c_j c_l |S_j \cup S_l| = 2T \|\psi_t\|_1 - \|\psi_t\|^2$ with $T = \sum_j c_j$ and $\psi_t = \sum_j c_j \mathbf{1}_{S_j}$ — form-independent and $O(m)$, exact (0.0) against the $K \times K$ double sum and against union sizes computed with no nestedness used (controls: one shuffled level set 3.2×10^{-2} , a 1% coefficient perturbation 4.8×10^{-3}). (2) The layer-cake counting lemma at a general base: $G(b) \leq 2b^2 \|v\|_\infty \|v\|_1 / \|v\|^2 + \varepsilon$ for *any* vector at all 6 preregistered bases; the classical dyadic 8 at $b = 2$ exactly, the base free (gain $4/b_*^2 = 3.7612$; the one-level-down cake breaks the domination on every draw). (3) The resolvent delocalisation bound: $\psi = \lambda R \psi$ (9.0×10^{-14} , an identity — no Davis–Kahan transfer anywhere) $\Rightarrow \|\psi\|_\infty \leq \lambda_{\text{up}} \max_j \|Re_j\|$ with Rayleigh–Ritz λ_{up} (1.0592–1.2883) and residual-paid certified column norms, within 1.061–1.337 of the truth; the lever: the certified columns sit 2.85–10.17 below the trivial ceiling $1/\lambda_{\text{min}}$, giving $\Gamma = 1.8356$ – 2.2797 against the useless $\sqrt{m}$. (4) The composite level lemma: $c_0^{\text{ap}} = 2b^2 \Gamma \min(1, \Gamma_1) + \varepsilon = 3.9042$ – 4.8488 (slack 1.986–2.418 over the measured constant, under the ceiling $16 = 4 \times \text{Maz'ya's Dirichlet } 4$), the certified window inequality $\text{cert_lam_max}(R) \leq c_0^{\text{ap}} \Psi_{\text{abs}}$ on 12/12 windows, chain loss 0.0423–0.1272 of the exact gap; the built-in no-go discriminator ($R = aa^\top + \varepsilon I$, $a_i = i^{-1/2}$: Γ grows $3.676 \rightarrow 6.798$, ratio $1.85 \geq 1.5$, $c_0^{\text{ap}} = 11.10$ exceeds every window value) fails exactly where it must, the theorem half surviving on every no-go size and the Dirichlet path-Laplacian control flat (1.0063). **Named limits as load-bearing content:** per instance on small windows — a finite list of certified window inequalities with an explicit maximum; nothing uniform in the zone index or in D , and explicitly *no* statement for all D : the asymptotic delocalisation of the Green columns (T146's remainder) stays *open*, typed open; only the sign-free / Green constant is bounded a priori — the energy-route σ_{tot} stays measured; every T146 exponent is a fit, sandbox. Fences: Maz'ya 1985 (strong-type half not used as a theorem anywhere) / Rayleigh 1877–Ritz 1909 / Cauchy–Schwarz / Miclo 1999 / Davis–Kahan 1970 (cited, deliberately not used) / Perron–Frobenius (not applicable to E , not used) / Fukushima–Ōshima–Takeda 1994 / Charikar 2000 / Goldberg 1984 / Gershgorin 1931 named classical; Weil 1952 cited, never used as a criterion; zero-firewall AST-checked; no marker moves. Machine-checked: [verification/v547_level_lemma_identities.py]. [E] (per-instance identities and certified inequalities)

PRIME.GREEN.SZEGO.IDENT.01 — the Green/Szegő identities of phase 2 (T147, T148). (1) The factorisation identity $\Gamma = \sqrt{m} \lambda_{\text{up}} \max_j \|Re_j\| = \sqrt{Q^*} \cdot S_w$ with $S_w = \lambda_{\text{up}} \|R\|_F$ purely spectral and $Q^* = m \max_j (R^2)_{jj} / \text{tr}(R^2)$ purely geometric — exact at 2.2×10^{-16} on 12 frame-A windows ($m = 26 \dots 285 \leq 300$) and 7 random positive definite forms; the certified factorisation (residual-paid column norms, Frobenius bracket) dominates the true Γ everywhere ($S_w = 1.1186$ – 1.5777 , $Q^* = 2.0879$ – 2.8634 ; controls $5.0 \times 10^{-3} / \geq 10^{-3}$). (2) The bottom-block trace identity $\sum_j (\Pi_B)_{jj} = |B|$ exact (3.3×10^{-16} ; $\theta = 10$, $|B| = 3$ – 5), hence $Q_B \geq |B|$ a theorem; the *measured* $Q_B/|B| = 1.5091$ – 1.7980 inside the preregistered band $[1, 2.2]$ against the Szegő/Widom sharp reference $2|B|$ (the rank- $|B|$ coordinate projector overshoots by ≥ 3.9). (3) The second-difference ℓ^1 ceiling: $|a_k| \leq \min(1, \|s_k\|_\infty \|L_0^p \psi\|_1 / \mu_k^p)$ for $p = 1, 2$ against the true coefficients everywhere, the m -free closed form $\|a\|_1 \leq 2\sqrt{\nu} + 1$ with $\nu = (m+1)^{3/2} \|L_0 \psi\|_1 / (2\sqrt{2})$, and $m \|\psi\|_\infty^2 \leq 2\|a\|_1^2$; Dirichlet calibration $\|a\|_1 = 1$ exact, $\nu \rightarrow \pi$ (controls: $\times 10^{-3}$ fails, a one-index μ shift breaks the identity at 1.5×10^{-1}). (4) The layer-cake S_w certificate: completed LDL^T inertia counts equal the direct spectral counts on all 96 (window, rung) pairs (Sylvester 1852; Bunch–Kaufman 1977), and the layer cake certifies $S_w \leq 1.3288$ – 1.8220 per window (true 1.1186–1.5777, slack ≤ 1.216 ; dropping the bottom layer breaks it everywhere). (5) The Liouville transform $m \|\psi\|_\infty^2 \leq \kappa_\Lambda m \|\hat{\varphi}\|_\infty^2 \leq 2\kappa_\Lambda \|a(\hat{\varphi})\|_1^2 \leq 2\kappa_\Lambda \text{ceil}(\hat{\varphi})^2$ with $\varphi = \Lambda^{-1/2} \psi$ and a-priori $\kappa_\Lambda = \max \Lambda / \min \Lambda = 1.3868$ – 2.5805 , stepwise on every bottom-block mode; the weight-class discriminator on a model section with a genuine order-2 zero ($f(0) = 0$, $f''(0)/2 = 1$ exact), ladder $m = 64 \dots 288$ at matched κ_Λ (mismatch 0.0013): the bounded-variation multiplier keeps ν flat (max/min 1.003) while the rough one (TV larger by 162.6) grows by 19.9 — the roughness, not the

conditioning; the first step $\times 10^{-3}$ fails provably (slack $\leq \kappa_\Lambda^2$). **Named limits as load-bearing content:** per instance on small windows — a finite list of certified window inequalities with an explicit maximum; nothing uniform in the zone index or in D , and explicitly *no* statement for all D : the one open lifting input ($\nu_L \leq F(\text{form})$ with F m -free; equivalently bounded $\text{TV}(\log \Lambda)$ or any weaker smoothness of the multiplier) stays *open*, typed open; the symbol-level order-2 hypothesis is *refuted* (T148: $f(0) < 0$ on every surface window) and enters nowhere as a theorem; σ_{tot} stays retired as a route; every T147/T148 exponent is a fit, sandbox. Fences: Kac–Murdock–Szegő 1953 / Widom 1958 / Basor–Ehrhardt 2009 / Böttcher–Silbermann (the mechanism’s address, never an authority) / Sylvester 1852 / Bunch–Kaufman 1977 / Euler 1735 / Rayleigh 1877–Ritz 1909 / Cauchy–Schwarz / Davis–Kahan 1970 (cited, computed vacuous, not used) / Gershgorin 1931 / Wilkinson 1968–Higham 2002 named classical; Weil 1952 cited, never used as a criterion; zero-firewall AST-checked; no marker moves. Machine-checked: [verification/v548_green_szego_identities.py]. [E] (per-instance identities and certified inequalities)

PRIME.GAUGE.PARITY.IDENT.01 — the gauge/parity identities of phase 2 (T149, T150).

(1) The gauge freedom: for *any* positive diagonal $\tilde{\Lambda}$ the whitening $\tilde{E} = \tilde{\Lambda}^{-1/2} A \tilde{\Lambda}^{-1/2}$ is an exact congruence, so $\lambda_{\min}(A) \geq \text{cert } \lambda_{\min}(\tilde{E}) \cdot \min_j \tilde{\Lambda}_j$ is a valid certified lower bound for every gauge (an identity plus one Rayleigh step; family maximum valid) — verified on all 48 gauge assemblies over 12 frame-A windows ($m = 26 \dots 285 \leq 300$); $\Gamma = \sqrt{Q^*} S_w$ holds *gauge-invariantly* at 2.4×10^{-16} ; the const gauge has $\text{TV}(\log \tilde{\Lambda}) = 0$ *exactly* with the certified entrywise sandwich $\sigma = 1.3868\text{--}2.5805$ (controls: a 1% Q^* perturbation breaks at 4.99×10^{-3} ; the completed-Cholesky floor refuses on the shifted indefinite form, 12/12). (2) The two-regime discriminator: the flutter smoother (moving geometric mean, half-width $m/16$) removes TV by $\geq 2.88\times$ while moving the Liouville-transported $\nu_{\tilde{L}}$ by ≤ 0.0088 — even the matched-amplitude rough re-gauge moves it ≤ 0.0046 (bar 0.01) — while the *untransported* functional jumps 0.41–9.76 under the same rough gauge: the roughness is in the form, not the multiplier (T149’s withdrawal of the TV hypothesis, wired per instance). (3) The explicit zone diagonal $\Lambda_r = c_0 - c_{M-1-2r} + \sum_{s \neq r} \max(c_{|r-s|} - c_{M-1-r-s}, 0)$ at residual 5.9×10^{-16} ; the exact split $c = c^{\text{arch}} + c^{\text{atom}}$ with $c^{\text{atom}} \leq 0$ entrywise and the closed atom budget $\|c^{\text{atom}}\|_1 \leq \sum_j \mu_j \leq 4B\sqrt{N}$, $B = 1.038821$ the verified maximum of $\psi(n)/n$ over the whole atom table (Chebyshev 1852, computed, never assumed; Abel summation); the archimedean section alone is *indefinite* on 12/12 (up to 5 negative eigenvalues, LDL^T) while the full section has zero — the atoms are co-responsible for the positivity, the additive perturbation reading structurally empty (control: dropping the positive part breaks at 4.25×10^{-1}). (4) The parity compression $U_-^T T_M(c) U_- = T - H$ and $U_+^T T_M(c) U_+ = T + H$ exact (3.9×10^{-16} , cross block 1.4×10^{-16}) on the 6 windows with $M \leq 300$: the reflection-odd section *is* the compression of the full symmetric Toeplitz section onto its antisymmetric parity sector, and the negative inertia of the sign-changing symbol ($f(0) = -29.2 \dots -6.3 < 0$, re-read from T148, never re-derived) sits entirely in the *even* sector, the odd sector counting zero (control: the sign-flipped basis misses by 3.99×10^{-1}). (5) The parity-sine calibration: the parity sines are exact eigenpairs of the parity Laplacian (corner 3 forced by antisymmetry; eigen-residual 2.3×10^{-13}), $\nu_k^P = \pi k^2$ on the exact model at $m = 2001$ to 3.0×10^{-5} , and the per-window ladder $\nu_k^P \leq Ck^2$ holds with certified $C = 11.70\text{--}17.28$ and the derived per-mode ceiling $m\|\psi_k\|_\infty^2 \leq 2(2\sqrt{C}k + 1)^2$ (controls: $\times 10^{-3}$ fails, a one-index shift misses by 0.75). **Named limits as load-bearing content:** per instance on small windows — a finite list of certified window inequalities with an explicit maximum; nothing uniform in the zone index or in D , and explicitly *no* statement for all D : the one open input after T150 (an m -free constant C in the odd-parity-sector ladder $\nu_k \leq Ck^2$ for a sign-changing-symbol Toeplitz section) stays *open*, typed open; $\text{TV}(\log \Lambda)$ stays withdrawn as a hypothesis (T149); σ_{tot} stays retired as a route (T148); every T149/T150 exponent is a fit, sandbox. Fences: Kac–Murdock–Szegő 1953 (parity sector) / Widom 1958 / Basor–Ehrhardt 2009 / Böttcher–Silbermann (the parity sectors as the algebra’s natural objects; the sign-changing symbol is exactly what Basor–Ehrhardt does not cover, and that gap is the open input) / Sylvester 1852 / Bunch–Kaufman 1977 / Chebyshev 1852 (verified on the table) / Abel / Weyl 1912 / Bauer–Fike 1960 / Davis–Kahan 1970 (cited, computed dead, not used) / Rayleigh 1877–Ritz 1909 / Cauchy–Schwarz / Gershgorin 1931 / Wilkinson 1968–Higham

2002 named classical; Weil 1952 cited, never used as a criterion; zero-firewall AST-checked; no marker moves. Machine-checked: [verification/v549_gauge_parity_identities.py]. [E] (per-instance identities and certified inequalities)

PRIME.ODD.SECTOR.IDENT.01 — the odd-sector identities of phase 2 (T151). (1) The grid step-over: the parity sines live on the *shifted* grid $\theta_k = 2\pi k/N$, $N = 2m+1$, which never contains $\theta = 0$, with $\mu_k^P = 2 - 2\cos\theta_k$ *exactly* (2.2×10^{-16} ; eigen-residual 2.3×10^{-13}); the symbol has $f(0) = -47.6 \dots -6.3 < 0$ (T148's honest negative, re-read, never re-derived) with a nonempty negative window $[0, \theta_c)$, and $\theta_c/\theta_1 = 0.371\text{--}0.485 \leq 0.9$ on *every* window of the battery (12 frame-A windows, $m = 26 \dots 285 \leq 300$): the odd grid *steps over* the negative window — odd-sector positivity is a grid fact. The honest negative beside it, promoted as content (T150's rest item 3 answered negatively): the symbol-only pencil floor $\kappa_{\text{sym}}(1 - \rho)$ is vacuous ($\rho \geq 1$) on 12/12 and the symbol moves by 0.45–3.27 relative across *one* mode spacing — the local model is a *matrix* statement, not a symbol statement (controls: the Dirichlet corner breaks the eigenpairs at 1.0, a one-index shift misses by 6.5×10^{-3}). (2) The certified bottom ladder: $\lambda_k(A) \leq S\mu_k^P$ for $k \leq 8$ by completed LDL^T inertia counts (Sylvester 1852; a certificate, never a sorted list), $S = 1.1019\text{--}6.4725$, the count equal to the direct spectral count on all 96 (window, k) pairs — order 2 in k , $f(0) < 0$ absorbed by the Rayleigh floor $L_P \geq \mu_1^P I$ (control: $S \times 10^{-3}$ certifies no eigenvalue below it). (3) The discrete Sobolev step: $\|\nabla\psi\|_2^2 = \langle\psi, L_0\psi\rangle - \psi_{\text{last}}^2$ *exactly* (7.6×10^{-15}), $\|\nabla\psi\|_2^2 \leq \langle\psi, L_P\psi\rangle$, and $\|\psi\|_\infty^2 \leq 2\|\psi\|_2\|\nabla\psi\|_2$ — licensed by the odd sector's virtual node $\psi_{-1} = 0$; calibration: the model pencil is (1, 1) (certified 1.1×10^{-12} at $m = 241$) and the Sobolev per-mode price is π to 5.1×10^{-8} at $m = 2001$ — the honest, m -free cost of leaving the ℓ^2 world (controls: the boundary term 4.8×10^{-2} , the Neumann undercount 0.50). (4) The closed archimedean minimum (T150's rest item 2 closed *and attained*): given the two per-window-checked shape properties ($c_i^{\text{arch}} < 0$, non-decreasing on $i \geq 1$), $\min_r \Lambda_r^{\text{arch}} = c_0^{\text{arch}} - c_{M-1}^{\text{arch}}$ *exactly* (residual 0.0; value 2.1719–3.4674), attained at $r = 0$ (control: the full atom-including kernel breaks the shape and misses by $\geq 8.7 \times 10^{-2}$). (5) The linear per-mode bound: with the certified pencil floor $\kappa = 0.1939\text{--}0.3599$ (one completed Cholesky per window, floor carried as fl/μ_1^P , direction re-read on the actual bottom eigenvalue) and $K_{\text{bot}} = S$, $b_k = 2m\sqrt{K_{\text{bot}}\mu_k^P/\kappa} \leq C_S k$ with $C_S = 12.081\text{--}26.953 \leq 2\pi\sqrt{K_{\text{bot}}/\kappa}$ on the bottom 8 modes of every window — *linear* in k where the quadratic ν ladder grew, every m -power cancelled, nothing additive anywhere; the bound dominates the truth and the parity Dirichlet energy on every read mode; the bottom pencil ratio $R = K_{\text{bot}}/\kappa = 3.7362\text{--}18.6607$ is the one scalar the chain still owes, typed *open* in m ; the discriminator: on the T145 no-go form the ratio explodes ($1.686 \times 10^3 \rightarrow 2.667 \times 10^4$, factor 15.8 over $m = 64 \dots 256$) while the exact parity model holds $R = 1$ to 1.1×10^{-12} (controls: the ladder $\times 10^{-3}$ fails, the inflated floor 1.5κ makes the Cholesky refuse on 12/12). **Named limits as load-bearing content:** per instance on small windows — a finite list of certified window inequalities with an explicit maximum; nothing uniform in the zone index or in D , and explicitly *no* statement for all D : the one open input after T151 (the m -freeness of the bottom pencil ratio $R = K_{\text{bot}}/\kappa$ for the odd parity sector of a sign-changing-symbol Toeplitz section) stays *open*, typed open — its flat trend ($x^{0.037 \pm 0.015}$) is a fit, sandbox; the symbol-only route stays certified vacuous; $\text{TV}(\log \Lambda)$ stays withdrawn as a hypothesis (T149); σ_{tot} stays retired as a route (T148); every T151 exponent is a fit, sandbox. Fences: Kac–Murdock–Szegő 1953 / Widom 1958 / Basor–Ehrhardt 2009 / Böttcher–Silbermann (the address of the symbol question — the computed answer: the pointwise symbol language does not extend to the bottom mode) / Weyl 1912 / Parlett / Hardy–Littlewood–Pólya 1934–Agmon 1965 / Sylvester 1852–Bunch–Kaufman 1977 / Rayleigh 1877–Ritz 1909 / Cauchy–Schwarz / Wilkinson 1968–Higham 2002 / Chebyshev 1852 named classical; Weil 1952 cited, never used as a criterion; zero-firewall AST-checked; no marker moves. Machine-checked: [verification/v550_odd_sector_identities.py]. [E] (per-instance identities and certified inequalities)

PRIME.RITZ.CEIL.01 — the fixed-size Ritz ceiling certificate of phase 2 (T154). (1) The direction lemma: for orthonormal Q and $W = Q^T A Q$, $\lambda_k(A) \leq \theta_k$ for *every* $k \leq \dim$ — Ritz values are *upper* bounds for the eigenvalues of the *same index* (Courant–Fischer 1920 / Cauchy 1829), verified against the exact spectrum on the certificate subspace of every window and on 24

random subspaces (one-sided excess 0.0) — so the fixed-size ceiling needs *no* residual argument; Temple 1928 / Kato 1949 is a floor device and is used nowhere in the module (control: the reversed reading is recognised as false on 12/12 windows, θ_k overshooting λ_k by ≥ 18.6 somewhere on every window). (2) The sixteen-column certificate: $\text{span}\{t_1..t_8\} + A^{-1}L_P \text{span}\{t_1..t_8\}$ has fixed dimension 16, independent of m (one Green step by Cholesky back-solves, no inverse formed), and its ceiling $K^F = \max_k \theta_k / \mu_k^P$ (8 completed Choleskys of matrices $\leq 8 \times 8$) equals 1.1019–1.5845 with $K^F \geq \max_k \lambda_k / \mu_k^P$ on every window and $|K^F / K_{\text{bot}} - 1| \leq 9.9 \times 10^{-7}$ against the declared cap 10^{-4} on 12/12 (K_{bot} the inertia-certified true ladder constant, count = direct count on all 96 pairs): the fixed-size ceiling *attains* the truth, retiring the size- m LDL^T counts from the ceiling step (controls: the parity sines alone overshoot 41.6–739; corrupting the Green columns breaks the attainment on 12/12; Kaniel 1966 / Paige 1971 quoted as the mechanism, never used as a bound). (3) The one-direction anatomy (angles measured, Björck–Golub 1973): the median of the seven aligned angles between the bottom eight of A and the bottom eight of L_P is 0.03 – 0.14° while the largest is 83.79 – 89.70° on every window of the battery (the 12 deepest in-cap frame-A windows, $n = 29 \dots 139$, $m = 96 \dots 291 \leq 300$ — declared, not tuned) — the misalignment is carried by *one* nearly orthogonal direction, and that direction *is* the collapse price $\lambda_1(A)/(t\mu_1^P) = 4.408$ – 7.042 , recovered in full per window by one completed Cholesky of $A - \gamma I$ (size m , certified, not fixed size, labelled as such; control: $A = L_P$ returns $K_{\text{bot}} = K^F = 1$, zero misalignment, price exactly $1/t$). (4) The no-go discriminator: on the T145 reconstruction $c(l) = 1/(1+l)$ the odd section is positive definite and the Schur floor survives ($t = 0.05$ on all 4 sizes) but K_{bot} explodes ($\times 35.4$) and the sixteen-column ceiling explodes *with* it ($\times 57.5$) over $m = 48 \dots 288$ — no manufactured closure; the Courant–Fischer direction is violated nowhere (16/16); the Dirichlet control (the Hankel half removed, arithmetic kept bit for bit) stops being positive definite ($\lambda_1 = -5.60 \dots -3.55$ certified on 3/3) — the reflection half is load-bearing for positivity. **Named limits as load-bearing content:** per instance on small windows — a finite list of certified window inequalities with an explicit maximum; nothing uniform in the zone index or in D , and explicitly *no* statement for all D ; the attainment is exhibited per window and *not* promoted to a theorem (T154’s no-go stress shows the certificate overshoots there by 1.12–2.19 — a property of the real section, not of the instrument); the two open inputs after T154 ($R1'$, an m -free floor for $\lambda_{\min}(B_{HH})$ on the bulk parity block; $R2'$, an m -free lower bound for the L_P floor on the complement of the eight bottom Ritz directions — arithmetic-free, worth 91–100% of the collapse price) stay *open*, typed open; every floor consumed is certified per window at size m ; PRIME.ODD.SECTOR.IDENT.01’s ladder stays as stated and nothing is re-opened. Fences: Courant–Fischer 1920 / Cauchy 1829 / Kac–Murdock–Szegő 1953 / Schur 1917–Haynsworth 1968 / Sylvester 1852–Bunch–Kaufman 1977 / Weyl 1912 / Temple 1928–Kato 1949 (a floor device, not used here) / Kaniel 1966–Paige 1971 (quoted, never a bound) / Björck–Golub 1973 / Wilkinson 1968–Higham 2002 / Chebyshev 1852 named classical; Weil 1952 cited, never used as a criterion; zero-firewall AST-checked; no marker moves. Machine-checked: [verification/v551_ritz_ceiling_certificate.py]. [E] (per-instance theorems and certified inequalities)

PRIME.WEIL.OPERATOR.01 — the localized-Weil-operator identification (research contract, [O]). The external operator shell for the $C = 1$ line (v618/v619), preregistered 2026-08-02 after the third external review. *Named reference frame (verified citations):* Suzuki, *Weil’s quadratic form via the screw function* (arXiv:2606.09096, June 2026) realizes the localized Weil quadratic form on every interval $[-a, a]$ as a canonical self-adjoint operator A_a via the screw function, proves the continuity of the lowest eigenvalue λ_a in a without assuming RH, and presents Yoshida’s criterion: $Q_W(v) > 0$ for all nonzero odd $v \in C_c^\infty(\mathbb{R})$ already implies RH; a first numerical realization is arXiv:2607.24830. *The lemma chain:* **(W1)** exact operator identification — the frame-A window bilinear form $\hat{A} = B - S$ is the Galerkin matrix of the odd restriction of A_a on the declared piecewise bases, exactly or up to an explicit rank-one transformation and a known normalization (no asymptotic “looks similar”); **(W2)** form density — the nested window spaces are form-dense in the odd form domain of A_a with Mosco/norm-resolvent convergence of the Galerkin forms; **(W3)** uniform discrete positivity — $\hat{A}_{a,h} \succeq 0$ for every a and every refinement h (the RH-hard step; the

$C = 1$ contraction reading of **v618** — the normalized arithmetic residue as an operator of norm $\leq 1/h$ — is the named candidate mechanism); (**W4**) passage to the continuum — W2 + W3 give $Q_W^a(v) \geq 0$ on all odd test functions for all a , hence RH via the localized criterion. *Kill tests:* (K1) the W1 identification fails beyond a window-dependent fit; (K2) a complete, numerically stable fresh window stably exceeds the frozen $C = 1$ bound (the **v619** complete-comb surface is the definitional domain); (K3) the odd-sector restriction loses the explicit-formula bookkeeping. *Fence:* a research contract, not a claim — no RH statement, no positivity claim beyond the measured surface; the zero-side representation stays diagnostic (a proof must run prime-side or through the screw function, avoiding the critical-line circularity). Status: **[O]** (preregistered; the audit and the citation verification are machine-checked in [\[verification/v624_external_lattice_audit.py\]](#)). *First executed slice (2026-08-02):* the *atomic half of W1 is closed* — the TFPT atom table is the prime measure of Suzuki's screw function, literally (positions log prime powers, weights $\Lambda(n)/\sqrt{n}$, exact; tent projection verbatim), and the first numeric Galerkin comparison measures a *non-scalar* conversion profile ($-0.064..-0.070$) — the preregistered rank-one/normalization freedom, now with data ([\[verification/v630_suzuki_contact.py\]](#)). On the geometric side the Hodge-chamber question is *measured*: all 67 complete windows land in the positive cone of the cover polarization lattice, on one sheet ([\[verification/v627_hodge_chamber.py\]](#)); P -canonicity and the chamber theorem stay open. *Second slice (2026-08-02, same day):* the non-scalar residual is *resolved* — it IS the zeta pole term. The Lerch collapse $[e^{-t/2}\Phi(e^{-2t}, 2, \frac{1}{4})]'' = 4e^{-t/2}/(1 - e^{-2t})$ (exact) gives the structure theorem $g''(t) = -2\cosh(t/2) - 4e^{-t/2}/(1 - e^{-2t})$ off the atoms (in the $\tilde{g}$ -normalization of this chain; see the fourth-slice erratum below — Suzuki's own smooth layer is $+\rho$): the chain-kernel smooth layer is $-4\times$ the TFPT arch density *minus* the pole block $2\cosh(t/2)$ (the $s = 0, 1$ pole weights — TFPT's separate rank-one, **v591**). The closed per-lag ratio law $-D[4 + (e^t + 1)(1 - e^{-2t})]$ is verified, and after pole subtraction the smooth conversion is the *scalar* $-4D$ (monotone to 1.0006); the atom-layer lattice constant is D^2 exactly. **W1 status: closed at the measured level on both halves** — $\hat{A}_{\text{arch}} = -(1/4D)$ Galerkin_{smooth} + pole rank-one, with the typed remainder (second-order discretization term, $d \leq 2$ boundary cells, L_0^2 projection) named for the theorem-level statement ([\[verification/v631_w1_dictionary.py\]](#)). *The implication map stays explicit:* W1 (now measured-closed) $\rightarrow$ W2 (form density, OPEN) $\rightarrow$ W3 (uniform discrete positivity, OPEN — the RH-hard step) $\rightarrow$ W4 (continuum passage, classical given W2+W3). Closing W1 does not move W3; no step is silently upgraded. *Third slice (2026-08-02, same day):* the typed W1 remainder *closes*, on three fronts. (i) **Boundary closure, exact:** the duplication identity $\psi(\frac{1}{4}) = -\gamma - 3\log 2 - \pi/2$ and the scheme conversion $k = \pi/4 + \frac{3}{2}\log 2$ are sympy-exact, so the two δ_0 conventions are one object in one scheme, and the exact boundary equation $A_{\text{arch}} = \frac{1}{4}g''_{\text{smooth}} - \frac{5}{4}(\log \pi - \psi(\frac{1}{4}))\delta_0$ verifies at relative residuals $< 10^{-22}$; the $d \leq 2$ cells become three computable constants and the second-order term is the derived, window-independent moment law $1 + 1/(6d^2)$ ([\[verification/v640_w1_boundary.py\]](#); the exact dictionary identities additionally machine-proved in Lean, `WeilDictionary.lean`). (ii) **Portability:** the frozen dictionary (closed formulas in D , no knob) executes unchanged on three fresh windows $h = 285/540/997$ — atom constant $D^2(\text{window})$ at machine precision — and the preregistered kill criterion is met nowhere ([\[verification/v641_w1_portability.py\]](#)). (iii) **Matrix level:** the full quadratic form closes at the operator level (16-mode block norm distance 2.7×10^{-3} fully derived, 6.3×10^{-6} after measure-level atom re-binning; the separated pole block numerically rank 1); the literal per-vector 1% bar fails only on two *typed* lattice residuals (near-cancellation denominators; tent-vs-B-spline atom profiles), certified as typed-residual checks ([\[verification/v642_w1_matrix.py\]](#)). *Fourth slice (2026-08-02, second round of the day): erratum + theorem + W2 start + W3 tool diagnosis.* **Convention erratum (corrected the same day):** the second and third slices read Suzuki's eq. (1.3) with Lerch coefficient -1 ; the paper's own §2.2 data lock $+1/4$ ([\[verification/v643_w1_theorem.py\]](#), C0.1). All slice-two/-three identities are correct identities of the chain kernel $\tilde{g} = g - \frac{5}{4}\text{Lerch}$, and every measured number transfers *verbatim* via the exact identity $c_{\text{gal}}^{\text{sm}}(\tilde{g}) = -4c_{\text{gal}}^{\text{sm}}(g)$ — only the labels change: Suzuki's own smooth layer is $+\rho$, the two-layer dictionary collapses to the *single* scalar $+1/D$ on both layers (sign-compatible with positivity), and the origin constant vanishes, $\kappa = 0$ exactly (the $-\frac{5}{4}L\delta_0$ term was an artifact of

the misreading; the v563 near-cell scheme *is* Suzuki's origin bookkeeping). **W1 is THEOREM-closed at the measure level** ([verification/v643_w1_theorem.py]): the projection lemma closes the last remainder — Suzuki's L_0^2 mean-zero condition is *automatic* on the u -side ($D : H_0^1 \rightarrow L_0^2$ a bijection), the window parity sector is a subspace of the form domain, $A_{\text{arch}} = -g''_{\text{smooth}}$ exactly at every lag (3.4×10^{-52}), and the full form equality holds at 1.28×10^{-10} on the common odd sector. **W2 is started** ([verification/v644_w2_form_density.py]): the classical FEM-density slice is verified at rate (with Q_W H^1 -bounded and H_0^1 a form core the density chain is classical, cited); Rayleigh–Ritz on the nested window spaces is monotone from above on all six sequences, and the λ_a scan is consistent with Suzuki Thm 1.3 ($\lambda_a = 0^+$ within $\sim 10^{-9}$, no sign statement); the Mosco/norm-resolvent remainder is *named*, not claimed. **W3 stays open, now with a tool diagnosis** ([verification/v648_sign_uncertainty.py]): the sign-uncertainty / LP-ceiling toolbox has a real, 25-digit-anchored dictionary to the critical strip (the $d = 1$ strip multiplier's phase slope is exactly $L = \log \pi - \psi(\frac{1}{4})$, the W1 δ_0 weight), but its quantitative mechanism is powered by the strip half-width and dies at $d = 1$ ($e^{-E^*} = 0.967$ where $< \frac{1}{2}$ is needed; $d \geq 21$ would be required; the window family's negative structure sits $5.2\times$ outside the maximal BCK ball): a BCK-type mass obstruction cannot certify $\hat{A}_{a,h} \geq 0$ — any W3 route through this toolbox must replace the dimension lever by a different large parameter; empirically the W3 surface is positive on all 67 complete windows (min $\lambda_{\min} = +8.26 \times 10^{-4}$, float-floor certified). **W4 unchanged**; no RH statement. *Fifth slice (2026-08-02, the closing bundle): the consolidated W2/W3 state.* **W2:** the envelope inequality holds *measured* a -uniformly — the total window symbol grows sub-log with family-uniform constants $s_{\text{tot}} \geq c_0 w^*(t) - C_0$, $(c_0, C_0) \approx (0.021, 0.055)$; the earlier “flat” read was a t -range artifact ([verification/v663_garding_envelope.py]); the Mosco/norm-resolvent preparation stands — the resolvent family is Cauchy with uniformly bounded $H_{\log}$ norms, and the eigen-scale collapse is typed: W2 routes must run through resolvent observables ([verification/v655_w2_mosco.py]); the two named lattice remedies for the Gårding drift are *refuted* — the drift lives in the total symbol, not in the lattice edge ([verification/v661_garding.py], [verification/v662_garding_edgeband.py]) — and the drift itself is *intrinsic* to the positivity problem, not Suzuki-frame-made: the external Baez–Duarte control frame shares it ([verification/v667_baez_duarte.py]). The remaining W2 steps are a Fejér spectral-density bound replacing the per-mode symbol read, plus the symbol-level proof. **W3:** reduced on the measured surface to *lock sign plus the uniform bound* $q \tan^2 \theta \leq 1 - \delta$ via the exact 2D formula $r_{\text{id}} = (1 - q \tan^2 \theta) / (1 - \tan^2 \theta)$ ([verification/v657_rid_alignment.py]; the margin-bridge correlation typed as a lock identification, [verification/v656_margin_link.py]); the resonance landscape is mapped *without positivity loss* — $\lambda_{\min} > 0$ on all 635 surveyed points, every $P > 1$ excursion a rotation artifact ([verification/v659_w3_landscape.py]; the risk survey [verification/v658_w3_uniform_bound.py]; the rotation predicate is typed honestly (necessary, not sharp; successor: block deflation; [verification/v660_theta_predicate.py]); and the *ground-truth recalibration* lands ([verification/v668_ground_truth.py]): on a proven RH analogue true positivity has *no uniform margin* — $\delta(K) \rightarrow 0$ exactly beyond the support-resolution depth — so shrinking margins on deeper windows are the *expected* behavior of a true positivity, and the alarm signal is only $\lambda_{\min} < -\text{floor}$, which has never been observed; conversely the machinery breaks by ~ 13 orders of magnitude the moment the Euler product is removed (the Epstein fire-wall), so the family positivity read does carry arithmetic information; the needles are frame-side ([verification/v667_baez_duarte.py]). *Kill criteria K1–K3 unchanged; no step silently upgraded; no RH statement.* *Sixth slice (2026-08-02, the Fejér-density round): the density plane closes; the needle search narrows threefold; the Li corollary.* **W2 — the density plane is closed** ([verification/v669_fejer_density.py]): the total window symbol *is* the Fejér-smoothed zero-counting density — the exact identity $s_{\text{tot}}(\cdot; a) = 2\pi(F_a \star dN)$ (Weil explicit formula on the tent test pair, verified 2.9×10^{-6} pointwise / 7.2×10^{-8} t -averaged against the Turing-certified comb) — and the *unconditional* chain RvM + Trudgian yields $\rho_{a,\delta}(t) \geq 0.306 \log(2+t) - C_0$ a -uniformly at $\delta \in \{4\pi, \pi a\}$ with 10/10 finite certificates on the zero-free grid (min margin +1.36). The honest floor: the literal $1/a$ and the plane-wave π/a widths sit *below* the RvM threshold $4\pi A_1 = 1.4074$ — per-mode symbol control is not certifiable this way at any height (the structural explanation

of the Gårding drift), so the remaining W2 piece is the packet-to-point translation below the floor. **W3 — the needle search is narrowed threefold:** the needles are *not low-rank* (the $(1+K)$ -block deflation fails honestly: leak median 1.000 at every $K \leq 8$, fidelity $p_{90} \approx 49\%$; [verification/v670_w3_block_deflation.py]), *not zero-comb-driven* (the Lehmer-pair link is a clean null: h^3 -partials all $p > 0.1$, placement control unremarkable; [verification/v671_lehmer_resonance.py]), and *not in the BD frame* ([verification/v667_baez_duarte.py]) — the mechanism search moves frame-/assembly-side. **New — the Li corollary and the E_4 lock:** the W1 window positivity *contains* 30/32 finite Li coefficients ($g_n = \frac{1}{2}e^{-|u|/2}L_{n-1}^{(1)}(|u|)$ derived exactly; band $2..4 \cup 6..32$ on $h = 1433$, pole-free hyperplane, odd sector certified by $\lambda_{\min} = +1.516 \times 10^{-3} > 0$; [verification/v672_li_corollary.py]), and the E_8 shell data and the ζ comb agree in *one* Li sequence ($\Lambda(E_4, s)$ completed with residues $\pm 1/240$ — the E_8 240 is the residue normalizer; $\lambda_n^L = 2\lambda_n^\zeta$ exact; budget usage 0.124; [verification/v673_li_e4.py]). *Kill criteria K1–K3 unchanged; no step silently upgraded; no RH statement. Seventh slice (2026-08-02, the consolidating contraction): the W2 end-state; the W3 contraction to the carrying form; two theorems and the honest typing of the wall.* **W2 — the end-state:** the density plane is closed ([verification/v669_fejer_density.py]); the Gårding statement now stands in *frame form with theorem constants* ([verification/v674_packet_garding.py]: the frame inequality $Q + C_0G \succeq c_0Y$ holds at every M with the v669 constants rebuilt zero-free, $c_0 = 0.3058$, margins $+1.52.. + 2.29$, a -uniform — while the packet *projection* norm honestly does not escape the $1/\log$ drift: typed residual with inverted expectation, the minimizer single-mode tight at 0.90); the Mosco mechanism is numerically complete (8/8 sequences, same module); the named remainder is *within-packet equidistribution* ($\theta \rightarrow 0.20$) = unconditional zero-gap information. **W3 — the contraction to the carrying form:** after the needle mechanism search *saturates* (all four assembly-side candidates miss: lag-quantization weakly significant $p = 0.007$ but too dense, phase gradient wrong sign, lock rotation cleanly killed, pole weight significant $p = 2 \times 10^{-4}$ but inverted — the needles are fourfold characterized as an emergent bulk/frame property; [verification/v675_needle_mechanism.py]), the contract W3 form becomes “*MARGIN-regime generic bound + needle risk map*” rather than a uniform bound with an exception predicate. The candidate mechanism $C = 1$ is demystified *and* grounded as a *quadrature theorem* ([verification/v676_c1_mechanism.py]: q_r = the exact zero-side reading of the lock profile, median residual 1.1×10^{-2} on 67 windows; h^{-1} = the DST normalization; the constant = the zero-free RvM density integral at ratio 1.005 — *no RH lever beyond the density, but computable*; $\sup \leq 1$ is not implied by the mean, the $\sin^2$ sampling budget is the precise remainder with factor-2 headroom; the v596 exponent corrected: -1.01 was flip-contaminated, clean -1.28). And the *structure theorem* ([verification/v677_w3_structure_theorem.py]) is the exact identity of the rung: window positivity $\Leftrightarrow$ alias positivity of the zero symbol (Cantoni–Butler compression + sandwich + master identity with $T_x \geq 0$ on the line) $\Leftrightarrow$ an off-line detector whose Epstein-lab census (exactly 12 off-line zeros, ζ_K control 0) predicts the form break at factor-2 level (ratio 0.803), with the threshold map $2\alpha s_{\min} \approx 1.4..2.6$ (the Ihara echo) — so *W3-on-the-family* is a theorem plus a detector certificate, while *uniform* W3 over all windows is the conjecture itself (Weil 1952 / Yoshida 1992, RH via the localized criterion), typed as such and *not* an RH statement. *Kill criteria K1–K3 unchanged; no step silently upgraded; no RH statement. Eighth slice (2026-08-02, the zero-gap round): the W2 end-state, with the last gap carrying a number.* **The frame chain is now fully unconditionally grounded:** the best explicit unconditional zero-gap theorem ([verification/v678_zero_gap_theorem.py]: the S -difference route with $S_{\text{bound}} = \min\{\text{Platt } |S| \leq 2.5167 \text{ to } 3.06 \times 10^{10}; \text{Trudgian 2014; Bellotti 2025 Cor. 1.5, asymptotic floor } 4\pi 0.10076 = 1.26619\}$, machine-verified on all 1999 Turing-certified comb gaps with 0 violations and min air $2.05\times$) makes the v674 frame packet floor *unconditionally* positive: adaptive bands $\delta_p = \kappa H_{\min}(b_p)$ hold $\geq \kappa$ ordinates per band unconditionally, and the theorem-shaped floor $0 < B_p \leq V_p$ holds on every packet (median worst-case discount 0.037) — zero-gap theorem + Fejér chain ([verification/v669_fejer_density.py]) ground the entire frame-Gårding statement in cited unconditional inputs. **The last gap (pointwise/projection form) carries a concrete number:** the packet projection ladder does not stabilize under adaptive re-partitioning either (typed residual with inverted expectation: 6/6 (κ, C) cells miss, the mini-

mizer stays single-mode at 0.898 — the obstruction is re-partition-invariant), and the pointwise main-lobe capture of a guaranteed zero *closes iff the explicit $S(t)$ constant A_1 improves from 0.10076 (Bellotti 2025) to $< 0.09017 = 1/(4a_0)$ — an 11.7% improvement of a named constant at the frontier of explicit analytic number theory, independent of RH*; the Platt branch bottoms out at $1.4183 > \pi/a_0 = 1.1331$, so no current chain reaches it at any height. This is the honest end-state of the W2 packet route: an unconditional frame form, a quantified external research target for the pointwise form, and within-packet equidistribution as the alternative input (W3/W4 territory). *Kill criteria K1–K3 unchanged; no step silently upgraded; no RH statement. Ninth slice (2026-08-03, the pinch-break and hole-closure round): the W2 end-state — the pointwise map at the anchor is gapless. The pointwise density floor at the anchor window is now a gapless split-type map* (comb-certified / scan-certified / abstract-Weil / unconditional-asymptotic), closing the 11.7% pinch and the $(2500, 7.27 \times 10^6)$ hole. **The pinch was a bookkeeping artifact** ([verification/v680_pinch_attack.py]): centered capture — the same zero-gap theorem spent two-sided, $\text{dist}(t, Z) \leq H_{\min}(t - 26)/2$, comb-verified on 6595 grid points with 0 violations — *doubles the threshold to $A_1 < 1/(2a_0) = 0.18034$, so the existing Bellotti constant 0.10076 passes with 79% headroom; and the Beurling–Selberg minorant eliminates A_1 from the counting chain entirely (the loss is exactly the π/a_0 minorant mass plus the explicit prime term $P = 2.534$; positive from $t^* = 1.11 \times 10^7$, every window including $a \rightarrow \infty$; Weil identity on the comb at 9.0×10^{-5}).* **The hole is closed** ([verification/v681_coverage_hole.py]): the exact prime term with a hierarchical Lipschitz certificate moves the abstract entry down a factor 72.5 to $t_x = 1.53 \times 10^5$, and a budget-certified Riemann–Siegel scan (223 949 zeros, Gabcke remainder, correctness 603/603 against the certified strip, found/expected 0.99774) carries $(2515, 1.56 \times 10^5]$ with floor 0.01664 — the statement: $s_{\text{tot}}(t; a_0) \geq 0.02259 \log(2 + t) - 0.5185$ for all $t \geq 10$, every region with named support and type (H1 honestly adjudicated: the Platt-constant chain reproduces the hole boundary exactly and cannot enter). **The W2 chain at the anchor:** density ([verification/v669_fejer_density.py]) + frame-Gårding ([verification/v674_packet_garding.py] / [verification/v678_zero_gap_theorem.py]) + pointwise ([verification/v681_coverage_hole.py]) — complete, with the mixed typing ledger explicit. *Remaining:* deep family windows $a > 4.43$ (Selberg-only entry), the formal Mosco writeup, the projection-norm form, $a \rightarrow \infty$ (= W3/W4 territory). *Kill criteria K1–K3 unchanged; no step silently upgraded; no RH statement. Tenth slice (2026-08-03, the great bundling): the master contract PRIME.UNIFPOS.01 and the five proof offensives. The master contract — the Uniform Positivity Theorem (PRIME.UNIFPOS.01, registered as a new contract).* Target statement, verbatim: *for every $a > 0$, Suzuki’s localized Weil operator B_a on $H_1^0(-a, a)$ is positive semidefinite, with the positivity derived from an operatoric/geometric representation whose definition uses neither Riemann zeros nor an RH-equivalent presupposition.* The machine-checkable intermediate form: $B_a = L_a + K_a$ with $L_a \succeq 0$ from a structural source and the relative bound $|\langle K_a v, v \rangle| \leq \theta \langle L_a v, v \rangle$ with $\theta < 1$ uniform — then $B_a \geq (1 - \theta)L_a \succeq 0$. **Annotated with the v682 verdict** ([verification/v682_lk_split_theta.py], LK-SPLIT-DIES): the two-sided form for smooth zero-free L is structurally dead — measured $\sup \theta$ per split 11.3–58.0 with every smooth L indefinite on the odd sector, and the death mechanism is a *positive* find: the extremal pencil direction sits at $t = \gamma_1 = 14.13$ (found from primes + digamma alone, no zero loaded), $\mu_{\max}$ follows the spike law $1 + 2\alpha/\Omega(t_{\text{peak}})$ (median dev 0.127), θ grows $\sim 2.47\alpha$ — *the deployed windows resolve individual zeros*, so the surviving directions are *one-sided* forms (the lower side alone — which must beat the pure-renaming floor $\theta = 0.998$ of the trivial spectral split) or *multi-resolution* splits (an L carrying the comb spikes at matched resolution; untested, the open lever). *Kill criteria:* (K1) a C_a/L_a defined through Riemann zeros = renaming; (K2) a contractivity statement that is itself RH-equivalent = renaming; (K3) any candidate binding at the trivial-split floor adds nothing beyond measured window positivity. **The rank-3 chain reaches theorem shape at the sign level:** [verification/v683_rank3_functionals.py] lifts the prime influence on the load-bearing 2×2 block into three exact functionals (4.9×10^{-16}) and adjudicates every entrywise route as class (c) (the T-B absorption margin demands $\sim 10^6 \times$ more than any per-entry input; no Cauchy–Schwarz win: 99.7% of atoms violate pointwise;

even a nonnegative pretentious measure breaks $\det S$); `[verification/v684_rank3_zeroside.py]` treats the one functional by the explicit formula ($\text{residuum} \leq 6.9 \times 10^{-6}$, 22 491 budget-certified zeros) and delivers the *unconditional* $\kappa_{\text{unc}} = 0.039\text{--}0.190 < 1$ (class (a); the pretentious escape blocked $\times 634$ by the classical zero-free strip); `[verification/v685_rank3_uniformity.py]` closes the uniformity quantifier: the symbolic envelope $K_{\text{env}} \leq 0.9798 < 1$ for all 62 complete windows with $a \geq 3.434$ plus five finite certificates $\kappa' \leq 0.11$ — $\det S > 0$ *unconditionally on the entire declared surface* (SURFACE-CLOSED; a surface theorem on the declared family, *not* uniform in all a ; T-B stays **prob:R1**). *[2026-08-03 prob:R1 update: T-B is now typed and largely closed. [verification/v692_rank3_lockgram.py] (7/7) types T-B(block) exactly: $\hat{A}_2 = \text{positive zero-Gram} + \text{psd rank-1 pole layer}$, the margin is the transverse zero mass (a sum of squares), on-line zeros only help (psd superadditivity) — “E kippt nicht” is a quantified partial-RH tail statement, neither a 2c-envelope question nor a W3 renaming. [verification/v693_rank3_density_close.py] (5/5) builds the cited explicit bounds (Platt–Trudgian 3×10^{12} ; Hasanlizada–Shen–Wong count; functional-equation halving; the explicit Ingham-form zero density, arXiv:2507.15184) into a sharpened safe-side envelope (sinh difference pairing, $V(0) = 0$, on-line subtraction; PEN down $\times 6.5\text{--}14.4$): 60/70 **complete windows close unconditionally-modulo-citations**; the exact open remainder is $h \in \{1359, 1721, 1868, 2018, 2093, 2472, 2475, 2630, 2656\}$ with $T^* \in [1.0 \times 10^{13}, 3.0 \times 10^{13}]$ plus $h = 5690$ at $T^* = 8.5 \times 10^{14}$; extension beyond the declared family grows $\sim e^{3.74a}$.] [2026-08-03 prob:R1 update II: [verification/v712_rank3_transverse_deck.py] (5/5) scalarizes the v692 margin identity along the transverse direction — the penalty pays only the transverse zero mass, median gain $\times 4.23$ on the open ten — and the census closes nine of them: 69/70 **complete family windows now close at the cited 3×10^{12}** (unconditional-modulo-citations); the single remainder is $h = 5690$ with $T^* = 1.6 \times 10^{14}$ (down from 8.5×10^{14}); the deck-triplication lever is honestly dead (triangle inequality: channel sups can only lose). T-B itself stays the conjecture.] **The kernel class collapses** (`[verification/v687_extremal_kernel.py]`, KERNEL-CLASS-TOO-NARROW): the pinning certificate $2\pi F_{a_0}(m\pi/a_0) = 0$ (sympy-exact) plus Shannon sampling collapse the entire admissible band-limited positive class onto $\{\lambda \text{ Fejér}\}$ with class supremum $\theta_{\text{req}} = 0.0901684 < 0.10076$ — no kernel optimization can beat the $S(t)$ constant; W2 stays closed via the v680/v681 route. **The W3 falsifier is constructive** (`[verification/v688_interpolation_detector.py]`, INTERP-FALSIFIER-CONSTRUCTIVE): every isolated off-line quadruple is detected from $2\alpha\delta \geq 1.974$ ($C_{\text{up}} = 0.987$), a $\times 2$ improvement over the v677 mode map, with masking fully adjudicated (46 broken + 2 provably sub-resolution of 48). **The geometric source opens a contract** (`[verification/v686_geometric_sos.py]`, PRIME.GEOMSOS.01): $\Lambda(n)$ reconstructed circle-free from E8 shell counting (7.1×10^{-31} , Satake data from counts, K1 passes), canonical dim-3 cover-SOS, K4 firing honestly; missing: the bundle H and the transport Φ (top-3 eigenvalue mass only 91.4%), milestones M1–M4 with kill criteria K1–K5. **The Hecke-SOS mechanism is extracted** (`[verification/v691_hecke_sos.py]`, HECKE-SOS-IHARA-MECHANISM-EXTRACTED): the target factorisation $A = B^*B + P$ exists *exactly* in the Ihara lab (B by Chebyshev recursion, no Cholesky; $P \succeq 0 \iff$ Ramanujan), the deployed ζ window form is exactly the sine/defect half of the canonical split (the part whose lab positivity is the RH analogue), the Euler mechanism is measurable (resonance lattice $2\pi j/\log 2$ exact, depth $\times 1.98$), and the missing ingredient is exactly named: Z1 — a self-adjoint geometric operator whose polynomial traces are the window moments (Hilbert–Pólya); Z2 would be RH itself. *Kill criteria K1–K3 unchanged; no step silently upgraded; no RH statement.**

The review contracts of the great bundling (registered 2026-08-03, brief). *E8.GAUSSIAN.CODE.01* — $E_8(\mathbb{Z}[i])/(1+i)$ is the information space of $\text{RM}(1,3)$: first slice *executed* by `[verification/v689_gaussian_code_bridge.py]` (26/26, GAUSSIAN-CODE-BRIDGE-EXACT: SNF-exact quotient $\mathbb{F}_2^4$, $240 = 15 \times 16$ census with the zero class provably empty, $\sigma =$ the $\text{RM}(1,3)$ family permutation with residual gauge of order 18, coordinate block $= F_1 + F_2 + F_3$; all must-fail controls fire); remaining: the theorem writeup. *[2026-08-03 addendum: the algebraic core is now formally verified — kernel-checked in Lean 4 (TfptCarrier/GaussianCodeBridge.lean, lake build green, no sorry/native_decide): the $\mathbb{Z}[i]$ -module with the SNF certificate (unimodular P, Q both directions), the 240-root census 15×16 with the zero class provably empty, μ_4 stability, and the σ*

family action with coordinate block = $F_1 + F_2 + F_3$.] **ARF.BOUNDARY.CODE.01** (registered 2026-08-05, [O]) — boundary code uniqueness: the non-circular selection chain boundary $\rightarrow$ code $\rightarrow E_8 \rightarrow \mathbb{F}_2^4 \rightarrow$ five slots *exists* with $g_{\text{car}} = 5$ as output ([[verification/v776_boundary_hamming_uniqueness.py](#)], 31/31, BOUNDARY-CODE-UNIQUE: census 135 $\rightarrow$ 30 $\rightarrow$ 14 $\rightarrow$ 2 $\rightarrow$ 1 without Hamming/ E_8 input; the flag datum is *one* deck-invariant $\mathbb{Z}_2$, the product of the four side orientations; Construction A of the selected code *is* E_8 ; the Arf selector is unique; packets 1 + 4). *The open contract form as registered:* derive evenness/self-duality of the mark bookkeeping from the seam axioms — the R1 *universe premise* (“the boundary realizes a doubly-even self-dual binary code on its 8 sides”) was the relocated and shrunk circularity, joined by the typed residuals R2 (deck placement is semantic, not selective) and R3 (the anchor naming $\bar{b}(F_\Sigma, A) = 1$ is part of the frozen selector, not a theorem). [2026-08-06 note, markers unchanged: all three residuals moved — R1 is replaced by R1' (the three physical axioms of **SEAM.CODE.TYPEII.01** below, [[verification/v799_seam_code_typeii.py](#)]); R2 is placed and R3 is named by the metaplectic S -lift (**SEAM.CLIFFORD.MODULAR.S.01** below, [[verification/v798_seam_clifford_modular_s.py](#)]).] [2026-08-06 round-22 note, markers unchanged: the attack surface sharpens — affine functions on the NS/R torsor reconstruct $\text{RM}(1, 3) = H_8$ with $\text{Stab}(\chi_{\text{NSR}}) = \text{AGL}(3, 2) = \text{Aut}(H_8)$ in its natural 8-point action ([[verification/v803_e8_jetcode_affine_nsr.py](#)]); a typed internal reconstruction, not a P2 removal — R1' unchanged.] No P2-removal claim; kills: a second boundary-legal code surviving the flagged selection, or a seam-axiom derivation forcing a non-doubly-even universe. No marker move. **SEAM.CODE.TYPEII.01** (registered 2026-08-06, [C]) — Type II from the seam axioms: at length 8 (four J -paired marks $\times$ two sides) the finite chain is exhaustively machine-verified over all 308,993 subspaces of $\mathbb{F}_2^8$ of $\dim \leq 4$ ([[verification/v799_seam_code_typeii.py](#)], 19/19, TYPEII-FORCED): mutual locality of the Construction-A boundary vertex words $\iff C$ self-orthogonal; integer conformal spin (trivial T sector phases) $\iff C$ doubly even; holomorphy/index one ($|\text{disc}| = 1$, S -closure of the single character) $\iff C$ self-dual — the survivor set is exactly the 30 $\hat{e}_8$ copies and no boundary-legal non-Type-II code exists. Chained to **ARF.BOUNDARY.CODE.01**, the full route seam axioms A1–A3 + deck + orientation bit $\rightarrow$ unique code $\rightarrow E_8 \rightarrow \mathbb{F}_2^4 \rightarrow g_{\text{car}} = 5$ stands with the single [C] residual R1': *the physical seam satisfies* A1 mutual locality (bose statistics / trivial monodromy), A2 integer conformal spin (modular T -alignment), A3 holomorphy/index one (a single modular-closed sector) — each a standard physical demand, not a coding-theory premise. Imported lattice-VOA/theta ingredients are classical, cited and machine-verified in every instance used; the OPE-linearity scope note is typed. Kill: a seam construction violating A1–A3 while staying boundary-legal. No P2 marker move. **SEAM.CLIFFORD.MODULAR.S.01** (registered 2026-08-06, partial state) — the missing Clifford bit as the metaplectic lift of modular S at $\tau = i$ ([[verification/v798_seam_clifford_modular_s.py](#)], 27/27, MODULAR-S-PARTIAL): *coset decision positive* — the per-binary-factor Hadamard lands exactly in the missing coset $\zeta_8 G_{31}$ with the exact fixed-point relations ($S^2 = I$, $T^2 = \text{Pauli parity}$, $(TK)^3 = \zeta_8 I$); the deck $J = iI$ is the metaplectic anomaly of the total symplectic Fourier (R2 placed) and $\zeta_8 = (1+J)/\sqrt{2}$ the real half-deck swapping the seam pair; the strict uniqueness census over all 46080 coset elements is *empty* — no Θ -real metaplectic involution fixes q^* , the Arf defect $q^* \mapsto q^* + \bar{h}(\cdot, c)$ is unavoidable — with exactly one G_{31} -conjugacy class of retyped survivors, and the σ -orbit Arf defects are exactly $\{A + F_\Sigma + F_i\}$ with XOR = the anchor A (R3 *named*, not fixed). *What remains:* derive the physical selection of the defect class (the anchor naming from seam data) or type the two-variant/defect residue as intrinsic gauge. Kill: a Θ -real metaplectic coset element fixing q^* (excluded here exhaustively) would falsify the defect law. No marker move. **CURVE.CODE.OUTERSPIN.01** (registered 2026-08-06, *decided* CANONICAL) — the outer bridge between the code duad side and the curve syntheme side ([[verification/v796_duad_syntheme_bridge.py](#)], 24/24, BRIDGE-CANONICAL): the Sylvester duality β with $\beta\sigma = \tau\beta$ and $\beta(q^*) = S^*$ (the unique $\langle \tau, \rho \rangle$ -fixed spread) exists and is unique up to the order-6 code joint stabilizer (one orbit including the τ/τ^2 swap); it resolves the v784 Arf mismatch canonically (q_Θ pulls back constantly to $q_{\text{even}}^* = q^* + \bar{h}(\cdot, A)$, the anchor shift), transports $\bar{5}/10$ and $3+2$ exactly, and sends t_A to $\rho = J$; χ_{NSR} transports as an honest exact two-variant torsor pinned by the v752 NS/R parity — that two-fold ambiguity is intrinsic and

typed, not open work. The `CURVE.CODE.2TORSION.01` inner-identification kill stands unchanged; no matter semantics (ROOTCLASS-MIXED). *[2026-08-06 round-22 note: the bridge sharpens into an incidence-operator statement — every census bridge factors exactly $N_{\text{code}} = P_{\Delta}^T N_{\text{curve}}^T P_{\Gamma}$, a transpose duality, not an isomorphism ([verification/v809_curve_code_doily.py]).]* No marker move. **E8.RAMIFIED.JETCODE.01** (registered 2026-08-06, [E]; carrying **E8.AFFINE.NSR.01**) — the jet-code normal form of the Gaussian compiler ([verification/v803_e8_jetcode_affine_nsr.py], 18/18 + 15/15, JETCODE-EXACT+AFFINE-NSR-EXACT): because 2 ramifies, $\mathbb{Z}[i]/(2) \cong \mathbb{F}_2[\epsilon]/(\epsilon^2)$ (unique iso; the split ring fails all 24 bijections) and $W = L/2L$ is free of rank 4 over the dual numbers; the deck is the jet unit $J = 1 + \epsilon$; among all 65536 sections of $0 \rightarrow \epsilon W \rightarrow W \rightarrow V \rightarrow 0$ there are zero μ_4 -equivariant splittings — non-splitness is the v782 no-origin obstruction; $240 = 15 \times 8 \times 2$ (root = address $\times$ error jet $\times$ orientation). Kill: a μ_4 -equivariant splitting, or a second iso class of the fiber ring. No marker move. **E8.SYNDROME.ALGEBRA.01** (registered 2026-08-06, dead as frozen) — the sharpest error-correction-hull hypothesis killed as preregistered ([verification/v805_e8_syndrome_algebra.py], 13/13 incl. the must-fire controls, SYNDROME-DEAD): max rank 222/256 over all four frozen phase conventions; the $\pm r$ proportionality obstruction structural for the whole diagonal-phase family; both post-hoc escapes score worse. Survives exactly: $16 + 240 = 256$, the 16 neutral kernels = the diagonal MASA, same-class products A_0 -valued over μ_4 . Successor criterion named: root-individual operator data beyond diagonal phases. The Construction-A Hamming theorem untouched. **CURVE.CODE.DOILY.01** (registered 2026-08-06, [E] + [O]) — the doily incidence identity ([verification/v809_curve_code_doily.py], 21/21, DOILY-EXACT): $NN^T = B + 2I$ entrywise on the K_6 doily (15 duads $\times$ 15 synthemes), spec = $\{9, 4^9, 0^5\}$; $N/3$ has singular values $\{1, 2/3, 0\}$ — the recovery base rate $2/3$ as the unique nontrivial singular value of the canonical duad-to-syntheme incidence; Petersen edges = synthemes 15/15; the Arf switch set T_6 = the six within-block duads. *Open target: the six-step composition identifying $(2/3)^6$ itself. [2026-08-06 round-23: decided — SIXSTEP-CLOCK-ONLY ([verification/v814_k5_sixstep_transport.py]): the rate arises canonically on three spatial routes (the doily alternation among them), but each carries a typed obstruction and sixth-power blindness makes per-step data mandatory — the deployed six is the clock exponent; the missing ingredient is a deployed shells-to-cusp isomorphism, named.]* Kill: any census bridge failing the transpose factorization. No marker move. **CARRIER.CUTCODE.01** (registered 2026-08-06, [E]; carrying **CARRIER.MOMENT.INCIDENT.01**) — the Arf split as the cut classification of K_5 ([verification/v810_cutcode_moment.py], 17/17 + 16/16, CUTCODE-EXACT+MOMENT-INCIDENT-PARTIAL): the cut map injective-isometric on all 256 cells; $16 = 1 + \bar{5} + 10$ is $(0|5) + (1|4) + (2|3)$; Clebsch $40 = 5 + 20 + 15$ over the Arf shells with within-10 = Petersen; roots/lines grade $80 + 160 / 20 + 40$ — Arf/cut grading, not matter; the master moment $\text{Tr}_{S+} X^2 = 4|X|^2 = 4h(E_8) = 120$ one symbolic identity. The frozen 210 decision: $\binom{10}{4}$ decorative (equivariant bijection killed by the exhaustive fixed-point obstruction; the partial $80 \rightarrow 5 \times 16$ miss-slot map and the Newton tie survive as audit). No marker move. **CARRIER.PETERSEN.RADIAL.01** (registered 2026-08-06, [E] + [H neu]; carrying **TRANSPORT.SIXTHROOT.01**) — the K_5 edge machine ([verification/v808_petersen_sixthroot.py], 19/19 + 16/16, PETERSEN-RADIAL-EXACT+SIXTHROOT-SPECTRAL-ONLY): charges $(-4, +1, +6) =$ the certified $SU(5)$ decuple; the edge graph is Petersen $\text{SRG}(10, 3, 0, 1)$; distance shells $(1, 3, 6)$ charge-pure; the flavor hexagon is the distance-2 shell edge-by-edge; spec(P^6) = $\{1, 64/729, 1/729\}$ is the deployed transport spectrum exactly; the 10-dim multiplicity reading ($\times 5 = g_{\text{car}}$ fast, $\times 4 = |\mu_4|$ slow) typed [H neu]. The sixth root: $B = \frac{1}{18}[[13, 1, 4], [1, 13, 4], [4, 4, 10]]$ unique symmetric doubly stochastic PSD in the corpus basis, $B^6 \text{ GL}(3, \mathbb{Q})$ -conjugate to but bit-equal to neither deployed form — the corpus freezes the spectrum, not the basis (the basis-freeze question is the named open). *[2026-08-06 round-23 dated correction + addendum ([verification/v814_k5_sixstep_transport.py]): the sixth-root census had missed the deployed v221 evaluation — T_{v221} , the exact spectral sum of v221's transfer $(1 - w)^6$, equals B^6 bit-exactly, so the verdict upgrades from SPECTRAL-ONLY to BITEXACT-vs-v221 and the basis-freeze question is answered on the deployed route (the frozen SPECTRAL-ONLY stands as the probe's adjudication; ABEL-DEAD correction precedent); and the [H neu] 10-dim reading now has a promotion-ready canonical PROPOSAL: $T_{10} = P_{10}^6$ in the Λ^2 basis passes*

all gates with the equitable $v774$ quotient = $(Q_{\text{Pet}}/3)^6$ bit-exact (TRANSPORT10-PROPOSED, no deploy claim).] Kill: a second PSD sixth root in basis, or a shell/charge mismatch. No marker move. **E8.QUARTIC.HALF.01** — G31 as the holomorphic μ_4 shadow of the E8 compiler: first slice executed by [verification/v690_quartic_half.py] (22/22, QUARTIC-HALF-ALIVE) with the honest correction: the vanishing half $\{2, 14, 18, 30\}$ is true but trivially μ_4 ; the substance is $F_8|, F_{12}|, F_{20}|, F_{24}|$ as a system of basic G31 invariants (Chevalley cited; the holomorphic restriction generates the full G31 ring); remaining: the symbolic degree-60 Jacobian factorization and the writeup. [2026-08-03 addendum: the μ_4 -orbit vanishing mechanism and the quartic power sum $\sum_{\alpha} \langle \alpha, x \rangle^4 = 576 (\sum_i x_i^2)^2$ over the explicit 240 roots are now kernel-checked in Lean 4 (TfptCarrier/QuarticHalf.lean, lake build green, no sorry/native_decide); the Chevalley basic-invariant step remains cited, not formalized.] **PRIME.W3.INTERPOLATION.01** — the alias-interpolation falsifier: first slice executed by [verification/v688_interpolation_detector.py] (constants above); remaining: the lemma as a paper lemma. [2026-08-03 addendum: the two named building blocks are closed near-proof by [verification/v694_interpolation_lemma_closure.py] (20/20, CLOSURE-BOTH-NEAR-PROOF): retention via an exact projection identity plus a closed-form $O(k^2)$ certificate ($r_f \geq 0.548$ on the family surface; straddle boundary case discovered, $k = 0$ branch secured), and the separation law form-corrected to the additive $\alpha^* = C_{\text{cell}}/\delta + C''/\Delta\gamma$ ($C'' \leq 0.59$, zero exceptions in 97 tests); the two open parent configurations detect at $\alpha^* = 7.90$; declared residues: the explicit Ingham constant chain, the Dirichlet-kernel overlap bound, the $C''(n)$ induction.] **PRIME.Z1.OPERATOR.01** — OPEN (Offensive 5 running): construct the self-adjoint geometric operator whose polynomial traces are the deployed window moments (the Hilbert–Pólya step of the extracted Ihara mechanism), without zeros as input; an operator defined through zeros or an RH-equivalent norm assumption = renaming; Z2 (the norm bound) would be RH itself — the factorisation localizes RH, it does not bypass it. [2026-08-03 update, the 5–5d series ([verification/v695_z1_trace_operator.py] / [verification/v696_z1_jacobi.py] / [verification/v697_z1_uvarov.py] / [verification/v698_z1_flow_recursion.py]): the contract stays OPEN, with the ground taken recorded honestly — (5) the measure comes from counting [E]: seam and orbifold are dead as Z1 spectrum (null-calibrated), while the E8 counting route delivers the Weil measure exactly (atoms 0.0, arch/pole via the closed Γ duplication split, 7/8 target spikes at $q = 0.000$); the comb sequence $c + \text{pole}$ is positive-feasible — the signedness of the deployed sequence is the pole subtraction. (5b) the canonical operator exists (CMV/Jacobi, all moments reproduced at 2×10^{-13}); no closed coefficient law yet, but the renaming clause is explicitly refuted: features on prime-power slots ($q = 0.000$), amplitude–mass link $r_{\log} = +0.76$, positivity load-bearing (170/171 vs 2865). (5c) masses: atoms are lag insertions (duality proven), exact transfer identity $\Delta\alpha = w_1/E$ (5.6×10^{-17}), and the inverted stabilization law — the Γ flow predicts the counting masses to $\sim 10\%$ (median ratio 1.026, corr 0.86). (5d) positions forced (windows 0.5–2 cells, jitter null 30/30, adversary –516 lags); shooting recovers the masses at the true positions to 0.11% median. Remaining: lookahead autonomy (the greedy-saturation failure of autonomous reconstruction) and the continuum reading of the operator.] [2026-08-03 update II, the montage and the moonshot: the contract stays formally OPEN — the continuum theorems are missing — but both remainders moved. (i) The lookahead question is decided ([verification/v702_z1_lookahead.py], 13/13): the Γ flow is a full-depth comb verifier ($913 \geq 898$, every fake dies ≥ 184 lags earlier) but not a generator — both prescribed lookahead routes reach 2–4 slots, and the per-slot information demand $g \sim 5\text{--}12$ bits is information-theoretic and parameter-robust (FLOW-VERIFIER-NOT-GENERATOR). (ii) The chain arc $S\text{--}A\text{...}S\text{--}G$ ([verification/v703_chain_tolerance_scaling.py] through [verification/v709_chain_zero_layer.py] plus [verification/v710_chain_two_stage_hecke.py] / [verification/v711_chain_firstbirth_scaling.py]) types the mass law honestly: tolerance induction dead (RH-grade), the arch is a deck-sector trace [E] ([verification/v705_chain_deck_sector.py], 3.4×10^{-16}), the flow α are the Schur parameters (10^{-40}), the section edge is an exact resolvent identity, and $G1 = [\text{E-edge}] + [\text{E-Hecke}] + [\text{M-rate}]$ with the corridor position ~ 0.53 as the named open selection question. (iii) The continuum candidate is assembled ([verification/v713_l1_montage.py], 16/16, L1-ASSEMBLED-MEASURED):

cover limit 6.2×10^{-8} , point masses at the D^2 rate, one UV slot, divergence exactly at $s \leq 1/2$ (the critical line as the boundary of the construction — a consistency signature, not a positivity statement); the missing theorems (self-adjointness, trace identification, positivity) are named. (iv) The atom layer is groupoid-internal (`[verification/v714_moonshot_hecke_groupoid.py]`, 19/19, MOONSHOT-GO; see PRIME.Z1.MOONSHOT.01 below). What is missing is exactly the continuum SÄTZE: until they exist, no marker moves.] [2026-08-03 update III, the evening round: the moonshot stages 2–4 plus the Satz-1 slice and the K collapse are done and machine-checked (see PRIME.Z1.MOONSHOT.01 below for the six modules `[verification/v716_moonshot_arch_glue.py]` through `[verification/v721_k1_node_capture.py]`). The Hilbert–Pólya candidate is now structurally complete at measurement level — an operator family, a state, an exact finite trace formula with a closed term dictionary to Weil 1952, and a measured spectral limit onto the zeros (100% of 377 at tol 0.25, rate -1.61 , SHA256 prediction firewall). The contract stays formally OPEN: the remaining hard substance is (L1) the identification of the vague limit at the line $s = 1/2$ (no mass loss — the single place the zero fluctuation enters) and (L2) the node convergence as a theorem (the capture half is proof-near, `[verification/v721_k1_node_capture.py]`; the $\sim 900\times$ super-resolution is the open K1b question); no marker moves, no RH statement.] PRIME.Z1.MOONSHOT.01 — OPEN (registered 2026-08-03, the groupoid program): derive the full Weil measure (atoms + arch + pole) as the trace layer of one geometric groupoid built on the $\mathbb{Z}[i]$ -E8 Hecke tower, with the arch layer glued (not direct-summed) and the E8 specificity established. Stage 1 (done): `[verification/v714_moonshot_hecke_groupoid.py]` (19/19, MOONSHOT-GO) shows the atom layer is groupoid-internal — the primitive degrees of the tower are exactly the Gaussian prime norms (maximality census; no prime projector anywhere: the primes come out of the tower), the log generator + cell normalizer de-divisorize the tower traces at 4×10^{-13} , and the σ descent reproduces 100 atom positions, 100 masses and 368 moments at deviation 0.00; declared ingredients (named as inputs, not derived): the $\sqrt{n}$ half-density and the σ descent. The honest fence (typed): the finite counting facts of stage 1 are E8-unspecific — the specificity question belonged to stage 2, which has since been executed. Stage 2 (done, 2026-08-03): `[verification/v716_moonshot_arch_glue.py]` (18/18, MOONSHOT-STAGE2-GLUED) — the arch glues: one object, one normalization (the free 3-scalar LSQ returns $\kappa = (1, 1, 1)$ to $< 10^{-12}$ on all five windows); the $1/4$ in $\operatorname{Re} \psi(1/4 + i\tau/2)$ is derived as the μ_4 fixed-sector offset; $\log \pi$ enters only through the declared self-dual (Tate) UV cell — the exact arch mirror of the declared $\sqrt{n}$ half-density; and E8 becomes forced at the gluing (the μ_2 tower, the wrong μ_4 class and the wrong deck twists all rip). Stage 3 (done): `[verification/v717_moonshot_state.py]` (21/21, STATE-ON-TRUNCATIONS) — the Weil functional is constructively a GNS vector state on every truncation (Levinson PD); the naive tower reading misses by $|\Delta|/W3\text{-margin} = 104\text{--}1563$ — the trace-formula boundary is the localized RH substance; KMS $\beta = 1$ (Bost–Connes critical) measured. Stage 4 (done): `[verification/v718_moonshot_spectral.py]` (10/10, SPECTRUM-CONVERGES-MEASURED) — the truncation spectra converge measurably onto the zeros: 100% of 377 frozen targets at tol 0.25 ($h = 1433$), rate -1.61 , monotone GNS K-ladder, SHA256 prediction firewall; the scramble control dies as a state. The Satz-1 slice (done): `[verification/v719_moonshot_traceformula.py]` (24/24, SATZ1-LEDGER-EXACT) — the finite trace formula is exact Gauss quadrature and the term dictionary to the classical Weil formula closes (pole/atoms/arch/UV); the E8 census over $\{\mathbb{Z}^8, E8\}$ (Mordell 1938): only the Gaussian E8 glues. The K collapse (done): `[verification/v720_moonshot_k2k3.py]` (14/14, K2K3-COLLAPSED-CLASSICAL) — the convergence items collapse onto classical analysis (tightness measured, unconditional tail majorant on $s > 1/2$, the boundary exactly $s = 1/2$); and `[verification/v721_k1_node_capture.py]` (14/14, K1-LEMMA-CAPTURE-ONLY) makes the capture half of the node question proof-near. Kill audit (criteria from the strategy council, adjudicated): (K1) does not fire — the gluing needs no zeros (SHA256-typed diagnostics); (K2) half-standing — the $\sqrt{n}$ half-density stays declared, its arch mirror $\log \pi$ is now typed as the self-dual normalization; (K3) does not fire at measurement level — the counterfactual census rips (the uniqueness theorem stays open); (K4) does not fire — positivity holds on every deployed window. The remaining substance, ordered: (L1) identification of the vague limit at the line $s = 1/2$ (no mass loss); (L2) node convergence as a theorem

(the $\sim 900\times$ super-resolution — the K1b question, exploration running); (L3) routine lemmas (tent/Jackson discretization, PNT tail majorant — both exhibited with constants). A research contract, *not* a claim; the contract stays OPEN; no marker move; no RH statement. [2026-08-03 update IV, the keystone round — the contract at its final state of the day. Eight further modules ([verification/v727_l1_identification.py] through [verification/v734_s1_canonical.py]) localize the remaining substance maximally. (L2) is resolved classically: the $\sim 900\times$ super-resolution is atom pinning of the raw measure — the memo's locality/symmetry mechanism dies on two preregistered bars ([verification/v728_k1b_superresolution.py], 11/13, K1B-ATOM-PINNING; lab rate $K^{-1.84}$, real correlate 0.64), and the explicit pinning lemma is certified with one-line proofs and 0 violations at 598 peaks/535 zeros including all 377 of the largest window ([verification/v730_strat2_pinning_lemma.py], 12/12); the certificate saturates at window width — sub-width certification needs atomicity beyond every window depth. The K1 diagram closes: capture theorem (v721) + pinning lemma (v730) + conditional gap rigidity ([verification/v731_strat2_gap_universality.py], 5/5, GAP-UNIV-KILLED for the strong measured form; a fixed-window conditional candidate remains via Lubinsky/Levin–Lubinsky/Avila–Last–Simon/Totik) + atom structure (= L1). (L1) is typed in four equivalent languages: determinacy of the limit moment problem is classical and the identification chain contains RH nowhere ([verification/v727_l1_identification.py], 11/11, L1-DETERMINED-WALL-TYPED); THE WALL is PD persistence of the window ladder — infinitely many PD windows suffice for RH, each window finitely decidable, measured 9/9; equivalently: moments (Hankel-PD) / state (Levinson-PD) / symbol (Fejér ≥ 0) / Hamiltonian (Krein $l_k > 0$ — [verification/v734_s1_canonical.py], 10/10, S1-CANONICAL-SUZUKI-BRIDGED: exact Krein correspondence at 1.6×10^{-15} , $H_h \geq 0$ measured on all nine windows, boundary phase = γ -side at scale exactly 1; Suzuki JFA 2021 as the fourth anchor, unconditional only for $\omega > 1$). The h -universal quantifier is the RH substance. Honest kills, fenced: naive reflection positivity is dead on the truncations — the FE symmetry buys exactly the atom-free Connes–Consani 2021 regime $|u| < \log 2$ and no more ([verification/v729_strat2_rp_universality.py], 5/5); the direct Sonin-compression transfer is dead ([verification/v732_strat3_sonin_prolate.py], 21/21 construction, SONIN-RANK-GROWS; the CC mechanism itself untouched, retuning forbidden). A first exact bridge into Connes' framework: the σ -descended groupoid maps by an exact weak functor candidate into the S -local periodic-orbit subalgebra ($S = \{\infty, 2, 3, 5\}$, 6859 convolutions at deviation 0, $\kappa \equiv 1$; [verification/v733_strat3_gate0_census.py], 13/13, GATE0-WEAK-FUNCTOR; full-algebra extension open). The literature anchor as the dual council reads it: the wall has three names — Weil 1952, Connes 1999/Connes–Consani 2021, Suzuki 2021 — with Meyer 2005 the proven boundary line (unconditional spectral realization without positivity); TFPT adds the computable zeta-free truncation framework with emergent prime places and per-window decidability. The contract stays OPEN; no marker move; no RH statement.] [2026-08-03 update V, the closing slice — the L1 route portfolio is complete. The state-transport route also stagnates ([verification/v735_strat3_ucp_inductive.py], 10/10 construction checks, UCP-STAGNATES): all eight adjacent GNS-ladder maps exist exactly as UCP quantile couplings — the Choi SDP reduces to a finite transport LP, unitality/state preservation to $3.1 \times 10^{-12}/1.2 \times 10^{-15}$, every Choi rank inside the sharp bound $h + h' - 1$ with log-log slope 0.991 (linearly controlled) — but the preregistered shift/grade covariance gate fails honestly (error tail not monotonically decreasing): mere state transport is not the missing L1 mechanism, and the maps are quantile Markov couplings, not $*$ -homomorphisms (multiplicativity defects $\sim 5\text{--}8 \times 10^{-3}$). The portfolio now reads: Sonin dead (v732), UCP stagnates (v735), Gate 0 weak functor (v733), Suzuki bridge a measurement (v734) — what remains for L1 is atomicity/identification substance, not transport machinery. The contract stays OPEN; no RH statement.] [2026-08-05, SHARPENED — the diagonal-Gram closure hands over (PRIME.GRAM.DIAGONAL.01 is closed per its own frozen stakes; the qf offensive [verification/v769_qf_contract_necessity.py] / [verification/v770_qf_spectral_bundle.py] / [verification/v771_qf_representation_census.py] / [verification/v772_qf_feshbach_effective.py] / [verification/v773_qf_cell_cocycle.py]). The sharpened target form: construct a self-adjoint geometric bulk operator with a canonical finite-dimensional boundary/threshold structure whose Weyl M -function delivers the qf block and

whose relative trace formula reproduces the full Weil formula. The measured handover content — mandatory constraints any Z1 candidate must carry: (i) the Hecke/E8 bulk with the archimedean lift ([verification/v714_moonshot_hecke_groupoid.py] / [verification/v716_moonshot_arch_glue.py]); (ii) the moving-edge threshold structure with its measured cadence — mode entries at $M \simeq 888/992/1108/1276$, ΔX widening $1.625 \rightarrow 2.625$ (spread 0.381, not regular at the 0.25 bar), avoided crossing 0.0039 at $M = 1240$ ([verification/v772_qf_feshbach_effective.py]); (iii) the ram-odd contact direction ($\cos \theta_1 = 0.997$ rung-stable, [verification/v771_qf_representation_census.py]); (iv) the settled per-function coupling levels (R2 family 0.225–0.358, R1 family 0.008–0.079, [verification/v770_qf_spectral_bundle.py]); (v) the cell-ordered prime transport in domain-preserving (Nevanlinna/Loewner) form — the exact Möbius/Redheffer cell cocycle of [verification/v773_qf_cell_cocycle.py]. Kills, active: an operator defined through zeros or an RH-equivalent norm assumption = renaming (unchanged); additionally, a candidate whose threshold structure cannot carry the measured widening cadence, or whose M -function block contradicts the settled qf coupling levels or the ram-odd contact direction, is dead on arrival. The contract stays OPEN; no marker move; no RH statement.] [2026-08-05, EXECUTED — the compactness split measured ([verification/v780_z1_compactness_trilogy.py], one module from three preregistered probes; gates $3/4 + 1/4 +$ legs $3/6$ with exactly the preregistered fails). The N2 GNS/Jacobi family carries the ram-odd contact ($\cos$ up to 0.9971), edge collisions at the source depth (0.0032 vs 0.0039) and settled couplings (14/14 P) natively — but not the widening cadence (its edge census is regular quadrature filling, the free-Jacobi type): the cell structure must be imported as boundary data. The finite boundary triple ($A = 2 - J_K$, imported Gram band) is exactly Herglotz through all typed transitions but dies as a whitened $K = M/2$ import — the Gram near-kernel packets are delocalized (cell-mass exactly 0.500 beyond the half-window), so whitening turns truncation loss into quasi-generic boundary noise (delivery 5.88; still arithmetic-specific: 5.9 vs GOE 13.0 vs random 132.5; the full-depth frame is provably out of reach by deepening, $e^{27.75}$ atoms). With the raw μ -weighted import the variable-edge Herglotz family is bounded (0.936) and equicontinuous (0.807) with summable $\sim$ rank-one entry poles at the arithmetic thresholds (c_{mass} 0.032 $\ll$ 1; rank ratios 0.80–0.87), while the delivery metric loses discrimination (typed UNRESOLVED-BY-THIS-METRIC) and the moment functionals do not settle (osc 22.1). The measured unification: this contract and PRIME.KMS.INDUCTIVE_STATE.02 share their compactness theorem (carried: Helly/Montel at finite level) and share their single remaining obstruction — state selection equals import-faithful boundary coupling (a coupling of the Gram band to the bulk that does not pass through the lossy half-window frame). The merged target is named Z1-COMPACTNESS (PRIME.Z1.COMPACTNESS.01); both rows stay, cross-referenced; any future module must attack the shared delta, not the two contracts separately. The contract stays OPEN; no marker move; no RH statement.]

The review master contract (PRIME.KMS.INDUCTIVE_STATE.01). OPEN (registered 2026-08-04, from the external review of the strategy portfolio): extend the finite Hecke functor of Gate 0 ([verification/v733_strat3_gate0_census.py], 13/13, GATE0-WEAK-FUNCTOR — exact on the periodic-orbit subalgebra, 6859 convolutions at deviation 0, $\kappa \equiv 1$) to a positive global KMS state ω at $\beta = 1$ on the closed groupoid algebra, with embeddings $\iota_h: A_h \rightarrow A_{E8}$ and $T_h = \omega \circ \iota_h$ — then Weil positivity follows for the dense test space, and with it RH. The five separation steps, each individually named and decidable: (1) algebraic extension from the periodic-orbit subalgebra to the full convolution algebra; (2) topological closability; (3) complete positivity; (4) the KMS condition at $\beta = 1$; (5) the Weil-limit identification. Kill criterion: complete positivity fails already on a finite extension — one finite counterexample retires the route. The contract names the gap; it does not close it. A research contract, not a claim; no marker move; no RH statement. [2026-08-04 update: the strong form is DEAD and the contract is refined. [verification/v740_kms_extension_switch.py] (8/8, KMS-EXTENSION-DEAD): $\beta = 1$ KMS is algebraically impossible on Gate-0's nonzero-grade Laurent unitaries — $T(u_p u_p^*) = 1$ versus $T(u_p^* \sigma_i(u_p)) = 1/p$, exact defects $1/2, 2/3, 4/5$ at $p = 2, 3, 5$ — and exact covariant $*$ -restrictions do not exist (0 of 141..1255 required boundaries); local extended Gram positivity survives (55,296 monomials/window). [verification/v741_kms_toeplitz_semigroup.py] (9/9,

KMS-TOEPLITZ-DEAD): the Bost–Connes semigroup Toeplitz correction repairs the algebraic KMS step completely (boundary-corrected identities exact; normalized bare error $10^{-2} \rightarrow 10^{-9}$ along the directed system), but the covariant inductive step stays dead (the preregistered UCP window covariance is nonmonotone). Separation steps (1) and (4) are thereby settled at the finite level; the surviving demand is restated below as *PRIME.KMS.INDUCTIVE_STATE.02*.]

The refined master contract: operator-system compactness (PRIME.KMS.INDUCTIVE_STATE.02). *OPEN* (registered 2026-08-04, refined from INDUCTIVE_STATE.01 after [verification/v740_kms_extension_switch.py] and [verification/v741_kms_toeplitz_semigroup.py]): keep the *proven* local blocks — the BC semigroup Toeplitz algebra per window (nonunitary isometries with boundary projections), the boundary-corrected $\beta = 1$ KMS identities (exact), extended Gram positivity on the factorized slices, and UCP state transport ([verification/v735_strat3_ucp_inductive.py]) — and state the single remaining global step purely as an *operator-system compactness/subsequence problem*: does the net of window states have a weak-* cluster point that is a positive KMS functional on the closed algebra, *without* demanding a covariant inductive system? *Kill criterion*: a finite obstruction to any cluster point (a uniform positivity violation along a subnet). Do *not* claim a covariant inductive system until a new canonical window map is found. *Literature anchor*: $\beta = 1$ KMS for the genuine Bost–Connes algebra is classical (Bost–Connes 1995); the TFPT burden is the compatibility of the glued window states with one global limit, not local KMS existence. A research contract, *not* a claim; no marker move; no RH statement. [2026-08-04 update: step (3) — *complete positivity* — is *SOLVED* on the factorized ramified 15-label Toeplitz layer. [verification/v756_kms_incidence_stinespring.py] (7/7, KMS-INCIDENCE-CP-COVARIANT): 105 explicit Kraus terms realize the incidence channel $K = B/7$ as a completely positive unital map, with all four compatibilities exact — σ -covariance at deviation 0, half-weight covariance $DU = 2^{-1/2}UD$ to 1.1×10^{-16} , $\beta = 1$ detailed balance to 6.9×10^{-18} , and Gate-0 compatibility to 3.3×10^{-16} . The local candidate state is $\tau_{\text{inc}} \otimes \tau_{\text{Toeplitz}}$ (channel-preserved). The remainder to a global state is exactly the compactness step of this contract: compatible incidence channels at every finite place and level, the interaction with non-label ramified multipliers and the arch lift, a canonical UCP map between the nine window systems, compactness/normality of the limit, and the Weil-functional identification.] [2026-08-04 update (the diagonal gram round): the compactness step now carries *MEASURED* handoff evidence and an abstract Lean closure. *Measured*: the ε -regulated resolvent is the admissible operator-system evaluation with ε -robust falling local defects ([verification/v766_handoff_bulk.py], 17/17, HANDOFF-BULK-CONVERGES); a target-independent growing-frequency PSD source Gram passes the finite ladder at slope -0.369 ([verification/v767_handoff_frequency_gram.py], 6/6, HANDOFF-GRAM-CONVERGES); the quadrature tail terminates algebraically (2.4×10^{-13} , uniform in the window), cross-window compatibility is closed at $\varepsilon = 10^{-3}$ on the deepest reachable ladder, and the source limit is identified with the deployed Weil functional on the frozen battery within 0.0485 ([verification/v768_handoff_tail_weil.py], TAIL-WEIL-PARTIAL+COMPAT-EPS3-CONVERGES). *Abstract*: the finite-stage Gram compactness/diagonal-extraction skeleton is machine-checked in Lean (*TfptCarrier/GramCompactness.lean*, 8 theorems). *THE SURVIVING OPEN OBJECT* is the q_f wall of [verification/v763_yosida_handoff.py] (QF-DEAD): the battery near-kernel pairing is non-Cauchy at honest maximal depth and survives orthogonalization by structural subordination — the weak-* cluster question of this contract now lives in exactly that bounded quantity; the round’s theorem form is registered as *PRIME.GRAM.DIAGONAL.01*. No marker move; no RH statement.] [2026-08-05 follow-up: the q_f object is adjudicated — q_f is representation gauge and the corner-native Gate 3’ is dead ([verification/v769_qf_contract_necessity.py]), the levels settle positive ([verification/v770_qf_spectral_bundle.py]), fixed-d Feshbach closes at a moving spectral edge ([verification/v772_qf_feshbach_effective.py]), and the cell cocycle is domain-only ([verification/v773_qf_cell_cocycle.py]): *PRIME.GRAM.DIAGONAL.01* is *CLOSED* per its own frozen stakes, and the compactness remainder of this contract hands over to the sharpened *PRIME.Z1.OPERATOR.01* (the Weyl M -function / boundary-threshold form with the measured q_f constraints). No marker move; no RH statement.] [2026-08-05, the unification measured

(`[verification/v780_z1_compactness_trilogy.py]`): the weak-* cluster demand of this contract and the operator demand of `PRIME.Z1.OPERATOR.01` are measured to be the same remaining object on the reachable surface. The cluster-existence half is carried: the variable-edge boundary family is bounded (0.936) and equicontinuous (0.807) on the frozen compacts with summable $\sim$ rank-one entry poles (c_{mass} 0.032) — every subsequence has weak-*/locally-uniform cluster points (Helly/Montel at finite level). The selection half is not carried (moment-functional tail oscillation 22.1), and its obstruction is the same measured object that fails the delivery legs: the lossy $K = M/2$ half-window import. State selection (this contract) = import-faithful boundary map (`OPERATOR.01`): one object viewed twice; the merged target is named `Z1-COMPACTNESS` (`PRIME.Z1.COMPACTNESS.01`); both rows stay, cross-referenced. No marker move; no RH statement.] [2026-08-05 add-on: the `Z1-COMPACTNESS` frame is now also the named limit frame of demand (2) of the new `PRIME.POSITIVE_DESCENT.01` contract below — the `GL1`-sector PSD persistence of the manifestly positive packet object is to be decided in exactly this weak-* cluster frame.] [2026-08-06 add-on: the CP extension gate (`[verification/v794_cp_extension_gate.py]`, `CPGATE-UNDECIDED-FALLING`) measures this demand sector-decorated: the $\chi = \text{GL1}$ instance of the four-sector window net is this contract's object (anchor 3.882×10^{-6} at $M = 1176$ reproduced on the same 10^8 comb), “state selection must settle” = “the sector margins never cross”; measured: `GL1` falls power-law ($\beta = -1.69$), χ_4 saturates at $\sim 1.5 \times 10^{-6}$ — the first measured sector floor.] [2026-08-06 round-23 note: the `SELECTION` half is now finite-level solved — Mosco form convergence + Friedrichs minimality on the `v762` dense core replace the failed moment selector on identical rungs (moment oscillation 22.08 vs Friedrichs resolvents 0.0004 / ledger-adjusted Weyl 0.0843 / raw μ -face 0.0000; cofinal-unique 2.5×10^{-5}), and the import diagnosis is completed at form level: nothing is whitened at the core (`[verification/v816_prime_mosco_selection.py]`, `MOSCO-SELECTS`; the new contract `PRIME.MOSCO.SELECTION.01` below). The remaining analytic object of this contract is the sector floor — restated as `PRIME.FLOOR.RATIO.01` below.] [2026-08-06 round-24 note: the floor contract is narrowed — the lock-block floor $\lambda\tau$ now carries a complete certified three-piece lower-bound skeleton on the deployed battery, closed at citation grade for all h up to $\alpha^* \approx 11$ (`[verification/v823_prime_lagrange_floor.py]`, `[verification/v824_prime_floor_skeleton.py]`); battery-relative and α -bounded, still open.]

The positive descent (`PRIME.POSITIVE_DESCENT.01`). *OPEN* (registered 2026-08-05, from the positive-protocol round; finite-level facts machine-checked in `[verification/v791_positive_descent.py]`, 30/30 incl. the five must-fire controls, verdict `DESCENT-PARTIAL`; the declared run-1 $\rightarrow$ run-2 calibration — the falsified linear state `F1`, the falsified all-PSD operator expectation `F2`, the corrected `C3` constant — is carried verbatim in the module provenance). *The finite-level facts*: (i) the packet GNS state on the extended positive register $\mathbb{N}[C_2] \otimes \mathbb{N}[\mathbb{F}_2^4] \otimes \mathbb{N}[\mu_4]$ is positive at every depth *by construction* (a sum of event squares; verified in exact rationals; the falling minimal sector is f_8^2 -weight dilution, structurally unable to cross zero), while the naive *linear* pushforward of the same nonnegative data is *not* a state (`F1`, quantified: -6.6×10^{-2} , depth-flat) — the squared/GNS form is the only positive aggregation found, and signedness is *observer projection*; (ii) the `GL1` character sector of the packet correspondence operator is *bit-identical* to the deployed Weil window (lag identity at 6×10^{-16}) and is its *unique* PSD sector of all 24 (margins $+5.3 \times 10^{-5} \rightarrow +1.2 \times 10^{-5}$ over $X = 4 \rightarrow 10$, reproduced read-only); (iii) every damped/twisted sector breaks because the register-trivial continuum cannot follow the atom leg (the pole-atom cancellation is exposed; localization: arch-only flat, arch+pole falling) — the breaking tensor factor is the *continuum* (e-slot) leg, not C_2 , $\mathbb{F}_2^4$ or μ_4 . *The contract therefore demands three objects*: (1) **sector-adapted continua** (the missing CP-functor data): for each character sector χ the arch/pole layer of the *twisted* channel (the explicit-formula continuum of the χ -twisted L -data; for the f_8 sector: a weight-4 archimedean factor, *no* pole), such that the sector window $T_{\text{cont}}^\chi + \sum_n \text{Re } \chi(x_n) Q_n$ is PSD at finite level — *finite-level falsifier*: exhibiting the f_8 -sector continuum and finding its window non-PSD kills the operator face for good; constructing it PSD promotes the functor from state level to operator level. (2) **sector-PSD persistence** (the `GL1` limit): the weak-* limit — in the `Z1-COMPACTNESS` frame of `PRIME.Z1.COMPACTNESS.01` /

[[verification/v780_z1_compactness_trilogy.py](#)] — of the GL1 sector functional stays in the cone as $X \rightarrow \infty$. This *is* Weil positivity, now typed as: “the sign sector of a manifestly positive packet object stays PSD” — the sign is never controlled directly, only the sector compression. (3) **the carrier intertwiner**: a positive identification, in the GNS limit, of the population register (χ_- readout = the damped Hecke ratio $a_n/\sigma_3(n)$) with the carrier register (χ_- readout = the unit-modulus sign); at finite level this step is *not* CP-trivial (the signed count fails positivity) — the renormalization/amplification limit is the single non-automatic step. *Stop conditions*: no fixed- d variants, no re-gating of the closed diagonal-Gram objects, no RH claim; demand (2) failing at any finite depth = DESCENT-DEAD for the route; demands (1) and (3) without (2) are decorative. *Inherited finite-level falsifier*: any rung where a non-GL1 carrier sector becomes PSD-binding while GL1 stays positive breaks the “GL1 is the unique carrier edge” reading measured here. A research contract, *not* a claim; no marker move; no RH statement. [2026-08-06 update, markers unchanged — the contract stays open for the floor: all three demanded objects are now delivered at finite level. (1-F) The demand sharpens to the continuum functor: $\chi \mapsto \Lambda_\chi = q(\chi)^{s/2} \prod_j \Gamma_{\mathbb{R}}(s + \mu_j(\chi)) L_{\text{an}}$ with (μ, q, pole) given by one closed rule; verified instances GL1 ($q=1$, pole), χ_4 ($q=4$), f_8 ($q=8$), $f_8 \otimes \chi_{-4}$ ($q=16$, Atkin-Li + Fricke), the conductor datum load-bearing at exactly $\ln 2$ and pinned from below by the PSD margins; the mirror-type sectors are structurally non-automorphic and carry no positivity demand ([[verification/v793_f8sector_conductor.py](#)], 15/15 + 16/16). (2-L/2-Z) The limit object acquires its exact form: per automorphic character the positivity of the sector Weil functional W_χ on the union test space V_∞ — $\inf_X \lambda_{\min}(T_{\chi, X}) \geq 0$ — with the net data a CP-extension theorem consumes measured (restriction exactness, interlacing, uniform bounds), and demand (2) at $\chi = \text{GL1}$ identified with the Z1-COMPACTNESS obstruction, sector-decorated; measured trend: no crossing, GL1/ f_8 /twist power-law falling, χ_4 saturating at the first measured sector floor ([[verification/v794_cp_extension_gate.py](#)], 17/17). (3) The carrier intertwiner exists at finite level in Stinespring form with exact Choi and all covariances, its four automorphic channels landing on the four sector windows from one CP map, and the F1 failure retyped as a negative Choi eigenvalue ([[verification/v801_prime_cp_intertwiner.py](#)], 29/29; the infinite-level demands live in PRIME.CP.INTERTWINER.01 below). The remaining analytic core, restated: the per-sector positivity floor (GRH-type, unproven) plus the intertwiner demands D1–D3. No RH statement.] [2026-08-06 round-23 note: selection and uniqueness are now both finite-level delivered — selection by Mosco+Friedrichs ([[verification/v816_prime_mosco_selection.py](#)], PRIME.MOSCO.SELECTION.01) and the unique positive limit by the master theorem ([[verification/v817_positive_descent_master.py](#)], TFPT.POSITIVE_DESCENT.MASTER.01: prime packet state defects summable at 0.701/doubling, $\hat{m}_2(p) = -1/15$ exact for every odd prime; positivity a separate hypothesis by control K_4). The floor itself reduces to the single ratio inequality $\rho = \tau/\tau_{\text{pnt}} > 0$ with a measured $h^{-3/2}$ envelope — the per-sector positivity floor is the one remaining analytic object, restated as PRIME.FLOOR.RATIO.01 below.] [2026-08-06 round-24 note: that remaining object is narrowed — exact Lagrange sum-of-squares identity, the certified fixed pair, the analytic fixed-pair bound, the family exhaustion, and the deep tail closed at citation grade for all h on the deployed battery ([[verification/v823_prime_lagrange_floor.py](#)], [[verification/v824_prime_floor_skeleton.py](#)]; the narrowed contract text below); battery-relative, α -bounded, still open.]

The carrier intertwiner (PRIME.CP.INTERTWINER.01). OPEN (registered 2026-08-06; demand (3) of PRIME.POSITIVE_DESCENT.01 solved at finite level in [[verification/v801_prime_cp_intertwiner.py](#)], 29/29, CP-INTERTWINER-EXISTS). *Object*: the finite-level carrier intertwiner $\Phi(a) = V^* \pi(a) V$ on the arrow $*$ -algebra $M_{15} \otimes \mathbb{C}[\text{tower}] \otimes (\text{packet slots at the odd places})$, V the 105-leg Kraus stack (unitarity $\sum_e E_{x_e x_e} = 7I$ exact), involution = arrow reversal, KMS half-weights $7^{-1/2}$ per leg / $n^{-1/2}$ universal. *Certified at finite level*: CP/unitarity (exact Choi, rank 105, eigenvalue 1/7); $\sigma/\text{deck}/\text{half-weight}/\beta = 1$ covariances exact; the Choi census on the generating family exact (integer Gram factorization); the four automorphic channels of the one map land on the deployed GL1 Weil window (machine precision) and the $\chi_4/f_8/\text{twist}$ $\Gamma_{\mathbb{R}}$ -rule windows (PSD on $X = 4..10$); the ramified conductor emptiness structural. *Demands (open)*: D1 the X -compatible dilation family and its

inductive limit (with the v794 margin trends as the honest obstruction data); D2 normal extension on the Z1-COMPACTNESS limit object (v780); D3 the limit identification of the GL1 channel with the critical-line Weil functional — *which contains RH and is not claimed.* [2026-08-06 round-22 note, D1–D3 unchanged: the intertwiner gains its register chart at the odd places — the affine Gray map, exactly deck-covariant with trivial μ_4 -cocycle ([verification/v804_prime_carrier_gray.py]) — and its 105 Kraus legs are a complete context protocol over the doily with the exact commutant $[B_{45}, B_{60}] = 0$ ([verification/v811_prime_kraus_doily.py]).] *Kill:* a finite positive packet whose required GL1 intertwiner fails the exact Choi census structurally. No marker move; no RH statement.

The Gray register chart (PRIME.CARRIER.GRAY.01). *Registered* 2026-08-06 ([verification/v804_prime_carrier_gray.py], 23/23, GRAY-INTERTWINER-EXISTS; one declared control repair carried verbatim). *Object:* the affine Gray encoding of $C_2 \times \mu_4$ into the R-class torsor, exactly respecting the v786 Packet480 kill (a group isomorphism is impossible — exponent 4 vs 2 — and not demanded). *Certified:* deck covariance exact with *trivial* measured μ_4 -cocycle; the residual a pure $\mathbb{Z}_3$ line twist on the σ -fixed classes; the extended Stinespring GL1 channel re-lands on the deployed Weil window at $\text{rel } 6.0 \times 10^{-16}$. *Demands:* D1 odd-place fiber charts; D2 the $\mathbb{Z}_3$ line twist as crossed-product structure; D3 the infinite-level CP extension (chained to PRIME.CP.INTERTWINER.01). *Kill:* a nontrivial cocycle under the frozen chart. No marker move; no RH statement.

The Kraus context protocol (PRIME.KRAUS.DOILY.01). *Registered* 2026-08-06 ([verification/v811_prime_kraus_doily.py], 21/21, KRAUS-DOILY-PROTOCOL; one declared report-text repair carried verbatim). *Certified:* the 105 Kraus legs = 15 diagonal + 90 off-diagonal over the doily $W(3, 2)$; the canonical $45 + 60 = \text{reversal quotient } (2:1 \text{ over the } 45 \text{ flags}) + \text{fiber expansion } (15 \times 4 = 60 = \text{the v783 stabilizer rays})$; the $Sp(4, 2)$ obstruction certifies no leg-subset partition (the literal partition is spread-relative = the ledger T3 census); the 45 transports quadratic ($P_x P_y = \pm P_{x+y}$), spanning $\text{End}(\mathbb{C}^4)$ at rank 16; the surprise $[B_{45}, B_{60}] = 0$ exactly — $K = \frac{3}{7}E + \frac{4}{7}R$ a direct-sum protocol with E a conditional expectation. *Measured:* the certified spread is *not* σ -invariant. *Named falsifier:* a σ -invariant spread reproducing the commutant split. [2026-08-06 round-23 addenda ([verification/v815_kraus_spread_commutant.py]): the falsifier pair is decided — σ acts on the six spreads as two 3-cycles with zero fixed spreads (outer-class forced; no invariant coarsening), but the direct-sum protocol is σ -stable as a family ($\sum_s B_{45}(s) = 4I + 2B$; per-orbit A_{orb} spectrum $\{0^5, 1^2, 4^5, 7^2, 9^1\}$); $[E, K] = 0$ extends exactly to channel/KMS-tower/odd places (the unique C_2 -character expectation per place) and the 600-dim register; the $\{K, \sigma\}$ commutant is 39-dim nonabelian with A_{orb} its canonical abelian 5-dim subalgebra (PRIME.AORB.REFINEMENT.01), window-silent beyond the K -spectrum; and the rule-A protocol set is canonically the 105 minimal dual checks of $[15, 10, 4]$ ([verification/v820_prime_kraus_rm24.py], PRIME.KRAUS.RM24.01).] No marker move; no RH statement.

The Lorentz spinor coordinates (PRIME.LORENTZ.SPINOR.01). *Registered* 2026-08-06 ([verification/v807_lorentz_nullselector.py], 22 checks with the three honest h -order FAILs encoded, LORENTZ-COORDS-REVEAL). *Exact:* $M(5, -3, 4) = 2(1, 2)^T(1, 2)$ — the compiler triple is a rank-one spinor square, Euclid $(2, -1) \mapsto (5, -3, 4)$. *The reveal:* the locking slope is strictly monotone in the window width α (66/66, Kendall 1.000, cross-ratio contraction CV 0.004) — the h -oscillation was α -jitter; the approach $2 - r \sim \alpha^{-0.23}$ glacial. *Named candidate:* the 1D monotonicity statement in α -order (the three h -order gates stay honest FAILs). *Kill:* a Kendall inversion under the frozen windows. No marker move; no RH statement.

The local-Clifford kill and the mod-8 completion (HECKE.LOCAL.CLIFFORD.01). *Registered* 2026-08-06, the analytic ordering *dead* as frozen ([verification/v806_hecke_local_clifford.py], 16/18 with the two FAILs the preregistered decider gates, controls 4/4, LOCAL-CLIFFORD-BREAKS; one declared bar re-freeze). All four $p \bmod 8$ class sectors fail PSD (signed class indicators carry no completed L -function; the trivial mod-8 component imprimitive); the margin ordering does *not* track the proven activation depths (Spearman +0.40); class 1 the unique positivity-completable class (Euler repair exact). The arithmetic multirate theorem untouched; the Clifford-activation

reading decorative analytically. *Positive side-result*: χ_8/χ_{-8} are two new PSD instances of the $\Gamma_{\mathbb{R}}$ conductor rule (exact parities) — the rule map covers the full mod-8 dual group. *Next falsifier*: primitive sector data, not class indicators. No marker move; no RH statement.

The flavor-filtration fingerprint (FLAVOR.GRAPH.FILTRATION.01). *Registered* 2026-08-06 ([[verification/v812_flavor_graph_filtration.py](#)], 16/16 incl. the four must-fire controls, FILTRATION-FINGERPRINT-ONLY). The $Q \rightarrow K \rightarrow L$ row budgets are exact, order-exact counts of the $K_{3,2}/\text{prism}/K_5/H_8$ incidence chain with level-wise additivity mirroring $L = K + Q$; *no* entry-level reconstruction from any frozen candidate; exhaustive false-positive rates 0.005–0.75% make the misses informative. Typed audit fingerprint — beside the audit identities, in no derivation. No marker move.

The P1 index-KMS identity (P1.INDEX.KMS.01). *OPEN* (registered 2026-08-06; [[verification/v813_p1_index_kms.py](#)], 13/13 incl. the two must-fire controls, P1-IDENTITY-CLOSED). *The identity*: $I \cdot \beta_{\text{angle}} \cdot c_3 = 4 \cdot 2\pi \cdot \frac{1}{8\pi} = 1$ exact in formal arithmetic, each factor independently sourced — the Jones index $I = 4 = |\mu_4|$ a c_3 -free theorem (zero occurrences gated in all four sources), $\beta_{\text{angle}} = 2\pi$ independently measured (v526; the radian pin is the one shared convention and cancels), c_3 the P1 axiom the identity *explains*, not derives. *Measured precursor*: $w_1 = w_3 = \frac{1}{4}$ exactly at every level with symmetric zero-sum offset $\pm 2^{-(l+1)}$ (the no-anomaly shape); the sea state does *not* equidistribute (dev 0.188). *The one open bridge*: the even modular-response normalization over the I deck sectors. *Kill*: any asymmetric sector offset. *Control*: index 2 gives $1/(4\pi)$, booked elsewhere by the corpus. P1 stays a declared input; no marker move.

Mosco selection (PRIME.MOSCO.SELECTION.01). *Registered* 2026-08-06 ([[verification/v816_prime_mosco_selection.py](#)], 23/23 guards+controls, 6/6 legs, MOSCO-SELECTS). *Statement*: selection by Mosco form convergence and Friedrichs minimality on the v762 dense core — the window-form chain q_X has exact nested recovery (structural, 1.2×10^{-12}), vanishing liminf defect (measured; the moving-edge family confirmed a genuine boundary channel, not an escape), Cauchy Friedrichs resolvents and Weyl functions *modulo the typed summable entry-pole ledger*, and cofinal-unique limits (2.5×10^{-5}): the unified compactness contract selects its state by form convergence, not by moment stability (moment baseline 22.08 reproduced exactly in-run). *Completed*: the import diagnosis at form level — the raw μ -weighted coupling of the core is exactly rung-independent, the whitened control reproduces the 5.8772 delivery failure exactly. *Remaining for theorem grade (typed)*: (i) Mosco-precompactness of $\{q_X\}$ on V_{∞} , (ii) identification of the limit form domain, (iii) entry-pole ledger summability at infinity. Positivity/sector floor stays a separate contract. No marker move; no RH statement.

The positive-descent master theorem (TFPT.POSITIVE_DESCENT.MASTER.01). *Registered* 2026-08-06 ([[verification/v817_positive_descent_master.py](#)], 14/14 incl. the four must-fire controls, MASTER-BOTH-INSTANCES; kernel-checked core `TfptCarrier/PositiveDescentMaster.lean`, the matrix positivity closure in `GramCompactness.lean`, cited). *The theorem (seven frozen hypotheses)*: finite operator systems with unital inclusions, positive states, CP transitions, finite character sectors, *summable* state defects, a sector-adapted closure datum, and a Mosco form family force a unique positive limit state with well-defined positive limit sectors; the core (telescoping + quantitative Cauchy tail; positivity closed; sectors commute) is kernel-checked. *Instance G_{net}* : hypotheses (1)–(5) measured (defects 0.267/doubling; the expectation CP via Choi; μ_4 evenness exact); open: normality/split. *Instance prime/packet* — *the new number*: state defects summable at 0.701/doubling $\approx 2^{-1/2}$ (the dilution law), with $\hat{m}_2(p) = -1/15$ exact for every odd prime; open: the sector floor (PRIME.FLOOR.RATIO.01, narrowed in round 24 by the certified floor skeleton v823/v824 — battery-relative, α -bounded, still open). *Control K_4* : summable defects with one non-positivity-preserving step give a non-positive limit — positivity is a separate hypothesis. The selection problem is eliminated on both sides. No marker move; no RH statement.

The floor as one ratio inequality (PRIME.FLOOR.RATIO.01). *OPEN* (registered 2026-08-06 — the reduced sector-floor contract; evidence [[verification/v818_sector_floor_attack.py](#)], 9/13 + 13/15 with exactly the preregistered honest FAILs, FLOOR-MECHANISM-PARTIAL + CAPTURE-BOUNDED

/ INGREDIENT-II-THEOREM-SHAPE / AMPLIFIER-MECHANISM). *Statement:* the sector positivity floor $\inf_X \lambda_{\min}(T_{\text{GLI},X}) \geq 0$ reduces, up to the measured $O(1)$ interlacing capture (median 1.53, range 1.04–4.36; the 2-mode lock block an $O(1)$ -faithful witness by Cauchy interlacing), to the *single* ratio inequality $\rho(X) = \tau(X)/\tau_{\text{pnt}}(X) > 0$ — the prime comb against its own smooth density — with the measured finite lower envelope $\rho \geq ch^{-3/2}$ (constant $c \approx 4.85$, envelope non-decaying, slope +0.14). *Certified/measured ingredients:* the direction $r(X)$ is owned by the density rotation law $dr/dt = -\kappa(w \cdot v_1)(w \cdot v_2)(1 + r^2)/(\lambda_1 - \lambda_2)$ (symbolic, sign hypothesis verified finite-level, 100% staircase sign-match, typed density-level by the Epstein control $s_E = 0.241$); the capture is angle-certified frame geometry ($\cos \theta$ median 0.990, scramble-robust — frame content, not arithmetic); the amplifier ρ is carried by the smallest prime powers (packet correlation +0.964, partially trend-driven, typed). *Kill:* the envelope failing at depth ($\rho h^{3/2}$ decaying below every positive constant) or the capture angle collapsing ($\cos^2 \theta < \frac{1}{2}$ at any rung). Selection and uniqueness are eliminated on both faces (PRIME.MOSCO.SELECTION.01, TFPT.POSITIVE_DESCENT.MASTER.01) — the floor is the single remaining analytic object of the prime programme. Interlacing runs floor $\Rightarrow$ margin (necessary, not sufficient); promotion of any ρ bound is *fenced* on the preregistered envelope + capture gates. [2026-08-06 round-24 narrowing — *what is now closed:* on the deployed 14-rung battery the lock-block floor $\det \hat{A}_2 = \lambda \tau$ has a complete three-piece uniform certified lower-bound skeleton — the *exact* Lagrange sum-of-squares identity over zero+pole rank-one carriers (machine wards $\leq 1.4 \times 10^{-9}$), the analytic fixed-pair bound $X_\infty^2(\alpha)(1 - C(\alpha)/h)$ with explicit $h_0(\alpha) \leq 500$ (h -uniform at every deployed α in-domain), the certified top-100 family exhaustion (0.93–0.97 of the floor, all carriers on-line $\leq 2 \times 10^4$), and the psd remainder closed at *citation* grade (product-sup envelope + Abel/RvM explicit constants; $T_{\text{ver}} = 3 \times 10^{12}$) for all h at the deployed α , with an h -free growth law ([verification/v823_prime_lagrange_floor.py], [verification/v824_prime_floor_skeleton.py]; PRIME.FLOOR.LAGRANGE.01 / PRIME.FLOOR.SKELETON.01 below). *What remains open — the contract:* the floor bound is *battery-relative* and α -bounded — (i) the citation-grade chain re-crosses X_∞^2 at the validity horizon $\alpha^* \approx 11.2$ (deeper α needs verified on-line *depth* beyond T_{ver} , never location); (ii) the fixed pair dies at the α -nodes of $\sin(\alpha\gamma_1)$, where the family limit (positive, minimum 1.62×10^{-4}) must take over — the family version of the analytic bound is not yet written; (iii) the product-sup Lipschitz certificate is a named elementary step, typed, not claimed; (iv) *nothing* bounds $\inf_X \lambda_{\min}$ on V_∞ — the $O(1)$ capture and the $\rho > 0$ demand at infinity remain exactly this contract. Kill criteria unchanged.] No marker move; NO RH claim.

The floor as an exact sum of squares (PRIME.FLOOR.LAGRANGE.01). *Registered* 2026-08-06 ([verification/v823_prime_lagrange_floor.py], 7/8 with the one preregistered-honest old-budget FAIL + 9/9, LAGRANGE-CONCENTRATED + PAIR-CERTIFIED; discovery probes `prime_lagrange_pair_probe.py` and `prime_lagrange_budget_probe.py`). *Certified (identity-grade wards):* every per-zero layer of the v692 master identity is exactly rank-one ($v_\gamma = 2\sqrt{w(\gamma)}(S_1, S_2)$), closed trigonometric reads, layer deviation $\leq 1.2 \times 10^{-16}$, and $\det(G_Z + P)$ is an exact sum of squares over the component pairs (Lagrange; ward $\leq 1.4 \times 10^{-9}$ on all 14 rungs); the pole is the universal non-collinear leg and the moving zero leg sits on the alias comb $\gamma \approx 2\pi k/D$. *Measured (per-rung, float-level):* $\lambda_{\min}(\hat{A}_2 - G_Z - P) \geq$ margin verified per rung eliminates the tail estimate — budget tightened a median 9.8 orders — and the certified interval $X^2(\text{pole}, \gamma_1)$ – BUD is strictly positive on 14/14 rungs, with the floor share growing as $h^{+1.06}$. *Honest typing:* the atom-side rank-one framing fails as frozen — the SOS realization lives on the zero+pole side; the psd of the remainder is verified numerically here, its citation-grade upgrade is PRIME.FLOOR.SKELETON.01. *Kill:* any ward failing on the frozen ladder. Feeds PRIME.FLOOR.RATIO.01 (narrowed). No marker move; no RH statement.

The floor-theorem skeleton (PRIME.FLOOR.SKELETON.01). *Registered* 2026-08-06 ([verification/v824_prime_floor_skeleton.py], 12/12 + 9/9, FLOOR-PARTIAL + TAIL-CLOSED-ALL-H; discovery probes `prime_floor_theorem_probe.py` and `prime_tail_envelope_probe.py`). *Uniform-verified on the deployed battery:* the analytic fixed-pair limit $X_\infty(\alpha) = 16\alpha\pi^2 \sin(\alpha\gamma_1) \sinh(\alpha/2)$ [bracket] with the explicit bound $L(h) = X_\infty^2(\alpha)(1 - C(\alpha)/h)$ for $h >$

$h_0(\alpha) = C(\alpha) \leq 499.6$ (zero sign flips; 25 α -nodes typed — h -uniform per α , not ladder-uniform in α); the certified top-100 family exhaustion (median 0.952 of the floor, carriers on-line $\leq 2 \times 10^4$; residue 0.015–0.049, non-growing; the analytic family limit positive across all nodes). *The tail closure*: the product-sup per-zero envelope + Abel/RvM with explicit unconditional Trudgian-grade constants close the psd remainder at citation grade on *all 14 rungs* at the fixed verified horizon $T_{\text{ver}} = 3 \times 10^{12}$ with margins 10^3 – 10^6 ; the growth law is h -free ($+1.98 \rightarrow +0.13$), the remaining growth α -only, validity horizon $\alpha^* \approx 11.2$ (the Vinogradov–Korobov region typed honestly useless). *Net*: $\lambda\tau \geq X^2(\text{pole}, \gamma_1) - \text{pert}(h) > 0$ at citation grade — no per-rung eigencheck, no zero-location input beyond the critical strip. *Honest boundaries*: battery-relative, α -bounded, necessary-side; the sup Lipschitz certificate a named elementary step; nothing bounds $\inf_X \lambda_{\min}$ on V_∞ . *Kill*: any closure margin failing on the frozen battery at the frozen horizon. Feeds PRIME.FLOOR.RATIO.01 (narrowed). No marker move; no RH statement.

The Reed–Muller identification (PRIME.PACKET.RM14.01). *Registered* 2026-08-06 ([verification/v819_prime_packet_rm14.py], 21/21 incl. the three must-fire controls, RM14-EXACT; Lean companion TfptCarrier/PackageRM14.lean, 30 theorems). *Certified*: the 15-label channel row hull is the punctured $\text{RM}(1, 4)^* = [15, 5, 7]$ (enumerator $1 + 15z^7 + 15z^8 + z^{15}$); $C_0 = [15, 4, 8]$ simplex; χ_{NSR} one codeword (7+8 its two sides, C_2 the affine bit); the self-reproduction cycle closes with exactly 1344 equivalences; the CSS code $[[15, 1, 3]]$ with dual $[15, 10, 4]$, logical distances (7, 3), triorthogonality $8/4/2$. *Typed [H]*: the ζ_8 /T-gate phase pointer (v798); the TFPT fingerprints ($14 = \dim G_2$, $3 = N_{\text{fam}}$, $4 = |\mu_4|$), audit-only. *Kill*: any census identity failing on the frozen frame. No marker move; no RH statement.

The canonical check rule and the plane frames (PRIME.KRAUS.RM24.01). *Registered* 2026-08-06, carrying PRIME.C2.PLANEFrames.01 ([verification/v820_prime_kraus_rm24.py], 16/16 + 14/14, KRAUS-RM24-RULE-CANONICAL + PLANEFRAMES-EXACT; one honest coset-census fix carried). *The honest double kill*: the literal leg-support claim is dead twice (weights $1/2/3 < \text{distance } 4$; $Sp(4, 2)$ orbits $15+90$ vs $45+60$ — no equivariant bijection exists). *Certified*: the canonical ω -built rule identifies the rule-A protocol set with the 105 weight-4 checks of $[15, 10, 4]$ exactly, $Sp(4, 2)/\sigma/\text{deck}/\text{KMS-covariant}$; the six Gram identities entrywise; $A_3 = 12 / B_3 = 16$ as per-point plane completion counts ($\sigma_3(3) = 28$, $a_3 = -4$, independent eta-product rebuild); the graded frame intertwiner $V^*V = \sigma_3(n)I$, $V^*GV = a_n I$ exact for $n = 3, 5, 7$. *Kill*: an equivariance mismatch of the ω rule under the frozen generators. No marker move; no RH statement.

The vacuum completion (PRIME.VACUUM35.01). *Registered* 2026-08-06 ([verification/v821_prime_vacuum35.py], 22/22 incl. the three must-fire controls, VACUUM35-EXACT). *Certified*: $140 = 105 + 35$; the Grams $22I+6J / 6I+J / 28I+7J$ entrywise with spectra $\{112, 22^{14}\} / \{21, 6^{14}\} / \{133, 28^{14}\}$; the E_7 completion $112 + 21 = 133$ as a matrix identity (the $E_7 \times A_1$ corpus reading a [C] fingerprint, anchor v170); Weitzenböck $H_{105}^T H_{105} = 2B^2 + 14I$ as the canonical label-side left factor (v758's continuum wall explicitly untouched); the $\pm B$ cancellation inside one Gram; the tight frame constant 28 on mean-zero; $\text{RM}(2, 4)$ dual-containing. *Typed [H]*: the vacuum/continuum reading — decided negatively by the closure row below. No marker move; no RH statement.

The vacuum-route closure (PRIME.CONTINUUM.UNSHORTEN.01, carrying PRIME.VACUUM.DILATION.01). *CLOSED* — *honest double kill* (registered and decided 2026-08-06; [verification/v822_prime_vacuum_dilation.py], 17/19 + 12/14 with the four FAILs being the preregistered PSD deciders themselves, controls $3/3 + 3/3$, UNSHORTEN-DEAD + DILATION-DEAD; first runs after freeze, no repairs). *The dead reading*: the archimedean/pole continuum as the $\text{RM}(2, 4)$ vacuum completion of the 15-label window objects — dead in *both* transcriptions: the one-slot bordering by the v758 mechanism verbatim and on scale (with the shape theorem: one slot induces only a rank-one pure- J coupling; the demanded $6I+J$ needs 35 separate rows), and the corrected target itself — the 35-row Stinespring dilation, shape fixed exactly, $\times 13.1$ distribution gain — still failing both sectors under both frozen normalizations. *The frozen fence (the round's quantitative product)*: any continuum re-entry through the RM check geometry is PSD iff its coupling satisfies $c \leq c^* = (2/21) \lambda_{\text{pencil}}(T_{\text{dep}}, T_{\text{cont}}^2)$; measured

deficits $\times 3 \times 10^7$ (GL1) / $\times 2 \times 10^2$ (f8) against the mass calibration — the reading dies *with the calibration*, not the geometry: the continuum's completion role is already exhausted inside T_{dep} (the explicit-formula balance), and the path-state construction is not buildable on this gating. The RM integer layer stays certified. No marker move; no RH statement.

The PD-persistence universal form (PRIME.PD.PERSISTENCE.01). *OPEN* (registered 2026-08-04): the universal-theorem form of the PD-persistence wall ([[verification/v727_l1_identification.py](#)], [[verification/v734_s1_canonical.py](#)]) with the important correction $\lambda_{\min} \geq 0$ instead of $\geq \delta$ — the Ihara lesson (FTR.IHARA.01): true RH analogues can have *exact zero modes after rank saturation*, so a uniform margin is stronger than necessary and the wrong instrument. *Target form*: cofinally infinitely many PD windows, established via a state / Gram representation / cone-preserving recursion — *not* via eigenvalue bounds. *The review blacklist, confirmed as permanent kills*: the raw symbol minorant, norm triangles, perturbation theory, position-blind estimates, the direct Sonin transfer ([[verification/v732_strat3_sonin_prolate.py](#)]) and naive reflection positivity ([[verification/v729_strat2_rp_universality.py](#)]) stay retired; further zero-matching records do *not* change the quantifier. A research contract, *not* a claim; no marker move; no RH statement. [2026-08-04 update (the incidence programme): the contract now carries a REGISTERED INDUCTION FORM, a CLOSED route, and a NAMED open object. (i) The induction form: the ramified Schur ladder on the canonical tower ([[verification/v755_simpler_schur_recursion.py](#)], 13/13, SCHUR-RECURSION-ALIVE) — $S_k \succeq 0$ measured on all 8 ram-odd stages ($n_k = 2^{2k+1}$, $\lambda_{\min}$ flat, long-double-hardened), with basis, Douglas range condition, Albert/Haynsworth step and cofinality all exact: the open proof need is exactly the structural Gram certificate $S_k = R_k^T R_k$ per step; the measured boundary data for any candidate R_k are recorded (the label part enters inverted, $\rho_k < 0$ with $|\rho_7|$ within factor 1.34 of 49/4 — cross-term structure, not an added block — and continuum vs atom sides are order-1 comparable). Honestly: for ζ data this is equivalent to tower PD — the wall in recursion shape. (ii) The closed route: the review's assembled certificate [[continuum root ⊕ Kraus](#)] is not mountable ([[verification/v758_simpler_certificate.py](#)], 8/8, CERT-CONTINUUM-ROOT-DEAD, the keystone negative): the arch+pol block is PSD on no stage (Levinson break at cell 47, exactly between the $n = 2$ and $n = 3$ cells, ladder-wide), the continuum Schur complement is no Haynsworth object, the additive split fails on every stage (the order-1 balance sits in the cross-term), and the rescue cascade is strictly cell-ordered — the wall IS the missing left factor. (iii) The named open object: the corrected target shape is the cell-wise cascade factorization $R = \prod(\text{cell factors})$ — rescue order $n = 2, 3, 4, 5, \dots$ across the channels — with the exact building blocks of the incidence programme ([[verification/v752_projective_hamming_incidence.py](#)], [[verification/v753_ramified_polarity.py](#)], [[verification/v754_ramodd_twostep.py](#)], [[verification/v756_kms_incidence_stinespring.py](#)]) as the label-side material. No marker move; no RH statement.]

The diagonal Gram closure theorem (PRIME.GRAM.DIAGONAL.01). *CLOSED* (registered 2026-08-04, the diagonal gram round; closed 2026-08-05 per its own frozen stakes — an honest negative adjudication, not a marker upgrade; the full closure cascade is the dated note at the end of this block). The target form, kept verbatim for the record: construct, for the countable dense test family $\{\phi_i\}$ of the canonical (D, X) tower ([[verification/v762_dense_weil_core.py](#)], 26/26, DENSE-CORE-CANONICAL: frozen enumeration, explicit stage map, admissibility located per class), PSD source matrices G_n without any target decomposition, zero data, or positivity test of the target, satisfying

$$\max_{i,j \leq n} |(G_n)_{ij} - W(\phi_i * \tilde{\phi}_j)| \leq 2^{-n};$$

then every fixed m -corner of W is a PSD limit of PSD matrices — Weil positivity on the rational span of the family, and the density step closes by the classical approximation argument. *The round's measured decision state*: Gate 1 positive — the handoff error is boundary/cutoff, not quadrature ([[verification/v759_handoff_fixed_window_resolution.py](#)], RESOLUTION-BOUNDARY); Gate 2 positive — the anti-alias identities are exact with sharp count $N_{f,\min} = 2M - 1$ ([[verification/v760_antialias_exact.py](#)], ANTIALIAS-CORRECTED) and the paired atom+pole remainder carries the explicit unconditional vanishing bound η_K with *no RH-strength input*

(`[verification/v761_atom_pole_abel.py]`, combined ABEL-PAIRED-BOUND); Gate 4 positive — the family is canonical (`[verification/v762_dense_weil_core.py]`); Gate 3 *negative on the canonical class* — the battery near-kernel pairing q_f is non-Cauchy at honest maximal depth and survives orthogonalization by structural subordination (`[verification/v763_yosida_handoff.py]`, QF-DEAD): *the q_f wall* — density and near-kernel avoidance collide on the same function class — is the named wall object; Gate 5 (the assembled diagonal certificate, the planned **v764**) was *not executed* — its precondition (theorem-capable bounds from Gates 2 and 3) failed via Gate 3; the red team is green (`[verification/v765_handoff_redteam.py]`, REDTEAM-GREEN-COMBINED, with the binding note that certificates must always carry *both* gate families — symbol/PSD and rate/ratio/identification). *Kill criteria that did NOT fire*: no RH-strength input was needed anywhere (the **v761** circularity flag and the **v763** KILL audit are machine-checked: no $1/\varepsilon$ bound, no PD-margin assumption); no spectral gap was assumed (exact zero modes stay admissible — the Ihara lesson). *The binding stop-list* (ten items, frozen): no atom/prime-layer positivity induction; no separate absolute atom/pole bounds; no 16-dimensional channel square sums; no exact neighboring-window embeddings; no unbounded A^{-1} observables; no Ward-drift signs; no rate fits as proof substitutes; no zero-matching records as progress metric; no positive continuum root as certificate factor; no cellwise PSD demands. A research contract, *not* a claim; no marker move; no RH statement. [2026-08-05 closure note — the route is CLOSED per its own frozen stakes. The final decision cascade, full tally: Gates 1/2/4 positive from the diagonal gram round (above); Gate 3 DEAD AT THE CORNER LEVEL — the audit pair (`[verification/v769_qf_contract_necessity.py]`, combined QF-GAUGE/GATE3PRIME-DEAD) first proves the full-ladder Cauchy demand on q_f was stronger than the contract needs (q_f is representation gauge: a deterministic ε -net subsequence of density 0.144 makes all 14 profiles Cauchy at the parent bars, and the contract's own E_Y corner objects are blind to q_f jumps, corr +0.038, median $\rho = 1.13$), and then the corner-native weakened Gate 3' dies on its own predeclared 1.6×10^7 comb ($X = 16.5625$): the corner increments fall to $X \sim 12.9$ and then rise sustained (+0.10..+0.34 per X unit at $\varepsilon = 10^{-3}$, 0/56 cells) — the **v764** precondition is refuted; Gate 5 / **v764** therefore was never executed. The q_f offensive's own line: gauge $\rightarrow$ corner rise (`[verification/v769_qf_contract_necessity.py]`) $\rightarrow$ settled-positive levels (`[verification/v770_qf_spectral_bundle.py]`, QF-SETTLES-POSITIVE: the band weights plateau at positive per-function constants, R2 family 0.225–0.358, R1 family 0.008–0.079; zero-bar depth $X^* \sim 227 = \text{never}$) $\rightarrow$ no fixed-d object (`[verification/v772_qf_feshbach_effective.py]`, FESHBACH-PARTIAL/EDGE-KEEPS-MOVING: the moving spectral edge — avoided crossing 0.0039 at $M = 1240$ on the 10^9 comb, widening non-regular entry cadence $\Delta X = 1.625/1.8125/2.625$, spread 0.381) $\rightarrow$ cocycle domain-only (`[verification/v773_qf_cell_cocycle.py]`, COCYCLE-DOMAIN-ONLY: exact identity and zero-breach domain transport, 0/6 convergence cells — structure without limit). What survives as theorem-shaped facts, each typed in its own module row: the exact anti-alias theorem (`[verification/v760_antialias_exact.py]`); the unconditional paired Abel bound (`[verification/v761_atom_pole_abel.py]`); the canonical dense family (`[verification/v762_dense_weil_core.py]`); the exact source-native domain-preserving monotone Möbius/Redheffer cell cocycle through mode entries (`[verification/v773_qf_cell_cocycle.py]`); the Kato bundle frame (`[verification/v770_qf_spectral_bundle.py]`); the Herglotz-certified exact Feshbach reduction (`[verification/v772_qf_feshbach_effective.py]`). NO new variants; the ten-item stop-list stays binding; the sharpened PRIME.Z1.OPERATOR.01 takes over with the measured handover content [2026-08-05: executed as the **v780** trilogy — compactness carried, selection the one shared open object, merged target Z1-COMPACTNESS]. Lean legacy (2026-08-05): the exact cell-cocycle core of the closed route is machine-checked as general Mathlib theorems over $\mathbb{R}$ — `TfptCarrier/CellCocycle.lean`, 18 declarations: the exact Banachiewicz/bordered-Schur update identity, PD cell block $\Rightarrow$ PSD corner correction $\Rightarrow$ monotone Loewner flow with domain preservation, bounded-monotone entrywise matrix convergence (the theorem names the missing analytic ingredient precisely — a uniform Loewner upper bound, which the finite ladder did not certify and which is not claimed), and the Redheffer/linear-fractional composition laws over arbitrary commutative rings. No RH statement.]

The structural upgrade: the canonical nested tower replaces the historical window

family (PRIME.TOWERNEST.01). The window quantifier of the PD-persistence formulation reduces *exactly* to the canonical nested (D, X) tower ([verification/v749_simpler_tower.py], 8/8, TOWERNEST-REDUCES): in X an ascending chain — X -nesting exact to 6.7×10^{-18} , so limit positivity is *one* limit object per chain — and in D dyadic downward inheritance, exact: PD fine *forces* PD coarse; every deployed window is a tower member at deviation 0.0. The window quantifier was an *artifact* of the historical selection: box step functions are dense, so tower-PD for $D \rightarrow 0$, $X \rightarrow \infty$ is equivalent to the Weil criterion (classical density step). PRIME.PD.PERSISTENCE.01 is therefore restated with the canonical truncation in place of the historical window family. *Honest limits:* incommensurable root- D chains connect only by approximate interpolation (the residual does not close the PD transfer), and the margins *fall* along the tower — the wall stands, in canonical nested form. *Companion diagnostics of the simplification trilogy:* the gluing identity is an identity-level *off-line detector* — it sees $\delta = 0.01$, sharper than the PD break at $\delta^* = 0.02$, and the envelope slope measures δ — while the rigidity thesis is honestly false (the identity holds unconditionally; [verification/v750_simpler_gluing.py], 5/5, GLUING-DETECTOR); and the component running is deterministic in direction with the exact $\mp u/2$ drift anatomy from $p^2 \equiv 1 \pmod{4}$ and a slope-Ward cancellation at 4×10^{-4} — the margin inherits *zero* deterministic drift — but the balance *sign* is zero content (budget $2.9 \times 10^4 \times$ the margin): the induction route is RH-circular at the balance level ([verification/v751_ward_monotone.py], 15/15, WARD-MONOTONE-MIXED). The cone-preserving layer recursion is retired by a one-liner: cone-preserving iff layer-positive, and no prime layer is ($c_0 = 0$; [verification/v743_schur_cone_recursion.py], 7/7, SCHUR-CONE-DEAD). No marker move; no RH statement.

The four fixed-size angle instruments (PRIME.ANGLE.INSTR.01). The instrument cores of the discovery parts T155 and T157, recomputed on the same declared surface as PRIME.RITZ.CEIL.01 (the 12 deepest in-cap frame-A windows, $n = 29 \dots 139$, $m = 96 \dots 291 \leq 300$; a no-go size ladder $m = 48 \dots 288$; parity and Dirichlet controls at $m = 64, 128, 256$; each statement with at least one mutation control that must fail loudly) — none of the four closes an angle, and none is promoted as anything but a per-instance instrument. (1) The 12×12 complement-floor certificate: $\min_{v \perp W} v^T L_P v \geq \mu_{13}^P - \lambda_{\max}(M^{1/2}(I - GG^T)M^{1/2})$ with $M = \text{diag}(\mu_{13}^P - \mu_k^P)$ and $G = T_{12}Q_W$ — an identity plus *one* certified Rayleigh bound on the high block, a theorem for every m and every subspace, direction checked: the certificate never exceeds the exact size- m complement bottom (12/12) and agrees to $1 - \text{ratio} \leq 1.01 \times 10^{-3}$ against the declared cap 2×10^{-3} on this small surface; exactness controls both ways ($W = \text{span}\{t_1..t_8\} \rightarrow \mu_9^P$; the *worthless* configuration $W = \text{span}\{t_2..t_{13}\} \rightarrow \mu_1^P$; both to 2.0×10^{-11} — the instrument reports its own worthlessness); a corrupted overlap row moves the certificate by ≥ 0.549 , only downward. (2) The sine-block confinement: from the certified pencil floor $A \succeq tL_P$ and the certified ladder ceiling $\lambda_1 \leq S\mu_1^P$ *alone*, $\|\gamma_H\|^2 \leq \lambda_1/(t\mu_{17}^P) \leq (S/t)/\rho_{17} = 0.0153\text{--}0.0245 < 1$ on 12/12 — the bottom eigenvector lives 97.5–98.5% inside the first 16 parity sines by a per-instance theorem (measured tail $6.7 \times 10^{-6}\text{--}3.0 \times 10^{-5}$; the discovery part’s measured “98 per cent” replaced by one line); negative control: the pencil ceiling κ in place of the ladder S is vacuous ($17.6\text{--}477 > 1$ on 12/12); mutation: the inflated floor κL_P is refuted at $e_1(A)$ by $3.3 \times 10^3\text{--}1.0 \times 10^5$. (3) The Schur-entry identity: $s = \mu_1^P t_1^T A^{-1} t_1 = (B^{-1})_{11}$ exactly (1.1×10^{-11}) and $(B^{-1})_{LL} S_L = I$ to 5.8×10^{-11} with $S_L = B_{LL} - B_{LH} B_{HH}^{-1} B_{HL}$ the *same* fixed-size 16×16 Schur complement the chain already forms, with the literal-entry direction $1/s \leq (S_L)_{11}$ on 12/12: $1/s = 4.0460\text{--}5.8962$ is an $O(1)$ number carried by one entry, so the R2'' loss ceiling $r \leq 1/(Ls)$ is a statement about *one* number; negative control: the inversion-free Cauchy–Schwarz ceiling is valid and useless, off by $250\text{--}7.27 \times 10^3$ (the cancellation in $b^T B_{HH}^{-1} b$ is nearly complete); mutation: a one-index shift breaks the identity by $1.55\text{--}2.15$. (4) The adaptive Lipschitz ceiling: interval bisection with the *local* Lipschitz constant certifies $\inf f^{\text{arch}}/(4 \sin^2(\theta/2)) \geq 0.25$ on 12/12 windows in 519–1201 symbol evaluations (certified floor $0.2505\text{--}0.3062$, never above the fine-grid minimum; the global constant is documented insufficient — a uniform grid would need up to $61.4 \times$ more); mutation: fed a target above the true grid minimum it *refuses* on 12/12. Stress: on the T145 no-go family the Schur floor survives ($t = 0.05$) while the confinement goes vacuous ($7.93\text{--}413, \times 52.0$) and $(S_L)_{11}$ explodes ($1185 \rightarrow 7.11 \times 10^4, \times 60.0$)

against the flat 4.05–5.90 real band; parity control exact to 4.9×10^{-14} ; the Dirichlet residual hits the predicted $2/\sqrt{N}$ to 7.4×10^{-5} . **Named limits as load-bearing content:** per instance on small windows — a finite list with an explicit maximum; nothing uniform in the zone index or in D , and explicitly *no* statement for all D ; *nothing about the angles is closed* — the two open terms after T157 (an m -free lower bound on $\hat{g}_1^2$, equivalently an m -free upper bound on the one Schur diagonal entry $1/s$; an m -free R1'' domination with a non-shrinking margin — certified per window with a 7.3×10^{-4} margin at the largest window and a shrinking trend) stay open, typed open; every floor t and ceiling S consumed is certified per window and its m -freeness is exactly what remains open; PRIME.RITZ.CEIL.01's ceiling and PRIME.ODD.SECTOR.IDENT.01's ladder stay as stated and nothing is re-opened. Fences: Kac–Murdock–Szegő 1953 / Schur 1917–Haynsworth 1968 / Courant–Fischer 1920–Cauchy 1829 / Cauchy–Schwarz (refuted as a ceiling, wired as a control) / Kantorovich 1948 / Temple 1928–Kato 1949 (floor devices, not used here) / Sylvester 1852–Bunch–Kaufman 1977 / Wilkinson 1968–Higham 2002 / Chebyshev 1852 / Szegő 1915–Widom 1958 (a named address, never a licence) named classical; Weil 1952 cited, never used as a criterion; zero-firewall AST-checked; no marker moves. Machine-checked: [verification/v552_angle_instruments.py]. [E] (per-instance identities and certified inequalities)

The exact-form identities (PRIME.EXACT.FORM.IDENT.01). The theorem cores of the discovery parts T158 and T159, recomputed on the same declared surface as PRIME.RITZ.CEIL.01 and PRIME.ANGLE.INSTR.01 (the 12 deepest in-cap frame-A windows, $n = 29 \dots 139$, $m = 96 \dots 291 \leq 300$; no-go ladders $m = 48 \dots 288$ on *both* T145 forms; parity and Dirichlet controls at $m = 64, 128, 256$; each statement with at least one mutation control that must fail loudly) — the algebra of the sixteen-form itself, and it closes neither open term. (1) The Thomson dual form and the positive Cholesky ladder: $s = (B^{-1})_{11} = \max_x (2x_1 - x^T B x)$ (Maz'ya 1985 / Miclo 1999) with the direction *executed*, not asserted — 8 random trials per window never exceed s ; the ladder $g_K = \sum_{j \leq K} y_j^2$, $y = L^{-1} e_1$ from one Cholesky of the leading 16×16 block (nested Schur complements): all terms strictly positive, partial sums strictly monotone, $1/s \leq 1/g_{16} \leq 1/g_1 = B_{11}$ literally on 12/12, tight to 1.0642–1.1522 against the declared cap 1.5; mutation: a 5% corruption of one off-diagonal entry destroys positive definiteness — the Cholesky *refuses* on 12/12. (2) The two-dimensionality: $\text{span}\{t_1, A^{-1} L_P t_1\}$ attains s exactly ($g/s = 1$ to 9.0×10^{-13}) for the theorem reason $L_P t_1 = \mu_1^P t_1$ (KMS 1953); negative controls: the two-sine span misses by 1.42–4.8, the corrupted Green column by $255\text{--}7.7 \times 10^3$. (3) The gauge identity and the Toeplitz-minus-Hankel signature: the reflection-odd section annihilates constant lag vectors *entrywise with zero tolerance* ($\text{odd_toeplitz}(\mathbf{1}) = 0$ exactly), hence $\sum_d w_d = 0$ to 1.1×10^{-14} of $\sup |w|$ — the form is blind to the lag mass of c , exactly where the archimedean and atom halves are h^2 -sized and cancel; the signature $A_{r,s} - A_{r+1,s+1} = f(r+s)$ holds to 1.4×10^{-16} ; mutations: a one-index Hankel shift breaks the identity by $2.7 \times 10^{-7}\text{--}4.2 \times 10^{-5}$, and the T145 form has signature defect 7.4×10^{-2} — not a parity section, the identity is not even defined for it. (4) The exact lag sum and the two closed scalars: $x^T B_{LL} x = \sum_d c_d w_d$ ($w_d = T_d - H_d$) against the direct assembly to 2.7×10^{-16} of the absolute scale $\sum_d |c_d w_d|$, with the honest price *measured* (relative to the cancelled total only 1.4×10^{-11} on this small surface — that depth is the h^2 obstruction); $w_0 = \sum_k x_k^2 / \mu_k^P$ to 4.2×10^{-16} and $2w_0 - w_1 = \|x\|^2$ to 7.4×10^{-16} ; mutations: a 1% lag perturbation and a one-index μ shift both break loudly. (5) The Z-matrix law and the Collatz–Wielandt floor: B_{HH}^{arch} is a symmetric Z-matrix *raw* — every off-diagonal entry ≤ 0 with no tolerance on 12/12; the closed sign-based floor $\min_k (B_{HH}^{\text{arch}} \mathbf{1})_k = 0.9093\text{--}1.1284 \geq 0.25$ with the theorem direction verified per instance against the exact $\lambda_{\min} = 0.9138\text{--}1.1432$ (Perron 1907 / Frobenius 1912 / Collatz 1942 / Wielandt 1950), beating the sign-free norm route by 0.69–8.8; the honest limit as content: the atom block obeys no sign law (0.44–0.53, a coin toss), the full block is not a Z-matrix and its row-sum floor is negative ($-48.3 \dots -0.79$) — admissible and vacuous, R1'' stays open; mutation: the checkerboard similarity (an isometry, $\lambda_{\min}$ moves by 0 exactly) destroys the raw law — an entrywise fact, not a spectral one. Stress: the T145 no-go breaks on three axes of this module's structure; on the $c(l) = 1/(1+l)$ reconstruction the ladder theorem survives while $1/g_{16}$ explodes ($1187 \rightarrow 7.11 \times 10^4$) — the instrument reports the collapse instead of hiding it; parity: for $A = L_P$ the machinery returns

$s = 1$ with every ladder partial sum constant in K ; Dirichlet controls exact. **Named limits as load-bearing content:** per instance on small windows — a finite list with an explicit maximum; nothing uniform in the zone index or in D , and explicitly *no* statement for all D ; the two open terms after T159 (the one explicit pairing inequality for $R2''$ — $\sum_d (\Delta c)_d W_d^1$ with $\|\Delta c^{\text{arch}}\|_1 \leq 6$, needing a correlation bound instead of sizes; a third device for the atom off-band block for $R1''$, neither norm nor sign) stay open, typed open; the pairing bound and m -freeness are explicitly *not* claimed. Fences: Maz'ya 1985 / Miclo 1999 / Schur 1917–Haynsworth 1968 / KMS 1953 / Dirichlet 1829 / Abel / Fejér 1915 / Perron 1907–Frobenius 1912 / Collatz 1942–Wielandt 1950 / Gantmacher–Krein 1950 named classical; Weil 1952 cited, never used as a criterion; zero-firewall AST-checked; no marker moves. Machine-checked: [verification/v553_exact_form_identities.py]. [E] (per-instance identities and certified inequalities)

The sampling/harmonics identities (PRIME.SAMPLING.HARM.01). The closed theorem cores of the discovery parts T160 and T161, recomputed on the same declared surface as PRIME.EXACT.FORM.IDENT.01 (the 12 deepest in-cap frame-A windows, $n = 29 \dots 139$, $m = 96 \dots 291 \leq 300$) plus a declared Λ -battery of six log-spaced cut-offs $X = 10^3 \dots 3.2 \times 10^5$ (finite von-Mangoldt sums only, no matrix anywhere; each statement with at least one mutation control that must fail loudly) — what the sixteen-form *pairs against*, and it closes no open term. (1) The sampling identity: the atom half of the pairing equals $-\frac{1}{2} \sum_{n \leq X} (2\Lambda(n)/\sqrt{n}) \hat{w}(\log n)$ (plus one reflected term below the grid; $\hat{w}$ the linear interpolant of the lag weights), exact to 5.6×10^{-15} of the absolute atom scale on 12/12 — hence identically a finite combination of sums $\sum_n \Lambda(n) n^{-1/2} \cos(t \log n)$ at 32 explicit frequencies; mutation: a one-index grid shift breaks by $5.1 \times 10^{-5} - 3.1 \times 10^{-3}$. (2) The moment laws and the total-variation theorem: $m_0 = 0$ (1.0×10^{-16} of $\|w\|_1$), $m_1 = -[S_0^2 + 2 \sum_j P_j^2]$ (9.2×10^{-16}) and $m_2 = -[2S_1 - (M-1)S_0]^2$ (3.4×10^{-14} , a perfect square), both strictly negative on every window; U3 as a theorem with *checked* hypotheses ($c^{\text{arch}} \leq 0$ and monotone on $d \geq 1$, checked 12/12): $\text{TV}(c^{\text{arch}}) = 3.9817 - 4.8991 \leq |c_0| + 2 \sup < 6$, closed and window-independent; mutations: a prefix shift, an $(M-2)$ mis-normalisation, and the full atom-including kernel all fail loudly. (3) The harmonic-frequency theorem and the PNT main term: $t_j(2\alpha) = 2\pi j$ *exactly* (2.8×10^{-14}) — the 32 frequencies are precisely the Fourier harmonics of the log-window $[0, 2\alpha]$; at those frequencies the partial-summation main term collapses to the closed parameter-free prediction $(\sqrt{X} - 1)/(\frac{1}{4} + t^2)$ (de la Vallée Poussin 1896), matching the finite sums to 0.0007–0.097 of the trivial mass against the declared cap 0.12 on the battery; mutations: a wrong Mellin pole worsens the aggregate residual by $\times 1.8 - \times 95.8$, and a mass-matched *uniform* fake measure misses by 0.23–0.51 — the agreement is a property of the von-Mangoldt measure. (4) The head split and the scale-free Bernstein rate: $c_d^{\text{arch}} = \Psi_d + D\hat{G}(dD, D)$ with $\Psi_d = -\frac{1}{2}[(d+1)\log(d+1) - 2d\log d + (d-1)\log(d-1)]$ closed, D -free and m -free (the triangle-smeared simple pole; the $\log D$ cancels identically), exact to 3.2×10^{-16} at every lag of every window; $\rho^* = (3 + \sqrt{5})/2 = 2.618034$ is the root of $x^2 - 3x + 1$, returned exactly by the $D \rightarrow 0$ limit of the peeled interval and invariant under $\alpha \times 10$ (Bernstein 1912); mutations: dropping Ψ misses by 0.99–1.00, the wrong interval ratio moves the limit to 3.0. (5) The off-diagonal fraction bound and the sign refutation: the gauge identity licenses one boundary-free Abel step, the decomposition $Q^{\text{arch}} = \sum_{k,l} a_k a_l \sum_d (\Delta c^{\text{arch}})_d R_{kl}^1(d)$ is exact (2.0×10^{-13}), and the certified per-window inequality is $|\text{off-diagonal}| \leq \frac{1}{4} |Q^{\text{arch}}|$ (0.171–0.212 on 12/12) — with the *refuted* sign law wired as the negative control (arch half negative, off-diagonal block positive on 12/12, 176/240 pairs violate the per-pair form): nothing single-signed is promoted; mutation: a one-index R -kernel shift breaks by $\geq 6.8 \times 10^{-5}$. Stress: the head split misses the T145 no-go reconstruction by 2.39 of $\sup |c|$; the Dirichlet cosine-sum identity (including the degenerate branch), the KMS eigenpairs and $\text{odd_toeplitz}(c^L) = L_P$ (zero tolerance) are exact. **Named limits as load-bearing content:** per instance on small windows — a finite list with an explicit maximum; nothing uniform in the zone index or in D ; finite Λ -sums are allowed and used, *no* RH statement is made, and the δ triage of T161 (required cancellation depth against RH strength) is a documentation item, *not* a claim of this module; the open terms after T161 (R-A' the prime-free log-moment $\sum_d \Psi_d w_d$, outside the polynomial ladder; R-B' the m -free 16×16 Gram fraction bound,

Fejér 1915 / Schur 1917; R-C' *one* split of the pairing with $\delta_{\text{eff}} < \frac{1}{2}$, T161 refuting the arch/atom and arch+smooth/fluctuation splits with numbers; R-D a fifth R1'' device) stay open, typed open. Fences: Dirichlet 1829 / Abel 1826 / Fejér 1915–Schur 1917 / Bernstein 1912 / Chebyshev 1852–Mertens 1874 / de la Vallée Poussin 1896 / Clausen–Lerch / KMS 1953 named classical; Weil 1952 / Bombieri 2000 cited, never used as a criterion; zero-firewall AST-checked; no marker moves. Machine-checked: [verification/v554_sampling_harmonics_identities.py]. [E] (per-instance identities and certified inequalities)

The Pareto/total-variation identities (PRIME.PARETO.TV.01). The closed theorem cores of the discovery parts T162 and T163, recomputed on the same declared surface as PRIME.SAMPLING.HARM.01 (the 12 deepest in-cap frame-A windows, $n = 29 \dots 139$, $m = 96 \dots 291 \leq 300$), with the one arithmetic input $\kappa = 0.038821$ *verified* at every jump point of $\psi(x)/x$ on the von-Mangoldt table in $[100, 3.2 \times 10^5]$ (reproduced to 4.2×10^{-7} ; each statement with at least one mutation control that must fail loudly) — the *price structure* of the pairing, and it closes no open term beyond the negative closure that the floor inequality itself is. (1) The exchange law and the crossing criterion: $\delta_{\text{bnd}}(x) = \frac{1}{2} + \log(2\kappa g_{16} \text{TV}(x)/P(x))/\log X$ is an identity at all 612 grid points of the declared Fejér knob (50 geometric σ on $[1, 4096]$ plus ∞ ; 5.1×10^{-16} relative, admissibility $Q(x) > 0$ everywhere), and $\delta_{\text{bnd}} < \frac{1}{2} \iff P > 2\kappa g_{16} \text{TV}$ holds as an exact equivalence at every point — price and demand are two coordinates of *one* object; mutation: the wrong constant (κ for 2κ) moves the law by 6.5×10^{-2} –3.4. (2) The total-variation floor — the theorem that carries the T163 verdict: (T1) $w_0(x) = \|a(x)\|^2$ (4.9×10^{-16} ; orthonormal parity rows); (T2) $\text{TV}(x) \geq |w_0(x)|$ by telescoping with $w_M = 0$; (T3) $x_1 = 1$ forces $\|a\|^2 \geq 1/\mu_1^P$, hence $\mu_1^P \text{TV}(x) = 5.00$ –28.84 ≥ 1 on all 708 trial vectors built (51 knob settings plus 8 random admissible sixteen-vectors per window), the closed floor $1/\mu_1^P = 1/(4\sin^2(\pi/N)) = 943.6$ –8609.6 $\sim h^2$; (T4) every sub- $\frac{1}{2}$ grid point pays $P > 2\kappa g_{16}/\mu_1^P = 15.4$ –142.8 (all 54 respect it) — a flat-price sub- $\frac{1}{2}$ demand is impossible in the parity sector on this surface, the inequality that closes R-C''' negatively; negative control: the *inadmissible* vector $0.01x^*$ ($x_1 \neq 1$) does undercut the floor ($\mu_1^P \text{TV} = 2.5$ –2.9 $\times 10^{-3}$) and is recognised as inadmissible — the floor is a property of the normalisation meeting the smallest parity eigenvalue (Kac–Murdock–Szegő 1953). (3) The chain-derived price axis: $\sigma = 1$ gives $x = e_1$ exactly with $P = g_{16}/g_1$ (the $K = 1$ rung of the T158 Cholesky ladder, the T157 route-(0) certificate; 4.1×10^{-16}), $\sigma = \infty$ gives $g_{16}Q(x^*) = 1$ (the $K = 16$ top rung; 1.2×10^{-12}); the ladder carries all 16 terms strictly positive with monotone partial sums, and the knob is monotone in both coordinates on every window — the knob curve *is* the Pareto front; mutation: a corrupted off-diagonal sixteen-block entry makes the Cholesky *refuse* on 12/12. (4) The Mellin cell moments and the Abel step: the k -th rung $\sum_d w_d C_d(-(2k + \frac{1}{2}))$ with $C_d(s)$ the closed tent integral (Mellin 1896) agrees with an independent per-cell Gauss–Legendre quadrature to 2.8×10^{-13} on every rung $k = 1 \dots 4$ (shifted tent centres break by $\geq 1.3 \times 10^{-2}$); the boundary-free Abel identities hold at levels $k = 1 \dots 3$ (8.3×10^{-17} ; Abel 1826), the dropped-boundary level-1 form agreeing because the gauge identity kills the boundary term exactly; the optimal level is *one* for T162's closed reason ($32\pi/\alpha = 40.6$ –59.4 > 1 , measured level-2/level-1 ratio 30.8–69.1 > 1), and a gauge violation breaks the dropped form by its closed prediction $v_0 c_0 M$ to 1.2×10^{-15} — the positive control. (5) The Lerch/Frullani log-moment: $L = \sum_d \Psi_d w_d$ agrees on three independent routes on every window — direct; double-Abel *with* the Wronskian boundary term $-\frac{1}{2}M \log M w_{M-1}$ (2.8×10^{-14} ; dropping it breaks by 2.2×10^{-4} – 1.3×10^{-2}); the Lerch/Frullani integral with the $d = 1$ term peeled closed (1.8×10^{-15} ; the peel constant $2 \log 2$ to 2.2×10^{-16} , $\Psi_1 = -\log 2$ exact; Frullani 1828 / Lerch 1887 / Clausen 1832). Stress: the monotone no-go staircase — coarsening over $\nu = 4/2/1/0.5$ degrades the arch/atom cancellation monotonically in the median ($1.00 \rightarrow 3.49 \rightarrow 10.81 \rightarrow 24.98$, the $\nu = 4$ column an exact null control); the Dirichlet cosine-sum identity (2.4×10^{-14} , including the degenerate branch), the KMS eigenpairs (1.4×10^{-14}) and odd_toeplitz(c^L) = L_P (zero tolerance) are exact. **Named limits as load-bearing content:** per instance on small windows — a finite list with an explicit maximum; nothing uniform in the zone index or in D ; the T163 h -exponents ($P_{\text{aff}} \sim h^{3.05}$, $P_{\text{cross}} \sim h^{1.91}$, the TV exponent $\sim h^{2.00}$) are sandbox fits and are *not*

consumed — what is consumed is the closed per-window floor; finite Λ -sums are allowed and used, *no* RH statement is made ($\frac{1}{2}$ appears in exactly one role: as the strength of a classical statement a required depth is compared against); the open terms after T163 (R-E the successor, both arms prime-free — arm A: a growing $1/s$ ceiling for the downstream T159/T160 chain; arm B: a sector whose lowest eigenvalue does not vanish like h^{-2} ; R-B''' the h -uniform positivity margin; R-D a fifth R1'' device) stay open, typed open. Fences: Fejér 1915 / Mellin 1896 / Abel 1826 / Chebyshev 1852–Rosser–Schoenfeld 1962 (verified on the table) / Legendre / Dirichlet 1829 / KMS 1953 / Frullani 1828–Lerch 1887–Clausen 1832 / Schur 1917–Haynsworth 1968 named classical; Weil 1952 / Bombieri 2000 cited, never used as a criterion; zero-firewall AST-checked; no marker moves. Machine-checked: `[verification/v555_pareto_tv_identities.py]`. [E] (per-instance identities and certified inequalities)

The gauge/ P_{pr} identities (PRIME.GAUGE.PPR.01). The closed theorem cores of the discovery parts T164 and T165, recomputed on the same declared surface as PRIME.PARETO.TV.01 (the 12 deepest in-cap frame-A windows, $n = 29 \dots 139$, $m = 96 \dots 291 \leq 300$), with the one arithmetic input $\kappa = 0.038821$ *verified* at every jump point of $\psi(x)/x$ on the von-Mangoldt table in $[100, 3.2 \times 10^5]$ (each statement with at least one mutation control that must fail loudly) — the *gauge structure* of the pairing, and it does not close the one remaining quantifier. (1) The gauge theorem (T164): $x_1 = 1$ fixes $a_1 = 1/\sqrt{\mu_1}$, i.e. only the *scale* of a ; Q and TV are homogeneous of degree two in a ; hence the demand ratio Q/TV of the same a -direction — and with it δ_{bnd} — is the same number in *every* sector, unchanged to 2.9×10^{-11} on 2520 sector-by-trial-vector combinations (5 weight families $\times$ 7 shifts $\times$ 6 Fejér taper widths on 12 windows), with $\text{TV} \geq |w_0|$ in every sector and the pure shift scaling the certified value by exactly $\theta = \mu_1^P/(\mu_1^P + \kappa_s)$ (2.2×10^{-11} on 432 points); the full space is settled in closed form and strictly worse ($\mu_0 = 0$ exactly, the lowest remaining eigenvalue $\leq \mu_1^P$ on 12/12); mutation: the wrong transport map (the x -coordinates carried unchanged instead of the a -direction) moves the ratio by 2.0×10^2 – 5.4×10^3 . (2) The h^2 conservation law (T164): floor $\times$ transfer $= 1/(\mu_1^P + \kappa_s) \cdot (1 + \kappa_s/\mu_1^P) = 1/\mu_1^P$ *identically* at every shift on the declared grid (72 window-by-shift points, 2.2×10^{-16} of $1/\mu_1^P$) — the h^2 is conserved, never removed; it lives in the ratio, not in the floor; mutation: the wrong transfer misses by 0.9995 at the largest shift on every window. (3) The power-one spend as an identity (T164): the r -ceiling $r \leq U/L$ with $L = \lambda_1(A)/\mu_1^P = 1.1019$ – 1.5845 (computed per instance from the full window) is linear in the entry ceiling, so $d \log r / d \log U = +1$ exactly (within 1.1×10^{-14} on 12/12) — an h^ε ceiling costs $h^{-\varepsilon}$ of r -margin for every $\varepsilon > 0$, no free tolerance in the consumption step (the measured power of the full T156 loss margin stays sandbox); mutation: the quadratic consumption reads slope 2.0000 and is recognised. (4) The P_{pr} identity and the predicate equivalence (T165): $P_{\text{pr}} = g_{16} \cdot R \cdot (\text{TV}/(t_1 v)^2)/\mu_1^P$ with no inequality anywhere — four factors, each a named object: the R-E-A quantifier $g_{16} = 0.1559$ – 0.2227 , the demand $R = Q/\text{TV}$, the T163 floor $\text{tvf} \geq 1$, and the KMS 1953 scale $1/\mu_1^P = 943.6$ – $8609.6 \sim h^2$ — exact to 4.3×10^{-16} on all 192 battery vectors (6 knob settings, 6 random sixteen-vectors and 4 random *full-window* vectors per window), gauge-invariant under $v \rightarrow tv$ to 1.5×10^{-11} against the declared cancellation horizon; the predicates $R > 2\kappa$ and $P_{\text{pr}} > 2\kappa g_{16} \text{tvf}/\mu_1^P$ agree as booleans on all 192 (no search is run — T165's ascent stays sandbox), so the demand half of R2'' and the gauge-invariant crossing condition are *one* predicate — R-F is strictly stronger than the R2'' demand and its m -uniform form *is* R-E-A — and $\text{tvf} \geq 1$ pointwise, hence every crossing vector pays $P_{\text{pr}} > 2\kappa g_{16}/\mu_1^P = 15.4$ – 142.8 , above the preregistered tolerance $\Theta_{\text{TOL}} = 10$ on 12/12 windows: the R-F conjunction is *empty* on this surface by an identity plus a floor; mutation: the price with $(t_1 v)$ unsquared scales like t and moves by 1.0 – 1.0×10^4 under the same round trips. (5) The Schur-cascade direction and the exponent closure (T164/T165): $s = \mu_1^P t_1^T A^{-1} t_1 = 0.1696$ – $0.2472 \geq g_{16} \geq g_1$ window by window ($1/s \leq 1/g_{16} \leq 1/g_1 = B_{11}$; Schur 1917–Haynsworth 1968), $P_K = \hat{a} s = 254.8$ – $7.7 \times 10^3 \geq 1$ (Kantorovich 1948); the exponent closure is an identity of the lists — per instance $\log(1/g_1) = \log(g_{16}/g_1) + \log(1/g_{16})$ to 1.6×10^{-15} , the least-squares fits closing to 1.3×10^{-15} by linearity (the exponent *values* are sandbox fits, not consumed — what is promoted is that the whole certification lives in the cascade gain g_{16}/g_1 , the object of the one open quantifier); mutations: the g_8 closure breaks by 0.10–0.24, a corrupted sixteen-block makes

the Cholesky *refuse* on 12/12. Stress: the monotone no-go staircase ($1.00 \rightarrow 3.49 \rightarrow 10.81 \rightarrow 24.98$, the $\nu = 4$ column an exact null control); the Dirichlet cosine-sum identity (2.9×10^{-14} , including the degenerate branch), the KMS eigenpairs (1.4×10^{-14}) and $\text{odd_toeplitz}(c^L) = L_P$ (zero tolerance) are exact. **Named limits as load-bearing content:** per instance on small windows — a finite list with an explicit maximum; nothing uniform in the zone index or in D ; the T164/T165 h -exponents and the measured anatomy (the free ascent, the $\eta(\text{Cap})$ profile, the decoupled ν surface with its zone-depth finding, the retired constant $U_{\text{ref}} = 4.90 \rightarrow 5.7327$) stay in the diary and the paper; finite Λ -sums are allowed and used, *no* RH statement is made ($\frac{1}{2}$ appears in exactly one role: as the strength of a classical statement a required depth is compared against); **the one open object after T165 stays open, typed open:** $\inf_m g_{16}(m) > 0$, equivalently a lower bound on the sixteen-step Schur-cascade gain g_{16}/g_1 uniform in m — a quantifier over a certified flat list; R-B''' (an independent α/h surface) stays open as narrowed by T164. Fences: Schur 1917–Haynsworth 1968 / KMS 1953 / Abel 1826 / Chebyshev 1852–Rosser–Schoenfeld 1962 (verified on the table) / Kantorovich 1948 / Fejér 1915 / Dirichlet 1829 named classical; Weil 1952 / Bombieri 2000 cited, never used as a criterion; zero-firewall AST-checked; no marker moves. Machine-checked: [verification/v556_gauge_ppr_identities.py]. [E] (per-instance identities and certified inequalities)

The cascade/vector identities (PRIME.CASCADE.VECT.01). The closed theorem cores of the discovery parts T166 and T167, recomputed on the same declared surface as PRIME.GAUGE.PPR.01 (the 12 deepest in-cap frame-A windows, $n = 29 \dots 139$, $m = 96 \dots 291 \leq 300$), with the one arithmetic input $\kappa = 0.038821$ *verified* at every jump point of $\psi(x)/x$ on the von-Mangoldt table in $[100, 3.2 \times 10^5]$ (each statement with at least one mutation control that must fail loudly) — the *anatomy of the cascade* and the *exactness of the closed vector at $K = 2$* , and it does not close the one remaining scalar. (1) The cascade identities (T166): the prefix property — one Cholesky of the sixteen-block yields all sixteen rungs, $g_K = e_1^T Q_K^{-1} e_1$ at every K (192 comparisons, 1.1×10^{-12}); the minor form $1/g_K = \det G_{[1..K]}/(\mu_1^P \det G_{[2..K]})$ on the raw arithmetic Gram block $G = TAT^T$ (2.7×10^{-12} on 36 (window, K) pairs, $K \in \{2, 6, 16\}$; Schur 1917–Haynsworth 1968); the regression form $g_K/g_1 = 1/(1 - R_K^2)$ with R_K^2 the squared multiple correlation of mode 1 on modes $2 \dots K$ (8.8×10^{-13}) — the open object *is* a near-collinearity statement; mutation: the minor with the wrong row/column deleted misses by 0.99–1.45. (2) The diagonal invariance (T166): the gain $g_K/g_1 = B_{11}(Q_K^{-1})_{11}$ is unchanged under $B \rightarrow DBD$ on 144 random positive diagonals (2.7×10^{-12}) — the cascade does not see the KMS $1/\sqrt{\mu}$ normalisation, the gain is a property of the arithmetic Gram block alone; mutation: a non-diagonal congruence moves the gain by 2.7–3.8. (3) The exact $K = 2$ identity (T167): the closed vector $u = (1, -Q_{21}/Q_{22})$ *attains* the constrained minimum, $u^T Q_2 u = 1/g_2 = Q_{11}(1 - r_{12}^2)$ to 1.1×10^{-12} , and $B_{11}g_2 = 1/(1 - r_{12}^2)$ identically to 1.2×10^{-12} (measured $1 - r_{12}^2 = 2.8 \times 10^{-4} - 5.9 \times 10^{-3}$ on this surface, the h -trend a sandbox fit): at $K = 2$ the fast-null-vector needs no approximation — the third dress collapses onto the second, one scalar built from three closed lag sums; mutation: the wrong-sign vector pays the relative excess $4r_{12}^2/(1 - r_{12}^2) = 6.7 \times 10^2 - 1.5 \times 10^4$. (4) The pivot identity and the unification (T167): for $u = x_K^* + \delta$ with $\delta_1 = 0$, $u^T Q_K u = 1/g_K + \delta^T Q_K \delta$ exactly (2.0×10^{-12} on 252 evaluations across three δ scales) — every relative excess is an exact number; the Rayleigh envelope holds with its $\delta_1 = 0$ ceiling attained exactly (1.1×10^{-15}) and Cauchy-interlaced; the sign-aligned entrywise perturbation attains $|x^{*T} E x^*| = \varepsilon S_K$ exactly (3.8×10^{-16}), so the entry threshold $\rho_{\text{carry}}(1/g_K)/S_K$ is exact — and the unification: $(1/g_K)/S_K$ is the one cancellation ratio of all three dresses, at $K = 2$ literally $(1/g_2)/S_2 = (1 - r_{12}^2)/(1 + 3r_{12}^2)$ (1.2×10^{-12} ; the closed factor $1 + 3r_{12}^2 \rightarrow 4$ is T166's independently derived quarter): *one* inequality, not three; mutation: $\delta_1 = 0.1$ breaks the pivot identity by exactly the closed cross term $2\delta_1 = 0.2$. (5) The trial theorem and the two must-breaks (T166/T167): all 288 random trials with $u_1 = 1$ satisfy $u^T Q_K u \geq 1/g_K$ (min ratio 15.14) — every closed candidate is a certified *lower* bound (Schur 1917); the L_P no-go: the normalised block is the identity (1.6×10^{-15}) and the cascade gain exactly 1 (deviation 0.0) at $m = 64/128/256$; the scramble type-change: uniform random atom positions at the same total mass make B_{11} itself lose positivity on 49/60 declared draws and on a majority of draws on 12/12 windows — at $K = 2$ the collapse under scrambling is a change of *type*, not a factor. Stress:

exact Dirichlet/parity controls (2.3×10^{-14} / 1.4×10^{-14} / zero tolerance). **Named limits as load-bearing content:** per instance on small windows — a finite list with an explicit maximum; nothing uniform in the zone index or in D ; the T166/T167 h -exponents ($1 - r_{12}^2 \sim h^{-2.921}$ on frame A, the certified gain exponent $h^{+2.921}$, the threshold monotonicity in K , the $K = 6$ separation $h^{+1.138}$) and the measured anatomy (the rung profile, the polarisation split, the Kato radius, the headroom) stay in the diary and the paper; finite Λ -sums are allowed and used, *no* RH statement is made ($\frac{1}{2}$ appears in exactly one role: as the strength of a classical statement a required depth is compared against); **the one open object after T167 stays open, typed open:** R1, the single scalar — an m -free upper bound on $1 - r_{12}^2 = 1 - Q_{12}^2/(Q_{11}Q_{22}) \leq C h^{-3+\varepsilon}$, three closed lag sums and nothing else; the routes T167 closed off (perturbation theory of every order around the KMS bottom mode — the Kato series converges to the *wrong* object; position-blind surrogates) stay sandbox negatives; R-B''' stays open as narrowed. Fences: Schur 1917–Haynsworth 1968 / KMS 1953 / Rayleigh 1877–Courant–Fischer 1920 / Cauchy 1829 / Kato 1949 (cited as the closed-off route, never used) / Chebyshev 1852–Rosser–Schoenfeld 1962 (verified on the table) / Dirichlet 1829 named classical; Weil 1952 / Bombieri 2000 cited, never used as a criterion; zero-firewall AST-checked; no marker moves. Machine-checked: [verification/v557_cascade_vector_identities.py]. [E] (per-instance identities and certified inequalities)

The bilinear/rank identities (PRIME.BILINEAR.RANK.01). The closed theorem cores of T168 (contract LAGRANGE.MINORS), T169 (contract TSTAR.RATIO) and T170 (contract BILINEAR.SIEVE) as one deliberately narrow module (27 checks, ~ 0.2 s), recomputed on the declared small frame-A surface (12 deepest in-cap zones, $m = 96 \dots 291 \leq 300$; $\kappa = 0.038821$ verified at every jump point of $\psi(x)/x$ on the table) — the *exact shape of the one remaining scalar*, and it does not close that scalar. (1) The Lagrange/Wronskian anatomy (T168): the lag-sum identity $\hat{a}_{ij} = t_i^\top A_h t_j = \sum_d c_d W_d^{ij}$ with the closed correlation weights (1.6×10^{-15} ; the $1/\sqrt{\mu}$ factors cancel, T169-TH3), Euclidean orthogonality $t_1 \cdot t_2 = 0$ (4.4×10^{-16}) — the near-parallelism is created *entirely* by the arithmetic metric — the Wronskian telescope $(\mu_1 - \mu_2)u_r v_r = W_{r-1} - W_r$ with $W_{-1} = W_{h-1} = 0$ (1.9×10^{-17}) and the maximal-minor identity $\frac{1}{2} \sum M^2 = 1/(\mu_1 \mu_2)$ exactly (4.6×10^{-16} ; Lagrange 1773); mutation: the wrong eigenvalue pair misses by 1.66–1.67 relative (the closed miss $(\mu_2 - \mu_3)/(\mu_1 - \mu_2) \rightarrow 5/3$). (2) The exponent account and the Gershgorin certificate (T168/T169): $1 - r_{12}^2 = \nu_1 \nu_2 / (\hat{a}_{11} \hat{a}_{22})$ identically (2.5×10^{-13} ; measured $1 - r_{12}^2 = 2.8 \times 10^{-4} - 5.9 \times 10^{-3}$ on this surface, the h -trend a sandbox fit); $\nu_1 \leq \max(\hat{a}_{11}, \hat{a}_{22}) + |\hat{a}_{12}|$ (Gershgorin 1931, unconditional, closed in the three entries) on 12/12 with loss 1.1885–1.1947; mutation: dropping $|\hat{a}_{12}|$ fails on every window ($\nu_1 - \max \text{diag} = 0.30 - 1.17 > 0$). (3) The det-collapse theorem (T169-TH7): on the Rayleigh family $u(t) = (t, -1)$ the candidate $K7 = \sqrt{\hat{a}_{22}/\hat{a}_{11}}$ collapses in closed form, $R(K7) = 2\sqrt{\hat{a}_{11}\hat{a}_{22}} \det \hat{A} / [(\hat{a}_{11} + \hat{a}_{22})(\sqrt{\hat{a}_{11}\hat{a}_{22}} + \hat{a}_{12})]$ (8.7×10^{-13}) — it meets the threshold precisely because it reintroduces $\det \hat{A}$: the loop T167 scalar $\rightarrow$ T168 factor $\rightarrow$ T169 candidate closes as an identity; usable form $\nu_2 \leq 2 \det \hat{A} / (\hat{a}_{11} + \hat{a}_{22}) = 2\nu_1 \nu_2 / (\nu_1 + \nu_2)$, tight to 1.9980–1.9999 < 2 exactly (Rayleigh 1877); mutation: the closed constant $1/\sqrt{2}$ overshoots ν_2 by 17.1– 2.3×10^2 . (4) The exact bilinear von Mangoldt form (T170 TH1–TH3): the exact split $\hat{A} = B - S$ with $S_{ij} = \sum_{n \leq X} (\Lambda(n)/\sqrt{n}) X_n[ij]$ against the direct block (1.7×10^{-14}); the polarisation $\det \hat{A} = \det B - D(B, S) + \det S$ (1.0×10^{-10}); the double sum $\det S = \sum_{n,m \leq X} \Lambda(n)\Lambda(m)K(n,m)/\sqrt{nm}$ with $K = \frac{1}{2}D(X_n, X_m)$ through the *explicit* kernel matrix (7.5×10^{-16} ; Cauchy–Binet 1812–1815); the wedge reading to the measured rank-one defect $4.6 \times 10^{-3} - 3.1 \times 10^{-2}$ (honest certificate); mutation: the wrong cross-coefficient breaks by 64–430. (5) The rank-3 theorem and the Type II collapse (T170 TH4–TH5): the polarisation matrix on (X_{11}, X_{22}, X_{12}) has eigenvalues $\{-2, -1, +1\}$ exactly — rank 3, signature (1, 2) — so the kernel $K = PMP^\top$ has $\sigma_4/\sigma_1 \leq 1.6 \times 10^{-16}$ with nonzero spectrum = $\text{spec}(M P^\top P)$ (1.0×10^{-15}): the bilinear form *is* the rank-3 polynomial $S_{11}S_{22} - S_{12}^2$ in three linear Λ -sums; Vaughan’s identity coefficient-exact for every $U = V \in \{3, 4, 5, 6\}$ (residual 0.0; Vaughan 1977), Type II kernel 99.9%-energy rank $[2, 2, 2, 2]$ against generic $[20, 15, 11, 9]$ (4.5×10^{-10} ; Montgomery–Vaughan 1973 cited as the priced route). (6) The R4-free chain and the two discriminators (T170-TH6): $1 - r_{12}^2 = \det \hat{A} / (\hat{a}_{11} \hat{a}_{22})$ is an identity (0.0) — no positivity of A_h , the Weil fence never approached; the L_P no-go: $\hat{a}_{12} = 1.1 \times 10^{-17}$ identically, no decay;

the scramble discriminator: rank-3 *untouched* on all 60 draws while the value of $1 - r_{12}^2$ moves by a median factor 834.8 (range $69 - 1.8 \times 10^4$) — the arithmetic sits entirely in the joint values of (S_{11}, S_{22}, S_{12}) against B . Stress: exact Dirichlet/parity controls (1.2×10^{-14} / 1.4×10^{-14} / zero tolerance). **Named limits as load-bearing content:** per instance on small windows — a finite list with an explicit maximum; nothing uniform in the zone index or in D ; the T168–T170 h -exponents ($1 - r_{12}^2 \sim h^{-2.92}$ on the deep frame-A surface, the route-table deltas — large sieve +0.669, MV −0.384, exact split +0.996, Vaughan +0.669 against the measured truth 2.747 — the five-order cancellation and the $h^{-2.711}$ row-angle trend) and the measured anatomy stay in the diary and the paper; finite Λ -sums are allowed and used, *no* RH statement is made ($\frac{1}{2}$ appears in exactly one role: as the strength of a classical statement a required depth is compared against); **the one open object after T170 stays open, typed open:** R1, now finally *classified* but not closed — an m -free unconditional certificate that the two rows of $\hat{A}$ become collinear at rate $h^{-3+\varepsilon}$, equally a joint near-degeneracy of the three explicit linear Λ -sums (S_{11}, S_{22}, S_{12}) against the archimedean block — a rate statement the module neither makes nor approaches; the T168/T169/T170 verdicts (MINORS-RESIST / RATIO-RESISTS / SIEVE-RESISTS) are sandbox verdicts. Fences: Lagrange 1773 / Cauchy–Binet 1812–1815 / Gershgorin 1931 / Rayleigh 1877–Courant–Fischer 1920 / Montgomery–Vaughan 1973–Vaughan 1977 (the priced route) / KMS 1953 / Chebyshev 1852–Rosser–Schoenfeld 1962 (verified on the table) / Dirichlet 1829 named classical; Weil 1952 / Bombieri 2000 cited, never used as a criterion; zero-firewall AST-checked; no marker moves. Machine-checked: [verification/v558_bilinear_rank_identities.py]. [E] (per-instance identities and certified inequalities)

The phase-2 capstone (PRIME.PHASE2.CAPSTONE.01). Promotion of discovery T171 (contract FINAL.MAP, verdict MAP-COMPLETE) as one module (35 checks, ~ 0.3 s): *the capstone verifies the map and claims nothing new* — the whole sixteen-link reduction chain of phase 2 as one machine-checked pass, the eight-route no-go battery, and the precision ledger, recomputed on the declared small frame-A surface (12 deepest in-cap zones, $m = 96 \dots 291 \leq 300$; $\kappa = 0.038821$ verified at every jump point of $\psi(x)/x$ on the table; the 16×16 low block positive definite on 12/12 — a numerical fact, never routed through the Weil criterion). (1) The chain, 13 THEOREM / 3 CERT: T152 Schur two-block (KMS 1.3×10^{-15} ; holds at $0.5\lambda_{\min}$, fails at $1.02\lambda_{\min}$ — sharp; Haynsworth 3.8×10^{-15}); T151 Sobolev $b_k \leq C_S k$, $C_S = (2\pi + 2)/\sqrt{N}$ (worst ratio 0.4816); T153 Ψ -collapse $\Psi = \sum_d c_d w_d = x^T G x$ (6.3×10^{-16} ; $\sum_d w_d = 0$ to 9.0×10^{-17}); T154 Ritz ceiling (interlacing; $g_{32}/s = 0.9520\text{--}0.9849$ from below); T155 12×12 floor [CERT] (holds at $(1 - 10^{-9})\lambda$, fails at $(1 + 10^{-4})\lambda$; $t = 0.2295\text{--}0.3719$); T156 $F(P, r)$ [CERT] ($P_K \geq 1$, $r_K \geq 0$, $L(16) = 6.6\text{--}15.6$); T158 Thomson/ladder (3.1×10^{-12} ; increments strictly positive; 36/36 trials below); T160 sampling (closed Dirichlet weights = correlation weights 1.1×10^{-15} ; atom half = sampled Λ -mass 9.9×10^{-15}); T163 TV floor/swap (Abel 1.8×10^{-11} ; overshoot $\times 5.2 \times 10^4\text{--}1.1 \times 10^6$); T164 gauge/quantifier (invariance 2.2×10^{-11} ; $\Psi(x^*) = 1/g_{16}$ to 7.0×10^{-12} ; 48/48 perturbed larger); T165 P_{pr} (1.1×10^{-16} ; = 1 at x^*); T166 cascade $1/(1 - R_{16}^2)$ (1.8×10^{-12} ; diagonal-invariant); T167 $K = 2$ exactness (6.8×10^{-13} ; milder $\times 1.62\text{--}5.20$); T169 det collapse R4-free ($\leq 4.2 \times 10^{-12}$; no positivity of A_h consumed, the Weil fence never approached); T170 rank-3 (J -eigenvalues $\{-2, -1, +1\}$ exact; $\sigma_4/\sigma_1 \leq 1.6 \times 10^{-16}$); T170 R1 shape [CERT] (the 2×2 Lagrange identity 3.6×10^{-16} ; $\sin \angle = 1.169 \times 10^{-4}\text{--}2.467 \times 10^{-3} \leq 10^{-2}$ per window; rate and quantifier OPEN). (2) The no-go battery, 8/8 failing as classified: size budget $\times 2.7 \times 10^3\text{--}1.6 \times 10^4$; symbol infimum $-47.6\text{--}-21.3 < 0$ on 12/12; sector change an invariance ($\leq 4.2 \times 10^{-12}$); perturbation $\times 9.8 \times 10^2\text{--}2.0 \times 10^4$ off scale (Kato 1949, the closed-off route); atom band dominates $\times 1.71\text{--}2.83$; sieve operator-norm on $\det S$ alone $\times 8.8 \times 10^3\text{--}7.5 \times 10^4$ over (by the rank-3 theorem a bound on $\det S$ alone cannot see the three-term cancellation; Montgomery–Vaughan 1973 / Vaughan 1977 the priced route); scramble $\times 679$ median (range $68\text{--}2.3 \times 10^6$) with rank-3 *untouched* on all 60 draws; L_P gain = 1 to 0.0. (3) The precision ledger: needed 2.752×10^{-4} at the deepest window ($m = 285$) — finer than the RH yardstick $h^{-0.5}$ by $215\times$ and the best unconditional $h^{-0.996}$ by $13\times$ on this surface; both self-similarity identities (6.8×10^{-13} / 3.1×10^{-13} , the t^* overshoot ≥ 0 everywhere, sign-agnostic). Mutations fail loudly (wrong regression row, wrong Thomson entry, off-by-one swap,

wrong polarisation coefficient — the 10^{-2} bar missed by orders). **Named limits as load-bearing content:** per instance on small windows — a finite list with an explicit maximum; nothing uniform in the zone index or in D ; the T171 trend fits (pooled sin-angle $h^{-2.33}$, frame-split $\nu = 4$: $h^{-1.86}$ / $\nu = 6$: $h^{-2.87}$, the no-go growth exponents, $\delta = +1.044$ against 3.0, the full-surface precision numbers 2.284×10^{-7} / $1.2 \times 10^5 \times$ / $3.1 \times 10^3 \times$ at $h = 1444$) are sandbox numbers on T171's two-frame surface and are *not* consumed; **zero new uniform-in- m statements are added — which is exactly why the capstone is promotable; the one open phase-2 object stays open, typed open:** R1 — an m -free unconditional certificate that the two rows of $\hat{A}$ become collinear at rate $h^{-3+\varepsilon}$, a joint near-degeneracy of the three explicit linear Λ -sums against the archimedean block; R2/R3 booked (T169), R4 off the path (T170-TH6); the T171 verdict MAP-COMPLETE is a verdict about the *map*, not about R1, and stays a sandbox verdict ($\frac{1}{2}$ appears in exactly one role: as the strength of a classical statement a required precision is compared against). Fences: Schur 1917–Haynsworth 1968 / KMS 1953 / Cauchy 1829–Courant–Fischer 1920 / Rayleigh 1877 / Abel 1826 / Dirichlet 1829 / Lagrange 1773 / Cauchy–Binet 1812–1815 / Gershgorin 1931 / Kato 1949 / Montgomery–Vaughan 1973–Vaughan 1977 / Chebyshev 1852–Rosser–Schoenfeld 1962 / Fejér 1915 named classical; Weil 1952 / Bombieri 2000 cited, never used as a criterion; zero-firewall AST-checked; no marker moves. Machine-checked: [verification/v559_phase2_capstone.py]. [E] (per-instance identities and certified inequalities)

The frame/deficit identities (PRIME.FRAME.DEFICIT.01). Promotion of the theorem cores of discoveries T172 (contract FRAME.BEYOND, verdict PARTIALLY-PORTABLE), T173 (FRAME.RATE, DEFICIT-VARIES) and T174 (CANCEL.IDENTITY, PARTIAL-CANCEL) as one module (24 checks, ~ 0.5 s): *the gauge route of phase 2, closed as algebra* — the companion to the capstone above, certifying what T172/T173/T174 established *about* the map's frame dependence, everything recomputed on a small declared surface (a 4×3 (anchor, h) rectangle of gap-blind frame-B cells, prime-power anchors $n = 41 \dots 557$ crossed with the declared ladder $h = (160, 226, 320)$, comb complete on every cell, positive definite on 12/12; one non-prime-power instance $n = 210$; one truncated-comb stress instance $n = 1009$; $\kappa = 0.038821$ verified at every jump point of $\psi(x)/x$; the builder's signature is $(n_{\text{zone}}, \alpha, M)$ — no frame label is an argument of any observable). (1) The identity transfer + the sieve-horizon discriminator: the KMS/Schur pair (3.2×10^{-16} ; quadratic-form pair 5.2×10^{-15}), the lag-sum / correlation identity (1.4×10^{-16} of the larger half; Abel 1826), the R4-free det collapse (at 0.020 of the conditioned bar $64\varepsilon/|1 - r_{12}^2|$) and the additive split (7.6×10^{-16}), re-derived *exactly* on a gap-blind frame-B instance *and* a non-prime-power-anchor instance — the links use no property of the frame rule or the anchor arithmetic, which is why they transfer; the truncated-comb instance turns *indefinite* ($\lambda_{\min} = -9.6 \times 10^4$) against 12/12 positive definite complete-comb cells, and the identity links keep holding there (none uses positivity): indefiniteness sits at the declared sieve horizon, not at a frame or zone. (2) The $q = 1 - s$ identity + the calibration: $hD = \alpha$ per cell (8.9×10^{-16}) forces the OLS slopes to obey $q = 1 - s$ to 2.9×10^{-14} (OLS is linear in its response); the gauge-degree account comes out as integers — $G2 = t/\lambda_{\max}(B_{LL})$ and the delivery $1 - r_{12}^2$ both $(d_{\text{amp}}, d_{\mu}) = (0, 0)$, $G1$ at $d_{\text{amp}} = 1$, $G3$ at $d_{\mu} = 1$ — so $G2$ is the unique doubly gauge-invariant arithmetic member of T173's family; $G3 = \mu_1^P G2$ identically (2.2×10^{-16} ; exponent $h^{-1.9955}$, the offset-2 explained); $(D/\alpha)^3 h^3 = 1$ a tautology (2.2×10^{-16}). (3) The blindness theorems + the additive limit: the amplitude $c \rightarrow \kappa c$ is a gauge of GAP, of $1 - r_{12}^2$ and of R over twelve decades (worst 0.026 of the per-cell conditioned bar; rebuilt from scratch 0.098); the scalar $\mu_1^P = 4 \sin^2(\pi/(2h + 1))$ — *the entire h^{-2} channel of the demand* — cancels exactly on both sides (0.001 of bar); the delivery is invariant under the full positive diagonal group (48 congruences spanning $e^{\pm 8}$, worst 0.036 of bar) while a non-scalar diagonal moves the demand by 0.23–1.40 (the asymmetry is the anatomy); neither the arch term nor the comb term alone has a positive definite low block on a single cell (0/12 + 0/12 against 12/12 for the sum): no factorisation $\hat{A} = (\text{frame}) \times (\text{arithmetic})$ exists, so T173's P6 is *not* promoted to a theorem — only its multiplicative half. (4) The KMS shape band (unconditional, declared $\varepsilon = (K\pi/(2h + 1))^2/6$): $|\sqrt{\hat{S}_k/k^2} - 1| \leq \varepsilon_{\text{cert}}$ on 12/12 and the k^2 -swap moves log GAP by ≤ 0.312 of the certified band $2 \log((1 + \varepsilon)/(1 - \varepsilon))$ (Rayleigh 1877; Schur 1917) while the delivery does not move at all. (5)

The placement discriminator: the value-multiset scramble destroys the positive low block on 24/24 declared draws and moves $|R|$ by a median factor 10^6 . Mutations fail loudly (the $d = 0$ halving dropped, the $N = 2h$ ladder, the off-by-one h , the comb-only rescale, the wrong universal profile). **Named limits as load-bearing content:** per instance on a small declared surface — a finite list with an explicit maximum; nothing uniform in the zone index or in D ; **the frame-free deficit is a measurement, not a claim of this row:** T174's $+0.1111 \pm 0.0222$ (5.0σ , 151 anchor clusters, cluster-robust; 0.4σ from T173's $+0.155 \pm 0.102$) stays a sandbox MEASURED number, as do all T172/T173/T174 h -exponents, the density anatomy and the ν /density collinearity; **the one open phase-2 object stays open, typed open:** R1, now stated frame-free — an m -free unconditional certificate that two explicit finite Λ -sum vectors become collinear at the demanded rate; the residual per-anchor scatter ($\chi^2/\text{dof} = 3.02$) and the sparse corner are sandbox items under test in T175 ($\frac{1}{2}$ appears in exactly one role: as the strength of a classical statement a required precision is compared against). Fences: Schur 1917 / KMS 1953 / Dirichlet 1829 / Rayleigh 1877 / Abel 1826 / Chebyshev 1852–Rosser–Schoenfeld 1962 / Wilkinson 1968–Higham 2002 named classical; Weil 1952 / Bombieri 2000 cited, never used as a criterion; zero-firewall AST-checked; no marker moves. Machine-checked: `[verification/v560_frame_deficit_identities.py]`. **[E]** (per-instance identities and certified inequalities)

Lemma 10 (G3–G4 — reflection positivity, fixed and integrated). *For each Θ -invariant g in the Calderón ball the physical boundary kernel $K_g = P_{\text{adm}}\Pi_{\text{phys}}\mathcal{N}_g^{TT}\Pi_{\text{phys}}P_{\text{adm}}$ is reflection positive, $\langle \Theta F, F \rangle_{K_g} \geq 0$ (G3); and $d\mu_{\text{Cal},\chi}$ is Θ -invariant, supported where G3 holds uniformly, so $\int \langle \Theta F, F \rangle_{K_g} d\mu_{\text{Cal},\chi} \geq 0$ (G4). **G4 is the decisive step:** RP after integrating over dynamical metrics, not merely at fixed background.*

Lemma 11 (G5 — gap dominance **[E]**). $2\|V_{\text{metric}}\|_{\text{rel}} < \Delta = 6 \log \frac{3}{2}$, with $\|V_{\text{metric}}\|_{\text{rel}} \leq 248c_3^2 = \frac{31}{8\pi^2} = 0.39262$, hence $2\|V\| = 0.7852 < 2.4328$ and $\Delta_{\text{eff}} \geq \Delta - 2\|V\| > 1.647 > 0$. *The numerical inequality is trivial (`[verification/v29_research_contract_certs.py]`); the work is the operator-norm bound (hard).*

Sugawara reading + atomic OS moment (`[verification/v171_os_moment_cluster.py]`). The G5 budget is the E_8 Sugawara constant: $248c_3^2 = \frac{31}{8\pi^2}$ with $31 = 1 + h^\vee(E_8)$ and $c((E_8)_1) = 248/31 = 8$, so gap dominance reads $|R^+(A_3)| \log(N_{\text{fam}}/|\mathbb{Z}_2|) > \frac{1+h^\vee(E_8)}{4\pi^2}$ — the A_3 transfer gap beats the E_8 Sugawara metric budget. Independently, the G3 reflection positivity is a *positive moment problem*: $\text{spec}(T) = \{1, (2/3)^6, (1/3)^6\}$ makes every two-point correlator $C(n) = A + B(64/729)^n + C(1/729)^n$ the moment sequence of a positive atomic measure, so each OS Hankel matrix factorises as a Vandermonde Gram and is positive semidefinite (a clean RP certificate; cluster constant 729/665, correlation length $\xi = 1/(6 \log \frac{3}{2}) = 0.4111$).

Theorem 1 (Decoupling theorem — physical low-energy claims do *not* need the full G_{metric}). *If the admissible sector is reflection-positive (G4) and the metric coupling is gap-dominated, $2\|V_{\text{metric}}\| < \Delta = 6 \log \frac{3}{2}$ (G5), then the admissible IR measure is stable under the ambient completion: a strictly positive effective gap $\Delta_{\text{eff}} = \Delta - 2\|V\| > 0$ protects the admissible sector from the (un-built) ambient/deep-UV, so every physical low-energy read-out (masses, mixings, α^{-1} , the $R+R^2$ regime) is independent of the open ambient measure. Symbolically*

$$\boxed{\text{RP} + 2\|V\| < \Delta \implies \text{admissible IR measure stable under ambient completion.}}$$

*Thus (G_{metric}) is not a diffuse “full quantum gravity missing” block: it is the sharply delimited ambient projective limit (G6), a strict-TOE certification target whose closure the falsifiable IR physics does not depend on. **Typing:** the inequality is machine-checked **[E]**; the operator-norm bound on $\|V_{\text{metric}}\|$ is the conditional input.*

Lemma 12 (G6–G7 — projective limit & Ward identities). *A projective system μ_{χ_n} with $\pi_{m,n*}\mu_{\chi_m} = \mu_{\chi_n}$, uniform moment bounds $\sup_n \int \|g\|_{H^s}^p d\mu_{\chi_n} < \infty$, tightness and regulator-independence yields $\mu_{\text{QG}} = \lim \mu_{\chi_n}$ on $\mathcal{G}/\text{Diff}$ (G6, the hardest step); the limit obeys the BRST Ward identities $\langle s\mathcal{O} \rangle = 0$, whence $\nabla^\mu T_{\mu\nu} = 0$ and the η boundary phase cancels the APS anomalies (G7).*

Gate 2 reduction: G6 is a *boundary projective limit*, not a bulk one ([verification/v76_gmetric_reduction.py])

The exact decoupling margin is $\Delta_{\text{eff}} = \Delta - 2\|V\| = 6\log \frac{3}{2} - \frac{31}{4\pi^2} = 1.648 > 0$, with $\|V_{\text{metric}}\|_{\text{rel}} \leq 248 c_3^2 = \dim(E_8) c_3^2 = \frac{31}{8\pi^2} (31 = 2^{g_{\text{car}}} - 1)$. **Holographic reduction:** because the seam is a *finite causal boundary*, the ambient *bulk* metric measure is reconstructed from a finite-codimension seam (Calderón) *boundary* measure — so G6 is a *boundary projective limit*. The conditional theorem is then

$$\text{RP}(\text{seam-Calderón boundary kernel}) + \text{tightness} \implies G_{\text{metric}} \text{ closes}$$

i.e. the dimension of the open problem drops from bulk QG to a finite seam-boundary measure.

Residual: the constructive boundary projective limit (constructive-QFT grade) — now discharged as a [C] redundancy (v369, keybox below). So Gate 2 is now *two clean tiers*: Tier A (IR physics) closed under RP+gap; Tier B (strict TOE) a [C] redundancy, no longer a diffuse “full QG missing”. [C]

The measure as one-loop fluctuations around the parameter-free saddle ([verification/v365_qg_oneloop_saddle.py], QGAMB.SADDLE.01)

With the saddle now parameter-free (v358/v359), the boundary measure of Tier B organises as a Gaussian one-loop fluctuation determinant whose ingredients are atom-fixed. **Fixed quadratic form [E]:** the fluctuation inverse-propagator stiffness is the saddle coefficient $1/c_3 = 8\pi$ — no free dimensionless dial, so the one-loop problem is *well-posed* (it was not before v358). **Gap-controlled convergence [E]:** the fluctuation operator obeys $M \geq \Delta_{\text{eff}} = 6\log \frac{3}{2} - \frac{31}{4\pi^2} \approx 1.648 > 0$ (v76/v337), so the Gaussian integral converges mode-by-mode with no zero/negative modes besides the saddle directions; on the OS spectrum $\{0, 6\log \frac{3}{2}, 6\log 3\}$ the regularised one-loop log-det $\text{Tr}' \log M = 6\log \frac{9}{2}$ is finite. **Conformal direction [C]:** the one wrong-sign mode ($c_{\text{conf}}(4) = -\frac{3}{2}$) is made convergent by the GHP contour ($\int e^{-|c|\phi^2} = \sqrt{2\pi/3}$) with IDG dressing (v334). **Tightness [E]:** $\chi = 1/(1 - (2/3)^6) = 729/665 < \infty$ (v330), so the one-loop measure has a projective limit on the admissible sector, with *zero* new dimensionless dials (v364). **Residual:** the full *non-perturbative* projective limit (G6 on the ambient sector) is now discharged as a [C] redundancy (v369, keybox below) — this sharpens C7 from “diffuse full QG” to a one-loop-controlled Gaussian problem around a parameter-free saddle and then to a certification object. [E]/[C]

The holographic route: the ambient measure is boundary-redundant, not missing dynamics ([verification/v369_qgamb_redundancy.py], QGAMB.REDUNDANCY.01)

There is a second, strategically stronger way to dispatch Gate 2 than building the measure: prove that the ambient sector carries *no* physical degrees of freedom relative to the holomorphic boundary algebra, so the measure QG.AMB.01 is a non-fundamental *certification* object — the role a coordinate choice plays in general relativity — rather than a missing piece of dynamics. The discriminators that make this *ambient-redundancy* statement well-posed are all in hand. **Trivial superselection [E]:** the number of DHR sectors of a chiral net is $|\det \text{Cartan}|$, which is 1 for E_8 (one primary, the vacuum; $\text{DHR}((E_8)_1) = \text{Vec}$) against 4 for the same- c rival $SO(16)_1$ — a holomorphic boundary net has *no* nontrivial superselection charge that a bulk could carry invisibly. **No torus degeneracy [E]:** the ground-state degeneracy on T^2 equals the number of anyons $= |\det K| = 1$ for E_8 , so there is no topologically protected bulk d.o.f. hidden from the boundary (the genus-1 obstruction of v344, in its redundancy reading). **Bounded reconstruction [E]:** the seam transfer deviation subspace contracts at the finite gapped Petz rate $(2/3)^6$ (v221), and the gap-decoupling margin $\Delta_{\text{eff}} = 6\log \frac{3}{2} - \frac{31}{4\pi^2} \approx 1.648 > 0$ (v76/v337) separates the un-built ambient sector from the admissible one. **The statement [C]:** with the Bisognano–Wichmann modular reconstruction map (v258/v309/v323), under SEAM.EQUIV.01 every gauge-invariant bulk observable is boundary-reconstructible, $\mathcal{O}_{\text{phys}}^{\text{bulk}}/\text{Diff} \cong \mathcal{A}_{\Sigma}^{\mu_4}$, and two ambient measures sharing the same boundary state are physically equivalent. **Residual [O]:** the construction consumes exactly two premises — SEAM.EQUIV.01 (the S3 lattice input, v366) and the BW intrinsicity of the raw collar (v329); it does *not* construct QG.AMB.01 but supplies the *alternative* route, so that Gravity-complete = Boundary-complete + Ambient-redundancy. [E]/[C]/[O]

The classical field equation, parameter-free: entanglement equilibrium gives the linearised Einstein equation with G from c_3 ([verification/v358_grav_entropy_equilibrium.py])

While Gate 2 (above) reduces the metric-sector *measure*, the classical metric-sector *field equation* is supplied directly by the entanglement first law $\delta S = \delta \langle K \rangle$ (Jacobson 2015; Faulkner et al.), carried out with TFPT's atoms. **Parameter-free coefficient [E]**. Equilibrium gives $G_{ab} = c_3^{-1} T_{ab}$ with $c_3^{-1} = 8\pi$ *fixed*: all four c_3 -roles are mutually consistent — the Unruh factor $1/(2\pi) = 4c_3$, the EH action $1/(16\pi) = c_3/2$, the Einstein coefficient $8\pi = 1/c_3$, the Bekenstein density $1/4 = 1/|\mu_4|$ — so *no free dimensionless Newton dial* enters. **Thermo = geo coincidence [E] (new)**. The first law requires the coefficient $2\pi/\eta$ (η the entropy density); the geometry gives $1/c_3 = |\mathbb{Z}_2| 2\pi \chi(S^2)$ (one-sided Gauss–Bonnet, v58); these are the *same* equation iff $|\mu_4| = |\mathbb{Z}_2| \chi(S^2) = 4$ — a TFPT atom identity (v216). So the thermodynamic (first-law) and geometric (Gauss–Bonnet) origins of c_3 are *one*. **The matter side, assembled [C] (closing J3)**. The ball modular Hamiltonian is the boost (Casini–Huerta–Myers; geometric by Bisognano–Wichmann, v323), $K_B = 2\pi \int_B w T_{00}$, $w = (R^2 - r^2)/(2R)$; the 3-ball weight integral is $\int_B w d^3x = 4\pi R^4/15$, so $\delta \langle K_B \rangle = \frac{8\pi^2 R^4}{15} \delta \langle T_{00} \rangle$ — the concrete matter boost flux of TFPT's stress tensor. **The entropy density, atom-fixed [E]**. The Bekenstein numerator $1/4 = 1/|\mu_4|$ (v57) and the entanglement normalisation = the central charge $c = g_{\text{car}} + N_{\text{fam}} = 8$ pin the dimensionless coefficient; the area-law form is established (v59). **Residual**: the full covariant extension is now *closed* (v359, next keybox); what remains is only (i) the Jacobson equation-of-state *status* (a thermodynamic derivation, not a from-action quantisation) [O] and (ii) the absolute scale v_{geo} (the Planck-area unit) [O], while the global ambient measure (Gate 2/C7) is now a [C] redundancy (v369) — *no free dimensionless parameter survives in the gravity coupling*.

And that boundary measure is a rigorously constructed net ([verification/v77_e8_conformal_net.py]). The level-1 central charges $c = \dim \mathfrak{g}/(1 + h^\vee)$ give $c(E_8)_1 = \frac{248}{31} = 8 = \text{rank } E_8$, $c(D_5)_1 = \frac{45}{9} = 5$, $c(A_3)_1 = \frac{15}{5} = 3$, and $c(D_5) + c(A_3) = 5 + 3 = 8 = c(E_8)$ is a *conformal embedding* $(D_5)_1 \times (A_3)_1 \subset (E_8)_1$ (coset $c = 0$). Now $(E_8)_1$ is the holomorphic $c = 8$ E_8 -lattice VOA / conformal net — one of the most rigorously constructed objects in mathematical QFT (Frenkel–Kac–Segal; Kawahigashi–Longo / Carpi). **So the conditional theorem becomes concrete: identify the seam-Calderón boundary measure with the $(E_8)_1$ lattice net** (carrier = $(D_5)_1 \times (A_3)_1$ subnet), and the gap $\Delta_{\text{eff}} > 0$ supplies the clustering/tightness — so G6 is imported into existing RCFT/conformal-net rigor rather than built from scratch. Residual = the net identification + bulk reconstruction. [E] (charges, embedding) + [C] (identification)

The full covariant field equation, parameter-free: fixed-volume equilibrium gives the Einstein tensor with conservation as an output ([verification/v359_grav_nonlinear_einstein.py])

The linearised result above (v358) extends to the *full* covariant equation by Jacobson's *fixed-volume* (maximal-vacuum-entanglement) equilibrium — no linearisation assumed. **Einstein tensor, not Ricci [E]**. Stationarity of the entanglement entropy at fixed volume forces the divergence-free combination: the 4D trace identity $g^{ab} G_{ab} = (1 - \frac{d}{2})R = -R$ (for $d = 4$) selects $G_{ab} = R_{ab} - \frac{1}{2}R g_{ab}$, so

$$G_{ab} + \Lambda g_{ab} = c_3^{-1} T_{ab}, \quad c_3^{-1} = 8\pi$$

with *both* coefficients TFPT-fixed: the Newton coupling $8\pi = 1/c_3$ (v358) and the cosmological constant Λ from α , $\rho_\Lambda = (3/4\pi^2)e^{-2\alpha^{-1}}$ (v60). **Conservation is an output [E]**. By Lovelock's theorem the only divergence-free, second-order, local symmetric tensor built from the metric is $a G_{ab} + b g_{ab}$, so $\nabla^a G_{ab} = 0$ forces $\nabla^a T_{ab} = 0$ — matter conservation is *derived*, not assumed. **So Gate B6 — the classical metric-sector field equation — is closed parameter-free at the local level. Residual**: the Jacobson equation-of-state *status* (a thermodynamic derivation, not a from-action quantisation) and the single unit v_{geo} stay [O], while the global ambient measure (Gate 2/C7, the constructive boundary projective limit) is now a [C] redundancy (v369) — no free dimensionless gravity dial remains. An *external candidate* for the missing action level is quantified in the next keybox (v473); it does not change this typing. [E]/[C]

The entropic-action bridge: an external candidate for the missing action level, quantified — nothing closes ([verification/v473_entropic_action_bridge.py])

The one honest **[O]** left in the classical sector — “equation of state, not a from-action quantisation” (v359) — now has an explicit *external* candidate: Bianconi’s entropic action $S_B = -\text{Tr}_F \ln(\tilde{G}\tilde{g}^{-1})$, the quantum relative entropy between the spacetime metric and the matter-induced metric (G. Bianconi, *Gravity from entropy*, Phys. Rev. D **111**, 066001 (2025); [arXiv:2408.14391](#); her Appendix C identifies $\Delta_{\tilde{G},\tilde{g}}^{1/2} = \tilde{G}\tilde{g}^{-1}$ with an Araki-type modular operator) — a local covariant action whose variation yields the Einstein equations plus an emergent nonnegative Λ at low coupling: exactly the missing variational layer, *if* the bridge identities hold. What is machine-checked today (GRAV.ENTROPIC.ACTION.01): **carrier Hodge count [E]**. Her $\Omega^{0,1,2}$ matter fiber, placed on the five-slot carrier $\mathbb{C}^5$ (*not* on 4d spacetime, where the fiber is $1+4+6=11$), counts $1+5+10=16=2^{g_{\text{car}}-1} = \dim S_{D_5}^+$ and Hodge-folds ($\star: \Lambda^1 \cong \Lambda^4$) onto $\Lambda^{\text{even}}\mathbb{C}^5$ — the carrier half-spinor’s own Pascal row (v2/v44); a vector-space identity *only*. **Coefficient match [E] (the one quantitative pin)**. Her weak-coupling Lagrangian $\mathcal{L} = 3\beta R - \alpha\mathcal{L}_m + \dots$ (the 3 = the three form blocks) matched to the TFPT EH normalisation $1/(16\pi G) = c_3/(2G)$ fixes her free constant *exactly*,

$$\beta'_B = \frac{c_3}{6} = \frac{1}{48\pi}$$

(the $6 = 3 \times 2 = p_2 = |R^+(A_3)|$ is recorded as a structural echo, not a proof). **Entropy potential = quadratic Λ [E]**. Her $\Lambda_G = \frac{1}{2\beta} \text{Tr}_F(\mathcal{G} - I - \ln \mathcal{G})$ is nonnegative and quadratic at the fixed point; with the TFPT displacement $\varepsilon_\star = e^{-\alpha^{-1}Q}$ it reproduces the closed Λ branch $\rho_\Lambda/\bar{M}_{\text{Pl}}^4 = (3/4\pi^2)e^{-2\alpha^{-1}}$ (v60), and the prefactor match yields the *exact* normalisation target $\text{Tr}_F Q^2 = 32c_3^4 = 1/(128\pi^4) = \frac{2}{3}\delta_{\text{top}}$ — a closed algebraic target for the two-form (pair) block ($\binom{5}{2}=10$), **[C]** until an explicit Q is constructed. **R^2 kill test [E] (pre-registered)**. The raw log expansion carries a curvature-squared coefficient of order $(c_3/6)^2$; the TFPT Starobinsky sector needs $\bar{M}_{\text{Pl}}^2/(12M_{\text{scal}}^2) = 1/(12c_3^7)$ (v36) — the exact mismatch is $3c_3^{-9} = 3(8\pi)^9 \approx 1.2 \times 10^{13}$: a direct identification of the raw entropic action with the TFPT R^2 sector is *false* without a mechanism (the scalaron as a light trace mode of her $\mathcal{G}$ -field generating c_3^{-9} , or a KMS-spectral renormalisation of the log action). **The bridge proper [C] (recorded, not proven)**. The compression conjecture $P_\Sigma(\tilde{G}\tilde{g}^{-1})P_\Sigma = \Delta_\Sigma^{1/2}$ — the physical Calderón compression of her relative metric operator equals the TFPT seam modular operator; if proven, S_B is a local volume representation of the TFPT relative spectral action $S_{\text{rel},\chi}$ (her log action is the all-scale-integrated heat kernel, by Frullani $-\ln A = \int_0^\infty \frac{dt}{t}(e^{-tA} - e^{-t})$, verified). Open with it: the operator-level Hodge items (Clifford action, Dirac intertwining, leaf involution, Calderón polarisation), and Lorentzian positivity of $\tilde{G}\tilde{g}^{-1}$ is *not* automatic (timelike gradients) — TFPT’s OS/reflection-positive route is the sturdier construction. **Residual [O] (unchanged)**. Nothing closes: the v359 equation-of-state typing stays **[O]**; adopting S_B would close it *only* after the compression conjecture *and* an R^2 mechanism; zero TFPT knobs are added.

The operator level, executed ([verification/v474_entropic_hodge_carrier.py]). Work packages 1 and 4-algebra of the bridge are now machine-checked on the finite carrier Fock space $\Lambda^\bullet\mathbb{C}^5$ ($\dim 2^{g_{\text{car}}}=32$). **Spinor structure exhibited [E]**: the ten CAR gammas satisfy $\{\Gamma_a, \Gamma_b\}=2\delta_{ab}$ exactly, the 45 bilinears span $\mathfrak{so}(10)$ and leak nothing from the 16-dim even subspace — S^+ is the even exterior algebra with its spinor structure, and Bianconi’s Hodge–Dirac operator is Clifford multiplication at symbol level ($c(\xi)^2 = -|\xi|^2$, symbolic). **The fold is the $5 \rightarrow \bar{5}$ conjugation [E]**: the traceless $\mathfrak{u}(5)$ weights of Λ^4 are the negatives of those of Λ^1 , so her fiber $1+5+10$ becomes the GUT half-spinor $16=1+\bar{5}+10$ — the SM carrier decomposition, not a dimension match; honest obstruction: $\Omega^{\leq 2}$ is *not* Clifford-invariant, so the identification must go through the fold, never through degree truncation. **The Q -target, decided [E]**: the uniform-block ansatz $\text{Tr} Q^2 = dk^2 c_3^4 = 32c_3^4$ with integer k admits *exactly* the supports $(d,k)=(2,4), (8,2), (32,1) = \{|\mathbb{Z}_2|, \text{rank } E_8, 2^{g_{\text{car}}}\}$, with the minimal uniform solution $q=c_3^2$ per Fock mode and the identity $\text{Tr} Q^2/\delta_{\text{top}}=32/48=2/3$; the two-form block 10 (the naive pair-sector reading) and the fiber 16 require irrational multipliers and are *excluded*. The physical choice among the three supports stays **[C]**; AP2 and the R^2 mechanism stay **[O]**.

The R^2 kill test, executed ([verification/v475_entropic_scalaron_gate.py]). On a maximally symmetric background the relative curvature operator has eigenvalues *exactly* $\{R; R/4 \times 4; R/6 \times 6\}$ (her Appendix-B flattening verified, incl. the 6×6 two-form block), so the exact vacuum action is $\mathcal{L}_B(R) = -[\ln(1-\beta R) + 4\ln(1-\beta R/4) + 6\ln(1-\beta R/6)] = 3\beta R + \frac{17}{24}\beta^2 R^2 + O(R^3)$ — the 3 re-derives her Eq. (45) and $\frac{17}{24}$ is the exact tensorial factor behind the pre-registered gap. With the pinned $\beta'=c_3/6$

both mass routes (Starobinsky matching and $f'/3f''$) give the raw entropic scalaron

$$m_{\text{raw}}^2 = \frac{4608\pi^2}{17} \bar{M}_{\text{Pl}}^2 \Rightarrow m_{\text{raw}} \approx 51.7 \bar{M}_{\text{Pl}}$$

— *trans-Planckian*, not light: the naive reading “the TFPT scalaron is the light trace mode of her $\mathcal{G}$ -field” is *dead* [E]; a viable mechanism must supply exactly $m_{\text{raw}}^2/m_{\text{TFPT}}^2 = \frac{72}{17}(8\pi)^9 \approx 1.69 \times 10^{13}$ in mass². The domain is not the obstruction (first log pole at $R=384\pi^2 \bar{M}_{\text{Pl}}^2$, trans-Planckian), and the Lorentzian caveat is now an explicit witness: a timelike gradient gives the Gg^{-1} eigenvalue $1 - \alpha v^2 \leq 0$ for $v^2 \geq 1/\alpha$ (ln undefined), while spacelike/Euclidean stay positive — the OS route is the sturdier construction. **Net verdict** [C]: of the three bridge interpretations, the entropic action covers the Einstein + Λ level (survives); the light-mode reading is dead; KMS-spectral renormalisation is the *only* surviving route to the R^2 sector. Nothing closes.

AP2, made well-posed ([verification/v476_entropic_compression_toy.py]). The compression conjecture itself is sharpened on an exactly solvable gapped chain. **The reading is decided:** on a *pure* bulk (the seam situation: pure bulk, thermal boundary) the covariance is a projection, so $f(x)=x/(1-x)$ is singular on its spectrum — the literal operator-side reading $P_\Sigma f(C)P_\Sigma$ is *ill-posed*, while the state-side $f(P_\Sigma C P_\Sigma)$ (Peschel) is finite: the identity can only mean *build Δ_Σ from the compressed relative metric*, which is exactly Bianconi’s own local construction — AP2 retyped, not weakened. **The mismatch is exactly second order** [E]: symbolically, $[C^n]_{AA} - (C_{AA})^n$ carries the cross-cut block twice (first order vanishes identically), so for any analytic f the two readings agree up to $O(\|C_{AB}\|^2)$ — the modular (entanglement) content *is* that quadratic correction. **Gap control:** at the bounded (covariance) level the mismatch obeys $d \leq x^2$ and falls with the gap, the decoupled cut agrees exactly — the two readings converge in the gap-dominated regime, where TFPT operates ($\Delta_{\text{eff}} > 0$, v76/v337). **Residual** [O] (**unchanged**): the continuum seam statement itself. No marker moves.

The surviving R^2 route, typed as one moment ([verification/v477_entropic_scaleflow.py]). The Frullani identity upgrades to a *scale-flow representation*: $-\ln a = \int_0^\infty \frac{dt}{t} (e^{-ta} - e^{-t}) = 2 \int_0^\infty \frac{dx}{x} (e^{-a/x^2} - e^{-1/x^2})$ — Bianconi’s log action *is* the flat scale-integral of relative heat-kernel actions, while TFPT’s $S_{\text{rel},\chi}$ evaluates *one* KMS scale with a weight f : the bridge between the two action principles is a choice of *scale measure*. In the moment dictionary (weight $w(t) dt/t$, μ_k its e^{-t} -moments) the coefficients read $3\beta R \mu_1 + \frac{17}{24} \beta^2 R^2 \mu_2$; the flat weight has $\mu_1 = \mu_2 = 1$ (= the raw v475 coefficients), and demanding $m^2 = c_3^7 \bar{M}_{\text{Pl}}^2$ forces *exactly*

$$\mu_2/\mu_1^2 = \frac{72}{17}(8\pi)^9 \quad \text{with} \quad \frac{4608\pi^2/17}{(72/17)(8\pi)^9} = c_3^7 \text{ (identically)}$$

— the v475 kill-test number *is* a moment ratio, and the “13 orders of magnitude” are not a bridge mismatch but a scale-measure datum: the flat measure fixes it wrongly, the TFPT KMS measure ($f_0=6(4\pi)^2/c_3^7$, v36) fixes it correctly — one KMS moment, two parametrisations, zero new dials. [C] (consistency, not derivation: the measure is TFPT input); the R^2 gate and the v359 typing stay [O].

The two remaining legs, first computable steps ([verification/v478_entropic_continuum_legs.py]).

Leg A (AP2 continuum): on the exactly solvable critical chain (infinite-volume kernel, 50-digit precision) the compressed interval state carries the conformal/BW structure quantitatively — the entanglement entropy follows Calabrese–Cardy with $c_{\text{est}} = 1.00089/1.00038/1.00021$ (monotone $\rightarrow$ the Dirac $c=1$; gapped control $\sim 6 \times 10^{-8}$), and its Peschel modular Hamiltonian organises on the CHM weight $\beta(x)=x(L-x)/L$ (corr 0.87 $\rightarrow$ 0.99, even-range bands vanishing identically, odd tail 1.8%): the state-side Δ_Σ (the reading *forced* above) flows to the CHM/Bisognano–Wichmann geometric form — *exactly* the object the entanglement-equilibrium derivation consumes (v323/v358); the bridge’s modular side and the gravity side meet in the continuum limit (a lattice exhibit, not a continuum proof — AP2 stays [O]; the natural continuum frame for the seam version is the *four-interval* multilocal modular geometry of the four marks, v480). *Leg B (the measure)*: the single-scale corollary is exact — $d\mu = \delta(t-t_0)dt$ gives $\mu_2/\mu_1^2 = e^{t_0}$, so the one-moment condition forces $t_0 = \ln \frac{72}{17} + 9 \ln(8\pi) = 30.461$, one dimensionless KMS time in closed form; the near-miss to $h(E_8)=30$ (difference 0.461) is explicitly *declined* per the no-free-pattern rule (v354/v355) — deriving the measure from the entropic structure stays [O]. [E]/[C]/[O]

The Simple-Current Extension Theorem — the central G_{net} closing statement [E]/[C]
 ([verification/v154_simple_current_theorem.py])

The conditional theorem can now be stated as one clean *simple-current extension*. Let $A = (D_5)_1 \otimes (A_3)_1$ be the carrier net and $L = \langle(1,1)\rangle \subset \text{Disc}(D_5) \oplus \text{Disc}(A_3) = \mathbb{Z}_4 \oplus \mathbb{Z}_4$ the isotropic μ_4 glue line ($q(k(1,1)) = k^2 \in \mathbb{Z}$). Then the local extension $B = A \rtimes L$ has

$$[B : A] = |L| = 4 = |\mu_4|, \quad c(B) = 5+3 = 8, \quad \mu(B) = \frac{\mu(A)}{|L|^2} = \frac{16}{16} = 1$$

so B is *holomorphic* (KLM μ -additivity), and a holomorphic $c=8$ chiral net is the lattice net of the *unique* even-unimodular rank-8 lattice — hence $B \cong (E_8)_1$ (the same- c rival $SO(16)_1$, $\mu=4$, is excluded). Every algebraic ingredient is machine-checked: the $\mathbb{C}[\mathbb{Z}_4]$ Longo Q-system ([verification/v125_glue_qsystem.py]), the $\mathbb{Z}_4$ -graded Frobenius E_8 hull ([verification/v143_graded_frobenius.py]), the Ramond glue sectors ([verification/v148_fock_census.py]). **Consequence:** G_{net} is internally *closed* if the seam–Calderón boundary net is *defined* to be A ; it stays the *one* external identification “the RP seam–Calderón net is A ” if the boundary net is to be *derived* from the seam kernel. Either way this is the single strict-TOE certification theorem still carrying weight — a clean extension statement, not “find a quantum gravity”. [2026-08-05 note, finite-level evidence for the identification ([verification/v779_gnet_gns_arf.py]): the inductive system — Haar isometry tower, commuting index-4 expectation squares, geometric state convergence — is exactly coherent; the finite Haag-duality census gives twisted-duality defect 0 with excess ratio $\dim B' / \dim A' = 4 =$ the local Watatani index; and the $\mathbb{C}[\mathbb{Z}_4]$ quasi-basis obstruction law $2^{-\ell}$ types the extension as exactly asymptotic at finite level. Typed finite evidence, never the continuum theorem; no marker move.]

And that one identification reduces to a single shared premise ([verification/v155_quasifree_boundary.py]). The identification splits (v110) into (a) “the seam Calderón involution is sheet-odd” — the double-cover deck, the same $|\mathbb{Z}_2|$ that halves $c_3 = \frac{1}{2 \cdot 4\pi}$ (structural) — and (b) “the seam 2-point kernel is quasi-free”. Part (b) is *not independent*: the boundary marginal of a reflection-positive *Gaussian* bulk is the compression PTP of the bulk covariance, and the compression of a contraction ($i\Gamma$ with spectrum in $[-1, 1]$) is a contraction — so the boundary/seam state is automatically quasi-free, with Wick inherited (every boundary correlator a Pfaffian of Γ_d). Hence (b) *follows* from “the seam bulk is a reflection-positive free field” — the *same* premise already load-bearing in the area law ([verification/v59_area_law_evidence.py]) and the EH mechanism ([verification/v150_replica_eh_model.py]/[verification/v151_bfk_split.py]; since exercised on the discretized collar with the seam’s own kernel and real replica sheets, v471 — the R3 kernel premise is discharged at the finite level, its residual retyping to the continuum scaling limit shared with SEAM.EQUIV.01, v336, plus the v152 anchor). **So the entire seam-side residual is one physical premise** — the seam bulk is a free RP field — shared by G_{net} , the area law and R3, and given it the chain Gaussian bulk $\Rightarrow$ quasi-free boundary $\Rightarrow$ one kernel = net (v113) $\Rightarrow$ carrier $A \Rightarrow (E_8)_1$ is exact. It is not reducible by a finite computation — it is the one place TFPT meets functional analysis.

The construction, exhibited and verified ([verification/v156_seam_net_construction.py]). The “ $B \cong (E_8)_1$ ” step is no longer an assertion — the chiral net is built explicitly and the identification *checked*. The Dirichlet-to-Neumann (Calderón) operator of the 2d Laplacian acts on a boundary mode e^{ikx} by the harmonic extension $e^{ikx-|k|y}$: $-\partial_n u|_0 = |k|u$, so for a free bulk the seam operator is the massless chiral dispersion $\omega_k = |k|$ that defines the free-fermion net on S^1 . Sixteen Majoranas give $c = 8 = 5+3$; the weight-1 currents are $248 = 120_{\text{NS}} + 128_{\text{R}}$ (the $SO(16)$ bilinears plus the Ramond spinor, Frenkel–Kac); and the holomorphic character is

$$\chi = \frac{E_4}{\eta^8} = q^{-1/3} (1 + 248q + 4124q^2 + 34752q^3 + 213126q^4 + \dots) = j^{1/3}$$

(verified by q -series), the 248 at level 1 being $\dim E_8$ — so the net *is* $(E_8)_1$. *What remains* is exactly (i) the free-bulk premise above and (ii) the operator-algebraic existence / uniqueness of the net generated by the seam Calderón kernel (a Kawahigashi–Longo-type net-construction theorem). Route 3 thus *exhibits and verifies* the target; the irreducible residual is the one premise + the standard net-existence step — where TFPT meets constructive QFT, now sharply located.

And the free-bulk premise itself reframes: freeness is a rigid fixed point, not a postulate

(`[verification/v157_rigid_fixed_point.py]`). Three facts click together. (i) The DtN/Calderón symbol is *universally* $|k|$: the boundary operator of *any* second-order elliptic bulk is an order-1 Ψ DO with principal symbol $|\xi|_g$ (Lee–Uhlmann), so the free chiral dispersion is bulk-detail-independent and any interaction lives in the lower-order (irrelevant) symbol. (ii) A holomorphic $c=8$ theory is *rigid*: every field has weight $(h, \bar{h}=0)$, an exactly-marginal deformation needs $(1, 1)$, and with $\bar{h}=0$ *no* $(1, 1)$ field exists — the fixed point is isolated, there is no continuous knob along which a non-Gaussian interaction could turn on. (iii) The *same* $|\mathbb{Z}_2|$ that halves the one-sided S^2 Gauss–Bonnet ($c_3 = \frac{1}{|\mathbb{Z}_2| 2\pi \chi(S^2)} = \frac{1}{8\pi}$) labels the sheet *and* is the Ramond/GSO projection giving $\mu=1$ ($248=120_{\text{NS}}+128_{\text{R}}$) — one object, three roles, so the geometry fixing c_3 is the projection making the net holomorphic. **Consequence:** premise (A) shrinks from “prove the seam is Gaussian” to “the gapped ($\Delta=6 \log \frac{3}{2} > 0$) boundary flow reaches the holomorphic fixed point” — given which, freeness is *forced* by rigidity + uniqueness ($(E_8)_1$, v83), not fine-tuned. A strictly smaller and simpler residual.

And the destination is a *proven* stable fixed point (`[verification/v158_fixed_point_stable.py]`). An exact operator-dimension count settles it: the chiral Majorana has $h=\frac{1}{2}$, the bilinear current $h=1$, the quartic $h=2$. The relevant window $0 < h < 1$ for bosonic deformations is *empty* (bosonic chiral operators have integer h , lowest 1) — nothing relevant flows the free theory away; the $h=1$ currents are chiral (v157), not $(1, 1)$ -marginal; and the lowest genuinely-interacting operator (the quartic, $h=2>1$) is *irrelevant*, as are all higher $2k$ -fermi operators ($h=k$). So the free chiral $c=8$ fixed point is *isolated and stable* against all interactions. Premise (A) thereby reduces to a *basin-of-attraction* statement with a proven stable target — the gapped ($\Delta=6 \log \frac{3}{2}$) flow lands here; the convergence is conformal-perturbation analysis, the destination is settled.

And the closure can be stated as one fixed-point theorem — which relocates (A) to a single internal premise (`[verification/v160_seam_gaussianity_from_pf.py]`). Phrase the argument internally: (i) a quasi-free state has *all* higher connected cumulants $\kappa_{2n}=0$ (the signed set-partition/Pfaffian inversion of $\langle \psi_1 \cdots \psi_{2n} \rangle = \text{Pf}(\Gamma)$ leaves only the degree-2 kernel — the v113 “one kernel = the net” property at the cumulant level); (ii) a kernel-preserving transport fixes the whole quasi-free state by Wick functoriality; (iii) a gapped primitive Perron–Frobenius transport has a *unique* fixed ray, so a non-Gaussian piece is impossible — if κ_{2n} were fixed it would be a *second* fixed ray (killing uniqueness), and if not fixed it sits in the gapped complement and decays to 0 (any gap > 0 suffices; the value $(2/3)^6$ only sets the rate). *The honest residual:* v56’s gapped PF uniqueness is established on the *three-dimensional* scalar transfer spectrum $\{1, (2/3)^6, (1/3)^6\}$ (the flavour gap / horizon Page-recovery rates), whereas the 16-Majorana seam state already carries $\binom{16}{4}=1820$ and $\binom{16}{6}=8008$ higher-cumulant directions — the Schwinger cone the transport must contract has dimension $\gg 3$, and uniqueness must hold there (not merely on the quasi-free sub-cone) for the argument to be non-circular. **So (A) is equivalent to one strictly smaller, internal, exact-testable statement:** the admissible TFPT seam transport is a *primitive, spectrally gapped, RP-cone-preserving* map on the *full* Schwinger cone, not only on the scalar readout spectrum of v56 — which the suite does *not* yet establish [E]/[O].

And that full-cone premise is now *discharged* to a single finite one-particle statement (`[verification/v161_one_particle_reduction.py]`). The lever is Bogoliubov second quantisation: a quasi-free transport is $T_\Sigma = \Gamma(t)$, the second quantisation of a one-particle contraction t on the 16-dimensional Majorana space H_Σ , and the Fock space grades by correlator degree, so $\Gamma(t)$ acts on the degree- m sector as the exterior power $\Lambda^m(t)$. Three exact consequences follow (all machine-checked): (i) $\text{spec}(\Gamma(t)|_{\deg m})$ is the set of products of m one-particle eigenvalues — the whole correlation spectrum is generated by the finite one-particle spectrum; (ii) **the cone gap equals the one-particle gap:** the sub-leading eigenvalue of $\Gamma(t)$ over *every* degree is $(2/3)^6$ (pad the largest contracted mode with fixed factors), so premise 5 on the cumulant space *is* premise 5 on the 16-dim space, the same number; (iii) the eigenvalue-1 sector is Λ^* of the fixed subspace — the disconnected Wick powers of K_Σ — so with a strict one-particle gap no independent connected cumulant is fixed, and an independent fixed cumulant *on* the fixed subspace is excluded by v113 (one polarisation $\Leftrightarrow$ one vacuum). Premise 1 (quasi-freeness) is itself *forced*, not assumed: the $120 = \dim \mathfrak{so}(16)$ weight-1 chiral currents *are* the Bogoliubov generators on the 16 Majoranas, and v158 makes every non-Gaussian deformation irrelevant. **So the three full-cone premises collapse to one finite, 16-dimensional statement:** the seam RG transport’s one-particle symbol is the gapped Calderón Bogoliubov map — fixed eigenspace = the rank-8 K_Σ polarisation (v113), sub-leading gap = $(2/3)^6$ (v56). Every *number* in it is already supplied by the suite; the one irreducible step is the physical identification that the actual seam RG generator equals this Bogoliubov map — the same operator-algebraic (Kawahigashi–Longo-class) statement as the net-existence residual of (A_2) , now with the infinite-dimensional cone eliminated [E]/[O].

And that finite statement carries no new open content: it factors into the two already-open items ([[verification/v162_seam_transport_identification.py](#)]). Take the one-particle symbol apart into its two data, the gap value (β) and the fixed space (α). (β) is forced: a μ_4 -equivariant operator is diagonal in the cusp basis — the deck average is the diagonal projection, $\sum_{k=0}^3 U^k X U^{-k} = |\mu_4| \text{diag}(X)$ for $U = \text{diag}(1, i, -i)$, and the commutant of U is exactly the diagonal algebra — so its spectrum is the cusp weights $\text{spec}(A_0^*) = \{0, \frac{1}{3}, \frac{2}{3}\}$ (the parabolic residue, [v115](#)), and the transfer eigenvalue $(1-\alpha)^{p_2}$ with $p_2=6=|R_+(A_3)|$ gives $\{1, (2/3)^6, (1/3)^6\}$, gap $6 \log \frac{3}{2}$ — every entry carrier data, no free knob; the lift to the 16 Majoranas is the established A_3 subnet grading ([v137](#)). (α) is the rank-8 Calderón polarisation of the $\omega_k=|k|$ free dispersion ([v113/v156](#)), the IR conformal free-fermion vacuum. **What is left** is only that the seam RG generator is μ_4 -equivariant and admissible — which is the R1 “seam clock” identification (the transfer operator as the seam’s gapped generator) — together with the A_2 net-existence theorem. Both are pre-existing open items, so premise (A) introduces *no independent open content*: it factors exactly into $A_2 + R_1$ with every number forced. The “free-bulk cloud” is gone; what remains is the same two identifications the programme already names — (A) is closed *conditionally* on them (a unification, honestly not an unconditional proof) [\[E\]/\[O\]](#).

And both residual identifications are pinned to their floor, so (A) has no independent open gate left ([[verification/v165_premise_a_closure.py](#)]). A_2 (net existence) is not open research but *assembly* of established mathematics: the 16 free Majoranas define a free-fermion conformal net (Buchholz–Mack–Todorov), the holomorphic $c=8$ lattice net is $(E_8)_1$ (Frenkel–Kac–Segal / Dong–Xu / Kawahigashi–Longo), the μ_4 extension is a Longo Q-system of index 4 ([v125](#)); the one TFPT-specific input — that the seam Calderón kernel is the free-fermion two-point function — is done because the DtN symbol is universally $|k|$ ([v156/v157](#)). So A_2 re-types [\[O\]](#) $\rightarrow$ [\[C\]](#) (closed by assembly). R_1 (the seam clock) reduces to **GATE.QGEO**: a μ_4 -equivariant flat gapped generator on the carrier is *unique* ($= A_0^*$, [v115/v116](#)), so its gap $(2/3)^6$ is forced once the seam generator lies in that class, and the residual is exactly one geometric identification — the seam collar boundary is the μ_4 -punctured sphere $\mathbb{P}^1 \setminus \mu_4$ — whose cohomological half is already established ([v137](#), $b_1=3=N_{\text{fam}}$). That is **GATE.QGEO**, an already-open realisation premise, *not* a new gate. [2026-08-03 **GATE.QGEO** note: quantitative progress on the continuum side via the quasi-free CAR route ([[verification/v715_qgeo_car_continuum.py](#)], 22/22, **QGEO-CAR-RATES-SUMMABLE**): all 6+1 deck/twist channels are Schatten-norm Cauchy with geometrically summable rates (N up to 3072/6144), the FH renormalization is fit-free lattice-derived (Barnes G , 8.8×10^{-7}), the RP cone is stable, and the majorant lemma (L1)–(L3) is named with constants and a classical proof route — named, not proven: the gate itself does not move.] [2026-08-04 **GATE.QGEO** note: the first proof block of the majorant lemma lands ([[verification/v745_qgeo_car_l1_sector_lemma.py](#)], 11/11, **L1-SECTOR-PROVEN-MODULO-ELEMENTARY**): (L1) is proven modulo elementary steps for all 6 undressed sectors on the fixed grid — the occupied-arc mode sums are exact geometric sums (exact summation replaces Euler–Maclaurin), the entire $1/N$ structure is the sublattice embedding offset, the leading term is the exact matrix $(iu/2MN)f'_q(s)$ (ratio $\rightarrow 1$ at 6×10^{-4}), $\gamma = 2$ is derived, the Taylor–Lagrange envelope holds with 39744+108 inequalities at 0 violations, and $r_1 = 1$ is sharp. The residues stay typed: (a) grid-sup [elementary], (b) L2 uniform FH/RH [hard-technical], (c) L3 uniformity [medium] — named, not claimed: the gate itself still does not move.] [2026-08-05 **GATE.QGEO** note: residue (a) is fully closed in the corrected $(1+\ln M)$ form ([[verification/v778_qgeo_gridsup_logsum.py](#)], 19/19 + 19/19, **GRIDSUP-SPLIT-CLOSED-LOG-LAW+FOURIER-LOGSUM-CLOSED**): the continuum entrywise supremum, the configuration- S_1 closure (explicit $B_{\text{conf}}(\varepsilon)$ via packing + $\zeta(2)/\zeta(3)$) and the refinement remainder (exact csc^3 integrals) close elementarily with zero census violations, while the leading part is honestly log-divergent under refinement — the literal [v715](#) clause “one $B(\varepsilon)$ for all grid refinements” is false at $r_1 = 1$ and is repaired to $\|D_N^{(M)}\|_1 \leq [B_0(q, \varepsilon) + (J(\varepsilon)/2) \ln(M+3)]/N$, certified at theorem level by the grid Fourier log-sum inequality $\|F_q^{(M)}(\varepsilon)\|_1 \leq c_0(q, \varepsilon) + (J(\varepsilon)/4) \ln(M+3)$, $J = 2|f'_q(\varepsilon)|$, with all constants exact (162-case census to $M = 6144$, zero violations; all 252 transfer cases verified; Araki/Powers–Størmer summability explicit). Honest: the measured slope is J/π^2 — the bound is valid, not sharp. Residues (b) and (c) stay open: the gate itself still does not move. 2026-08-06 architectural note ([[verification/v802_gnet_martingale.py](#)]): the martingale route bypasses (b)/(c) for the limit-existence step of the G_{net} continuum strand (summable state-norm defects suffice) and relocates them into the typed normality step — a reallocation, not a proof of (b)/(c).] By the No-Unit Theorem ([v153](#)) the irreducibles stay exactly $\{\pi, v_{\text{geo}}\}$ — closing (A) adds none. **Final accounting**: premise (A) = (A_2 assembly, [\[C\]](#)) + (R_1 =**GATE.QGEO**, already open) + the $\{\pi, v_{\text{geo}}\}$ irreducible core (a theorem) — it carries zero independent open gates and ceases to be its own gate [\[E\]/\[O\]](#).

And the two structural residuals — net existence and full-cone reflection positivity —

are now *discharged to a theorem*, leaving the seam realisation as the one irreducible open premise ([`verification/v175_net_existence_full_cone.py`]). The CAR second-quantisation functor $\Gamma(t) = \bigoplus_m \Lambda^m(t)$ lifts the one-particle seam contraction t to the *complete* Fock space $\Lambda^\bullet(\mathbb{R}^{16})$ ($\dim 2^{16}=65536$); by the exterior-power spectrum theorem every eigenvalue of $\Gamma(t)$ is a subset product of $\text{spec}(t)$, so checking all 2^{16} products shows $\Gamma(t)$ is a *contraction* (full-cone RP), with eigenvalue-1 multiplicity $2^8=256$ (the wedges of the rank-8 K_Σ polarisation) and sub-leading eigenvalue exactly the one-particle gap $(2/3)^6$ — so full-cone RP/gap reduce *for every* m (not only $m \leq 6$, `v161`) to the one-particle contraction (Shale–Stinespring), *discharging the v160 scope premise by a theorem*. And A_2 is now an *assembled and verified* certificate: the $(E_8)_1$ character $E_4/\eta^8=1+248q+4124q^2+34752q^3+213126q^4+1057504q^5$ (level-1=248), the embedding $248=120+128$, and the E_8 Cartan matrix even with $\det 1$ (the unique rank-8 even unimodular lattice, `v83`) — the net *exists* by cited theorems. **Both residuals then collapse to the same input**: that the actual seam Calderón kernel is this gapped one-particle contraction on $\mathbb{P}^1 \setminus \mu_4$, i.e. QGEO.REALIZE.01, whose finite half is proven (`v168`). That realisation half is a constructive-geometry/AQFT statement, *not* closeable by a finite computation, and is left honestly [O] — the single irreducible structural premise; net existence and full-cone RP are [E].

The No-Unit Theorem — v_{geo} is an irreducible metrology primitive [E]/[O] ([`verification/v153_no_unit_theorem.py`])

v_{geo} is not an open gap. Every load-bearing compiler datum — $g_{\text{car}}, |\mu_4|, N_{\text{fam}}, \text{rank } E_8, \alpha^{-1}$, the determinant ladder $(3, 4, 8, 20)$, $c_3 = \frac{1}{8\pi}$, φ_0^{ret} — is mass-dimension 0 and *invariant* under a unit rescaling $L \mapsto \lambda L$, whereas a mass / energy / $1/G$ carries nonzero dimension. A map from dimension-0 inputs to a dimension- $d \neq 0$ output is simultaneously invariant and covariant: impossible unless the unit is introduced. **So a dimensionless boundary compiler provably cannot select an absolute scale** — v_{geo} is a *forced input*. The three apparent residuals then collapse onto the one unit, differing only in mass-dimension:

$$U_{\text{point}} \sim v_{\text{geo}}, \quad \frac{1}{G} \sim v_{\text{geo}}^2, \quad m/\mu = e^{3/4} \text{ (a pure ratio),}$$

so the irreducibles stay exactly {one dimensionful unit, π }. *Re-typing*: v_{geo} is an *irreducible metrology primitive* (a declared unit, like choosing a metre), not a missing derivation.

Sharpened across gravity, and the final tally ([`verification/v364_vgeo_sharpen.py`]). After the parameter-free gravity (`v358/v359/v361`) the *gravity* sector adds no new scale either: the Einstein coefficient $1/c_3 = 8\pi$ and the Λ prefactor $(8\pi)^2 \cdot 48c_3^4 = 3/(4\pi^2)$ are dimensionless, so $1/G \sim v_{\text{geo}}^2$ and $\Lambda \sim (\text{dimensionless}) v_{\text{geo}}^4$ both reduce to v_{geo} times atoms. Hence v_{geo} is the *single* dimensionful input of the *complete* theory (matter and gravity), and the final tally is $\{v_{\text{geo}}, \pi\}$ with *zero* dimensionless dials — a $\sim 26 \rightarrow 1$ reduction against the Standard Model. ($1/G$ stays UV-sensitive (Sakharov, `v68`), so it is a metrology readout, not a clean prediction — consistent with v_{geo} being the anchor.)

The torsor closure of the interface ([`verification/v725_phys_vgeo_torsor_audit.py`], V GEO . TORSOR . 01, 21/21). The v_{geo} interface is now *structurally closed* as an $\mathbb{R}_+$ scale torsor, machine-audited on the corpus: (1) the complete export table $O_i = r_i v_{\text{geo}}^{d_i}$ with machine-derived dimensionless r_i ($\bar{M}_{\text{Pl}}, 1/G, \Lambda, U_{\text{point}}, m/\mu$ at $d = 1, 2, 4, 1, 0$); (2) the dimension matrix has *rank* 1 — exactly one independent scale (conditional [A] on the flavor block via the declared seam=Planck identification, named, not hidden); (3) *all* consistency conditions are homogeneous under $L \mapsto \lambda L$ — the machine form of this theorem; (4) the leave-one-anchor-out cross-calibrations are compatible (G vs ρ_Λ : the same $\bar{M}_{\text{Pl}}$ to 0.11%, `v274`); (5) the *calibration theorem* in torsor language: one external reference selects the point in the torsor, everything else is prediction. The closure is the *calibration form* — no absolute-scale derivation (impossible per this theorem; that is the content, not the defect); the one remaining dimensionless scale ratio is *named*, $H_{\text{EW}} = \ln(\bar{M}_{\text{Pl}}/v_{\text{EW}}) = 37.1776$ (naive $c_3/\varphi_0^{\text{ret}}$ ladders honestly excluded); the [E]/[O] typing of this keybox is unchanged.

Theorem G — the closure statement [O] (target)

The relative spectral boundary kernel over the diffeomorphism-quotiented metric sector admits a regulator-independent, reflection-positive projective-limit measure μ_{QG} . Its Ward identities imply covariant conservation, and its low-curvature expansion is the TFPT $R + R^2$ seam action

$$f_R R_{\mu\nu} - \frac{1}{2} f(R) g_{\mu\nu} + (g_{\mu\nu} \square - \nabla_\mu \nabla_\nu) f_R = \bar{M}_{\text{Pl}}^{-2} T_{\mu\nu} + O(R^3/M^4),$$

so the closed covariant equation becomes the controlled IR shadow of the full measure. *Status*: the local covariant field equation is parameter-free (`v358/v359`); Theorem G remains open only for the

projective-limit ambient measure μ_{QG} ($G6$, the deepest item). $G5$ certified; the regime equation closed [verification/v28_gravity_fR.py].

Research Contract — CELEST.SEAM.01 (the celestial and twistor continuum route)

Reader’s note. A compact synthesis of this contract’s executed work packages (WP1–WP5d, both WP5d stages, + the WP5e α , β , γ , δ_2 , ε_2 and ε_1 stages + the three back-reaction milestones M1–M3) — the narrative arc from the μ_4 clock to the $(E_8)_1$ boundary shadow — is given as a dedicated section in `tfpt_3_e8_audit_bootstrap` (“The celestial and twistor continuum route”); this contract remains the full technical reference (typing fence, kill tests, work-package statements). The bridge from this contract’s compiler-side constructions to reconstructed Minkowski physics is *not* part of this contract: it is the separate central contract WOIT.OS.TWISTOR.01 below — everything in CELEST.SEAM.01 is preparation for it.

The object diagram (read this first). The route is one chain of objects; each arrow is a separately typed claim. Where mathematics ends and physics begins is visible at a glance: the last two arrows are [O].

Arrow	Status	Claim / modules	Kill test
$(\Sigma, \mu_4, \rho, \Theta) \rightarrow \mathbb{P}^1 \setminus \mu_4$	[E]	SEAM.MARKS chain: four marks, clock, cross-ratio 2, $\tau=i$ pillowcase (v168/v214/v216/v180; bit reduction v506/v507/v510/v512)	a fifth mark, a non-order-4 clock, or a marks-preserving root without flag transitivity
$\mathbb{P}^1 \setminus \mu_4 \rightarrow \mathbb{C}^2 / \mathbb{Z}_4$	[E]	CELEST.WP1.01/CELEST.WP2.01: the glue is the flat $\mathbb{Z}_4$ monodromy on the A_3 ALE, clock = Kähler $U(2)$ phase, clock ² = deck (v492/v493)	equivariant-sector closure failure (3112/6720 control) or a surviving shape modulus
$\mathbb{C}^2 / \mathbb{Z}_4 \rightarrow \mathbb{PT} / \Gamma$	[E]/[C]	CELEST.WP5E.*: equivariant twistor uplift — anomaly ledger, level-from-flux, axion slot, back-reacted Ω_N , twisted KS measure, a_0 uplift, the δ_1 decision, the measure decision (v505/v509/v511/v513/v514/v515/v516/v517/v518/v520)	the preregistered level kill (“inflow demands $k \neq 1$ ”); the M1–M3 kills were evaluated by v515/v516/v517 and did not fire; the δ_1 kill <i>fired on the derived measure</i> (v518), and the declared/derived measure question is since <i>decided at probe level</i> in favour of the declared reading under the typed premises TP-1..TP-4 (v520) — the residual [O] is the constructive w_m derivation
$\mathbb{PT} / \Gamma \rightarrow \mathcal{A}_{\text{hol}}$	[O]	the <i>interacting</i> holomorphic algebra (global BCOV+SDYM quantisation; WP5e proper; target of WOIT.OS.TWISTOR.01; CFT-side anchors v497–v504)	anomaly mismatch at the Costello–Li grade
$\mathcal{A}_{\text{hol}} \xrightarrow{\text{OS}} \mathcal{A}_{\text{Mink}}$	[O]	OS reconstruction with the real structure Θ — the WOIT.OS.TWISTOR.01 target (the free-collar RP/OS witnesses v379/FORM.SEAM.MMST.01 are the <i>free</i> anchor only; the α stage v519 pins Θ itself and free RP on the seam system — kill test (1) discharged at the free level, live on $\mathcal{A}_{\text{hol}}$)	the seven kill tests of WOIT.OS.TWISTOR.01 below

Definition (the group extension, exactly). The shorthand “ $\mathbb{PT} / \mathbb{Z}_4$ ” used in this contract is precise but deserves unpacking, because *two different* order-4 structures are in play and only one of

them is quotiented:

- $\Gamma_{\text{ALE}} = \langle \text{Deck} \rangle \cong \mathbb{Z}_4 \subset SU(2)$, the deck group $(z_1, z_2) \mapsto (iz_1, i^{-1}z_2)$: *triholomorphic*, preserves the holomorphic symplectic form Ω ; this is the $\mathbb{Z}_4$ that is *quotiented* ($\mathbb{C}^2/\Gamma_{\text{ALE}}$ is the A_3 singularity $XY = Z^4$, and $\mathbb{P}\mathbb{T}/\Gamma$ means the quotient by Γ_{ALE} acting on the twistor space of the ALE) [E] ([verification/v492_celestial_z4_orbifold.py]).
- $\rho = \text{Clock} = \text{diag}(i, 1) \in U(2)$: a Kähler, *not* triholomorphic isometry ($\det \rho = i$ rotates Ω by the μ_4 generator — the WP1 S5 computation); it is *not* in Γ_{ALE} and is *not* quotiented — it *normalises* Γ_{ALE} and survives the quotient as the residual clock symmetry [E] (v492).
- The global object is the extension generated by both: since $\rho^2 = \text{Deck}$ exactly (v492/v506), $\langle \Gamma_{\text{ALE}}, \rho \rangle = \langle \rho \rangle$ is *cyclic* — projectively $\mathbb{Z}_4$ (the clock on the sphere) with ρ^2 generating the deck $\mathbb{Z}_2$ there, and on the spin/fermionic level the canonical $\mathbb{Z}_8$ tower ($V^2 \propto U$, $V^4 \propto (-1)^F$, nonsplit; $8 = 2|\mu_4|$) [E] (v506/v507).

The action table, entry by entry ([E] = arithmetically established at the current stratum, [O] = a quantisation question):

Object	Γ_{ALE} (Deck)	ρ (Clock)	Status / witness
$K_{\mathbb{P}\mathbb{T}}$ (canonical class)	trivial	weight via $\det \rho = i$	[E] (incidence-forced column weights $(+1, +1, -1, -1)$, v514)
Ω (hol. symplectic)	preserved	rotated by i	[E] (v492 S5; back-reacted Ω_N closed-form with $(2\pi i)^2$ -integral periods and forced charge 4, v515)
BCOV fields ($O(-2)$ tower)	orbifold sectors	character series $P_0..P_3$	[E] ledger / [O] quantisation (v514)
open-string fields (SDYM E_8)	glue-equivariant sector	sector rotation $j \mapsto j+1$	[E] ($D[d] = 60(d+1)/64(d+1)$, v492) / [O] interacting
boundary states ($(E_8)_1$ shadow)	μ_4 sector split	clock phase on $ s\rangle$ etc.	[E] character/GNS level (v497–v500) / [O] as an actual net

The A_3 role table (no silent identifications). Several rank-3 A_3 ’s and external gauge factors circulate near this contract; the following table states, for each pair, either the connecting functor (with its witness) or an explicit *not identical*:

Object	Role and relations
A_3^{family}	the family factor in the lattice $D_5 \oplus A_3$ ($\mathfrak{su}(4)$ -flavour; three families from its exponents, v128/v137). Related to A_3^{ALE} by the McKay correspondence of <i>types</i> only (both are type A_3); not identical as realisations — one is a sublattice of E_8 , the other the singularity type of $\mathbb{C}^2/\mathbb{Z}_4$. The bridge that <i>does</i> connect them is the Kronheimer–Nakajima quiver (v479: same ADE datum, different functors).
A_3^{ALE}	the ADE type of $\mathbb{C}^2/\mathbb{Z}_4$ ($XY = Z^4$); its resolution carries the three exceptional spheres (v492/v493).
three exceptional spheres	the Coxeter/Picard–Lefschetz structure of the resolved A_3^{ALE} : the clock acts as a Coxeter element of $W(A_3)$ on H_2 (eigenvalues $\{i, -1, -i\}$, v493); they carry the three twisted axion slots (v505) and the lockstep fluxes (v509).
D_5^{carrier}	the carrier factor ($g_{\text{car}} = 5$); glued to A_3^{family} by the μ_4 Lagrangian glue into E_8 (v92/v125); no role on the ALE side beyond the zero-mode algebra $\mathfrak{g}_0 = D_5 \oplus A_3$ (v492).
Woit’s $SU(3)_{\text{color}}$	<i>external programme reference only</i> (Euclidean Twistor Unification: colour as the rank-three quotient bundle on $\mathbb{P}\mathbb{T}$). Not identical — no identification claimed ; in particular $A_3^{\text{family}} \not\cong SU(3)_{\text{color}}$ (an $\mathfrak{su}(4)$ flavour factor is not a colour gauge bundle), and TFPT’s colour lives in the SM readout of the carrier, not on $\mathbb{P}\mathbb{T}$.
Woit’s $SU(2)_{\text{weak}}$	<i>external programme reference only</i> (the internal spin factor of the Euclidean twistor construction). Not identical — no identification claimed ; whether an internal $SU(2)$ of that kind can be realised on $\mathbb{P}\mathbb{T}/\Gamma$ is exactly kill test (6) of WOIT.OS.TWISTOR.01.

Hypothesis (as corrected by WP1). *The seam is the glue-equivariant $\mathbb{Z}_4$ -orbifold sector of the celestial chiral algebra on the A_3 ALE space $\mathbb{C}^2/\mathbb{Z}_4$; the μ_4 clock is the Kähler $U(2)$ phase $\text{diag}(i, 1)$ whose square is the deck group (spin bridge $\mathbb{Z}_8$, $8 = 2|\mu_4|$ — the $c_3 = 1/(8\pi)$ seam winding integer); and the associated twistorial bulk is the self-dual sector of TFPT gravity.* This is the fourth research contract, alongside (U_{wall}) , (G_{metric}) and **CONTRACT.F.01**: a numbered chain of work packages with pre-registered kill tests, *not* a claim. The method is the celestial-chiral-algebra (CCA) construction of Bittleston–Homans–Sharma on Eguchi–Hanson space ([arXiv:2305.09451](#), JHEP 09 (2023) 008 — the $\mathbb{Z}_2$ case, refined here to $\mathbb{Z}_4$ with a flat connection at infinity in the Kronheimer–Nakajima / heterotic-on-ALE sense), inside the celestial-holography programme (Strominger, PRL **127**, 221601: $w_{1+\infty}$; Costello–Paquette–Sharma, PRL **130**, 061602: Burns holography, the $SO(8)$ WZW₄ model; Bittleston–Sharma–Skinner, CMP **403** (2023): quantizing the non-linear graviton). The gauge-algebra admissibility gate is already passed in the literature: Costello’s one-loop axionic-consistency analysis ([arXiv:2111.08879](#)) admits the exceptional algebras *including* E_8 for SDYM on twistor space (Okubo: no independent quartic Casimir). The spin bridge in the hypothesis is no longer a bookkeeping convention: on the 16-Majorana seam the NS deck implementation is *forced* to order 4 ($U^2 = (-1)^F$ exactly, nonsplit $\mathbb{Z}_4$) and the quarter-shift lifts to a canonical $\mathbb{Z}_8$ tower with $V^2 \propto U$ — “ $8 = 2|\mu_4|$ ” fermionically explained ([\[verification/v506_seam_clock_rigidity.py\]](#), **SEAM.CLOCK.RIGIDITY.01**); and the nonsplit class is arrangement-*sensitive*: the edge (silver) arrangement splits ($U_e^2 = +1$, zero roots), so the fermions *measure* the alignment bit as a Fidkowski–Kitaev-type extension class ([\[verification/v507_seam_bit_origin.py\]](#), **SEAM.BIT.ORIGIN.01**) — and that class is nonsplit *iff* the deck acts freely (all 17 seam-circle involutions, **v510**), which the covering deck does by topology.

WP1, executed and verified: the E_8 μ_4 -glue is the flat $\mathbb{Z}_4$ monodromy of the equivariant CCA on the A_3 ALE space — verdict B ([\[verification/v492_celestial_z4_orbifold.py\]](#), **CELEST.WP1.01**)

Everything in this box is exact arithmetic (sympy, no floats). **Inner grading / flat monodromy [E]:** the **v128** glue grading ($240 = 52+64+60+64$, additive on all 6720 root pairs, dims with Cartan $(60, 64, 60, 64)$, $\mathfrak{g}_0 = D_5 \oplus A_3$ carrier = 60) is *inner*: $h = (2, 2, 2, 2; 0^4)$ reads the glue class mod 4 on all 240 roots, so the glue is $\text{Ad}(\exp(2\pi i h/4))$ — an order-4 torus element, i.e. a *flat $\mathbb{Z}_4$ monodromy* in the Kronheimer–Nakajima sense; the A_3 -side detector $h' = (0^5; 1, 1, 1, -3)$ reads the *same* class, so the glue charge is the diagonal $(1, 1)$ of the discriminant group $\mathbb{Z}_4 \times \mathbb{Z}_4$ — the **v92/v125** Lagrangian glue. **The spatial side [E]:** $\mathbb{C}^2/\mathbb{Z}_4$ under the deck $(z_1, z_2) \mapsto (iz_1, i^{-1}z_2)$ is the A_3 singularity $XY = Z^4$ (invariant ring exact; Hilbert series $(1+t^4)/((1-t^2)(1-t^4))$; Molien table verified). **The equivariant sector closes [E]:** the glue-equivariant SDYM(E_8) modes ($j + m \equiv 0 \pmod{4}$) form a Lie subalgebra with graded dimensions $D[d] = 60(d+1)$ (even d) / $64(d+1)$ (odd d) — uniform per parity *only* because $\dim \mathfrak{g}_0 = \dim \mathfrak{g}_2 = 60$ and $\dim \mathfrak{g}_1 = \dim \mathfrak{g}_3 = 64$: the **v128** dimensions are exactly the consistency condition. The zero modes ($d=0$) are $\mathfrak{g}_0$ alone — *the unbroken algebra on the orbifold is the carrier $D_5 \oplus A_3 + \text{Cartan} = 60$* ; the asymptotic density is $62/248 = 1/4 = 1/|\mathbb{Z}_4|$ (the orbifold index), and the $\mathbb{Z}_2$ halfway sector reproduces the Eguchi–Hanson towers $120/128 = SO(16)$ adjoint/spinor — μ_4 refines the sheet $\mathbb{Z}_2$ exactly as $4 = 2 \times 2$ (**v125**). **Character level [E]:** Θ_{E_8} splits into the four glue-coset thetas (shell 1 = $(52, 64, 60, 64)$ re-derived from the *lattice*); the four sector characters Θ_{C_j}/η^8 sum *exactly* to $\chi_{(E_8)_1} = 1+248q+4124q^2+34752q^3$ (the **v377** series); the sector conformal weights are $h_{D_5} = (0, \frac{5}{8}, \frac{1}{2}, \frac{5}{8})$, $h_{A_3} = (0, \frac{3}{8}, \frac{1}{2}, \frac{3}{8})$ — the **v92/v125** discriminant form $(5x^2+3y^2)/8 \pmod{1}$ — and the glue diagonal has *integer* $h = (0, 1, 1, 1)$: the **v125** isotropy is the locality of the extension to $(E_8)_1$. **The critical correction [E] (why verdict B, not A):** the A_3 deck $\text{diag}(i, i^{-1})$ induces on the celestial sphere $z = z_1/z_2$ only $z \mapsto -z$ (projective order 2, the sheet flip) — it is *not* the order-4 clock $z \mapsto iz$. The exact bridge: any $SU(2)$ lift of the clock has order 8 (spin group $\mathbb{Z}_8$, which would quotient to A_7), and (spin clock)² = the deck generator *exactly*, with $|\mathbb{Z}_8| = 8 = 2|\mu_4|$ — the $c_3 = 1/(8\pi)$ winding integer. The order-4 clock *is* realised, as the $U(2)$ phase $\text{diag}(i, 1)$: projective order 4, fixed points $\{0, \infty\}$, free μ_4 orbit (all **v180** clock properties), normalising the deck (so preserving the glue flat connection), with $\det = i$ rotating the holomorphic symplectic form by the μ_4 generator (a Kähler, *not* triholomorphic isometry — it rotates the twistor sphere by $\pi/2$). **Gravity side [E]:** clock-invariance selects from the 3-parameter versal A_3 family $XY = Z^4 + a_2 Z^2 + a_1 Z + a_0$ *exactly* the 1-parameter family $XY = Z^4 + a_0$, whose four branch points are *one* μ_4 orbit with cross-ratio 2 — the seam marks (**v179** clock orbit; **v168/v214**

cross-ratio $2 \Rightarrow j=1728$ pillowcase). **Negative controls [E]:** the false spatial action $\text{diag}(i, i)$ is killed three ways ($\det = -1 \notin SL(2, \mathbb{C})$, no CY/ALE resolution; Poisson closure fails in 14/14 nonzero cases; conjugation symmetry broken), the false glue (60/64 blocks swapped) breaks additivity on 3112/6720 root pairs, and of all 24 relabelings exactly $\{\text{id}, j \mapsto -j\} = \text{Aut}(\mathbb{Z}_4)$ are additive — the grading is rigid. **Verdict [C]:** the celestial and glue $\mathbb{Z}_4$ structures match under *one* precisely named extra identification — $\text{clock}^2 = \text{deck}$ on the sphere (the sheet- $\mathbb{Z}_2$ spin bookkeeping) — so WP1 lands at verdict B (match modulo one named identification), close to but honestly short of A. [E]/[C]

Typing and non-circularity. Admissible *inputs* are the finite seam/glue data already in the suite — the μ_4 Q-system glue (v92/v125), the graded hull (v128), the seam rigidity and pillowcase geometry (v168/v214), the Möbius clock (v180) — plus the WP1 [E] arithmetic ([verification/v492_celestial_z4_orbifold.py]). *Not* admissible as an input: the *continuum existence* of the holomorphic $(E_8)_1$ net on the seam — that is precisely the SEAM.EQUIV.01 target, and assuming it would make the contract circular. *Scope fence:* the twistorial bulk of this contract is the *self-dual* sector only, never full Einstein gravity — the classical field-equation route (entanglement equilibrium, v358/v359) and the entropic-action bridge (GRAV.ENTROPIC.ACTION.01, v473) are separate, non-competing statements about the full equation. SEAM.EQUIV.01 stays [O] throughout; nothing in this contract moves it.

Work packages.

- **WP1 (done, verdict B) [E]/[C]:** the structural $\mathbb{Z}_4$ arithmetic of the keybox above, executed and machine-verified as CELEST.WP1.01 [verification/v492_celestial_z4_orbifold.py].
- **WP2 (done, verdict B) [E]/[C]:** the clock-invariant deformation theory, executed and machine-verified as CELEST.WP2.01 [verification/v493_celestial_wp2_clock_deformation.py] (sympy exact, 47 checks). Both targets landed, *sharpened*: (i) a_0 is the pure *scale* of the seam configuration — the binary-quartic invariants of $w^2 = Z^4 + a_0$ are $I = 12a_0$, $J = 0$ *identically*, so $j = 1728$ and the $\tau=i$ pillowcase shape (v168/v214) is *frozen* for every $a_0 \neq 0$ (no shape modulus survives clock-invariance; the negative controls $a_1 Z \Rightarrow j=0$ and $a_2 \neq 0 \Rightarrow j=1556068/81$ show the test has teeth); (ii) the three resolved spheres ($3 = \text{rank } A_3 = N_{\text{fam}}$) carry *exactly* the three nontrivial μ_4 characters $\{i, -1, -i\}$ under the clock, which acts on H_2 as a *Coxeter element* of $W(A_3)$ (char $x^3 + x^2 + x + 1$, order $h(A_3)=4=|\mu_4|=N_{\text{fam}}+1$) and *is* the Picard–Lefschetz monodromy of the family ($a_0 \rightarrow e^{2\pi i} a_0$ gives the 4-cycle); the surviving deformation direction is the weight-1 χ_1 -Fourier diagonal $(1, i, i^2)$ — *not* the naive invariant diagonal ($\det(A-1) = -4$: no invariant cycle) — and all three sphere volumes move in lockstep ($|\Pi_j| = \sqrt{2}t$). The BHS deformed-algebra pattern transfers from $\mathbb{Z}_2$ to $\mathbb{Z}_4$: the fibre bracket $-4\text{Nambu}(XY - Z^4 - a_0)$, anchored at the $a_0=0$ orbifold, closes with corrections exactly *linear* in a_0 at the $\mathbb{Z}_4$ wrap, conserving the μ_4 grade (no sector leak — the WP1 equivariant sector deforms consistently), with a_0 in the weight-0 slot of BHS’s c^2 . Verdict B via three named [C] identifications (-4Nambu as *the* $k=4$ CCA bracket; period = root difference; the seam-scale reading via $\text{clock}^2 = \text{deck}$); the full deformed twisted-sector OPEs and the $W(\mu)$ -type scaling limit for $k=4$ stay [O]. The v216 [O]-residual is *typed*, not moved: given the order-4 clock, the deformation freezes the $(2, 2, 2, 2)$ modulus to the square automatically — a relocation of the same order-4 carrier input, not a new derivation (since v510 that input reads: the square-modulus datum $\tau = i$; its position half is topology; since v512 it reads equivalently: one discrete symmetry-lift bit, flag transitivity $\iff \tau = i$; and since v521/v525/v529 free OS positivity, the whole mark-decorated free-plus-twist state class and the first genuinely interacting FK toy are documented as the *eighth*, *ninth* and *tenth* side-blind tests — the bit is not derivable from free RP/Θ existence, from any Wick-computable twist decoration or from the FK interaction class, and remains genuine discrete input; since v528 it is the twist-class choice, in principle interferometrically readable; and since v534 it carries the first *positive* selection datum — reflection positivity on the interacting mark family keeps exactly the $\delta = \pi/2$ member alive, toy-fenced).

- WP3 (done, verdict B) [E]/[C]:** the Green–Schwarz alignment test, executed and machine-verified as CELEST.WP3.01 [verification/v495_celestial_wp3_gs_alignment.py] (exact Fraction/sympy, 25 checks). The arithmetic side is sharp: the Okubo coefficient $\text{Tr}_{\text{adj}} X^4 / (\text{Tr}_{\text{adj}} X^2)^2 = 5/(2(\dim \mathfrak{g} + 2))$ is *derived* as a polynomial identity for all 8 algebras on Costello’s list (with a sharp $\mathfrak{sl}_4$ negative control: $5/32$ vs $3/32$), the closed form $\tilde{\lambda}^2 = 10h^\vee/(\dim + 2) = h^\vee + 6$ holds exactly across the Deligne series, and for E_8 the axion coefficient is *exactly* $\tilde{\lambda} = 6$ (unit-trace) resp. $\lambda_{\text{fund}} = 1/10$ (adjoint-trace), so the GS coupling $\kappa = \lambda_{\mathfrak{g}}/(8\pi\sqrt{3})$ satisfies $(\kappa/c_3)^2 = 12 = |\mu_4|N_{\text{fam}}$ resp. $1/300$ — exact anchor rationals, with the anchor decomposition $h^\vee = 30 = |\mathbb{Z}_2|N_{\text{fam}}g_{\text{car}}$, $\varphi(30) = 8 = \text{rank}$, $\tilde{\lambda} = |\mathbb{Z}_2|N_{\text{fam}}$, $\lambda_{\text{fund}} = 1/(|\mathbb{Z}_2|g_{\text{car}})$ (a byproduct: the printed $\lambda_{\mathfrak{so}_8}^2 = 3/2$ in Costello’s appendix A is a factor-2 slip; his own weight content gives 3, Okubo-consistent). κ/c_3 itself is *irrational* ($2\sqrt{3}$): only the squared alignment is exact. **The look-elsewhere caveat is part of the result [C]:** the same squared-rational alignment holds for *all eight* algebras on the list ($8/8$ — the test as such has zero selective power), $\tilde{\lambda}$ -integrality passes $2/8$ (shared with $\mathfrak{sl}_3$), and the only $\mathfrak{e}_8$ -selective single test is $g_{\text{car}} = 5 \mid h^\vee$ ($1/8$); the conjunction that isolates $\mathfrak{e}_8$ is post hoc, not preregistered. *Alignment survives; selectivity does not* — the c_3 -connection of the celestial route is convention-level compatibility plus genuine λ -arithmetic, *not* $\mathfrak{e}_8$ -selective evidence, and never a derivation of c_3 from celestial data.
- WP4 (done, verdict B(ii)) [E]/[C]:** the type mismatch, executed head-on and machine-verified as CELEST.WP4.01 [verification/v496_celestial_wp4_character_shadow.py] (exact integer/Fraction/sympy, 25 checks). The $(E_8)_1$ character is *not* a conformal block of the celestial $E_8[\mathbb{C}^2]$ S-algebra in its own (jet) grading — the obstruction is localised *three* independent ways: (a) the CP grading gives spin $1 - d/2$, unbounded below, and 248 is never a jet-tower dimension ($60(d+1)/64(d+1)$, $d \leq 100$); (b) the cumulative generator count is quadratic ($31s^2 + 92s + 60$, $31 = k + h^\vee$), so the jet Fock grows as $n^{2/3}$ against the character’s $n^{1/2}$ (Meinardus poles $s=2$ vs $s=1$; the ratio f_n/χ_n increases strictly for $n = 1..12$); (c) the level-2 null ideal of $(E_8)_1$ deletes *exactly* $27000 = 30^3 = (h^\vee)^3$ out of $\text{Sym}^2(248) = 1 + 3875 + 27000$ (the character keeps $4124 = 1 + 248 + 3875$) — and has no jet analogue. **But the boundary/period reading holds exactly at the current stratum:** the zero-mode slice is the 60 vacuum-sector currents, the single-line jet slice cycles the four sector counts $(60, 64, 60, 64)$, one full μ_4 period of loop energies sums exactly to 248 (the single-particle level of the $248q$ stage), and the glue-diagonal weights $(0, 1, 1, 1)$ are integers — while the free loop Fock at level 1 counts $897266 \gg 248$, i.e. the rational truncation must be *imposed in a limit*. That is precisely the MMST scaling-limit shape of SEAM.EQUIV.01 (v336/v449: the rational net exists in the *limit* of the lattice collar, not in the pre-limit algebra): WP4 is hereby *retyped* from “character as conformal block” to “character as boundary/limit shadow”, and the constructive limit question (which limit, which topology, convergence) passes to WP5 — where WP5a has since answered the “which limit” part at the counting level (the coefficientwise χ_w limit below), WP5b has constructed the deleting operator, WP5c the limit state whose GNS kernel contains the ideal, WP5d the local-net legs (the α -stage two-interval index chain and the β -stage split + strong-additivity witnesses — all three KLM ingredients of complete rationality on the lattice), and WP5e- $\alpha/\beta/\gamma/\delta_2/\varepsilon_2/\varepsilon_1$ have anchored *both* faces of the twistor uplift, closed its exchange sub-branch, executed the full-tensor ledger, executed the level-from-flux dial and built the bulk-axion slot (the CFT-side prefactor + level pinning; the equivariant twistor skeleton with the index bridge and the rigid $32 T_3$ residual; the exchange no-go with certificate; the full-tensor confirmation of the collapse with the one cubic d -symbol channel that opens; the lockstep theorem making “one level” a theorem of clock invariance, with the sector-counter dial pinning $k = 1$; the equivariant $O(-2)$ slot construction with $\tilde{\lambda} = 6$ triply pinned and the first GH/ A_3 back-reaction step), and all three quantised back-reaction milestones M1–M3 have since been executed (the closed-form Ω_N with $(2\pi i)^2$ -integral lockstep periods and forced source charge $4 = |\mu_4|$, v515; the twisted KS measure on the declared completion reading — per-sector Okubo squares, the $32 T_3$ cancelled, the $\psi = 64$ slice supplied without

the cubic d -channel, **v516**; the a_0 uplift coupled to the centre count on four scales, **v517**), leaving the global BCOV quantisation (WP5e proper) as the remaining open piece — the BCOV *measure question* is since decided at probe level: the δ_1 chain itself was decided by **v518** (kill under the derived measure, all three testers fail), and **v520** has since decided *which* of the two exact measures the BCOV integral produces — single-valuedness of the one-loop lattice factor is *derived* from two independent constructive sources (F-independence of the moduli integral + the Quillen pairing) under the typed premises TP-1..TP-4, forcing the declared **v516** completion reading, and **v523** has since *derived* the w_m normalisation itself at probe level ($1/\det_j$ = the Atiyah–Bott/zeta-determinant fixed-point factor, computed from three independent sources under the typed premises TP-REG/TP-Q/TP-NUM/TP-CH); the residual **[O]** narrows to the global BCOV integral beyond the fibre zero-mode factor (throughout, $\mathbb{P}^T/\mathbb{Z}_4$ is the quotient by Γ_{ALE} , normalised by the clock $\langle \rho \rangle$ — the definition block above; the ε_1 slot itself is executed; the quantised milestones M1–M3 preregistered in **v514** are all executed via **v515/v516/v517** — the **v514** fence is fully worked off).

- **WP5 (the summit) — subdivided; WP5a–WP5d (both WP5d stages) plus the WP5e $\alpha, \beta, \gamma, \delta_1, \delta_2, \varepsilon_2$ and ε_1 stages and the back-reaction milestones M1–M3 are landed.** The twistorial construction as a *second continuum route* for SEAM.EQUIV.01 (Costello–Li grade), carrying the WP4 constructive-boundary-limit question. Subdivision WP5a–WP5e:

- **WP5a (done) [E]/[C]:** “the null ideal from the limit”, executed and machine-verified as CELEST.WP5A.01 [verification/v497_celestial_wp5a_null_ideal.py] (exact integer/Fraction arithmetic, 34 checks). Three results make the WP4 limit reading precise. (i) *Stabilisation theorem*: the one-parameter family χ_w of graded Fock characters on the chiral jet generators (loop energy m , radial number r ; $E_w = m + wr$ in quarter units $u = q^{1/4}$) *contains* the chiral jet grading as its $w=2$ member (generator counts 64, 120, 128, 180, 192, 240, 256, 300 per degree), and its u^n coefficient equals the quarter-moded loop Fock for *all* $w \geq n+1$ — strictly larger for every $w \leq n$ (verified $n \leq 8$, $w \leq 10$): the WP4 “boundary-limit shadow” is a precise *coefficientwise* limit $\varepsilon = 1/w \rightarrow 0$ with explicit stabilisation threshold $w = n+1$, not a slice. (ii) *The null ideal is derived, not cited*: Freudenthal multiplicities + the Weyl dimension formula + generic character peeling give $\text{Sym}^2(248) = V(2\theta) \oplus V(\omega) \oplus V(0) = 27000 + 3875 + 1$ with peeling residual exactly zero (independent Weyl-orbit cross-check); $27000 = 30^3 = (h^\vee)^3$ exactly; level-1 integrability ($\langle 2\theta, \theta^\vee \rangle = 4$, $\langle \omega, \theta^\vee \rangle = 2$) leaves the vacuum as the only level-1 primary ([C] Kac, used only through its counting consequence); level-2 quotient $31124 - 27000 = 4124 = 1 + 248 + 3875$ exactly. (iii) *Two routes, one number*: the μ_4 theta-split sector sum at $q^2 = (1036, 1024, 1040, 1024)$ gives the *same* 4124 — the untwisted null-ideal quotient and the boundary/twisted lattice presentation count one space (q^3 : 34752). Negative controls: $SO(16)_1$ through the same pipeline gives *four* components ($5304 + 1820 + 135 + 1$), $5304 \neq 14^3$ (the $h^\vee{}^3$ identity is not generic), quotient $2076 = \Theta_{D_8}/\eta^8$ at q^2 (the recipe validated on a second algebra), and block weights $(0, \frac{1}{2}, 1, 1)$ that *cannot* fuse into one local character; only $P = 4 = |\mu_4|$ periodisation reproduces the 248 layer; swapped glue breaks the zero-mode anchor. *Honest limit [O]*: the limit does *not* generate the truncation (the loop Fock counts $897266 \gg 248$ at integer level 1) — WP5a fixes the ideal’s *size and location* quantitatively and gives the celestial route the same two-step shape as MMST (limit + maximal ideal); the operator content is exactly WP5b–e, of which WP5b–WP5d and the WP5e $\alpha, \beta, \gamma, \delta_2$ and ε_2 stages are executed below.
- **WP5b (done) [E]/[C]:** “the singular vector as an operator”, executed and machine-verified as CELEST.WP5B.01 [verification/v498_celestial_wp5b_singular_vector.py] (exact integer/Fraction arithmetic, 53 checks, deterministic) — *success on the preregistered criterion*: the deleting object exists and is explicit. $|s\rangle = (E_{-1}^\theta)^2|0\rangle$ (weight 2θ , level 2 = 8

- quarter units = q^2 , an *integer* level) is constructed in a machine-built Chevalley/Frenkel–Kac basis (cocycle asymmetry on all 57600 pairs; the $[e_\alpha, e_{-\alpha}]$ sign *forced* by Jacobi, $\text{SGN} = -1$; κ derived, $\kappa(\theta^\vee, \theta^\vee) = 2$), and $J_1^a|s\rangle = 0$ is machine-verified for *all* 248 generators with the case classification 190 (first bracket) / 57 (second) / 1 ($a = F^\theta$, the central-term cancellation — the *only* case that sees the level k); plus $J_2^a|s\rangle = 0$, $E_0^a|s\rangle = 0$, exact weight and Shapovalov norm 0. *Level dial*: the F_1^θ coefficient is $2(1-k)$ — nonzero at $k=0$ and $k=2$, zero exactly at $k=1$: without the central extension the deletion operator does not exist. μ_4 compatibility: glue class $j(\theta) = 1$ (machine-built h -adapted chamber, $\langle\theta, h\rangle = 5$, height $29 = h^\vee - 1$), clock phase $i^{2j} = -1$, class(2θ) = 2 sheet-even (WP5a-consistent) with the deleting $\theta\text{-}\mathfrak{sl}_2$ crossing the sheet-odd classes (1, 3); 8 quarters = q^2 via the per-period dictionary. The module it generates is *the* ideal: weight 2θ has multiplicity 1 in the level-2 Fock, and the direct $\mathfrak{g}_0$ -lowering orbit BFS from $|s\rangle$ reproduces the Freudenthal multiplicities of $V(2\theta)$ exactly through depth 4 ($27000 = (h^\vee)^3$, quotient 4124; [C] Weyl complete reducibility beyond depth 4). Negative controls separate honestly: the level-1 current state and generic level-2 states are *not* singular; at $k=2$ the *cube* is (the $(E^\theta)^{k+1}$ pattern is generic Kac level mechanics); $SO(16)_1$ has the same singular vector but keeps three extra level-1 primaries ($h = \frac{1}{2}$ breaks one-block fusion) — the singular-vector mechanism is level-1 *generic*, the *one-block closure* is the E_8/μ_4 -specific part. *Handover to WP5c* (answered below): in the twisted quarter-slot moding of the χ_w limit Fock, two sector- C_1 modes never sum to 8 quarters (minimum $6 = q^{3/2}$) — the per-period dictionary (v496), not the per-slot identification, carries $|s\rangle$ to q^2 ; exactly the GNS/limit-state question (kernel $\supseteq$ ideal) that WP5c answers.
- **WP5c (done)** [E]/[C]: “the GNS limit state”, executed and machine-verified as CELEST.WP5C.01 [verification/v500_celestial_wp5c_gns_limit_state.py] (exact integer/Fraction arithmetic, 35 checks, deterministic) — *success on the preregistered criterion*: the family ω_w exists and its limit has the null ideal in its GNS kernel. ω_w is the quasi-free state with loop sector = the affine $k=1$ vacuum n -point functions (via the machine-determined compact anti-involution $\theta(e_\alpha) = -e_{-\alpha}$, the unique anti-automorphism sign on all 61504 basis pairs) and radial sector = oscillator pairings x^{wr} (the exact Gibbs regulator, x -adic); *the WP5a threshold holds at the state level*: $\omega_w = \omega_\infty \bmod x^{N+1}$ for all $w \geq N+1$, sharp at $w = N$. The core is fully machine: the *complete* exact level-2 Gram — 31124 monomials in 9361 weight blocks — has rank 4124 exactly (the preregistered target), kernel $27000 = V(2\theta)$ *weight by weight* (Freudenthal cross-check on all 9361 weights), every block PSD, rank table per Weyl orbit $(0, 0, 1, 8, 44)$, level-1 rank 248 positive definite (the current layer survives). μ_4 : the clock descends to GNS with level-2 rank split $(1036, 1024, 1040, 1024) = \Theta_{C_j}/\eta^8$ at q^2 — the WP5a two-routes identity at the *state* level — and $|s\rangle$ is the zero vector of $\text{GNS}(\omega_\infty)$, resolving the WP5b twisted-slot tension. The KILL branch is probed and not triggered: a CCR obstruction shows *no* w -uniform state can damp the radial modes (the family formulation is *necessary*), and the wrong constructions are caught (family $E'_w = mw+r$ erases the 248 layer, $710955 \neq 248$; no damping keeps $897266 \neq 248$); k -dial and D_8 controls separate honestly ($k=2$: $\langle s|s\rangle = +4$, no ideal in the kernel; D_8 : one block of four, $h = \frac{1}{2}$). [C] quasi-free extension to the full $\ast$ -algebra; Kac positivity beyond level 2.
 - **WP5d — both stages (done)**: α [E]/[C], β [E]/[C] (**continuum uplift** [O]): “the local net”, approached from the lattice. The α stage — the KLM *two-interval index* as the operational Haag-duality discriminator ($(E_8)_1 \Rightarrow \mu=1$, one sector; $SO(16)_1 \Rightarrow \mu=4$) — is executed and machine-verified as CELEST.WP5D.01 [verification/v501_celestial_wp5d_two_interval_index.py] (39 checks; Gaussian lattice machinery ED-validated to 10^{-15} , exact Fractions for the arithmetic). Measured on the 16-layer seam carrier (critical Majorana ring, $c_- = 8$): the fermionic two-interval

- mutual information is *extensive* (the $\mu=1$ reference; c fit 0.5000, residual $\rightarrow 0$ with N), while the sector-summed (orbifold) prescription pays exactly one classical bit — the $\ln 2$ plateau at machine precision ($|\Delta_2 - \ln 2| = 1.1 \times 10^{-15}$ at $N=512$), so $[F : F_{\text{even}}] = 2$ and $\mu_{\text{gauged}} = 4$, the entropic twin of the **v490** parity census (two independent lattice witnesses); the orbifold also breaks two-interval complementarity ($S(E) \neq S(E')$, the direct duality-failure witness), with the complementary-pair budget bounded by $\ln 4 = \ln \mu(SO(16)_1)$ as a *double-limit* statement. The condensation arithmetic is anchored at both measured ends: $\det \text{Cartan}(D_5) \cdot \det \text{Cartan}(A_3) = 16$, $\text{KLM/Longo-Rehren } 16/4^2 = 4/2^2 = 1$, $\Sigma d_i^2 = (4, 4, 1)$, $\theta_v = 1$ exactly at $\nu = 2c_- = 16$ (rivals $\neq 1$) — so $\mu = 1$ after condensation and the preregistered KILL (“ μ -offset $\neq 0$ after condensation”) is *not* triggered; the $\nu=1$ control shows a non-removable offset ($\theta_v \neq 1$: the discriminator has teeth), the trivial phase shows none, and the wrong sector sum loses the full $\ln 2$. **[C]** the continuum dictionary “offset = $\ln$ index” (KLM; Longo–Xu; Casini–Huerta–Magan–Pontello). **The β stage (done) [E]/[C] — split + strong additivity, the two remaining KLM legs** — is executed and machine-verified as CELEST.WP5DB.01 [verification/v504_celestial_wp5d_klm_completeness.py] (37 checks; Gaussian lattice machinery ED-validated, exact GF2/integer algebra for the exact legs; the $U(1)$ /Dirac number-conserving subnet as the preregistered negative control — the current-net analogue where strong additivity famously *fails* in the continuum). *Strong additivity is algebraically exact on the lattice*: with the shared boundary Majorana (“touching = share the point”), $\text{Even}(A) \vee \text{Even}(B) = \text{Even}(A \cup B)$ *exactly* (GF2 spans full $64/64$, $256/256$, $512/512$, $1024/1024$; literal matrix rank $32/32$), while disjoint intervals give exactly *half* (rank $16/32$, index 2 — the missing sector is odd $\otimes$ odd, the single $\ln 2$ bit of the α stage, localised at the split point); the $U(1)$ /neutral algebras do *not* generate the union even with the shared site (gaps $2/10/52$, growing). The entropic witness uses the honest discriminator *bounded vs divergent*: the $\mathbb{Z}_2$ /orbifold touching deficit stays $< \ln 2$ with the Ising approach $(\ln 2 - \Delta_2) \sim N^{-p}$, $p = 0.2444$ vs $2\Delta_\mu = \frac{1}{4}$ (the naive “defect $\rightarrow 0$ ” expectation is *false* at the sharp lattice split — bounded $\Leftrightarrow$ finite index, Longo–Xu), while the $U(1)$ control *bursts* $\ln 2$ from $L = 128$ on and grows as $\frac{1}{2} \ln \text{Var } Q_A$ (Klich–Levitov slope 0.10134 vs $1/\pi^2 = 0.10132$: infinite index). The split property is witnessed quantitatively: the coupling-block singular values fall exponentially at the elliptic-nome rate $\pi K(1-x)/K(x)$ to 1.3–2.0% over $d = 16..512$, $\sigma_1 \sim x^{0.5044}$ (vs $\frac{1}{2}$), trace norm summable — and the orbifold *inherits* the same ladder exactly at machine level (P_A flips $C \rightarrow -C$, σ_k identical to 8.3×10^{-17} ; the even-bilinear coupling Gram is the second compound $\Lambda^2 C$ with singular values $\{\sigma_a \sigma_b\}$ — Longo heredity, **[C]**). The Pimsner–Popa bound is a one-line *identity*, $E(a) - a/2 = PaP/2$, so $\lambda = \frac{1}{2} = 1/[F : F_{\text{even}}]$ exactly, attained in exact integers ($16384 + 2048$ monomial sweeps, 0 violations; random pencil $\min \lambda = 0.5000000000$); the two-interval group gives $\lambda_{E_4} = \frac{1}{4} = 1/\mu$ exactly, and the index is pinned by *two independent routes* ($e^{\Delta_\infty} = 2.0000000000 = 1/\lambda_{\text{PP}}$; $e^{2\Delta_\infty} = 4 = 1/\lambda_{E_4} = \mu$); the $U(1)$ control has $\lambda = 1/(m+1) \rightarrow 0$. *With the α stage, all three KLM ingredients of complete rationality — split, strong additivity, finite μ — are now witnessed on the lattice.* The continuum uplift stays **[O]** and is honestly fenced: the theorem “finite-group orbifolds of completely rational nets are completely rational (hence split + strongly additive)” is *Xu’s theorem* (algebraic orbifold CFTs) — cited, *not* claimed; the continuum proof for the concrete seam quotient net on the collar (**v336/v449**) and for the interacting condensed $(E_8)_1$ net remains WP5e / Costello–Li territory.
- **WP5e (subdivided): α stage (done) [E]/[C] — the CFT-side prefactor + level pinning; β stage (done) [E]/[C] — the equivariant anomaly ledger on twistor space, including the honest a_0 refutation; γ stage (done) [E]/[C] — the sphere-axion pairing check, a rigid *negative* result with certificate; δ_2 stage (done, intermediate verdict, exact) [E]/[C] — the full-tensor ledger: the**

collapse confirmed, one cubic door opens; ε_2 stage (done, verdict B) [E]/[C] — **the CPS level-from-flux dial, with “one level” lifted to a theorem; the global BCOV quantisation stays [O].** The α stage is executed and machine-verified as CELEST.WP5E.ALPHA.01 [verification/v502_celestial_wp5e_prefactor_level.py] (exact sympy/Fraction arithmetic, 33 checks). Both exactly computable consistency anchors of the uplift hold. (i) *The $q^{-1/3}$ prefactor is exact μ_4 vacuum-energy bookkeeping:* the clock is *inner* ($h = (2, 2, 2, 2, 2; 0^4)$) and the A_3 detector $h' = (0^5; 1, 1, 1, -3)$ both read the glue class mod 4 on all 240 roots; $(h, h) = 20$, $(h', h') = 12$, $\text{sum } 32$), so the induced twist on the 8 torus bosons is $\theta = 0^8$ in *every* sector — a *shift* orbifold, not a rotation orbifold — and each sector carries the same $8 \times (-\frac{1}{24}) = -\frac{1}{3} = -c/24$ (Hurwitz- ζ exact, $E_b(\theta) = -\frac{1}{24} + \theta(1 - \theta)/4$); the sector ground exponents are $-\frac{1}{3} + (0, 1, 1, 1)$ with h_j = the discriminant form, an identity holding via spectral flow ($j^2(h, h)/32$, mod 1) *and exactly* via the $10 + 6 = 16 = 2^{g_{\text{car}}-1}$ Majorana free fermions of the WP5d seam carrier (R-NS shift $n/16$: $\frac{5}{8}, \frac{3}{8}, 1$); the four sector characters sum to $E_4/\eta^8 = q^{-1/3}(1, 248, 4124, 34752)$; the rotation reading *fails* (the geometric A_3 deck gives $\frac{3}{16} \neq \frac{3}{8}$ — the WP1 “clock $\neq$ deck” lesson at the Casimir level). (ii) $k = 1$ *is forced three independent ways:* the current condition $h(J) = k = 1$ (the $\mathbb{Z}_4$ extension closes onto the E_8 adjoint $60+64+60+64 = 248$ only at spin 1); the conformal embedding ($D_5+A_3 \subset E_8$: $47k^2+219k-266 = 47(k-1)(k+266/47) = 0$; $D_8 \subset E_8$: $128k(1-k) = 0$); and the central charge $248k/(k+30) = 8 \Leftrightarrow 240(k-1) = 0$ (= the prefactor itself); plus the WP5b singular-vector dial ($\|(E_{-1}^\theta)^m 0\|^2 = m! k(k-1) \cdots (k-m+1)$, singular vector exactly at grade $k+1$; $k = 1$ deletes $V(2\theta)$: $31124 - 27000 = 4124$). *Honest sharpening:* glue- h integrality $h(J^j; k) = k(0, 1, 1, 1)$ holds for *all* $k = 1..8$ and fixes nothing — the originally conjectured integrality route is dead; the three other dials carry. Controls: $D_8/SO(16)_1$ has the *same* $q^{-1/3}$ prefactor but $h = (0, \frac{1}{2}, 1, 1)$ (the discriminator is glue- h integrality, not the prefactor); $\mathbb{Z}_2$ or μ_4 rotation twists break the common prefactor; wrong $k \in \{2, 3, 4\}$ fails all five dials coherently. **The β stage (done) [E]/[C]** — **the equivariant anomaly ledger on twistor space** — is executed and machine-verified as CELEST.WP5E.BETA.01 [verification/v505_celestial_wp5e_beta_equivariant_ledger.py] (exact sympy/Fraction arithmetic, 47 checks). The twistor side is pushed as far as exact arithmetic reaches: the Atiyah–Bott/Lefschetz fixed-point skeleton of the one-loop box anomaly on $\mathbb{C}^2/\mathbb{Z}_4$ with the μ_4 -glue monodromy. (i) *The AB ledger is exact:* denominators $\det(1 - g_j^{-1}) = (2, 4, 2)$ with Dedekind sum $5/4 = (|\mathbb{Z}_4|^2 - 1)/12$; equivariant characters $\text{tr}_{g_j}(\text{Ad } E_8) = (248, 0, -8, 0)$ by two independent routes; invariant average $60 =$ the carrier; Frobenius 61568. (ii) *Okubo per sector, the honest sharpening:* only the *invariant* sector is Okubo-quadratic ($Q^{(0)} = 36\langle x, x \rangle^2$, $36 = \tilde{\lambda}_{e_8}^2$ — the v495 identity re-derived through the glue grading); the twisted sectors carry irreducible T_5/T_3 content, and the AB-weighted sum cancels the D_5 quartic *exactly* while leaving the *rigid* residual $32T_3$ (no admissible reweighting fixes it; the graded GS exchange is rank-obstructed in sectors 1–3) — twisted-sector closed strings must carry the rest [O]. (iii) *The index bridge* (the centrepiece): $f(m) = \frac{1}{4} \sum_j (i^{jm} - 1)/\det_j = \text{ch}_2(T_m) = -(C^{-1})_{mm}/2$ *exactly* — the fixed-point ledger and the McKay/Kronheimer intersection ledger are the same arithmetic; the glue-bundle ch_2 defect is the integer -78 by *both* routes (classes $(-24, -30, -24)$). (iv) *The level dials say $k = 1$ geometrically:* the lattice current count is 240 at $k=1$ and exactly 0 at $k = 2, 3, 4$; the embedding residual $47k^2+219k-266 = (0, 360, 814, 1362)$; $h(J; k) = k$ is integer for *all* k (integrality alone fixes nothing — honest, exactly as on the CFT side); one scale $\Rightarrow$ one level (v493 periods; since the ε_2 stage a *theorem* of clock invariance, v509). (v) *The honest refutation:* the clock-invariant modulus $a_0 \in O(8)$ fills the BSS graviton slot $O(2)$, *not* the axion slot $O(-2)$ (Hamiltonian weights $X = -2 \neq +2 = Y$; mismatch $4 = |\mu_4|$, typed [C] as a coincidence) — “the theory brings its own GS axion as a_0 ” is *refuted*; what does match: the three $H^2(\text{ALE})$ classes carry exactly the three twisted-sector Coxeter

characters $\{i, -1, -i\}$ (bijection, no invariant class), so the bulk axion must come from the $O(-2)$ tower field itself (construction [O]). Controls: $\text{diag}(i, i)$ breaks the ledger four independent ways; $SO(16)$ glue gives characters $(120, 0, +120, 0)$ and defect $-30 \neq -78$ with a *failing* bulk Okubo (independent quartic Casimir); $k = 2$ dies on the closure dial ($-31/48 \neq -\frac{1}{3}$) while naive integrality dials stay integer. The preregistered kill does *not* fire on the equivariant skeleton. **The γ stage (done) [E]/[C] — the sphere-axion pairing check, an honest rigid negative result** — is executed and machine-verified as CELEST.WP5E.GAMMA.01 [verification/v508_celestial_wp5e_gamma_pairing.py] (exact sympy/Fraction arithmetic, 27 checks). Question: can the three twisted sphere axions η_m (Coxeter eigencharacters $\{i, -1, -i\}$ on $H^2(\text{ALE})$, the β -stage slot bijection) cancel the rigid $32 T_3$ residual ($A_{\text{fix}} = 9P_1 - 30P_2 - 15P_3 + 32 T_3$) by Green–Schwarz-type *exchange* with sector-compatible quadratic vertices? *No, with a certificate.* (i) *Vertex-space theorem*: the $W(D_5) \times W(A_3)$ -invariant quadratics on the 8-dim glue Cartan are *exactly* $\text{span}\{s_5, s_3\}$ (dim 2) and the invariant quartics *exactly* $\text{span}\{P_1, P_2, P_3, T_5, T_3\}$ (dim 5) — complete by Weyl nullspace arithmetic over all bidegrees; the *product theorem* is the master kill: the product of any two invariant quadratics has identically zero T_5/T_3 content, so every exchange image — any axion count, any selection rule, any couplings — is annihilated by Φ_{T_3} while $\Phi_{T_3}(A_{\text{fix}}) = 32 \neq 0$. (ii) *Selection enumeration*: the strict two-index $\mathbb{Z}_4$ rule collapses (the class bilinears are nonzero only for $j' = -j$, forcing $m = 0$: the sphere axions have *no* invariant bilinear Cartan vertex at all); the generous twist-insertion channels $E_{13} = (16, -96, 144, 0, 0)$, $E_{22} = (16, 32, 16, 0, 0)$ give a rank-2 exchange matrix with $\text{rank}([M | A_{\text{fix}}]) = 3$ and annihilator certificate $(\Phi_{T_5}, \Phi_{T_3}, \Phi_P)(A_{\text{fix}}) = (0, 32, 72)$ — two independent obstructions, the second driven by the side discovery $K^{(0)} = -15 K^{(2)}$ (the even-sector quadratics are parallel); $\ker M = 0$ forces the unique non-disturbing coupling $c = 0$ (the sphere axions are rigidly *silent*), and the sector-0 Okubo mechanism is undisturbed ($\Phi_P(Q^{(0)}) = 0$). (iii) *Naturalness dissolves*: ch_2 -natural and Atiyah–Bott-weight couplings both carry certificate $(0, 0)$ against the required $(32, 72)$ — the failure is scale- and coupling-independent (the ansatz space misses the target identically, not by an unnatural factor). Controls: the flipped rule changes the enumeration but not the T_3 kill (Φ_P is rule-dependent, Φ_{T_3} is not); $SO(16)$ has *no* sphere partners (odd McKay classes empty) *and* uncanceled T_5 $((15, -18, -15, -4, 40) — E_8$ is doubly special); D_8 has no T_3 structure at all $(16 T_8 + 12 s_8^2)$. Number fences typed [C] only: $32 = (h, h) + (h', h') = 20 + 12$, $72 = 2\tilde{\lambda}_{e_8}^2$. The β -stage slot bijection is untouched and the preregistered level-kill still does *not* fire — what dies is the exchange *realisation* of the pairing. **The ε_2 stage (done, verdict B) [E]/[C] — the CPS level-from-flux dial** — is executed and machine-verified as CELEST.WP5E.EPS2.01 [verification/v509_celestial_wp5e_eps2_level_from_flux.py] (exact sympy/Fraction arithmetic, 28 checks). Target: $k = 1$ from flux quantisation on the lockstep spheres, following the Costello–Paquette–Sharma Burns-holography mechanism (PRL **130**, 061602; arXiv:2306.00940). (i) *The CPS skeleton is replicated exactly*: the S^3 period of the brane back-reaction 3-form is $(2\pi i)^2 N$ (pullback density $-\sin 2\theta$, their eq. 3.33 — large-gauge invariance quantises N); the exceptional-sphere Kähler flux is $2\pi N$ exactly; the boundary level magnitude is $2N$ via the symplectic-boson double contraction ($N = 1, 2, 3$), Dynkin normalisation $T(\text{vector}) = 1$ — “level = flux quantum $\times$ Dynkin index”, *linear* in the flux. (ii) *The pairing matrix is pinned, not postulated*: the β -stage ch_2 ledger + integrality + unimodularity + effectivity force $P_{mi} = c_1(T_m) \cdot [\Sigma_i] = \delta_{mi}$ by complete enumeration (row candidates $8/6/8$; 48 unimodular solutions; with $P \geq 0$ exactly 2: identity + the A_3 diagram flip), so the glue fluxes are $F_i = (64, 60, 64) = \dim \mathfrak{g}_i$ (total $188 = 248 - 60$, flip-invariant) and every charged root current carries *exactly one* flux quantum through exactly one sphere. (iii) *The level formula, honest*: “level = total open-string flux” is *killed by the lockstep test itself* ($k_i = (64, 60, 64)$ unequal — F_i is the flavour multiplicity, the brane

matter content); what works is $k = \text{flux per adjoint current} = F_i / \dim \mathfrak{g}_i = (1, 1, 1)$, anchored non-circularly by the lattice current count $(240, 0, 0, 0)$ for $k = 1..4$ and the embedding index $j(D_5 \subset E_8) = j(A_3 \subset E_8) = 1$ ($60 = 2h^\vee$). The ord-4-vs-level-1 tension is *resolved*: the clock order counts *fractional flux sectors*, not the level ($\mu = 16 = \text{ord}^2$; the μ_4 glue diagonal is isotropic and Lagrangian; KLM $16 \rightarrow 16/4^2 = 1$, zero residual fractional flux), and a *new dial* pins the value: $\#\text{primaries}((E_8)_k) = (1, 3, 5, 10, 15, 27)$ for $k = 1..6$ from the affine marks — exactly *one* sector iff $k = 1$, independent of the CPS transplantation. (iv) *The lockstep theorem* (the lifting of the β -stage “one scale $\Rightarrow$ one level” to a *theorem*): clock invariance forces $a_3 = a_2 = a_1 = 0$ sharply, the four branch points are one μ_4 orbit, $|\Pi_j| = \sqrt{2}t$ is *equal* on all three spheres, the monodromy acts as the pure phase i and $\det(A - 1) = -4 \neq 0$ (no invariant flux vector) — with the CPS flux-level dictionary, $k_1 = k_2 = k_3$ is a *theorem* of clock invariance; the new falsifier: the clock-forbidden family $Z^4 - Z$ (smooth, disc -27 , branch points not a μ_4 orbit) gives *zero* lockstep chains over all 24 orderings. Honest limit: uniform flux doubling keeps the lockstep — the theorem proves *equality*, not the value; the value 1 comes from the current/sector counters and the CPS minimal quantum $N = 1$. Controls separate honestly: $k = 2$ dies on the closure dials (count 0, residual 360, prefactor $-31/48$, $\#\text{prim} 3$) while the lockstep dial honestly does *not* fire; $SO(16)$ glue leaves spheres $1/3$ flux-dark with condensation stuck at 4; the A_2 form has $\mu = 12$ (no Lagrangian subgroup exists at all) and Coxeter order $3 \neq 4$. Typed honestly: the CPS dictionary itself on $\mathbb{P}T/\mathbb{Z}_4$ stays **[C]** (their geometry is the $\mathbb{C}^2$ blow-up with one (-1) -sphere and $\mathfrak{so}(8)$ at negative level — not the A_3 ALE); the type-I B-model back-reaction on $\mathbb{P}T/\mathbb{Z}_4$ stays **[O]** only in its δ_{1d} derivation face (all three milestones M1–M3 have since been executed, **v515/v516/v517**; the δ_1 chain itself has since been *decided* by **v518** — kill under the derived measure — and the declared/derived measure question has since been decided at probe level in favour of the declared reading under the typed premises TP-1..TP-4, **v520**; the residual **[O]** is the constructive w_m derivation). **The δ_2 stage (done, intermediate verdict, exact) [E]/[C] — the full-tensor ledger: the collapse is real, and one cubic door opens** — is executed and machine-verified as CELEST.WP5E.DELTA2.01 [[verification/v511_celestial_wp5e_delta2_tensor_ledger.py](#)] (exact Kostant/Weyl weight arithmetic + sympy/Fraction, 41 checks). The γ -stage collapse was a *Cartan-restricted* statement; δ_2 recomputes the ledger on the full adjoint tensor structure. (i) *Structure*: $\mathfrak{g}_0 = \mathfrak{d}_5 \oplus \mathfrak{a}_3$ is *semisimple* with no $\mathfrak{u}(1)$ factor (class-0 roots split $40+12$, root span rank 8); the twisted sectors factorise multiplicity-free into minuscule Weyl orbits $\mathfrak{g}_1 = (16_s, 4)$, $\mathfrak{g}_2 = (10, 6)$, $\mathfrak{g}_3 = (\mathfrak{g}_1)^*$, each irreducible; and the *innerness theorem* decides the collapse wholesale: the grading element h lies in the Cartan of $\mathfrak{g}_0$, so every $\mathfrak{g}_0$ -invariant tensor carries total sector charge $0 \bmod 4$ — the γ -stage collapse is a *full-tensor* statement and holds across arities (all 15 non-neutral trilinear sector triples have $\text{Hom} = 0$: the sphere axioms have no invariant partner operator at any arity ≤ 3). (ii) *The Hom tables*: bilinear invariants exist only at $j' = -j$ (dims $2/1/1/1$, the Killing blocks); the neutral trilinear multisets carry dims $(3, 2, 2, 1, 1)$, all cross-sector trilinears are pure brackets (antisymmetric in their repeated legs) — the *only* totally symmetric trilinear on all of $\mathfrak{e}_8$ is the $\mathfrak{su}(4)$ d -symbol ($\mathfrak{so}(10)$ has no cubic Casimir: $\text{Inv}(S^3 45) = 0$ protects T_5). (iii) *The new channel*: the quartic projection of the d -symbol exchange is $Q_{dd} = (0, 0, -\frac{1}{240}, 0, \frac{1}{60}) = \frac{1}{60}(T_3 - P_3/4)$ by two independent routes (Killing propagator $2h^\vee = 60$ from the roots; hand route $|x^2 - (\text{tr } x^2/4)\mathbf{1}|^2/60$), and $\Phi_{T_3}(Q_{dd}) = \frac{1}{60} \neq 0$ — the γ -stage master kill (the product theorem) covers *quadratic* vertices only and does not extend to cubic ones: the T_3 direction is reachable through the d -symbol channel, while brackets, tadpoles and mixed $d \times f$ die exactly and no T_5 analogue exists. (iv) *The new certificate*: $M = [E_{13}, E_{22}, E_{00}, Q_{dd}]$ has rank 3, augmented rank 4 — the pairing in the charge reading remains *obstructed*, but with the genuinely *weaker* certificate $\{\Phi_{T_5}, \psi = \Phi_P - \Phi_{T_3}/4\}$, $\psi(A_{\text{fix}}) = 72 - 8 = 64$; and

the relaxed reading (charge bookkeeping dropped on the bilinear P -block) becomes *exactly solvable*: $A_{\text{fix}} = -u + 8v + 2w + 1920 Q_{dd}$ (residual 0) — where the γ stage found $T_5 = T_3 = 0$ persisting even relaxed, the d -channel closes the T_3 gap with small integer coefficients. Number fences typed **[C]** only: $64 = \dim \mathfrak{g}_1 = 2^6$; $1920 = |W(D_5)| = 8 \cdot 240$ — and the 1920 fence has now been stress-tested and *fails as a derivation claim* (`[verification/v513_celestial_dterm_nonderivation.py]`, `CELEST.DTERM.NONDERIV.01`): the $1920 = |W(D_5)|$ reading is look-elsewhere-loaded (11/924 pre-declared catalog expressions hit 1920, vs. 8/924 for the control target 1800) and convention-contingent (only 1 of 5 normalisations produces 1920: Gell-Mann gives 120, $\mathfrak{su}(4)$ -Killing 256, plain trace 32, ch_3 -normalised 69120); the charge-legal flux pairings give 2048/1800 — they bracket and miss — while the only flux hit is the charge-forbidden (1, 2) pairing; quantisation delivers only lattices ($60\mathbb{Z}$, with 1920 the 32nd multiple; strict ch_3 units exclude it); and the twisted cubic index $32i$ clashes in class provenance with the true quartic 32 (classes 1/3 vs. 0/2: same number, disjoint mechanism). The convention-stable **[E]** core is the factorisation $c_d = \Phi_{T_3}(A_{\text{fix}}) \times 2h^\vee(E_8) = 32 \times 60$; the fence stays **[C]**, the physical generation of the 32 stays **[O]** — no marker moves. Controls: the Killing control replicates the γ -stage channels and A_{fix} exactly; $SO(16)/D_8$ has *no* symmetric cubic ($\text{Inv}(S^3 \text{adj}_{D_8}) = 0$), no odd sectors and no A_3 block — the δ_2 channel cannot even be posed (E_8/μ_4 doubly special); the false $\mathfrak{g}_0 = \mathfrak{d}_5 \oplus \mathfrak{a}_2 \oplus \mathfrak{u}(1)$ inflates every Hom dimension ($(5, 2, 2) \neq (2, 1, 1)$, $\text{Inv}(S^3 \mathfrak{g}_0) = 6 \neq 1$) and the false sector content $(16_s, \bar{4})$ kills every pairing. What stays open after δ_2 **[O]**: the arity ≥ 4 ledger, the *physical* justification of the cubic GS term ($c_d = 32 \times 60$, the convention-stable core; the $1920 = |W(D_5)|$ reading is look-elsewhere-loaded and convention-contingent, **v513**; ch_2 -naturalness for cubic vertices *undecided* — no canonical geometric datum exists for a cubic vertex), and the burden on δ_1 is *sharpened*: it must supply either the $\psi = 64$ slice or the selection-rule relaxation — a burden since *discharged* by the M2 milestone: the declared KS measure supplies the $\psi = 64$ slice exactly (**v516**), so the cubic d -channel is not needed ($c_d \text{ free} = 0$) and its physical-justification question stays dissolved — the δ_1 decision (**v518**) does *not* revive it (no branch fires under the derived measure). **The δ_1 stage (done, DECIDED — kill under the derived measure, the $\mathbb{Z}_2$ anchor a method boundary) [E]/[C]** — is executed and machine-verified as `CELEST.WP5E.DELTA1.01` (`[verification/v518_celestial_wp5e_delta1_derived_kill.py]` (exact $\mathbb{Q}(\zeta_8)/\text{phase arithmetic} + 30\text{-digit kernels}$, 30 checks; the $\delta_{1b}/\delta_{1c}/\delta_{1d}$ probe chain consolidated). The exploration stage had left the verdict undecided with four exact sharpenings — (i) the strict holomorphic q^0 reading *refuted at the $\mathbb{Z}_2/\text{Eguchi-Hanson anchor}$* (a method boundary, not a kill of the contract); (ii) the forced leading (T_5, T_3) ratio 4:3; (iii) the contact term = the *modular completion* of the Atiyah–Bott data (a Harvey–Moore-type τ -integral, orbit split $15 = 12 + 3$); (iv) massive coset levels $n \geq 2$ mandatory. The τ -integral has now been *evaluated under a derived measure*, and the route is decided. Four exact results: (i) *the 16-component Weil completion closes*: the discriminant module of $D_5 \oplus A_3$ is $\mathbb{Z}_4 \times \mathbb{Z}_4$ with $q = (5x^2 + 3y^2)/8$, Gauss sums $2\zeta_8^5 \times 2\zeta_8^3 = 4 = \sqrt{16}$ (signature 0 mod 8 = rank E_8), the exact Weil relations hold in $\mathbb{Q}(\zeta_8)$, the diagonal and anti-diagonal Lagrangians are the two E_8 gluings, and the 16-vector theta S -covariance drops the naive-4-rule residual from $2.91 = O(1)$ to $\sim 10^{-39}$; (ii) *the obstruction is a finite μ_4 character, identified*: the dressed gauge blocks $M = G[a, b] \eta^{-8}$ carry a multiplier system whose $SL(2, \mathbb{Z})$ relation defects are (1, 1, 1) exactly on all 15 pairs (a *character* of the orbit stabilisers $\Gamma_1(4)/\Gamma_0(2)^\pm$, not a genuine 2-cocycle), on $\Gamma_1(4)$ explicitly $\lambda(\gamma) = i^{2B+C/4}$; (iii) *the cancellation exists and is the twisted fibre block $f_1 f_3$, not the three sphere axions*: $G[a, b] = f_1 f_3$ is an exact identity, the T-fix mechanism is one line of exact phase arithmetic ($(-1) \times e(-1/6) = e(1/3) = \chi_4(T)$), and the exact cancellation table reads $\text{none} \rightarrow 4$, $f_1 f_2 f_3 \rightarrow 4$, $f_2 \rightarrow 6$, $f_1 f_3 \rightarrow 1$ (both orientations), wrong weights $\rightarrow 4$ — the strict

solutions are exactly χ_4 (dims (3,3)) and χ_{10} (dims (1,1)), certified on the dressed functions; (iv) *all three preregistered testers fail* under both derived solutions (T_5 fractions 0.5/0.83 instead of 0; $W_{13} = 0$ instead of $W_2:W_{13} = 4:3$; spreads 1.05/1.47 against $-A_{\text{fix}}$; no $\psi = -64N$ slice), and the honest (N_1, N_2) orbit rebalance rescues nothing in the positive cone — a genuine KILL on the derived surface. Controls: the $\mathbb{Z}_2/\text{EH}$ anchor (gauge-only residual order 2, cancelled to 1 by the same fibre dressing), $SO(16)$ (order-4 supply structurally absent), wrong form/wrong signature (break exactly). Fences [C]: the $f_1 f_3 = \text{KS-weight identification}$ (deck weights (1,3) = the twistor fibre weights forced by the v514 incidence ledger — exact match, *not* a complete BCOV derivation), the kernel-family convention, the discrete χ_k choice (both evaluated, both fail). TENSION, stated honestly (the mandatory fence): the DECLARED completion reading (v516/M2) delivers the $\psi = 64$ slice and cancels $32 T_3$; the DERIVED chiral measure (this module) fails all three testers — both are exact; the sharp open question was which of the two is the true BCOV measure (the declared reading is supported by the δ_1 modular-completion finding but not derived; the derived measure rests on the $f_1 f_3$ identification). Neither result is hidden behind the other. That question is since *decided at probe level* by v520 in favour of the declared reading (single-valuedness derived from F-independence + the Quillen pairing under the typed premises TP-1..TP-4; the kill of this module is *sharpened*: the column-canonicalised χ_4 member is boundary-consistent yet tester-dead); and the constructive BCOV derivation of the w_m normalisation itself is since *executed* by v523 (verdict ERFOLG; the residual [O] narrows to the global BCOV integral beyond the fibre zero-mode factor). **The ε_1 stage (done) [E]/[C] — the $O(-2)$ bulk-axion slot** — is executed and machine-verified as CELEST.WP5E.EPS1.01 [verification/v514_celestial_wp5e_eps1_axion_slot.py] (exact sympy/Fraction arithmetic, 34 checks, verdict B). Three results, each exact. (i) *The slot is a construction*: on $\mathbb{P}T' = \text{Tot}(O(1) \oplus O(1) \rightarrow \mathbb{P}^1)$ the pushforward ledger $\pi_* O(-2)$ gives per spacetime degree d exactly $(d+1)^2$ classes = the exact wave-operator nullspaces, and the *equivariant* bookkeeping under the v492 action $(\mu_1, \mu_2) \mapsto (i\mu_1, i^{-1}\mu_2)$ (incidence-forced column weights (+1, +1, -1, -1)) closes *block by block* for all $d \leq 6$ and all four characters; the character series are $P_0 = 1 + 3t^2 + 15t^4 + 21t^6 + 45t^8 + \dots$, $P_1 = P_3 = 2t + 8t^3 + 18t^5 + \dots$, $P_2 = 6t^2 + 10t^4 + 28t^6 + \dots$, the $d = 0$ slot has multiplicity (1, 0, 0, 0) — *the bulk axion survives the projection* — and the Molien invariant ring is $(1 - t^8)/((1 - t^4)^2(1 - t^2))$: $Z \in O(2)$, $X, Y \in O(4)$, one relation in degree 8 = the a_0 weight (the v492/v493 bridge); the E3 bijection is sharpened per character (twisted minimal content $(2t, 6t^2, 2t)$, no degree-0 mode, = the Coxeter eigenvalues, $\det(A - 1) = -4$), and the graviton control separates a_0 (the $O(+2)$ slot starts invariant only at fibre degree 4 with multiplicity $3 = \{X, Z^2, Y\}$). (ii) $\tilde{\lambda} = 6$ *is triply pinned*: Okubo $\text{Tr}_{\text{adj}} X^4 = 36\langle x, x \rangle^2 = (6\langle x, x \rangle)^2$ on the 240 glue roots ($36 = h^\vee + 6$, $\tilde{\lambda} = 6 = |\mathbb{Z}_2|N_{\text{fam}}$); the measure chain on $\mathbb{P}T/\mathbb{Z}_4$ cancels the quotient factor $\mu = 1/4$ *exactly* (vertex $^2 \mu^2$, propagator $1/\mu$, anomaly μ : $\tilde{\lambda}_{\text{eff}}^2 = 36$ for arbitrary μ), with the wrong bookings quantified and *excluded* (anomaly-only $\Rightarrow \tilde{\lambda} = 3$, missing propagator renormalisation $\Rightarrow \tilde{\lambda} = 12$); and the flux side (minimal CPS quantum $N = 1$, a single exchange channel exactly at $k = 1$, $(\kappa/c_3)^2 = 12$ exact). (iii) *The first honest back-reaction step (Gibbons–Hawking A_3)*: clock-invariant four-centre configurations form exactly two branches (axis 4 = pure resolution or *one* free μ_4 orbit = pure deformation); the free orbit gives $\prod_p (Z - i^p z_0) = Z^4 - z_0^4$, i.e. $XY = Z^4 + a_0$ with $a_0 = -z_0^4$ — the v493 family *re-derived from the geometry*, with the a_0 monodromy = one clock step; the charge- k GH point is exactly $\mathbb{R}^4/\mathbb{Z}_k$ ($k = 4$: the seam boundary $S^3/\mathbb{Z}_4$); the periods are $\Pi_j = 4\pi t_0(i - 1)(1, i, i^2)$ with equal moduli (lockstep, the v509 counter-check from geometry) and the Coxeter clock re-derived ($A^4 = 1$, $\det(A - 1) = -4$, $\Pi \cdot A = i\Pi$); the CPS “ $N \log$ ” analogue carries source charge $4 = |\mu_4|$, with the honest sharpening that the Ricci-flat ALE has asymptotic log coefficient exactly 0 (the CPS log is an exceptional-locus statement;

Burns contrast $\det g \neq 1$), and the multipole selection rule $m \equiv 0 \bmod 4$ stores $-a_0$ in the first symmetry-breaking $(4, \pm 4)$ harmonic. Controls: **508** ($\tilde{\lambda}^2 = 12$ irrational; perfect squares in the Deligne series only $\{9, 36\}$), $\text{diag}(i, i)$ (Veronese cone, no hypersurface), $k = 2$ (three channels). **[C]**: conditional on Costello's flat- $\mathbb{P}^1$ matching $\lambda_g^2 E = A$ (the measure chain is classical bookkeeping). **[O]** (the honest fence): the *quantised* BCOV one-loop coefficient on $\mathbb{P}^1/\mathbb{Z}_4$ and the twisted channels $(32T_3)$ — preregistered as M1–M3 below; all three have since been executed (**v515/v516/v517**), the δ_1 chain has since been *decided* by **v518** (kill under the derived measure), the measure question has since been decided at probe level by **v520** for the declared reading, and the w_m normalisation has since been *derived* at probe level by **v523**; what stays **[O]** is the global BCOV integral beyond the fibre zero-mode factor. **WP5e proper [O] (the global BCOV quantisation)**: the BCOV/Kodaira–Spencer theory on $\mathbb{P}^1/\mathbb{Z}_4$ itself (quantisation, non-fixed-point contributions, the twisted closed-string measure). The exchange sub-branch is *closed* by the γ stage, the full-tensor ledger (δ_2), the level-from-flux dial (ε_2) and the bulk-axion slot (ε_1) are *executed*, and δ_1 is *decided* by **v518** (kill under the derived measure); the named remaining target, with preregistered success/kill criteria: **(the measure question — the δ_1 successor; decided at probe level, ERFOLG-A)** the δ_1 τ -integral had been evaluated on *both* sides: the M2 milestone (**v516**) *delivers* the coefficient $(9, -30, -15, 0, 32)$ -completion exactly on the *declared* measure (every sector the same Okubo square, the $\psi = 64$ slice supplied), while the *derived* chiral measure (**v518**: blockwise $SL(2, \mathbb{Z})$ covariance solved for, the μ_4 multiplier character cancelled by the twisted fibre block $f_1 f_3$) fails all three preregistered testers — both are exact, and the stated tension was the sharp remaining question: *derive* from the BCOV integral on $\mathbb{P}^1/\mathbb{Z}_4$ which of the two measures is the true one. That decision is executed and machine-verified as **CELEST.WP5E.MEASURE.01** [[verification/v520_celestial_wp5e_measure_decision.py](#)] (the δ_{1e}/δ_{1f} consolidation, 39 checks, ERFOLG-A on the preregistered decision layer): the exact invariant subspace of the dressed 16-component Weil system is $\dim_{\mathbb{Q}}\{v : \rho(T)v = v, \rho(S)v = v\} = 8 = 4 \times 2$ with *every* kernel vector in the $\mathbb{Q}(\zeta_8)$ -span of $\{e_H, e_{H'}\}$ (both theta = E_4 = the unphased trace), and single-valuedness of the physical one-loop lattice factor is *derived* from two independent constructive sources rather than postulated — (*source 1, the F-lemma*) every strict chiral closure at physical weight carries a nontrivial character ($\chi_4(T) = e(1/3)$, $\chi_{10}(T) = e(5/6)$, exact; pointwise covariance of the canonical column-matched member certified at 3.9×10^{-16} with $|G| \geq 59.6$), and an exact change of variables forces $\int_{\gamma_F} G d\mu = \chi_4(\gamma) \int_F G d\mu$, so fundamental-domain independence forces $\int = 0$ for *every* chiral integrand: the **v518** route was never a well-defined nonvanishing moduli integral (the kill is *sharpened*); (*source 2, the Quillen pairing*) the doubled hol $\times$ antihol transports satisfy $|t\bar{t} - 1| < 8.3 \times 10^{-40}$ on all 15 pairs ($|\chi_4|^2 = 1$ exactly vs one-sided $\chi_4^2 = e(2/3) \neq 1$), the doubled 256-dim system closes *strictly* at trivial character and physical weight (nullspaces $(28, 17) \geq 11$) containing the unphased diagonal *and* the physical columns $w_b \otimes \bar{w}_b$, while the hol $\otimes$ hol control is *empty* $(0, 0)$; the forced unphased numerator then reproduces the **v516** Okubo squares *levelwise* inside the **v518** scaffold (exact 4-design on every level $n \leq 8$, J -weighted spreads $\sim 5 \times 10^{-41}$, $\psi(\text{skeleton}) = +0.2302 = -\psi(\text{contact})$ — the J -weighted mirror of ± 64), the declared reading also wins the column-canonicalisation discriminator (the physical AB column is contained *exactly* in the χ_4 family, canonicalising (N_1, N_2) , yet the canonical member fails every tester) and the $\mathbb{Z}_2/\text{EH}$ anchor (all three derived instantiations fail; only the declared reading hits $9\langle x, x \rangle^2 = (1/4) \times 36$ with the scale tooth $c = 1/2 = |\mathbb{Z}_2|h^{A_1}$). Typed premises **[C]** (the mandatory fence): TP-1 (F-independent moduli integral), TP-2 (nonvanishing — it must source $\psi = 64$), TP-3 (the **v518** kernel convention), TP-4 (the Quillen structure of BCOV F_1). The residual **[O]** was the constructive BCOV derivation of the w_m normalisation itself — the success branch (“the BCOV derivation

lands on the declared completion reading, carrying the forced 4:3 leading ratio”) has fired at probe level, the kill branch (“it lands on the derived chiral measure”) is dead — and that derivation is since *executed* (verdict ERFOLG) as CELEST.WP5E.WM.01 [verification/v523_celestial_wp5e_wm_derived.py]: the $1/\det_j$ weight is COMPUTED from three independent sources — the equivariant mode ledger of the fibre $\mathbb{C}^2$ (closed form $1/((1 - qi^j)(1 - qi^{-j}))$ exact at every order $n \leq 120$, regular at $q = 1$ with Abel value $(1/2, 1/4, 1/2) = 1/\det_j$, fixed-point splitting at every truncation level for the phased AND the v520-forced unphased trace, so $\text{contact}_j = (Q^{(0)} - Q^{(j)})/\det_j$ is computed, not declared), the zeta/Quillen route ($\det_\zeta = 4\sin^2(\pi j/4) = \det_j$ exactly with the unique real positive holomorphic section via the $SU(2)$ conjugation pairing, and the δ_{1f} block constant term $1/\det_b$), and the consistency chain (the v516 numbers reproduced one by one: $w = 4h$, $c = 4 = |\mu_4|$, squares $(18, 9, 18)$, total $45\langle x, x \rangle^2$, $\psi \pm 64$) — with the negative controls behaving ($1/\det^2$ and $1/|1 - i^j|$ break the chain quantifiably, the $\mathbb{Z}_2/\text{EH}$ anchor passes at $c = 2 = |\mathbb{Z}_2|$, the $SO(16)/D_8$ kill fires, $\text{diag}(i, i)$ breaks the zeta = AB identity itself, the $k = 3, 5$ weights wander correctly) and the typed premises TP-REG/TP-Q/TP-NUM/TP-CH as its own mandatory [C] fence; the residual [O] narrows to the global BCOV integral on $\mathbb{P}T/\mathbb{Z}_4$ beyond the fibre zero-mode factor (the resolved-ALE route of v514/v517 not built); **(the cubic GS term — dissolved)** the $\psi = 64$ slice is *delivered* by the declared KS measure (v516); the cubic d -channel is *not needed* (c_d free = 0) — its physical-justification question (the honest v513 target $c_d = 32 \times 60$) stays dissolved: the δ_1 decision (v518) does not revive the channel (no branch fires under the derived measure); a ch_2 -type naturalness datum for *cubic* vertices still does not exist; **(the back-reaction milestones, preregistered as M1–M3 in v514; all three executed — the v514 fence is fully worked off)** the type-I B-model *back-reaction* on $\mathbb{P}T/\mathbb{Z}_4$, which would upgrade the ε_2 CPS dictionary from [C] to derived: **M1 (done, SUCCESS)** [E]/[C] the A_3 Ω_N (the Beltrami/Kodaira–Spencer deformation sourced by the glue currents on the three lockstep spheres; preregistered success = a closed-form Ω -deformation whose $S^3/\mathbb{Z}_4$ periods are $(2\pi i)^2 \times$ an integer flux vector with the lockstep phases $(i - 1)(1, i, i^2)$; kill = unequal sphere fluxes or a non-integer period — then the CPS transplant *and* the ε_2 level dial die) — executed and machine-verified as CELEST.WP5E.M1.01 [verification/v515_celestial_wp5e_m1_omega_n.py] (exact sympy, 30 checks): the residue form $\omega_2 = dX \wedge dZ/X$ is *derived* (three ambient routes; orbifold-cover factor $4 = |\mu_4|$; clock phase i), the twistor family closes as $XY = Z^4 + 4t_0^2\lambda^2Z^2 - t_0^4(1 - \lambda^4)^2$ ($e_1 = e_3 = 0$ identically; seam fibre $\lambda = 0 =$ the v493 family) with the CY-compatible clock lift $(Z, \lambda) \mapsto (iZ, -i\lambda)$, $\Omega \rightarrow +\Omega$, and $\Omega_N = \Omega_0 + \sum_p N_p K_p$ is *closed form* (CPS/Bochner–Martinelli kernels on the four centre twistor lines) with every $S^3/\mathbb{Z}_4$ period $(2\pi i)^2$ -integral, the seam-fibre phase vector $2\pi i t_0(i - 1)(1, i, i^2)$ with clock covariance $\Pi \rightarrow i\Pi$, the flux vector $N(1, 1, 1, 1)$ forced uniform (clock orbit + K_4 connectivity) and the lens geometry *forcing* the source charge $4 = |\mu_4|$ ($N \in 4\mathbb{Z}$); the 12 conifold nodes sit exactly on the 8 eighth roots of unity ($8 = 2|\mu_4|$); honest fence: on the clock-forbidden $Z^4 - Z$ family the $(2\pi i)^2$ quantisation *also* holds — integrality alone is *not* the discriminator, the discriminator is the lockstep phase structure ($0/24$ vs $8/24$) plus the clock forcing of equal fluxes; neither kill fired; **M2 (done, SUCCESS on the declared completion measure — verdict B)** [E]/[C] the twisted KS measure (quantise the $O(-2)$ tower on $\mathbb{P}T/\mathbb{Z}_4$ with the ε_1 mode ledger and compute the one-loop box coefficient per sector; preregistered success = the bulk sector reproduces $\tilde{\lambda}^2 = 36$ *and* the twisted sectors pair with the three sphere axions cancelling the rigid $32 T_3$ residual; kill = a leftover independent T_3 -type quartic in any sector) — executed and machine-verified as CELEST.WP5E.M2.01 [verification/v516_celestial_wp5e_m2_twisted_ks_measure.py] (exact sympy/Fraction, 23 checks): the *declared* completion contact term $\text{contact}_j = (Q^{(0)} - Q^{(j)})/\det_j$ (the twisted-sector loop contributes the *unphased*

sector trace with the g^j zero-mode normalisation) carries the exact completion-weight identity $w_m = \sum_j (1 - i^{jm}) / \det_j = (0, \frac{3}{2}, 2, \frac{3}{2}) = 4h_m = |\mu_4|h_m = -4\text{ch}_2(T_m)$ — the three sphere axions pair through their *own* McKay ch_2 charges, *no free scale, no fit*; the locks are parameter-free ($T_5 = 0$ for *any* scale, ratio $h_2:h_1 = 4:3 =$ the δ_1 -forced leading ratio *reproduced*, the T_3 budget forcing $c = 4 = |\mu_4|$ uniquely); every twisted channel becomes the *same* perfect Okubo square $36\langle x, x \rangle^2 / \det_j$ ($T_5 = T_3 = 0$ in every sector — the KILL does not fire), the total is $45\langle x, x \rangle^2 = \frac{5}{4} \times 36\langle x, x \rangle^2 = \text{Dedekind} \times \text{Okubo} =$ the unique quartic-free weighting of the β -stage rigidity theorem; both γ -stage certificates are killed ($\Phi_{T_3} : 32 \rightarrow 0$, $\Phi_P : 72 \rightarrow 0$) and the δ_2 slice $\psi = 64$ is *supplied exactly* — no cubic d -channel needed ($c_d \text{ free} = 0$); controls: wrong scale leaves $T_3 = 32 - 8c$, the shuffle breaks T_5 and T_3 , $SO(16)$ keeps $T_5 = 20/T_3 = -40$ (the KILL fires there: E_8 doubly special), $\text{diag}(i, i)$ degenerates, the $\mathbb{Z}_2/\text{EH}$ anchor passes at scale $2 = |\mathbb{Z}_2|$. Mandatory fence [C]: the completion reading (the loop of twisted states carries the *unphased* sector trace with the same zero-mode normalisation) is *declared*, supported by the δ_1 modular-completion finding, *not* derived from the BCOV integral within that module — the $\delta_{1b}/\delta_{1c}/\delta_{1d}$ chain has since been executed and decided by v518 (the *derived* chiral measure fails all three testers), the declared-vs-derived measure question has since been decided at probe level by v520 for the declared reading (typed premises TP-1..TP-4), and the w_m normalisation has since been derived at probe level by v523 (the residual [O] narrowing to the global BCOV integral beyond the fibre zero-mode factor); **M3 (done, SUCCESS)** [E]/[C] the a_0 uplift (lift the $(4, \pm 4)$ GH multipole to the Beltrami differential and compute the induced Kähler-potential correction; preregistered success = a log-type correction with coefficient tied to the source charge $4 = |\mu_4|$; kill = decoupling from the centre count) — executed and machine-verified as CELEST.WP5E.M3.01 [verification/v517_celestial_wp5e_m3_a0_uplift.py] (exact sympy, 23 checks): the uplift object is the generalized-Legendre-transform kernel $\chi = \log P_4$ — the *log of the M1 family polynomial* — well-typed (the $O(2)$ section is a null coordinate: *any* kernel is harmonic) and matching the V-ledger through the exact residue identity $\partial_x[\text{transform}(\log \eta_p)] = 1/r_p$ (flux -4π per centre, source charge $4 = |\mu_4|$); the log-type correction arrives on *four coupled centre-count scales*: (i) the asymptotic kernel $\log \chi = 4 \log \eta + a_2/\eta^2 + (a_0 - a_2^2/2)/\eta^4 + \dots$ with coefficient $4 = |\mu_4|$, the seam-fibre first correction *exactly* $a_0/\eta^4 =$ the $(4, \pm 4)$ multipole (exact m -grading, no $\log \times$ power terms — the ε_1 honest zero preserved); (ii) the GLT tower $p_{4k} = 4(-a_0)^k$ with the $n \equiv 0 \pmod 4$ selection rule; (iii) the exceptional-locus $\log \chi(0) = \log a_0 = 4 \log t_0 + i(4\varphi_0 + \pi)$ (response 1 at the locus vs power-law at infinity); (iv) the period response $d \log \Pi_j / d \log a_0 = 1/4 = 1/|\mu_4|$ uniformly, integrating to the monodromy $i =$ one Coxeter clock step (the WP2 monodromy reproduced from perturbation theory), with a_0 -rigid $\mathbb{Z}_8$ node support and topological $(2\pi i)^2$ fluxes; controls: the $(4, 0)$ multipole is clock-invariant, the $\mathbb{Z}_2/\text{EH}$ analogue reads $2 = |\mathbb{Z}_2|$ on *every* dial, $k = 3, 5$ orbits move the coefficient with the centre count ($O(6)/O(10)$ slots), the forbidden family fails both dials ($p_3 = 3$, $e_4 = 0$) — the KILL (decoupling) does not fire; fence [C]: the GLT dictionary ($\eta_p \log \eta_p =$ the GH Kähler-potential kernel, Lindström–Roček / Ivanov–Roček); [O]: the full nonlinear Kähler potential of the resolved A_3 ALE. Success (unchanged): partition function $= E_4/\eta^8$ *including* the $q^{-1/3}$ prefactor *derived from the twistor side*; kill: an anomaly mismatch at the Costello–Li grade — neither the CFT-side dials of v502, nor the equivariant skeleton of v505, nor the exchange no-go of v508, nor the flux/sector dials of v509, nor the full-tensor ledger of v511, nor the axion-slot construction of v514, nor the M1–M3 back-reaction milestones of v515/v516/v517, nor the δ_1 decision of v518, nor the measure decision of v520 trigger it (all computable dials say $k = 1$, “one level” is now a theorem, the bulk axion survives the projection, the back-reacted Ω_N carries the forced integral lockstep flux, the twisted KS measure lands on the unique quartic-free weighting, the a_0 uplift

couples to the centre count, the δ_1 kill is a statement about the derived measure, not about the algebra, and the measure decision selects the declared reading at probe level), but the global quantisation is untested.

Honest status line: WP5a–WP5d (both WP5d stages), the WP5e α , β , γ , δ_1 , δ_2 , ε_2 and ε_1 stages and the M1–M3 back-reaction milestones are exact-arithmetic [E] constructions (plus ED-validated lattice witnesses in WP5d) with [C] named identifications (“radial decoupling = collar thickness $\rightarrow 0$ ”; Weyl complete reducibility beyond BFS depth 4; the Kac mechanics beyond level 2; the quasi-free extension; the KLM/Longo–Xu/CHMP index dictionary; Longo split heredity; Kac generation of the maximal submodule for $k \geq 3$; the WP5e- β dictionaries — $\text{CS}(\rho) \bmod 1 = \text{discriminant weight}$, $32 = (h, h) + (h', h')$, $-30 = -h^\vee$, the Ω -contraction; the WP5e- γ number fences $32 = 20+12$, $72 = 2\tilde{\lambda}_{e_8}^2$; the WP5e- δ_2 number fences $64 = \dim \mathfrak{g}_1$, $1920 = |W(D_5)|$ — the latter typed look-elsewhere-loaded and convention-contingent by v513; the WP5e- ε_2 CPS dictionary on $\mathbb{P}T/\mathbb{Z}_4$; the WP5e- ε_1 conditionality on Costello’s flat- $\mathbb{P}T$ matching; the M1 global kernel patching, Hitchin small resolution and CPS brane dictionary, v515; the M2 *declared* completion reading of the twisted KS measure, v516; the M3 GLT dictionary, v517); *WP5a–WP5d plus WP5e- $\alpha/\beta/\gamma/\delta_1/\delta_2/\varepsilon_2/\varepsilon_1$ plus M1–M3 are landed — all three KLM ingredients of complete rationality carry lattice witnesses, both faces of the inflow are anchored, the exchange sub-branch is closed with a certificate, the collapse is confirmed full-tensorially with one cubic door open, “one level” is now a theorem of clock invariance with the sector-counter dial pinning $k = 1$, the bulk-axion slot is a construction with $\tilde{\lambda} = 6$ triply pinned, the back-reacted Ω_N is closed-form with $(2\pi i)^2$ -integral lockstep periods and the source charge $4 = |\mu_4|$ forced, the twisted KS measure (on the declared completion reading) cancels the $32T_3$ residual and supplies the $\psi = 64$ slice without the cubic d -channel, the a_0 uplift couples to the centre count on four scales, and the δ_1 chain is decided — kill under the derived measure, tension with the declared reading stated — WP5e proper (the global BCOV quantisation) is the single remaining open piece, with the continuum uplift of the WP5d lattice witnesses honestly fenced as Xu’s theorem (cited, not claimed); the exchange sub-branch is closed by v508, the full-tensor ledger executed by v511, the level-from-flux dial executed by v509, the axion slot executed by v514, the M1–M3 back-reaction milestones executed by v515/v516/v517 (the v514 fence fully worked off), the δ_1 chain decided by v518, the measure question decided at probe level by v520 (the declared completion reading wins: single-valuedness derived from F-independence + the Quillen pairing under TP-1.TP-4; the derived chiral measure was never a well-defined nonvanishing moduli integral), the w_m normalisation itself derived constructively at probe level by v523 (Atiyah–Bott mode ledger + zeta/Quillen determinant + consistency chain, typed premises TP-REG/TP-Q/TP-NUM/TP-CH), and the named remaining target is the global BCOV integral beyond the fibre zero-mode factor (the cubic-GS-term question stays dissolved: v518 does not revive the d -channel) — SEAM.EQUIV.01 stays [O] and nothing in WP5a–WP5e-M3, the δ_1 decision or the measure decision moves it.*

Kill tests (pre-registered). (K1) the equivariant sector could have *contradicted* the $52+64+60+64$ split (uniformity of $D[d]$ needs exactly $\dim \mathfrak{g}_j = (60, 64, 60, 64)$) — *survived by WP1 and WP2*: the false-glue control shows the test has teeth (3112/6720 violations), and WP2 adds the deformation side (a shape modulus surviving clock-invariance, or a_0 -corrections leaking between μ_4 sectors, would have fired — neither occurred; the clock-variant controls fail in two independent ways each). (K2) if one-loop associativity of the celestial E_8 theory demanded extra fields that TFPT does not possess, the contract dies (Costello’s axionic mechanism is the first pass: E_8 is admissible; the TFPT-specific field content is the open half). (K3) — *evaluated by WP3* ([[verification/v495_celestial_wp3_gs_alignment.py](#)]): the trigger (“no exact alignment between the GS coefficient and the c_3 normalisation”) does *not* fire — exact alignment exists ($(\kappa/c_3)^2 = 12$ resp. $1/300$). But the look-elsewhere battery *demotes the scope of K3 itself*: the alignment format passes 8/8 across Costello’s whole list, so surviving K3 certifies compatibility

plus λ -arithmetic, *not* $\mathfrak{e}_8$ -selective evidence — alignment survives, selectivity does not. **(K4)** — *evaluated by WP4* ([[verification/v496_celestial_wp4_character_shadow.py](#)): K4 fires only against the *exact*-block reading (the character is provably not a conformal block in the jet grading); it does *not* degrade the contract to “ E_8 is an admissible gauge algebra for celestial SDYM”, because the sector arithmetic holds exactly — WP4 lands at the boundary-limit reading (verdict B(ii), the SEAM.EQUIV.01 scaling-limit shape), with the constructive limit handed to WP5 — and made precise there by WP5a ([[verification/v497_celestial_wp5a_null_ideal.py](#)): the coefficientwise χ_w limit with threshold $w = n+1$ and the derived ideal), with the deleting operator itself constructed by WP5b ([[verification/v498_celestial_wp5b_singular_vector.py](#)): $|s\rangle$ explicit, level dial $2(1-k)$), the limit state by WP5c ([[verification/v500_celestial_wp5c_gns_limit_state.py](#)): the ideal in the GNS kernel, level-2 rank 4124 exactly), the two-interval index chain by WP5d- α ([[verification/v501_celestial_wp5d_two_interval_index.py](#)): μ -chain $16 \rightarrow 4 \rightarrow 1$ anchored at both measured ends), the split + strong-additivity legs by WP5d- β ([[verification/v504_celestial_wp5d_klm_completeness.py](#)): all three KLM ingredients of complete rationality witnessed on the lattice; the continuum uplift is Xu’s theorem, cited not claimed), the CFT-side anchors of the uplift itself by WP5e- α ([[verification/v502_celestial_wp5e_prefactor_level.py](#)): the $q^{-1/3}$ prefactor as exact μ_4 vacuum-energy bookkeeping, $k = 1$ forced three ways), the equivariant twistor skeleton by WP5e- β ([[verification/v505_celestial_wp5e_beta_equivariant_ledger.py](#)): the index bridge $f(m) = \text{ch}_2(T_m)$ exact, glue defect -78 two routes, geometric $k = 1$ dials, the rigid $32 T_3$ residual, and the honest a_0 refutation — the kill branch does not fire on the skeleton), the exchange no-go by WP5e- γ ([[verification/v508_celestial_wp5e_gamma_pairing.py](#)): the sphere-axion pairing by quadratic-vertex exchange refuted with the three-functional certificate $(0, 32, 72)$ — the $32 T_3$ burden moves to the twisted contact terms, δ_1 binary), the full-tensor ledger by WP5e- δ_2 ([[verification/v511_celestial_wp5e_delta2_tensor_ledger.py](#)): the collapse confirmed full-tensorially by the innerness theorem, the cubic d -symbol channel opening T_3 with the weaker certificate $\psi(A_{\text{fix}}) = 64$ and the relaxed solution $c_d = 1920 Q_{dd}$ -coefficient — typed by the v513 negative certificate as $c_d = 32 \times 60$ with the $|W(D_5)|$ reading look-elsewhere-loaded), and the level-from-flux dial by WP5e- ε_2 ([[verification/v509_celestial_wp5e_eps2_level_from_flux.py](#)): the lockstep theorem — “one level” as a theorem of clock invariance, falsifier $Z^4 - Z$ at $0/24$ — plus the sector-counter dial $\#\text{prim}((E_8)_k) = 1 \Leftrightarrow k = 1$), the bulk-axion slot by WP5e- ε_1 (v514: the equivariant $O(-2)$ slot construction, $\tilde{\lambda} = 6$ triply pinned, the first GH/ A_3 back-reaction step), the back-reacted Ω_N by the M1 milestone (v515: closed form, $(2\pi i)^2$ -integral lockstep periods, forced source charge $4 = |\mu_4|$), the twisted KS measure by the M2 milestone ([[verification/v516_celestial_wp5e_m2_twisted_ks_measure.py](#)): the completion-weight identity $w = 4h = -4 \text{ch}_2$, per-sector Okubo squares, the $32 T_3$ cancelled and the $\psi = 64$ slice supplied on the declared completion reading — the M2 kill did not fire) and the a_0 uplift by the M3 milestone ([[verification/v517_celestial_wp5e_m3_a0_uplift.py](#)): the log-type correction coupled to the centre count on four scales, monodromy i — the M3 kill did not fire), and the δ_1 chain decided by the consolidated δ_1 module ([[verification/v518_celestial_wp5e_delta1_derived_kill.py](#)): the derived chiral measure fails all three preregistered testers — kill under the derived measure, the declared/derived tension stated), and the measure question decided at probe level by the consolidated δ_{1e}/δ_{1f} module (v520: single-valuedness derived from two constructive sources, the declared reading wins under TP-1..TP-4), and the w_m normalisation derived constructively by the WM module ([[verification/v523_celestial_wp5e_wm_derived.py](#)): the $1/\det_j$ fixed-point weight computed from the Atiyah–Bott mode ledger, the zeta/Quillen determinant and the consistency chain — the WM kill did not fire), leaving the global BCOV quantisation (WP5e proper, narrowed to the global BCOV integral beyond the fibre zero-mode factor) as the single remaining open piece.

Honest success estimate. The probability of *full* success (WP5e: a second continuum route to SEAM.EQUIV.01) remains low — it is constructive holography at the Costello–Li grade. The *partial*-findings expectation is now realised and *exceeded*: WP1–WP4, all five WP5 constructive milestones (both WP5d stages), the WP5e α , β , γ , δ_1 , δ_2 , ε_2 and ε_1 stages and the M1–M3 back-reaction milestones are executed with sharp structural results (verdicts B, B, B, B(ii); WP5a–WP5d passed;

WP5e- α pins the CFT side, WP5e- β the equivariant twistor skeleton, WP5e- γ closes the exchange sub-branch with a certificate, WP5e- δ_2 confirms the collapse full-tensorially and opens the one cubic door, WP5e- ε_2 makes “one level” a theorem and pins its value by the sector counter, WP5e- ε_1 builds the bulk-axion slot with $\tilde{\lambda} = 6$ triply pinned, M1 delivers the closed-form back-reacted Ω_N with forced charge $4 = |\mu_4|$, M2 delivers the twisted KS measure on the declared completion reading (the $32T_3$ cancelled, the $\psi = 64$ slice supplied, no cubic d -channel), M3 delivers the a_0 uplift coupled to the centre count on four scales, and the δ_1 decision (v518) closes the chain with an honest kill under the derived measure, the measure decision (v520) selects the declared reading at probe level, and the w_m derivation (v523) computes the fixed-point weight from three independent sources; *WP5e proper — the global BCOV quantisation, narrowed to the global BCOV integral beyond the fibre zero-mode factor — is the single remaining open piece* — the WP1 critical correction (clock $\neq$ deck; clock² = deck), the WP2 sharpenings (the clock *is* the Picard–Lefschetz/Coxeter monodromy of the seam deformation; a_0 pure scale with the $\tau=i$ shape frozen), the WP3 non-selective alignment (exact λ -arithmetic, honestly typed by the 8/8 look-elsewhere table), the WP4 boundary-limit localisation of the character mismatch (three exactly-located obstructions, one surviving limit structure), the WP5a null-ideal derivation (stabilisation threshold $w = n+1$; $27000 = (h^\vee)^3$ derived; quotient = sector sum = 4124; $SO(16)_1$ contrast), the WP5b singular-vector construction (the deletion operator explicit, case tally 190/57/1, level dial $2(1-k)$, one-block closure typed E_8/μ_4 -specific), the WP5c limit state (the ideal in the GNS kernel, the full 9361-block Gram at rank 4124, the family proved *necessary* by a CCR obstruction), the WP5d- α two-interval index (the $\ln 2$ plateau at 10^{-15} , the μ -chain $16 \rightarrow 4 \rightarrow 1$ measured at both ends), the WP5d- β KLM completion (strong additivity exact on the lattice with the shared boundary point; the bounded-vs-divergent entropic discriminator with the Ising $\frac{1}{4}$ -exponent approach against the diverging $U(1)$ control; the elliptic-nome split ladder with exact orbifold inheritance; Pimsner–Popa $\lambda = \frac{1}{2}$ and $\lambda_{E_4} = \frac{1}{4} = 1/\mu$ with integer attainment — all three KLM ingredients witnessed, Xu’s continuum theorem cited not claimed), the WP5e- α prefactor/level pinning (the $q^{-1/3}$ prefactor as exact μ_4 vacuum bookkeeping; $k = 1$ forced three ways; the glue-integrality route honestly retired), the WP5e- β equivariant ledger (the index bridge $f(m) = \text{ch}_2(T_m)$ exact with the -78 defect; the rigid $32T_3$ twisted residual; the honest a_0 refutation with the three sphere classes filling the three twisted axion slots), the WP5e- γ exchange no-go (the invariant vertex spaces exactly $\dim 2/5$; the product theorem; the certificate $(0, 32, 72)$; naturalness dissolved scale-independently — an honest negative result that converts the pairing question into the binary δ_1 contact-term test), the WP5e- δ_2 full-tensor ledger (the innerness theorem confirming the collapse across arities; the unique symmetric survivor = the $\mathfrak{su}(4)$ d -symbol with $Q_{dd} = \frac{1}{60}(T_3 - P_3/4)$ opening the T_3 channel; the weaker certificate $\psi(A_{\text{fix}}) = 64$ and the exactly solvable relaxed reading with $c_d = 32 \times 60$, the $1920 = |W(D_5)|$ fingerprint honestly retired as look-elsewhere-loaded by the v513 negative certificate), and the WP5e- ε_2 level-from-flux dial (the CPS skeleton exact; the pairing matrix pinned by complete enumeration with fluxes $(64, 60, 64)$ and one quantum per current; the naive “level = total flux” killed by the lockstep test itself; the lockstep theorem with its Z^4-Z falsifier; the sector counter $\#\text{prim}((E_8)_k) = 1 \Leftrightarrow k = 1$), the WP5e- ε_1 axion slot (the equivariant Penrose ledger closing block by block; the $d = 0$ slot surviving the projection; $\tilde{\lambda} = 6$ triply pinned with the wrong bookings 3 and 12 excluded; the GH/ A_3 step re-deriving the v493 family and the Coxeter clock from geometry), the M1 back-reacted Ω_N (the residue form derived with cover factor $4 = |\mu_4|$; the closed twistor family with its CY-compatible clock lift; every $S^3/\mathbb{Z}_4$ period $(2\pi i)^2$ -integral with the forced uniform flux vector and the lens-forced charge $4 = |\mu_4|$; the honest integrality-is-not-the-discriminator fence), the M2 twisted KS measure (the completion-weight identity $w = 4h = -4\text{ch}_2$ with no free scale; every twisted channel the same perfect Okubo square; both γ -stage certificates and the δ_2 $\psi = 64$ slice supplied exactly; the declared completion reading honestly fenced [C], v516) and the M3 a_0 uplift (the GLT kernel $\chi = \log P_4$ with the log coefficient $4 = |\mu_4|$ on four coupled centre-count scales, the period response $1/4$ integrating to the Coxeter monodromy i , v517) and the δ_1 decision (the 16-component Weil completion closing exactly with the E1.5 residual $2.91 \rightarrow \sim 10^{-39}$; the μ_4 multiplier obstruction identified as a *character* of the orbit stabilisers by the exact koboundary test; the cancellation achieved by the twisted

fibre block $f_1 f_3 = G$ — not the three sphere axioms; all three preregistered testers failing under both derived solutions with the honest (N_1, N_2) rebalance rescuing nothing; the declared/derived measure tension stated as the mandatory fence, v518) and the measure decision (the exact invariant subspace = $\text{span}\{e_H, e_{H'}\}$ forcing the unphased E_4 numerator once single-valuedness is derived from F-independence + the Quillen pairing; the consistency circle to v516 closed levelwise; the typed premises TP-1..TP-4 named, v520) — exactly the kind of sharp, falsifiable refinement the contract format is for. Status: CELEST.SEAM.01 [O] (research contract, WP5e proper the single remaining open milestone); CELEST.WP1.01 [E]/[C] (executed); CELEST.WP2.01 [E]/[C] (executed); CELEST.WP3.01 [E]/[C] (executed); CELEST.WP4.01 [E]/[C] (executed); CELEST.WP5A.01 [E]/[C] (executed); CELEST.WP5B.01 [E]/[C] (executed); CELEST.WP5C.01 [E]/[C] (executed); CELEST.WP5D.01 [E]/[C] (executed, α stage); CELEST.WP5DB.01 [E]/[C] (executed, β stage — the continuum uplift stays [O], Xu's theorem cited); CELEST.WP5E.ALPHA.01 [E]/[C] (executed, α stage — the CFT-side prefactor + level pinning); CELEST.WP5E.BETA.01 [E]/[C] (executed, β stage — the equivariant anomaly ledger; the global BCOV quantisation itself stays [O]); CELEST.WP5E.GAMMA.01 [E]/[C] (executed, γ stage — the exchange no-go with certificate; the δ_2 ledger since executed by v511, the twisted contact terms δ_1 since decided by v518 — kill under the derived measure, tension with v516 stated); CELEST.WP5E.DELTA2.01 [E]/[C] (executed, δ_2 stage — the full-tensor ledger, intermediate verdict: collapse confirmed by innerness, the cubic d -symbol channel opens T_3 with the weaker certificate $\psi = 64$; the arity ≥ 4 ledger stays [O]; the $\psi = 64$ burden since discharged by M2/v516 — the cubic-GS question stays dissolved, the δ_1 decision (v518) does not revive it); CELEST.DTERM.NONDERIV.01 [E] (executed, the c_d negative certificate — the $1920 = |W(D_5)|$ fence typed look-elsewhere-loaded and convention-contingent, the convention-stable core $c_d = 32 \times 60$; the fence stays [C], the generation of the 32 stays [O]); CELEST.WP5E.EPS2.01 [E]/[C] (executed, ε_2 stage — the level-from-flux dial, verdict B: “one level” a theorem, the sector counter pins $k = 1$; the CPS dictionary on $\mathbb{P}T/\mathbb{Z}_4$ typed [C], the type-I B-model back-reaction milestones M1–M3 all since executed via v515/v516/v517); CELEST.WP5E.EPS1.01 [E]/[C] (executed, ε_1 stage — the $O(-2)$ bulk-axion slot construction with $\tilde{\lambda} = 6$ triply pinned and the first GH/ A_3 back-reaction step; the M1–M3 milestones it preregistered are all since executed; the BCOV-integral derivation stays [O]); CELEST.WP5E.M1.01 [E]/[C] (executed, back-reaction milestone M1, SUCCESS on the preregistered criterion — the closed-form Ω_N with $(2\pi i)^2$ -integral lockstep periods and the lens-forced source charge $4 = |\mu_4|$; honest fence: integrality alone is not the discriminator; M2–M3 since executed via v516/v517); CELEST.WP5E.M2.01 [E]/[C] (executed, back-reaction milestone M2, SUCCESS on the preregistered criterion *on the declared completion measure* — the completion-weight identity $w = 4h = -4\text{ch}_2$ with no free scale, every twisted channel the same perfect Okubo square $36\langle x, x \rangle^2 / \det_j$, the $32 T_3$ cancelled, both γ -stage certificates and the $\psi = 64$ slice supplied exactly, no cubic d -channel needed; the completion reading stays [C]-declared within that module — the δ_1 chain since decided by v518, and the completion reading since *derived at probe level* from two independent constructive sources (F-independence + Quillen pairing) under the typed premises TP-1..TP-4 by v520; the residual [O] is the constructive w_m derivation); CELEST.WP5E.M3.01 [E]/[C] (executed, back-reaction milestone M3, SUCCESS on the preregistered criterion — the a_0 uplift on the GLT kernel $\chi = \log P_4$ with the log coefficient $4 = |\mu_4|$ coupled to the centre count on four scales, the period response $1/4$ integrating to the Coxeter monodromy i ; the GLT dictionary stays [C], the full nonlinear Kähler potential stays [O]); CELEST.WP5E.DELTA1.01 [E]/[C] (executed, δ_1 stage — DECIDED: kill under the derived measure; the 16-component Weil completion closes exactly, the μ_4 multiplier obstruction is a character (koboundary defects $(1, 1, 1)$, $\lambda(\gamma) = i^{2B+C/4}$ on $\Gamma_1(4)$), the cancellation is the twisted fibre block $f_1 f_3 = G$, and all three preregistered testers fail under both derived solutions; the $f_1 f_3 = \text{KS-weight identification}$ and the kernel-family convention stay [C]; the declared-vs-derived BCOV measure question — the stated v516 tension — was the named [O] and is since decided at probe level by v520 for the declared reading, sharpening this kill; the residual [O] is since narrowed by v523 to the global BCOV integral beyond the fibre zero-mode factor); CELEST.WP5E.MEASURE.01 [E]/[C] (executed, the measure decision — ERFOLG-A on the preregistered decision layer: single-valuedness derived from two independent constructive sources

(F-independence + the Quillen pairing), the invariant subspace exactly $\text{span}\{e_H, e_{H'}\} = E_4$, the consistency circle to v516 closed levelwise, the $\mathbb{Z}_2/\text{EH}$ anchor and the column canonicalisation both selecting the declared reading; typed premises TP-1..TP-4 stay [C]; the constructive BCOV derivation of the w_m normalisation itself — the named residual [O] — is since executed by v523); CELEST.WP5E.WM.01 [E]/[C] (executed, the constructive w_m derivation — verdict ERFOLG per the frozen preregistration: the $1/\det_j$ fixed-point weight COMPUTED from three independent sources (the equivariant mode ledger with Abel value $(1/2, 1/4, 1/2)$; the zeta/Quillen determinant $4\sin^2(\pi j/4)$ with the unique real positive section; the δ_{1f} block constant term $1/\det_b$), the v516 chain reproduced number by number ($w = 4h$, $c = 4 = |\mu_4|$, squares $(18, 9, 18)$, total $45\langle x, x \rangle^2$, $\psi \pm 64$), all negative controls behaving (the $\mathbb{Z}_2/\text{EH}$ anchor at $c = 2 = |\mathbb{Z}_2|$, the $SO(16)$ kill, the $\text{diag}(i, i)$ break, the $k = 3, 5$ wandering); typed premises TP-REG/TP-Q/TP-NUM/TP-CH stay [C]; the global BCOV integral beyond the fibre zero-mode factor stays [O]); SEAM.EQUIV.01 untouched, stays [O].

Research Contract — WOIT.OS.TWISTOR.01 (the Osterwalder–Schrader twistor bridge)

Typing. This is the actual bridge from compiler to physics — everything in CELEST.SEAM.01 is preparation for it. It is a research contract [O] (ledger row WOIT.OS.TWISTOR.01: Open, research contract), *not* a claim; nothing below asserts that the construction exists. The external programme it engages — Woit’s Euclidean Twistor Unification (arXiv:2104.05099) — is a *named reference frame* for the shape of the target, never a confirmation in either direction (the Woit-bridge paragraph in tfpt_3 states exactly what is missing).

Input. $X_+ = (\mathbb{PT}/\Gamma)_+$ (the positive-time half of the quotiented projective twistor space; $\Gamma = \Gamma_{\text{ALE}} \cong \mathbb{Z}_4$, normalised by the clock — the definition block of CELEST.SEAM.01); $\mathcal{A}_{\text{hol}}$, the *interacting* open+closed twistorial algebra ($\text{SDYM}(E_8)$ open strings + BCOV closed strings — the WP5e-proper object, whose free/equivariant skeleton is pinned by v492–v517); ρ , the order-4 clock $\text{diag}(i, 1)$; Θ , the anti-linear real structure induced by the seam reflection; and $\mu_{\text{BCOV}+\text{SDYM}}$, the interacting functional (the global BCOV+SDYM quantisation on $\mathbb{PT}/\Gamma$). *Precision (i), from the α stage:* “induced by the seam reflection” means the seam-circle *reflection*, not the deck/covering involution — the deck (whose free-ness is topology, v510) furnishes the $(-1)^F$ Kramers class $\Theta_t^2 = (-1)^F$, not the OS Θ ; the seam-circle reflection furnishes $\Theta_{\text{Fock}}^2 = +1$ ([verification/v519_woit_theta_rp_free.py]). *Precision (ii):* on the RP side the μ_4 marks sit at the *bond midpoints* of the 16-Majorana seam circle (the half-offset/NS-natural placement, 4 sites per mark quadrant) — the cut through the sites fails reflection positivity exactly ([verification/v519_woit_theta_rp_free.py]). *Precision (iii), from the β_1 stage:* “gauge-invariant subalgebra” at the free level means the GSO/fermion-parity $\mathbb{Z}_2$ — the only θ -compatible part of the μ_4 tower (the free-fermion shadow of the E_8 glue: $(E_8)_1 = \text{even NS of } SO(16)_1 + \text{R spinor}$); the clock is the euclidean rotation itself (*timelike*: all 16 seam mirror axes invert it, none commutes), and its equivariant statement moves to β_2 — OS quotient first, then the clock as reconstructed transfer operator ([verification/v522_woit_beta1_gso_gauge.py]).

Target theorem. $\Theta^2 = 1$ and $\Theta\rho\Theta = \rho^{-1}$; the gauge-invariant algebra is reflection-positive; and the Osterwalder–Schrader quotient produces $(\mathcal{H}, \Omega, U(\mathcal{P}_+^\dagger), \mathcal{A}_{\text{Mink}})$ with: a positive Hilbert metric, positive energy, local causality, *one chiral fermion generation without mirror doubling*, the TFPT charge lattice, an internal $SU(3) \times SU(2) \times U(1)$ action, and the $(E_8)_1$ seam net as an *actual* boundary net (not only as a character).

Kill tests (pre-registered, all seven live). (1) Θ is incompatible with the clock (no anti-linear Θ with $\Theta^2 = 1$ and $\Theta\rho\Theta = \rho^{-1}$ exists on $\mathcal{A}_{\text{hol}}$). (2) Reflection positivity fails after gauge fixing (the gauge-invariant subalgebra of $\mu_{\text{BCOV}+\text{SDYM}}$ is not RP). *Named toy threat (no status change):* on the first genuinely interacting 16-Majorana Fidkowski–Kitaev seam toy (SEAM.INT.FKTOY.01, [verification/v529_seam_interacting_toy_fk.py]) Kill-Test 1 does *not* fire (Θ exists with $\Theta^2 = +1$ and clock-tower inversion), but Kill-Test 2 *fires at toy level* — RP breaks for every $g > 0$ by

an interference mechanism, with the *straddle law* (RP fails exactly on quartet-straddled cuts, stays PD on quartet-avoiding ones; 24/24) as a concrete hurdle for the interacting stages (γ and beyond; the free/quotient-level β_3 is since executed, v565). Fence: one toy / one interaction class / [C] flat-band parent — this narrows the route (every $\mathcal{A}_{\text{hol}}$ must protect RP against exactly this mechanism), it does not close it. *The hurdle, executed as a selector (SEAM.STRADDLE.CONE.01, [verification/v534_seam_straddle_cone.py]; no status change)*: on the equivariant interacting mark family (independent couplings on the four mark quartets, both signs, every admissible cut — including the $\pi/2$ straddled clock axes the 24-entry law had not covered) reflection positivity survives for *exactly one* member $\times$ sign: $\delta = \pi/2$ with positive coupling, PD on all four cuts over $g \in \{1/32..8\}$ with the minimum eigenvalue lifted *away* from the RP boundary; every asymmetric member dies. The straddle law refines (asymmetric straddling kills, symmetric straddling protects), the literal leading-order cone formalization is dead (the protection is nonperturbative), and the threat paragraph gains its complement: a passing interaction class *exists* and its unique survivor is the alignment bit — the same member that carries the v528 twist-class order parameter. Toy-level evidence for a dynamical origin of the bit, not a derivation; the kill tests stay formally live on $\mathcal{A}_{\text{hol}}$. (3) The reconstruction produces a vector-like mirror generation (chirality is lost in the OS quotient). (4) The Penrose transform reaches only free / self-dual states (the interacting extension does not exist at the Costello–Li grade). (5) The reconstructed net has the right character but the wrong OPE / fails net equivalence (character-level agreement without operator-algebraic agreement). (6) The internal $SU(2)$ remains a spacetime factor (no genuine internal weak factor separates from the Euclidean rotation group). (7) The four μ_4 marks are not incidence-compatibly extendable over spacetime (the mark data of the seam cannot be propagated through the incidence relation).

The α stage — executed (*done*; the free/equivariant milestone WOIT.THETA.FREE.01, [verification/v519_woit_theta_rp_free.py]). The real structure *exists*, and free reflection positivity picks the *same* family. Exactly *two* families of anti-linear structures on $\mathbb{C}^2$ normalise the clock: family D ($z \mapsto \mu\bar{z}$, $|\mu| = 1$) satisfies $\Theta\rho\Theta = \rho^{-1}$ *exactly* with $\Theta^2 = +1$ (-1 impossible: $M\bar{M} = \text{diag}(|\mu|^2, 1)$ — Kramers-free), inverts the deck and reflects the seam circle (2 cut points); family A ($z \mapsto \mu/\bar{z}$) centralises the clock projectively and *never* inverts it — it *defines* the euclidean section (Woit’s ρ_{tw} side: $\rho_{\text{tw}}^2 = -1$, no real points, replicated exactly on $\mathbb{C}^4$), while family D is the OS conjugation (σ_{std} , real points $\mathbb{RP}^3$). Mark compatibility pins $\mu \in \mu_4$ (a 4-element torsor, 2 clock orbits); the $\mathbb{Z}_8$ spin plane has no phase leaks; in exact $\text{Cl}(16)$, $\Theta_{\text{Fock}} = U_r \circ K$ has $\Theta_{\text{Fock}}^2 = 2^7 > 0$ (normalised $+1$), inverts the Fock clock tower ($V \mapsto 4096 V^{-1}$) and normalises the deck, while the deck-induced candidate has $\Theta_t^2 = (-1)^F$ — the v510 split/nonsplit dichotomy is the $\Theta^2 = +1$ vs $(-1)^F$ dichotomy. Free RP holds on the bond cut with *no* degree truncation (one-particle Gram (8,0,0) PD, even $\deg \leq 2$ sector (29,0,0), complete $N = 8$ half-sided algebra PD; twist $\eta = +i$ forced), the site cut and the clock-centralising family fail RP exactly, and the anti-chiral state flips the odd sector to negative definite (the free shadow of kill test 3). **Kill test (1) therefore does *not* fire at the free/equivariant level — and stays formally *live* on the interacting algebra $\mathcal{A}_{\text{hol}}$; kill tests (2)–(7) are untouched. No marker moves: the contract stays [O].**

The β_1 stage — executed (*done*, verdict UNDECIDED per the frozen preregistration; the typed milestone WOIT.BETA1.GSO.01, [verification/v522_woit_beta1_gso_gauge.py]). “The clock is time-like, GSO is the gauge datum.” The preregistered clock-tower reading of “gauge” (the $\mu_4/\Gamma_{\text{ALE}}$ equivariance averaged *before* the OS quotient) is structurally the *wrong* operationalisation: the one-step clock insertion $T_1 = \langle \theta(e_a), \alpha_S(e_b) \rangle$ violates Hermiticity *exactly* (witness entry $-i/(8\sin(5\pi/16))$ where the conjugate transpose demands $+i/(8\sin(5\pi/16))$); the pairing is complex-symmetric: 745 matching, 96 anti-matching, 0 violations — “OS-symmetric” is strictly weaker than Hermitian, so “positivity after gauge fixing” is not even well-posed for the clock reading). The mechanism is typed exactly: *all* 16 dihedral reflection axes of the NS seam circle invert the clock lift ($RSR^{-1} = S^{-1}$), none commutes — the μ_4 clock is Woit’s euclidean rotation, and the θ -compatible (gaugeable) part of its $\mathbb{Z}_8$ Fock tower is exactly the 2-torsion $\{1, (-1)^F\}$ = the GSO/fermion-parity $\mathbb{Z}_2$ (Hermiticity census over all powers; α^4 = the 2π rotation of the seam = fermion parity). Under that corrected typing **gauge-fixed RP holds exactly**: the

parity-projected bond-cut Gram is $(29, 0, 0)$ PD at $N = 16$ ($\deg \leq 2$, min eigenvalue 1.78×10^{-6} at 40 digits) and $(8, 0, 0)$ PD on the complete $N = 8$ half algebra (no degree truncation); the site-cut defect *survives* gauge fixing $((7, 9, 6) —$ the bond placement of precision (ii) is gauge-invariant information); family A stays indefinite $((17, 12, 0)$ vs family D's $(29, 0, 0)$: the α -stage triangulation survives); the Ramond (wrap +1) “true $\mathbb{Z}_4$ ” lift fails state invariance (the $\text{NS}/\mathbb{Z}_8$ tower is forced); and the Kramers class stays trivial on the invariant sector ($\Theta_t^2 = \text{id}$ on $\mathcal{A}^G$, Θ restricts with $\Theta^2 = +1$). **Kill test (2)’s free shadow does *not* fire** (no legitimate Hermitian gauge-fixed form has a negative direction), and kill test (1) stays discharged also gauge-invariantly; both stay formally live on $\mathcal{A}_{\text{hol}}$. Transparency (documented in the module): the first frozen run scored 6/13 — the clock-reading checks failed exactly as the mechanism predicts — and the preregistered deg-2 clock-invariant dimension 30 was a combinatorial error (correct orbit census $28 + 4 = 32$); the verdict follows the frozen logic (Hermiticity failure $\Rightarrow$ structural-failure branch $\Rightarrow$ UNDECIDED). β_1 *proper* (the clock-equivariant statement) is re-routed through β_2 . No marker moves: the contract stays [O].

The β_2 stage — executed (*done*, verdict SUCCESS per the frozen preregistration, [C]-typed per contract precision (iii); the milestone WOIT.BETA2.OS.01, [verification/v524_woit_beta2_os_quotient.py]). “The OS quotient of the free system made explicit.” $\mathcal{H}_{\text{phys}} = \mathcal{A}_+/\mathcal{N}$ is *explicit and nondegenerate* at both levels: at $N = 16$ ($\deg \leq 2$) the 37×37 bond-cut OS Gram is exactly Hermitian and parity-block-diagonal with inertia $(37, 0, 0)$ PD (min eigenvalue 1.7801×10^{-6} at 40 digits) and null space $\{0\}$, $\dim 37 = 29 \oplus 8$; at $N = 8$ the *complete* half algebra gives $(16, 0, 0)$ PD with $\dim 16 = 8 \oplus 8 = 4^2$ — compact euclidean time reconstructs a *thermal* (KMS) representation, with the exact certificate $\sin^2(3\pi/8) - \sin(\pi/8)\sin(5\pi/8) = 1/2$. The euclidean rotation becomes a Klein–Landau *local symmetric semigroup*: $\tau_k = \omega(\theta(a)\alpha_k(b))$ is exactly Hermitian on every shrinking domain D_k ($k = 1..7$, dims $29, 22, 16, 11, 7, 4, 2$), the chain identities are exact and the vacuum is fixed; the positivity pattern *is* the α -stage site/bond dichotomy (even steps PSD via the exact square identity $T(2j) = A_j^* A_j$, odd steps indefinite with exactly zero one-particle diagonal — the one-step transfer is *not* positive: the free chirality datum, kill-test-3 shadow sharpened). **The clock is implemented on $\mathcal{H}_{\text{phys}}$ as the positive self-adjoint transfer step $T^{N/4}$** $((11, 0, 0)$ PD, min 6.7×10^{-5} ; compressed spectrum $[0.0195, 1.6414]$ with vacuum eigenvalue exactly 1; spectral projections certified at $\sim 10^{-40}$ — exactly the calculus the non-Hermitian pre-quotient average of v522 could not have), and at $N = 8$ the compressed clock spectrum is *exactly* $\{1, \sqrt{2} - 1\} = \{1, 1/\delta_S\}$ in both parity sectors (the silver axes of the v519 μ_4 torsor returning as the clock eigenvalue); the reconstructed rotation group $U(s) = e^{isH}$ is unitary with the group law — per precision (iii) the [C]-operationalisation of the contract phrase “the clock acting unitarily”. The v522 non-Hermiticity is *resolved*: the census $(745, 96, 0)$ is reproduced, every anti-matching entry is a wrap overlap and every overlap-free entry matches — the pre-quotient failure was exactly the domain/wrap artifact. The pre-declared KMS deviation arrives with exactly the declared witnesses: *no contraction* on the compact circle (edge-mode expansion $C(1)/C(3) = 1 + \sqrt{2} = \delta_S$ exactly; $\det(G - \tau_4) < 0$) — a typed KMS finding, not a β_2 failure (the contract demands no contraction). GSO/ Θ : the grading survives; θ_{cut} and the deck candidate do *not* act on $\mathcal{H}_{\text{phys}}$ (support censuses); the *perpendicular* torsor mirror descends anti-unitarily ($\eta_{\perp} \in \{\pm i\}$) with $\Theta_{\text{phys}}^2 = +1$ on *every* sector — the quotient is Kramers-free — and $\theta_{\text{cut}} \circ \theta_{\perp} = \alpha_{N/2}$ exactly (the torsor closes onto the deck). Controls: the site cut is indefinite $((3, 3, 1)/(2, 2, 4) —$ the contract kill branch fires for the site placement), family A admits no quotient $((4, 4, 0))$, the anti-chiral state flips to $(8, 8, 0)$ (quotient existence itself selects the chiral orientation), the wrong semigroup direction is non-Hermitian exactly off-domain (all mismatches wrap-overlap). **Kill tests (1) and (2) are strengthened and the (3) shadow is sharpened; none fires** — all seven stay formally live on $\mathcal{A}_{\text{hol}}$. Transparency (documented in the module): run 1 scored 18/20 — a sympy simplify failure on a *true* $\pi/16$ product formula ($\sin(7\pi/16) - \sin(5\pi/16) = 2\cos(3\pi/8)\sin(\pi/16)$), fixed by a Laurent-polynomial certificate; no criterion, tolerance or expectation changed. No marker moves: the contract stays [O]; β_3 is executed next (below).

The β_3 stage, executed (WOIT.BETA3.DUALITY.01, [verification/v565_woit_beta3_pt_duality.py]; verdict SUCCESS per the frozen preregistration; no status change). The $\mathbb{PT} \leftrightarrow \mathbb{PT}^*$ duality is *constructed as a compatibility class* and typed against σ_{std} , with the contract kill branch decided *empty by algebra*. **(i) The class:** a duality is a nondegenerate Hermitian form Φ on $\mathbb{C}^4$ ($\sigma(Z) = Z^\dagger \Phi$, mapping $\mathbb{PT} \rightarrow \mathbb{PT}^*$); clock compatibility has the entrywise residual $(\bar{\lambda}_j \lambda_k - 1) \Phi_{jk}$ and forces the four cross-eigenspace entries to zero exactly: two 2×2 Hermitian blocks, $16 \rightarrow 8$ real parameters (the deck adds nothing; the wrong clock lift $\text{diag}(i, i, 1, 1)$ breaks the split). **(ii) The dichotomy and the forced (2, 2):** pairing against Woit's Euclidean structure ρ_{tw} via $M_{\text{tw}}^\dagger \Phi M_{\text{tw}} = \varepsilon \bar{\Phi}$ splits the class exactly in two. The *Euclidean* branch ($\varepsilon = +1$; $\sigma_{\text{std}} = \mathbf{1}$ lives here) has seam-line norm $a(|z|^2 + 1)$: no null point, no OS cut, ever — exactly the section-defining role Woit assigns it. The *Minkowski* branch ($\varepsilon = -1$) has seam-line norm $a(|z|^2 - 1)$: **the null locus meets the seam line exactly in the seam circle** — the OS cut is *forced*, not imposed — and the second block is exactly minus the conjugate of the first (charpoly mirrored), so the **total signature is (2, 2) for every nondegenerate member**: Woit's Minkowski signature is a theorem of clock + ρ_{tw} pairing, not an input. **(iii) The kill is empty:** every compatible duality inverts clock *and* deck exactly (family D); a clock-centralising (family A) duality would need $\rho_4^2 = \text{scalar}$, but $\rho_4^2 = \text{diag}(-1, 1, -1, 1)$ — no solution; and family A independently has no quotient ((4, 4, 0), the v524 control). **(iv) The identification:** every seam-preserving Minkowski member induces *one* projective conjugation $z \rightarrow -\bar{z}$ ($\mu = -1$, fixed points $\pm i$), perpendicular to σ_{std} 's $z \rightarrow +\bar{z}$; under the declared RP-side placement (v519 precision (ii); exact rational- π arithmetic, offsets $\mp \pi/2$) that member is $\theta_\perp$ — the mirror whose quotient descent v524 certified: Θ_{phys} is the shadow of the $\mathbb{PT} \leftrightarrow \mathbb{PT}^*$ duality, anti-unitary, Kramers-free ($\Theta^2 = +1$ per sector), time-reversing the semigroup, closing with the σ_{std} -type cut onto the deck ($\theta_{\text{cut}} \circ \theta_\perp = \alpha_{N/2}$; upstairs: the clock conjugation exchanges $\mu = +1 \leftrightarrow \mu = -1$). *Role separation complete:* $\sigma_{\text{std}} \sim$ the OS cut (the metric), the Φ -twisted Minkowski duality $\sim$ the descending mirror. Free/quotient level only; $\mathcal{A}_{\text{hol}}$ untouched; all seven kill tests stay live there; the declared conventions (seam line, seam preservation, placement, torsor relabeling) are cited, not invented. No marker moves: the contract stays **[O]**; γ is the next milestone.

Thermal follow-on (no status change). The free OS quotient of β_2 is KMS, and its temperature normalisation closes onto the Nariai/BH chain as the *third leg* of c_3 (SEAM.THERMAL.KMS.01, [verification/v526_seam_thermal_kms_nariai_bridge.py]: β measured = N at both levels; $\beta_{\text{angle}} = 2\pi = 1/(4c_3)$, $T_{\text{seam}} = 4c_3$; modular Hamiltonian closed; class-1 anti-numerology selection; non-compact $\Rightarrow T = 0$). The silver clock eigenvalue is explained and demystified in parallel (SEAM.CLOCK.SILVER.01, [verification/v527_seam_clock_silver_spectrum.py]: structure SUCCESS, connection KILL). No marker moves.

The β/γ roadmap (named milestones, each with success and kill criteria). (β_1) Θ on the gauge-invariant subalgebra + gauge-fixed RP on the equivariant SDYM(E_8) sector — success: the v519 OS form extends to the gauge-invariant subalgebra with positivity after gauge fixing; kill: kill test (2) fires (RP fails after gauge fixing). *Executed via v522 (above): UNDECIDED under the frozen preregistration — the clock reading is time-like (wrong operationalisation), the gauge datum is the GSO $\mathbb{Z}_2$, gauge-fixed RP holds under that typing, neither kill fires; the clock-equivariant statement moves to β_2 .* (β_2) The OS quotient of the free system made explicit — success: $(\mathcal{H}, \Omega)$ constructed from the v519 bond-cut form with the clock acting unitarily; kill: the quotient degenerates (zero or indefinite completion). *Executed via v524 (above): SUCCESS, [C]-typed per precision (iii) — $\mathcal{H}_{\text{phys}}$ explicit and PD at both levels, the clock a positive transfer step with spectral calculus (exact $N = 8$ spectrum $\{1, \sqrt{2} - 1\}$) and a reconstructed unitary rotation group; the pre-declared KMS deviation carries exactly the silver witnesses; neither kill fires; β_3 is next.* (β_3) The $\mathbb{PT} \leftrightarrow \mathbb{PT}^*$ duality typed against σ_{std} — success: Woit's Minkowski conjugation ($\mathbb{PT} \rightarrow \mathbb{PT}^*$) identified with the family-D conjugation on the quotient; kill: the duality forces a clock-centralising (family A) structure instead. *Executed via v565 (above): SUCCESS — the kill branch is empty by algebra (every clock-compatible duality is family D; family A would need $\rho_4^2 = \text{scalar}$), the OS cut and the (2, 2) signature are forced on the Minkowski branch, and the unique induced seam member is $\theta_\perp$, the v524 Θ_{phys}*

carrier. (γ) The chirality theorem + the mark incidence — success: the odd-sector definiteness flip of v519 upgraded to “one chiral generation without mirrors” on the reconstructed net, and the four μ_4 marks extended incidence-compatibly; kill: kill tests (3)/(6)/(7) fire. *The FREE slice is executed via [verification/v608_gamma_free_slice.py] (9/9, GAMMA-FREE-SLICE-LANDED): on the explicit v524 net the mirror state’s odd sector is strictly negative definite (sector-exact flip — every mirror fermion vector has negative OS norm, so no mirror mode survives reconstruction: kill test (3) cannot fire at the free level), the chiral generation is multiplicity-free (clock spectrum exactly $\{1, \sqrt{2} - 1\}$), and the mark transport exists (τ_2 is the mark- $A \rightarrow B$ transport, Hermitian PD: kill test (7) does not fire at the free level). γ itself stays open on $\mathcal{A}_{\text{hol}}$ (kill tests (3)/(6)/(7) formally live there); kill test (6) is untouched by the free slice. The INTERACTING toy slice is since executed via [verification/v615_gamma_toy_interacting.py] (11/11, GAMMA-TOY-LANDED): on the RP-surviving interaction class (the alignment bit $\delta = \pi/2$, $s = +$, v534) the mirror state’s odd sector is strictly negative definite on every admissible cut and every coupling $g \in \{1/32..8\}$ (bounded away from zero — kill test (3) cannot fire on the surviving class), the mark transport α_4 is Hermitian PD at every g (kill test (7) does not fire at the interacting toy level), the odd transfer spectrum stays multiplicity-free (honest finding: strong-coupling near-doubling at $g = 8$ without closing), and the RP-dead member has no gamma home — the dynamical bit selection and the chirality data co-locate on the same member. Fence: one toy, one interaction class; γ proper stays open on $\mathcal{A}_{\text{hol}}$. Kill test (6) has since received its first (kinematic) slice via [verification/v616_su2_internal_kinematic.py] (17/17, SU2-KINEMATIC-SEPARATED): on $\mathbb{C}^4 = \mathbb{H}^2$ the internal $su(2)$ (the right quaternion action, with $\rho_{\text{tw}} = \text{right-}j$ verbatim) preserves every twistor fiber, commutes exactly with the spacetime quaternion action and intersects it in $\{0\}$ — the internal factor separates kinematically, the clock factorizes exactly as $\text{left-}e^{i\pi/4} \circ \text{right-}e^{i\pi/4}$, the ALE deck is purely spacetime-side, and the complex choice breaks the internal $SU(2)$ to $U(1)$ with the v519 mark torsor as its μ_4 ; the DYNAMICAL half of kill test (6) (a genuine gauge action on the reconstructed net) stays open and formally live on $\mathcal{A}_{\text{hol}}$.*

Scope fence (explicit non-claims of the celestial/twistor branch). Self-dual is *not* full 4D physics. Until and unless this contract closes, the celestial/twistor branch claims *none* of the following: both helicities (the anti-self-dual completion), generic scattering amplitudes, local matter fields, full Einstein dynamics (beyond the self-dual sector), electroweak symmetry breaking, confinement. The classical field-equation route (entanglement equilibrium, v358/v359) is a separate, non-competing statement about the full equation; SEAM.EQUIV.01 and its route split (SEAM.EQUIV.MMST.01/SEAM.EQUIV.TWISTOR.01) are stated in their own rows and are not moved by anything here. Status: WOIT.OS.TWISTOR.01 [O] (research contract; the α stage is executed — WOIT.THETA.FREE.01 [E], v519 — with kill test (1) discharged at the free level only; the β_1 stage is executed — WOIT.BETA1.GSO.01, v522, verdict UNDECIDED per the frozen preregistration, with kill test (2)’s free shadow not firing, the gauge datum typed as the GSO $\mathbb{Z}_2$ and contract precision (iii) added; the β_2 stage is executed — WOIT.BETA2.OS.01, v524, verdict SUCCESS, [C]-typed per precision (iii), with the OS quotient explicit and PD at both levels, the clock a positive transfer step with spectral calculus and a reconstructed unitary rotation group, and the pre-declared KMS deviation carrying exactly the silver witnesses; the β_3 stage is executed — WOIT.BETA3.DUALITY.01, v565, verdict SUCCESS, with the kill branch empty by algebra, the OS cut and the (2, 2) signature forced on the Minkowski branch of the duality class, and the unique induced seam member identified with the v524 Θ_{phys} carrier; no marker move).

Certifiability and recommended order

Building block	Machine	Hardness
W_{wall} enumeration ($C_U^{(1)}$)	Lean / Python / Wolfram	easy (<i>done</i>)
D_4 fixed-locus polynomial ($C_U^{(2)}$)	Sage / Mathematica	medium
selector uniqueness ($C_U^{(3)}$)	interval arithmetic	medium
SNF, determinants, spectra	Lean	easy–medium
principal-symbol ellipticity (G1)	xAct / Cadabra / Lean	medium
gap inequality (G5)	Lean	easy (<i>done</i>)
operator-norm bound (G5)	analysis + numerics	hard
projective limit (G6)	not primarily machine	very hard
Ward identities (G7)	BV/BRST formal + checks	hard

Recommended order.

Selector-triangle pairings $\rightarrow v_{\text{geo}}$ scale anchor $\rightarrow G_{\text{net}}$ inclusion theorem

The flavor interface (historically U_{wall}) is now reduced to the *selector-triangle pairings* (the dual normal pair (d, n) forcing R columnwise, v136/v139); it is finite, falsifiable and the cheapest visible upgrade. The v_{geo} scale anchor is the one dimensionful input (same nature as $1/G$). G_{net} — the index-4 seam-net inclusion certifying the metric sector — is more fundamental but a *programme*, not a single proof: heavy machinery, with the projective limit the genuine summit. F_{transfer} (source $\rightarrow$ pole / relic / cosmology) is the remaining downstream interface.

F_{transfer} **is a typed functor, not a bag of open topics.** It is standard physics fed TFPT *source* data, $F_{\text{transfer}} = F_{\text{observable}} \circ F_{\text{threshold}} \circ F_{\text{RG}}$, with exactly four interfaces:

Interface	Map	Status	Source data $\rightarrow$ observable
F_{pole}	source $\rightarrow$ pole masses	[E] / [C]	Koide 53/54 = $a^T(R+Q)\mathbf{1}/(2\mathbf{1}^T R a)$ is an exact operator factor [E] (the missing Sheet-Diamond corner); the pole-transfer interpretation is [C] ([verification/v183_koide_f_corner_transfer.py])
$F_{\text{Boltzmann}}$	CP source $\rightarrow \eta_B$	[C]	$\tilde{m}_1=m_3/A_\Lambda$ anchors the washout, but $M_1=M_R\varphi_0^4$ only relocates the free input (M_R is the seesaw scale, not a compiler power), so η_B stays a <i>sharper scenario</i> , not a closure ([verification/v184_etaB_anchored_boltzmann.py])
F_{relic}	$(f_a, m_a, \theta_i, N_{\text{DW}}) \rightarrow \Omega_a$	[C]	the determinant-line axion can be all of DM, but only on the $\theta_i \approx 170^\circ$ hilltop where the relic is exponentially sensitive — a viable scenario ([verification/v185_axion_relic_solver.py])
F_{QCD}	$(\alpha_s, b_3, \Lambda_{\text{QCD}}, \text{lattice}) \rightarrow m_p/m_e$	[O]	a QCD matching contract ($b_3=-7$ a carrier output), <i>not</i> a compiler number — the rejected 1836 near-formula is the discipline working

A machine guard enforces the typing: the four families must always carry a [C]/[O] marker and must never be promoted to a primitive [E] compiler prediction ([verification/v187_ftransfer_laws.py]). Exact *algebraic* sub-parts (the Koide factor, $b_3=-7$) may be [E]; the *physical* prediction never is.

The functor contract CONTRACT.F.01, four axioms ([verification/v213_ftransfer_functor.py]). Beyond the typing guard, the functor itself satisfies four checkable structural axioms, each instance a *discrete* compiler kernel [E] composed with an *external* continuous solver [C]: (1) **μ_4 -deck equivariance** — the discrete kernels are carrier/deck invariants $\{|\mu_4|, N_{\text{fam}}, g_{\text{car}}, A_\Lambda\}$, and the Koide multiplier $\lambda_2=(2/3)^6=64/729$ is the μ_4 -deck transfer eigenvalue (v82/v54); (2) **Plücker preservation** — the dimensionless readouts are Plücker readings ($53=a^T(R+Q)\mathbf{1}$, $\text{Pl}(F)=(1, 22, 30)$; $\|\text{Pl}(K)\|_1=11$); (3) **positivity/stochasticity** — $\text{spec } T=\{1, (2/3)^6, (1/3)^6\}$ is positive (RP/OS),

$\kappa_f \in (0, 1)$; **(4) external modules explicit** — the continuous content (pole transfer, the flavored Boltzmann network, the finite- T $\chi(T)$ ODE, the lattice C_N) is named and external, never a compiler number. So `CONTRACT.F.01` is the third research contract alongside (U_{wall}) and (G_{metric}): a typed functor, *not* a bag of open topics — but a consolidation, *not* a closure. [2026-08-05 dated note — the jet route closes, the fibered refinement stands (`[verification/v777_ftransfer_clock_jets.py]`): the executed external-clock contract `FTRANSFER.CLOCKS.01` (below) fires its preregistered $K1$ kill sympy-exactly — the transported Schwarzians are $\{-\Delta^2/2, -\Delta^2/2, 0\}$, $\Delta = 6 \log(3/2)$, so no common continuous clock exists for the four jets. The surviving architecture is measured: F_{transfer} is a fibered functor — one shared discrete PGL_2 base (the anchor cross-ratio of all four channels is $4/3$ exactly, symbolically and through the frozen g_{*s} table at $\text{dev } 6.5 \times 10^{-4}$; the deck generator is the unit parabolic σ -translation, $\text{tr}^2 - 4 \det = 0$ — the `v632` lattice-explicit discrete half) with channel-specific projective connections whose obstruction is the constant seam cocycle, values exactly $\{0, \pm \Delta^2/2\}$ (antisymmetry and the cocycle identity exact; $d/d\sigma$ identically zero); the classification matches the predeclared marker — thermal/proper-time clocks $\{\tau, \beta\}$ on the $-\Delta^2/2$ coset, the RG clock $\{\log \mu\}$ on the 0 coset, the relic channel degenerate and honestly unclassified. The functor axioms above stand; the jet/continuous-clock unification demand is replaced by the fibered form, with the Schwarzian differences stored as the obstruction cocycle, not fitted away. The one named proof-grade next step — derive the coset assignment from KMS /modular clock typing — is executed one round later and is a theorem (`[verification/v792_ftransfer_kms_coset.py]`, 18 checks, $KMS\text{-}COSET\text{-}THEOREM$, registered as `FTRANSFER.KMSCOSET.01`): $S = -(\ln \lambda)^2/2$ with λ the PGL_2 multiplier of the flow's time-1 holonomy per clock e -fold; the classifier uses no Schwarzian and no derivatives (non-circularity gated on its own source), the blind assignment classifies the four deployed charts exactly as the fibered functor carries them, the unblinding is sympy-exact, and the controls fire (fake KMS caught; affine invariance; the `v578` Möbius clock as the positive case) — the fiber classification is derived, not case-inspected. The four observables stay $[C]/[O]$; no marker move.]

A preregistered cross-ratio check for the four solvers — designed, and since executed on its discrete half. The design is machine-checked in `[verification/v566_parabolic_anchor_selfcode.py]`, the discrete-half execution in `[verification/v571_cr_discrete_corners.py]`. The disc -7 norm line $\alpha_{t+1} = \alpha_t + 2$ (`v533`) equips the transfer path with a projective invariant: $CR(\alpha_{-2}, \alpha_{-1}; \alpha_0, \alpha_1) = \frac{4}{3}$ exactly — and, honesty first, $\frac{4}{3}$ is the *universal* cross-ratio of any arithmetic progression, so the test content is not the number but the *discipline*: if the four solvers (J, K, C, F) export canonical one-dimensional projective states y_J, y_K, y_C, y_F — declared *before* any evaluation — then Möbius transport along the path forces $CR(y_J, y_K; y_C, y_F) = \frac{4}{3}$, and the fourth state is determined fit-free by the first three (y_F reconstruction identity). Negative control with teeth: the raw determinant path $(2, 4, 14, 32)$ gives $CR = \frac{28}{25} \neq \frac{4}{3}$ — the already-known norm sequence cannot pass. *The discrete half is now executed*: the four corners are the in-repo kernels $J, K, C, F = M(1, t) = R + Q \text{diag}(1, t, t)$, $t = -2.1$ (`v224`), and the corpus's one canonical declared solver reading — the Koide anchor response $\ell(M) = a^T M \mathbf{1}$ (`v183`) — reads $(26, 35, 44, 53)$ on them: $CR = \frac{4}{3}$ exactly, and the fit-free reconstruction *forces* $y_F = 53$ from $(26, 35, 44)$ alone — the Koide numerator is Möbius-determined by the three upstream corners (a new exact cross-corner identity); honesty theorem stated out loud: ℓ is affine on the path (step $9 = N_{\text{fam}}^2$), and *every* linear functional of the kernel passes — the check separates linear from nonlinear exports (the dominant-eigenvalue export fails at 1.3222 , the determinants at $\frac{28}{25}$, both as they must) **[E]**. Status: the *continuous* half (the external solvers exporting their own states) stays open exactly as this contract fences it; contract markers unchanged **[C]**. *The frozen export protocol for the continuous half* (design **[X]**, preregistered here): each instance must export, *before* evaluation, the canonical one-dimensional datum of its own native flow — the fixed-point multiplier of its contraction (cf. the native rates of `v425`: $F_{\text{pole}} (2/3)^6$ exact in-corpus, $F_{\text{Boltzmann}}/F_{\text{relic}}/F_{\text{QCD}}$ external) — as a projective state on the path; only then is $CR = \frac{4}{3}$ tested, with no renormalisation after the fact. *Executed* (`[verification/v575_cr_continuous_uniflow.py]`): the in-repo native exports form the quadruple $\{(2/3)^6, (2/3)^6, 1, -7\}$ — the repetition is *forced* by the single-flow theorem

(v425: F_pole and F_Boltzmann are literally the same seam contraction) — and a repeated quadruple gives $\text{CR} \in \{0, 1, \infty\}$ in every corner assignment (theorem, plus all 24 permutations): the continuous half can never pass *natively*; it closes onto the single-flow reduction, verdict UNIFLOW-DEGENERATE. The four independent external data are the dimensionful anchors (v_{geo} , M_R , C_p , θ_i) — [O] walls, not projective states — so a future continuous test needs genuinely *external* anchor-level exports; the meaningful cross-ratio content was and remains the discrete half. Both halves of the check are now executed [E]/[C]. *The jet-level successor is executed natively too* ([verification/v578_native_jets.py]): when point values degenerate, the Möbius-invariant *Schwarzian* of the exported trajectories is the right jet test — and natively, the exact v99 Koide trajectory has $\{q, t\} = -\Delta^2/2$ in wall time (constant, trajectory-independent; F_Boltzmann identical by the single-flow theorem) while being Möbius in its *own* exponential clock $u = e^{\Delta t}$ (Schwarzian 0), and the 1-loop QCD flow is Möbius in its own RG time already (Schwarzian 0): the native quadruple $\{-\Delta^2/2, -\Delta^2/2, 0, \text{degenerate}\}$ shares *no* wall-time jet invariant — each instance is *projective in its own clock*, the obstruction is the clock, and the clocks are anchored by the external data. Verdict MOEBIUS-IN-OWN-CLOCK; a future joint jet test needs external anchor-level jets; markers unchanged [E]/[C].

The discrete→dynamic lens: F_transfer is the readout end of the one principle ([verification/v303_ftransfer_dynamics.py], FR.DYNAMICS.01). The typing above is sharpened by a *dynamical* reading. The main-branch mechanism is a gapped, positivity-preserving relaxation to a *unique* attractor (the local-averaging update $m \leftarrow (A+2I)/4m \rightarrow$ the E_8 marks, gap $6 \ln(3/2)$). All four F_transfer instances share that *same shape*: [E] F_pole (Koide) *is* the seam dynamics — the source→pole/branch relaxation is the Möbius map with fixed-point multiplier $F' = (2/3)^6 = \lambda_2$, the seam-clock subleading eigenvalue, so $|F'| < 1 \Rightarrow$ a unique attractor at the *seam* rate (v82); [C] F_Boltzmann (η_B) is a gapped positive contraction (washout $\kappa_f \in (0, 1)$ toward the unique balance, an H-theorem); [C] F_relic (axion) freezes at an adiabatic fixed point (the comoving number relaxes to a constant); [E] F_QCD (m_p/m_e) is the 1-loop RG flow with the carrier $b_3 = -7$, whose Gaussian UV fixed point is the attractor (asymptotic freedom, Λ_{QCD} generated). *Honest boundary*: only F_pole has the seam rate $(2/3)^6$ exactly; the other three share the *shape* with external rates (thermal, cosmological, RG). So F_transfer is not a bolt-on but the downstream manifestation of the one discrete→dynamic principle — the compiler fixes the discrete seed + integer kernel, a gapped flow carries it to the observable — and the theory’s simplicity is preserved precisely because the external rates are honestly fenced (v187), not because they reduce to the seam. (Even F_pole’s flow time is non-integer, v93: the relaxation is the *continuous* semigroup $e^{t \log F}$ generated by the discrete operator — exactly the diffusion-semigroup reading of the main branch.)

The single-flow reduction (DYN.TRANSFER.UNIVERSAL.01). The dynamic lens is sharpened to a precise reduction ([verification/v425_dyn_transfer_universal.py]): the shared “shape” is literally the *one* native flow of the theory — the seam modular/recovery semigroup (v238/v240, gap $6 \ln(3/2)$) — restricted to each sector, extending the static universal-gap principle (v383) to a dynamic one. [E] the v99 Koide generator $\dot{q} = (\Delta/N_{\text{fam}})(q-2)(q-5)$ linearised at the attractor $q=2$ has rate $-\Delta$, so the time-1 contraction is $(2/3)^6$ (the recovery eigenvalue, $q=5$ unstable); [E] the Boltzmann washout is that same contraction, with the CP phase $4\pi/3$ native (v320); [E] the QCD rate is $b_3 = -7$. So the transfer *dynamics* is native; only the *anchors* stay external — the IR pole scale and M_R are v_{geo} -class, C_p is the lattice $O(1)$, and F_{relic} ’s cosmological initial condition θ_i is the one genuine wall. A reduction, *not* internalisation: no new premise, and the firewall (v187) holds — the observable stays [C].

Both thermal-time routes are dead: the external clock contract stands (FTRANSFER.STT.01 / FTRANSFER.COCYCLE.01). The radical counter-idea to the external-clock wall — Connes-style thermal time, “the four clocks = four modular flows of one state” — is now machine-killed on *both* checkable routes. [X] *Strong thermal time* ([verification/v723_phys_modular_clock.py], 11/11, STT-KILLED + INTERNAL-PAIR-CONFIRMED): a modular generator is self-adjoint, hence diagonalizable — but the F_QCD 1-loop clock map $u \mapsto u + 2b_0$ is *parabolic* and not diagonalizable

(single Jordan block), and the symbolic lemma “exp of a real-diagonalizable matrix is diagonalizable” closes the door: no constant-rescaled modular flow produces the QCD clock; the pole multiplier is confirmed *exactly* $(2/3)^6 = 64/729$ (hyperbolic, dimensionless $\Rightarrow$ internal; F_Boltzmann the same element by v425). [X] *Conditioned modular flows* ([verification/v724_phys_t3b_modular_flows.py], 12/12, T3B-DEAD): with target-free graph projectors, *one* global KMS scale $\Delta = 6 \log(3/2)$ and the Connes cocycle composition exact to 1.6×10^{-15} along J–K–C–F, the four conditioned flows come out *loxodromic/complex* — the frozen v578/v632 solver classes (hyperbolic / parabolic / degenerate) do *not* arise from conditioned modular flows of one internal mother state (the NEEDS-LINDBLAD escape is honestly refused). *Typed consequence*: the external clock contract is *confirmed* and sharpens to the anchor census {one unit = Λ_{QCD} (v_{geo} -class by No-Unit v153), one angle θ_i , one lattice $O(1)$ C_p } — T3’s clock wall and T1’s unit wall are *the same wall*, counted once. No marker moves; the functor typing above is unchanged.

The frozen external-clock contract (FTRANSFER.CLOCKS.01) — preregistered 2026-08-04, executed 2026-08-05. Since both thermal-time routes are dead, the clocks *must* be external — and without clocks fixed in advance every jet statement is arbitrarily reparametrizable. The contract therefore freezes them (design byte-frozen 2026-08-04 in experiments/ftransfer-clocks/hypotheses/ftransfer_clocks_v1.yaml; typed [X], a search-target design, *not* a claim). *The four frozen clocks, exactly one physical clock per solver*: F_pole = proper time τ (the on-shell/pole-mass phase clock; frozen gauge: one v99 seam unit = one proper-phase e-fold); F_Boltzmann = inverse temperature β (FIRAS-anchored adiabatic chart; one washout seam unit = one β e-fold; the z -linear alternative is recorded as *excluded*); F_QCD = RG time $t = \ln(\mu/M_Z)$ ($\overline{\text{MS}}$, 1-loop $b_0 = 7$ on the frozen window $[m_t, 1 \text{ TeV}]$; identity gauge); F_relic = cosmological e-folds $\ln a$ (Planck 2018 background; the flow is frozen, the jets *degenerate* — the channel is scored only on kills and dictionary validity). *The preregistered prediction* (FTC.P1): under the frozen affine dictionaries (slopes $-1, -1, +1, -1$ into the common chart $\sigma = \ln(\mu/M_Z)$) *one* constant-coefficient Riccati equation carries all non-degenerate transported jets — equivalently, the transported Schwarzians of F_pole/F_Boltzmann/F_QCD are *equal*; the seam channels transport to $-\Delta^2/2 = -2.959235170076978$ exactly, so the prediction forces the transported QCD Schwarzian onto that value. *Three kills*: K1 NO-COMMON-CONNECTION (symbolic inequality of the transported Schwarzians; on kill the executor must exhibit the forbidden reparametrization per failing pair); K2 FREE-REPARAM-NEEDED (any post-freeze change of a gauge or dictionary; the YAML sha256 is recorded by the executor); K3 SECOND-CLOCK-NEEDED (any channel export needing two time variables). *The honest prior, preregistered*: the *expected* outcome is the K1 kill — affine transport scales a Schwarzian by slope² and cannot move 0 (the QCD value) onto $-\Delta^2/2$ (the seam value); a K1 kill would finalize the v578 diagnosis at the *physical-clock* level and close the jet route of CONTRACT.F.01; ONE-CLASS would be the surprise. *Executed* (2026-08-05, [verification/v777_ftransfer_clock_jets.py], *one module from two probes*, 24 + 14 checks, 0 failures): the K1 kill fired *exactly* as the frozen prior expected — verdict NO-COMMON-CONNECTION. The freeze guards were byte-verified at run time (YAML sha256 880224f7...; the two data tables — Saikawa–Shirai 2018 g_{*s} , PDG 2024 $\alpha_s(Q)$ — were curated and SHA-frozen *before* the first run); all four dictionary-validity legs pass (Koide scheme check 571 $\times$; washout entropy-chart drift 1.15×10^{-4} ; QCD overlay max dev 0.83%; relic onset slope dev 0.0602); the transported Schwarzians are sympy-exactly $\{-\Delta^2/2, -\Delta^2/2, 0\}$ — no tolerance involved — and the K1 duty is discharged: the pass-enabling reparametrization $h = \text{Möbius}(e^{\pm\Delta\sigma})$ is exhibited and is non-affine, exactly the freedom the freeze forbids; K2/K3 stay quiet. The same module carries the surviving architecture (the fibered refinement — see the CONTRACT.F.01 dated note above). [2026-08-06 completion (FTRANSFER.KMSCOSET.01, [verification/v792_ftransfer_kms_coset.py], 18 checks, KMSCOSET-THEOREM): the coset assignment of these clock charts is now a theorem from KMS/modular typing — $S = -(\ln \lambda)^2/2$ from the time-1 PGL_2 holonomy multiplier per clock e-fold, with a Schwarzian-free, derivative-free classifier (the detailed-balance multiplier cross-ratio as the discrete KMS signature), blind assignment and exact unblinding; the three-probe chain executor $\rightarrow$ fibered groupoid $\rightarrow$ coset theorem is whole.] Nothing here upgrades CONTRACT.F.01: the four observables

stay [\[C\]](#)/[\[O\]](#); no marker move; no RH statement (unrelated surface; discipline note). So the visible residual is one geometric postulate (QGE0.SYM.01), one unit gauge (v_{geo}), and this one typed transfer functor — everything else is out-of-sample data.

External works built upon

The TFPT discrete core (the two axioms, the E_8 glue, the integer skeleton) is self-contained and machine-checked. Its *bridges* to continuum/AQFT physics, to gravity and to the lattice/singularity picture build on established mathematics, listed here by topic; the body refers to these by author name.

Modular theory and algebraic quantum field theory.

- M. Takesaki, *Tomita's Theory of Modular Hilbert Algebras and its Applications*, Lecture Notes in Math. **128**, Springer (1970).
- J. J. Bisognano, E. H. Wichmann, *On the duality condition for a Hermitian scalar field / for quantum fields*, J. Math. Phys. **16** (1975) 985–1007; **17** (1976) 303–321.
- K. Osterwalder, R. Schrader, *Axioms for Euclidean Green's functions* I, II, Comm. Math. Phys. **31** (1973) 83–112; **42** (1975) 281–305.
- S. Doplicher, R. Haag, J. E. Roberts, *Local observables and particle statistics* I, II, Comm. Math. Phys. **23** (1971) 199–230; **35** (1974) 49–85.
- R. Haag, *Quantum field theories with composite particles and asymptotic conditions*, Phys. Rev. **112** (1958) 669–673; D. Ruelle, *On the asymptotic condition in quantum field theory*, Helv. Phys. Acta **35** (1962) 147–163.
- R. Longo, *Index of subfactors and statistics of quantum fields* I, II, Comm. Math. Phys. **126** (1989) 217–247; **130** (1990) 285–309.
- Y. Kawahigashi, R. Longo, M. Müger, *Multi-interval subfactors and modularity of representations in conformal field theory*, Comm. Math. Phys. **219** (2001) 631–669.

Conformal nets, vertex operator algebras and lattice constructions.

- I. B. Frenkel, V. G. Kac, *Basic representations of affine Lie algebras and dual resonance models*, Invent. Math. **62** (1980) 23–66; G. Segal, *Unitary representations of some infinite-dimensional groups*, Comm. Math. Phys. **80** (1981) 301–342.
- T. J. Osborne, A. Stottmeister, *Conformal field theory from lattice fermions*, Comm. Math. Phys. **398** (2023) 219–289, [arXiv:2107.13834](#), [doi:10.1007/s00220-022-04521-8](#). (This is the seam scaling-limit theorem the TFPT continuum leg cites — the Koo–Saleur $\rightarrow$ Virasoro convergence, $c = \frac{1}{2}$, and the smeared WZW-current Theorem 5.1. The internal TFPT code-token “MMST” and the legacy claim-ids SEAM.MMST.* / filenames v*_mmst_* denote *this* Osborne–Stottmeister theorem.)
- V. Morinelli, G. Morsella, A. Stottmeister, Y. Tanimoto, *Scaling limits of lattice quantum fields by wavelets*, Comm. Math. Phys. **387** (2021) 299–360, [arXiv:2010.11121](#), [doi:10.1007/s00220-021-04152-5](#); and *Operator-algebraic renormalization and wavelets*, Phys. Rev. Lett. **127** (2021) 230601. (The operator-algebraic (wavelet/Wilson–Kadanoff) renormalization *framework* the Osborne–Stottmeister theorem above builds on.)
- M. S. Adamo, Y. Moriwaki, Y. Tanimoto, *Osterwalder–Schrader axioms for unitary full vertex operator algebras*, [arXiv:2407.18222](#) (2024).
- M. S. Adamo, L. Giorgetti, Y. Tanimoto, *Rational and non-rational two-dimensional conformal field theories arising from lattices*, [arXiv:2506.01008](#) (2025); *Representations of conformal nets associated with infinite-dimensional groups*, [arXiv:2508.07109](#) (2025).
- P. Naaijken, D. Penneys, B. Wallick, *(commuting-projector reflection positivity / LTO-RP)*, [arXiv:2605.10693](#).

Lattices and quadratic forms.

- J. H. Conway, N. J. A. Sloane, *Sphere Packings, Lattices and Groups*, 3rd ed., Grundlehren der math. Wiss. **290**, Springer (1999).
- C. L. Siegel, *Über die analytische Theorie der quadratischen Formen* (the Minkowski–Siegel mass formula), Ann. of Math. **36** (1935) 527–606.

Singularities and geometry.

- E. Brieskorn, *Beispiele zur Differentialtopologie von Singularitäten*, Invent. Math. **2** (1966) 1–14.
- J. Milnor, *Singular Points of Complex Hypersurfaces*, Ann. of Math. Studies **61**, Princeton Univ. Press (1968).

Spectral geometry, gravity and thermal time.

- A. H. Chamseddine, A. Connes, *The spectral action principle*, Comm. Math. Phys. **186** (1997) 731–750.
- A. Connes, C. Rovelli, *Von Neumann algebra automorphisms and time–thermodynamics relation in general covariant quantum theories*, Class. Quantum Grav. **11** (1994) 2899–2918.
- R. M. Wald, *Black hole entropy is the Noether charge*, Phys. Rev. D **48** (1993) R3427–R3431.
- H. Casini, M. Huerta, R. C. Myers, *Towards a derivation of holographic entanglement entropy*, JHEP **05** (2011) 036, [arXiv:1102.0440](#).




Figure 3: The structural-residual reduction chain ($v175 \rightarrow$ the certification round). The naive “build a quantum-gravity measure” shrinks, one machine-checked step at a time, from a vague measure to one falsifiable physical statement — *is the raw seam $(E_8)_1$ at $\tau = i$?* (SEAM.EQUIV.01; its conformal-deck face QGEO.SYM.01 is a corollary, v335). The closing arc ($v276$ – $v302$) pins the shared fact and derives the gap $6\ln(3/2) > 0$; the certification round discharges the extension leg to the crossed-product package (v469), the R3 graph→geometry bridge to Kronheimer 1989 (v479) and exhibits the four-interval μ_4 mechanism (v480). Every step above the bedrock is a theorem, a Lean proof or an established citation; the residual — the cited continuum scaling-limit existence (v336) — stays honestly [O].

F_{transfer} : one shape, four rates — a gapped relaxation to a **UNIQUE** attractor




Figure 4: **The discrete→dynamic lens on F_{transfer}** ([verification/v303_ftransfer_dynamics.py]). All four transfer instances share *one* shape — a gapped, positivity-preserving relaxation to a *unique* attractor, the same Perron-Frobenius / H-theorem / RG-fixed-point structure as the main-branch update $m \leftarrow (A+2I)/4m \rightarrow$ the E_8 marks. *Only* F_{pole} (Koide, gold) runs it at the *seam* rate: its Möbius multiplier is $F'=(2/3)^6=\lambda_2$, the seam-clock subleading eigenvalue. $F_{\text{Boltzmann}}$ (η_B , thermal washout), F_{relic} (axion, cosmological freeze) and F_{QCD} (m_p/m_e , the RG flow to the Gaussian UV fixed point with the carrier $b_3=-7$) share the *shape* with *external* rates. So **F_{transfer}** is the readout end of the one principle, not a bolt-on; the theory's simplicity is preserved because the external rates are honestly fenced (v187), not reduced to the seam.