

The saturated dual resolvent

A specialist dossier on the critical bordered block $R^\dagger$

Stefan Hamann

August 30, 2026

Abstract

On a family of explicit finite discrete measures, a particle-hole duality identifies a two-measure subordination inequality with the spectral condition $R \succ \frac{1}{2}I$ on a dual hole kernel. A rank-one border of that kernel produces a single object $R^\dagger$ whose positivity $R^\dagger \succ \frac{1}{2}I$ is equivalent to the original inequality *and* a terminal budget condition, and whose critical three-by-three Schur block carries both. A constrained equilibrium problem on the same dual weight places the binding pair of nodes astride a saturated hard-edge block. All identities below are finite linear algebra; all numerical values are a frozen census on a finite family. No asymptotic theorem is proved. After the two-rank inertia cut, the graph-resolvent dictionary, and the edge-balance chain, the two remaining holes close at the same kind of object: an unsigned cancellation of von Mangoldt data (Section 12). Round 427 audited the campaign numbers (Section 13). The current architecture is an *open program* in the Krein–Langer class $\mathcal{S}_\kappa$ (Section 14): bulk membership $\mathcal{S}_{\leq 1}$ and edge membership $\mathcal{S}_0$ via a Redheffer pivot, with two quantifier weakenings. The older minimal-inertia and saturation/RHP questions are retained as Problems 1 and 2. This document is a problem statement for specialists, not a claim, and nothing in it is evidence for or against the Riemann hypothesis.

Status and grades. Every assertion is marked EXACT (finite identity, including rational arithmetic), THEOREM-GRADE (finite-matrix theorem on measured inputs), LEAN-VERIFIED (sorry-free in `rh/lean/`, kernel-checked), CENSUS (measured on a frozen finite family; no extrapolation), HYPOTHESIS (unproved analytic statement), REFUTED (named internal route closed by a sealed probe), or OPEN PROGRAM (named target, no existence proof). The construction of the underlying measures (μ, ν) is given in full in the companion note [4]; the present dossier does not repeat it. Uvarov-transform language, the discrete-hard-edge placement, and the Lubinsky-class exclusion are recorded there (§frozen-placement, §frozen-dualhole) and are cited, not duplicated. Machine records: sealed probes of rounds 356, 359, 360, 362, 363, 366, 367 and 369–372 (repository `experiments/tfpt-discovery/`); Lean transcription round 373 (`RH/Haynsworth.lean`, `RH/DualResolvent.lean`); graph-resolvent consolidation round 412 (`RH/GraphResolvent.lean`); edge-balance consolidation round 426 (`RH/EdgeBalance.lean`); campaign audit round 427; selected-sequence reopening round 428; head-identity round 429. Companion notes: `campaign_audit.tex`, `qn_reopened.tex`, `zloc_head.tex`.

1 Setup

Fix a prime power z (the window anchor). A triangular (“tent”) test function of half-width Δ , determined by z and a fixed oversampling parameter, samples the von Mangoldt comb and an archimedean kernel, producing an even trigonometric polynomial f — the symbol of a circulant. Folding the symbol onto $[-1, 1]$ by $x_j = \cos \theta_j$ yields a signed discrete measure on the cosine grid, split by sign into two positive measures μ and ν with disjoint supports in $[-1, 1)$ and S atoms in total [4, construction]. Write $N = \lceil S/2 \rceil$ and $Y = \text{supp } \nu$. Across the whole family the

grid is unculled, so $S = 2N - 1$ [exact]: half-filling is Borodin’s complementary rank condition, not an extra hypothesis.

The running instance is $z = 16$: $S = 367$, $N = 184$, $|Y| = 104$. The binding pair of ν -atoms sits at grid (fold) indices $(2, 4)$, the two rightmost odd nodes of the cosine lattice. The inequality under study, denoted L^* , asks

$$\int p^2 d\nu < \int p^2 d\mu \quad \text{for every real polynomial } p \neq 0 \text{ with } \deg p < N.$$

Equivalently, the ν -sampling matrix E_N in the μ -orthonormal frame satisfies $\lambda_{\max}(E_N) < 1$. An augmented form $L^\dagger$ adjoins a rank-one border (a budget-normalized smooth test functional) and asks $\lambda_{\max}(E_N + bb^T/B) < 1$. Both statements are open for the unbounded family. On every tested window the inequalities hold, with spectral margins decaying as a power of N ; structurally destroyed comparison inputs fail already at a small fraction of the critical degree [4]. No statement here claims either inequality.

2 Definition of $R^\dagger$

Lift (μ, ν) to the positive measure $\eta = \mu + \nu$ with weights $u_j = |\tilde{w}_j|$ on the union $X = \text{supp } \eta$. Let P_X be the monic node polynomial of X . Borodin’s reciprocal weight [1] is $u_j^\vee = 1/(u_j P'_X(x_j)^2)$. On this cosine grid the dual weight has the closed form

$$u_j^\vee \propto \frac{c_j(1 - x_j)}{|f(\theta_j)|}, \quad (1)$$

where c_j is the folding weight ($c_j = \frac{1}{2}$ at the left endpoint, else 1): the original weight carries $+\log|f|$, the dual weight $-\log|f|$ [exact, three evaluation routes]. Let $\Pi_{N-1}^{u^\vee}$ be the rank- $(N-1)$ Christoffel–Darboux projection kernel of $u^\vee$ on X , dressed by $\sqrt{u^\vee}$, and let R be its restriction to Y .

Let b be the border functional in the μ -orthonormal frame, $B > 0$ the budget, $\beta = b/\sqrt{B}$, and v the node-side image of the border under the μ -Christoffel–Darboux kernel ($v_j = \sqrt{\nu_j/B} \int K_\mu(y_j, t) d\sigma(t)$). Write $\gamma = \beta^T \beta$ and $\tilde{v} = Dv$, with D the diagonal sign matrix of P'_X .

Definition 1 (Bordered dual resolvent). *The augmented dual kernel is the $(|Y| + 1) \times (|Y| + 1)$ matrix*

$$R^\dagger = \begin{pmatrix} R^{-1} & \tilde{v} \\ \tilde{v}^T & 1 + \gamma \end{pmatrix}^{-1}. \quad (2)$$

The extra coordinate is the border; the leading principal block is a rank-one Sherman–Morrison deformation of R (Proposition 2(A4) below).

Route (i) — adjoining a virtual atom and taking the $(S+1)$ -point Borodin kernel — is available on a single-atom toy border (half-filling shifts $S = 2N - 1 \mapsto S^\dagger = 2N$, dual rank $N - 1 \mapsto N$) and agrees with (2) up to sign conjugation [exact, rational arithmetic]. On the real windows the border is a signed smooth-comb measure with S atoms, every one of which collides with a grid node, and is not a point evaluation (representability spread 56.8 on the running instance). The bordered-resolvent form is therefore the general object.

3 Exact duality formulae

Write $\text{margin} = 1 - \lambda_{\max}(E_N)$ and $\text{margin}^\dagger = 1 - \lambda_{\max}(E_N + bb^T/B)$.

Proposition 1 (Particle-hole dictionary [1]). *At half-filling $S = 2N - 1$,*

$$L^* \iff \lambda_{\max}(\mathbf{E}_N) < 1 \iff R \succ \frac{1}{2}I,$$

with the spectral map margin $= 2 - 1/\lambda_{\min}(R)$ [theorem-grade; absolute deviation $\leq 9.9 \cdot 10^{-10}$ on 85 windows]. The L -ensemble transformation $Q = \mathbf{E}_N(I + \mathbf{E}_N)^{-1}$ is exact [rational arithmetic on a 5-atom model; double precision $\leq 7.6 \cdot 10^{-12}$ on 85 windows], and combined with Borodin's complementation $\Pi_N^u = D(I - \Pi_{N-1}^v)D$ yields $\mathbf{E}_N = D(R^{-1} - I)D$ on Y .

Live census: $\lambda_{\min}(R) - \frac{1}{2}$ ranges from $+4.2 \cdot 10^{-5}$ (running instance: $\lambda_{\min}(R) = 0.500041882$) down to $+5.9 \cdot 10^{-11}$ on the deepest tranche — all positive, no counterexample. Comparison inputs with destroyed source structure violate $R \succ \frac{1}{2}I$ structurally ($\lambda_{\min}(R) - \frac{1}{2} \in [-0.500, -0.083]$). Honest limit: at the deepest instances the double-precision resolution of the dual route ($\sim 5 \cdot 10^{-10}$) exceeds the margins, so the hole coordinate is exact algebra, not a sharper certification instrument.

The pair identities $c = \varepsilon_1 \varepsilon_2 (R^{-1})_{12}$, $p = 2 - (R^{-1})_{11}$, $q = 2 - (R^{-1})_{22}$ identify the two-point Uvarov block of the companion note with the $(1, 2)$ principal minor of $2I - R^{-1}$ [theorem-grade]. The absolutely continuous hard-edge class is excluded at census grade: the rescaled pair positions $a_j = 2N^2(1 - y_j)$ are family-constant against $\pi^2 j^2/4$ on all 85 windows (maximum deviation $0.33/N$), while the kernel correlation ρ_K decays with fitted exponent 1.4222, contrary to Lubinsky's Bessel limit at fixed microscopic arguments [3], [4, §frozen-placement].

Proposition 2 (Augmented dictionary). *Let $Z = \begin{pmatrix} R^{-1} & \tilde{v} \\ \tilde{v}^T & 1+\gamma \end{pmatrix}$, so $R^\dagger = Z^{-1}$, and write $M^\dagger = R^\dagger - \frac{1}{2}I$. The following are finite-matrix identities [theorem-grade; rational arithmetic on toys, graded double precision on 75 windows of the prime-generated family plus two character-twisted families of 42 windows each].*

- (A1) *The augmented node Gram $G^\dagger$ is the sign-conjugate of $(R^\dagger)^{-1} - I$: the particle-hole complementation extends by one row.*
- (A2) *margin $^\dagger = 2 - 1/\lambda_{\min}(R^\dagger)$ (deviation $\leq 1.1 \cdot 10^{-11}$ shallow, $\leq 5.5 \cdot 10^{-10}$ at the deepest computed row).*
- (A3) *$L^\dagger \iff R^\dagger \succ \frac{1}{2}I$, two-sided. Both truth directions occur on real windows: six character-twisted rungs with terminal ratio $q_N > 1$ have $\lambda_{\min}(R^\dagger) - \frac{1}{2} < 0$ (e.g. $-9.8 \cdot 10^{-5}$), exactly as the identity demands.*
- (A4) *Sherman–Morrison on the Y -block: $R_{YY}^\dagger = R + (R\tilde{v})(R\tilde{v})^T/(1 + \gamma - \tilde{v}^T R\tilde{v})$, and $R_{bb}^\dagger = 1/(1 + \gamma - \tilde{v}^T R\tilde{v})$ (deviation $\leq 1.2 \cdot 10^{-14}$ on every computed row, including the deepest). The border is a rank-one insertion at dual-resolvent level.*
- (A5) *$R^\dagger \succ \frac{1}{2}I$ if and only if $R \succ \frac{1}{2}I$ and $q^\dagger < 1$, where $q^\dagger = \gamma + v^T(I - \mathbf{E}_N)^{-1}v$ is frame-invariant (equals the squared dual border norm $u^T H^{-1}u/B$) and coincides with the standing terminal coordinate. The dictionary $1 - q^\dagger = (5/7)(1 - q_N)/B$ holds to relative $6.7 \cdot 10^{-6}$.*
- (A6) *The Schur complement of $M^\dagger$ onto the border coordinate equals $(1 - q^\dagger)/(2(1 + q^\dagger))$ globally, and equals the same value inside the local 3×3 block of Proposition 3 (quotient $\leq 1.0 \cdot 10^{-11}$).*
- (A7) *Cauchy interlacing: $\lambda_k(R^\dagger) \leq \lambda_k(R) \leq \lambda_{k+1}(R^\dagger)$. The border can only lower the bottom of the dual spectrum (margin $^\dagger \leq$ margin). On the running instance, $0.500041459 \leq 0.500041882$.*

A formal reconstruction, independent of the dual coordinate, is already machine-checked in Lean: $L^\dagger \iff L^* \wedge \text{Terminal}$ and $L^* \wedge \text{Terminal} \Rightarrow$ positivity of every bordered window matrix

through the cap (RH/Augmented.lean, RH/Closure.lean; axiom audit in RH/Audit.lean, consuming only the two named canonical holes). The dual-resolvent dictionary itself is sorry-free since round 373: $L^\dagger \iff R^\dagger \succ \frac{1}{2}I$ under the μ -ONB whitening (RH/DualResolvent.lean) [lean-verified]. Haynsworth additivity on the two-rank and mixed J -forms is likewise sorry-free (RH/Haynsworth.lean; Section 8). What is *not* formalized, and not proved analytically, is the pair of census premises (P1)–(P2) of Problem 1 for the unbounded family — and, a fortiori, $R^\dagger \succ \frac{1}{2}I$ itself.

4 The critical three-by-three block

Partition $Y = \{y_1, y_2\} \cup C$, with $\{y_1, y_2\}$ the binding pair (folds 2, 4) and C the rest of Y . Write $M = R - \frac{1}{2}I$ and S_N for the 2×2 Schur complement of M onto the pair. Then [exact]

$$M \succ 0 \iff M_{CC} \succ 0 \text{ and } S_N \succ 0,$$

and S_N is a ratio of Sylvester-bordered minors of $\det M$ (the rest block sits inside the border; a pair-only truncation is false by 4.07 nats on the running instance and exactly on the rational toy). The off-diagonal of the dual resolvent $A^{-1} = (I - 2R)^{-1}$ is the Casoratian of the two consecutive dual orthonormal polynomials, resolvent-dressed (discrete IKS identity) [theorem-grade]. The diagonal entries $(A^{-1})_{kk}$ are *not* Casoratian data: their cross share $(A^{-1})_{12}^2 / ((A^{-1})_{11}(A^{-1})_{22})$ has median 0.702 on 85 windows (range 0.46–0.84). No standalone explicit prefactor turns the Casoratian into $\det S_N$ at the sealed bars.

Proposition 3 (Binding). *On the 74 resolvable windows of the family $(\lambda_{\min}(R) - \frac{1}{2} > 10^{-9})$,*

$$\text{bind} = \frac{\lambda_{\min}(S_N)}{\lambda_{\min}(R) - \frac{1}{2}} \in [1.0003, 1.0605], \quad \text{median } 1.0058$$

[census]. *After the rank-one border, the 3×3 Schur block $S_3^\dagger$ of $M^\dagger$ on (pair, border) satisfies*

$$\text{bind}^\dagger = \frac{\lambda_{\min}(S_3^\dagger)}{\lambda_{\min}(R^\dagger) - \frac{1}{2}} \in [1.0003, 1.0608], \quad \text{median } 1.0056$$

[census]. *The in-block border Schur equals the global border Schur of (A6). The rank-one deformation does not move the pair structure: on the running instance the cross share of $R^\dagger$ is 0.6990 against 0.6973 for R ; ladder medians 0.708 against 0.702.*

On the running instance the three eigenvalues of $S_3^\dagger$ are $(4.16 \cdot 10^{-5}, 4.90 \cdot 10^{-4}, 1.76 \cdot 10^{-2})$: the bottom mode is the L^* pair channel, the top mode is the terminal border channel. The critical eigenvector’s border mass is $2.5 \cdot 10^{-5}$ there, and $5.8 \cdot 10^{-4}$ at the near-terminal window where $\text{margin}^\dagger / \text{margin} = 0.8716$. The giant operator behaves, at the critical point, like this 3×3 block.

5 The pair straddles the saturation edge

Two independent finite objects describe the dual ensemble’s occupation of the cosine grid.

Exact occupation. Let $o_\eta(x_j)$ (resp. $o_{\text{dual}}(x_j)$) be the diagonal of the rank- N η -projection (resp. the rank- $(N-1)$ dual-hole projection). Borodin complementation on the diagonal gives

$$o_\eta(x_j) + o_{\text{dual}}(x_j) = 1 \tag{3}$$

at every union node, with trace ward $\sum_j o_{\text{dual}}(x_j) = N - 1$ [exact; rational arithmetic on the toy, $\leq 2.2 \cdot 10^{-11}$ live]. This is the finite counterpart of an equilibrium density versus its upper constraint, with no solver and no asymptotics. On the prime-generated family, fold 1 (the rightmost node) is the unique occupation anomaly: $o_{\text{dual}}(x_1)$ lies 0.25–0.33 below the bulk median on all 85 windows, while the pair folds 2 and 4 are flat (deviation ≤ 0.10).

Obstacle problem. Minimize the discrete energy

$$E(m) = \sum_{j \neq k} m_j m_k \log \frac{1}{|x_j - x_k|} + \sum_j m_j (-\log u_j^\vee)$$

over $0 \leq m_j \leq 1$, $\sum m_j = N - 1$. The box constraint is the Baik–Kriecherbauer–McLaughlin–Miller upper constraint [2]; dropping it piles mass ≥ 92 times the cap onto single nodes. A deterministic projected first-order solver (FISTA plus exact box-simplex water-filling, sealed iteration counts, KKT partition gates) produces, on four sealed instances including the running instance, a dual-void block equal to folds $\{1, 2, 3\}$ [census]. By (3) this is the primal saturated block at the right hard edge.

Proposition 4 (Straddle, census). *The binding pair (2, 4) straddles the saturation-block edge 3 | 4: fold 2 lies inside the saturated zone, fold 4 outside. Auxiliary character-twisted families on the same grid have a different partition — they occupy fold 1, with no fold-1 occupation anomaly on $0/42 + 0/42$ windows — and a sign-scrambled comparison input is disordered at the edge while failing $R_{CC} \succ \frac{1}{2}I$ ($\lambda_{\min}(R_{CC}) - \frac{1}{2} = -0.4962$) with the pair block itself still positive. The identity chain is world-blind algebra; the partition, and the positivity, are not.*

The asymptotic equilibrium-measure theorem (BKMM band and saturation limit shape for the field $-\log u^\vee$) is not proved [hypothesis, open]. The finite partition is a measurement about this family.

6 The observed exponent

On the 57 calibration windows a Theil–Sen fit of $\log \lambda_{\min}(S_N)$ has slope -3.334 , hitting the standing margin exponent 3.332 at distance 0.002 and rejecting an alternative reading 1.65 at distance 1.684 [census]. The diagonal dual-resolvent entries carry the same exponent ($|(A^{-1})_{11}|$ slope $+3.365$, $|(A^{-1})_{22}| + 3.292$, $|(A^{-1})_{12}| + 3.316$). Weight-side Christoffel exponents (1.79 and 0.38 families) miss by a wide margin: they do not carry the resolvent diagonals.

A dimensionless combination $t_{\text{geo}} = (\lambda_{\min}(R) - \frac{1}{2}) \sqrt{(A^{-1})_{11}(A^{-1})_{22}}$ is margin-locked: slope -0.009 on the 57, range $[0.222, 0.287]$ on all 74 resolvable windows [census]. Exact algebra identifies t_{geo} with a function of $\det(S_N/\varepsilon)$ and the cross share; family-constancy is a local-parametrix-class census, not a Gamma-function derivation. The Gamma parametrix of [2, Thm 2.7, Rem 2.8] adjacent to a saturated band remains the natural analytic candidate [hypothesis, open].

An adjugate identity $\det S_N \cdot 4[(A^{-1})_{11}(A^{-1})_{22} - (A^{-1})_{12}^2] = 1$ [exact] implies that the wander of $\log \det S_N$ is the wander of $-\log((A^{-1})_{11}(A^{-1})_{22})$ (fine-structure correlation $+0.9835$), with the cross-share term five times subleading. The leading common wander lives in the diagonal resolvent data and cancels algebraically in the minor.

7 Relative error-operator form

Two constant-matrix targets fail the census and force a relative statement.

Rest block. Cauchy interlacing on the principal submatrix R_{CC} , and the Schur-complement eigenvalue inequality, give

$$\lambda_{\min}(R_{CC}) - \frac{1}{2} \geq c(\lambda_{\min}(R) - \frac{1}{2}) \quad \text{with } c = 1 \quad (4)$$

as a finite theorem, gated in both directions on the rational toy and on 74 + 84 resolvable windows [theorem-grade]. On the prime-generated family the measured ratio lies in [8.8, 54.2] (median 20.3). On the character-twisted families the ratio reaches the interlacing line: minima 1.0 on 11 of 84 windows. Any analytic rest-block clause is a theorem only at $c = 1$; a comfortable factor 8.8 does not transfer.

Leading coefficient. The schematic $S_N = N^{-\alpha}(M_0 + O(N^{-\epsilon}))$ with M_0 a constant positive matrix is too rigid. The condition number $\kappa_S = \lambda_{\max}(S_N)/\lambda_{\min}(S_N)$ ranges in [6.3, 30.5] (median 13.9), a range ratio $4.88 > 1.5$: the margin direction collapses (bind median 1.0058) but the second eigenvalue wanders by an $O(1)$ factor [census]. The honest form is projective: one controls $S_N/\lambda_{\min}(S_N)$ in a bounded set of positive-definite matrices, or equivalently the error operator $E = P_{\text{true}}P_{\text{model}}^{-1} - I$ of a local Riemann–Hilbert parametrix in a *relative* operator norm on the critical block, not as an additive constant remainder.

No bound on E is proved [open]. Equation (4) at $c = 1$ is the only theorem-grade error control presently available, and it is finite interlacing, not an asymptotic remainder. After rounds 367–373 this relative form is the *stronger* target: the primary question no longer needs a remainder that tracks $N^{-\alpha}$ (Section 8).

8 The minimal inertia form (rounds 367–373)

The schematic $S_N = N^{-\alpha}(M_0 + O(N^{-\epsilon}))$ of Section 7 is a margin asymptotic. The two-rank inertia cut replaces it by two coarser statements on the rest block $A_0 = R_{N-3} - \frac{1}{2}I$ and the 2×2 Schur matrix $K_2 = I + U^T A_0^{-1}U$, with U the last two dual Christoffel–Darboux columns.

Analytic discount. Write $\text{ind}_-(A)$ for the number of strictly negative eigenvalues of a real Hermitian matrix A . It is enough to prove, uniformly on the admissible family,

$$\text{ind}_-(A_0) \leq 1 \quad (P1)$$

$$\det K_2 \leq -c_0 < 0 \quad \text{on the branch } \text{ind}_-(A_0) = 1. \quad (P2)$$

(P1) is an integer mode count. (P2) is the sign of an $O(1)$ 2×2 source. An asymptotic argument need only beat a *fixed* signature reserve, not the shrinking $N^{-3.33}$ spectral margin of Section 6.

Census on the 74 resolvable windows ($\lambda_{\min}(R) - \frac{1}{2} > 10^{-9}$) [census, round 367]: overload $\text{ind}_-(A_0) \geq 2$ is 0/74; the dichotomy is 45 exact-one (nneg-1) against 29 already positive-definite at rank $N - 3$ (P1 vacuous). On the nneg-1 branch, $-\det K_2 \in [1.157, 1126.389]$, census floor 0.50 holds 45/45, Theil–Sen slope of $\log(-\det K_2)$ against $\log N_w$ is +0.543 (flat against the margin rate -3.332), and the restatement correlation against log margin on the 57 calibration windows is -0.0064 . The implication $(P1) \wedge (P2) \Rightarrow R \succ \frac{1}{2}I$ holds on all 74 resolvable rows; it is sufficient, not necessary (the vacuous-29 already have $A_0 \succ 0$).

Theorem 1 (Two-rank Haynsworth). *Let A_0 be real Hermitian and invertible with $\text{ind}_-(A_0) = 1$, and let $K_2 = I_2 + U^T A_0^{-1}U$. If $\det K_2 < 0$, then $A_0 + UU^T \succ 0$ [lean-verified, RH/Haynsworth.lean, sorry-free].*

Finite real algebra only; the census premises (P1)–(P2) on windows are *not* asserted. The Lean statement is `haynsworth_two_rank`. On the hole kernel, $A_0 + UU^T = R - \frac{1}{2}I$, so Theorem 1 converts (P1)–(P2) into $R \succ \frac{1}{2}I$. The $R^\dagger$ dictionary of Proposition 2(A3),(A5) then lifts to $L^\dagger$ once the terminal Schur $q^\dagger < 1$ is available; both directions of $L^\dagger \iff R^\dagger \succ \frac{1}{2}I$ are sorry-free in `RH/DualResolvent.lean`.

Mixed 3×3 form. The border is not a third unaugmented CD column. At dual-resolvent level the Sherman–Morrison insertion of Proposition 2(A4) does supply a third column with an explicit signature. Write $s = R\tilde{v}$, $\text{den} = 1 + \gamma - \tilde{v}^T R\tilde{v}$, and

$$U_{\text{mix}} = \begin{pmatrix} u_{N-3} & u_{N-2} & s \\ 0 & 0 & -1 \end{pmatrix}, \quad J = \text{diag}(1, 1, 1/\text{den}).$$

Then [theorem-grade, round 369; Fractions residual exactly 0 with $\text{den} = 23/30$; live residual $\leq 1.04 \cdot 10^{-14}$]

$$M^\dagger = \text{blkdiag}(A_0, -\tfrac{1}{2}) + U_{\text{mix}} J U_{\text{mix}}^T. \quad (5)$$

The generalized Haynsworth identity for invertible Hermitian J ,

$$\text{In} \begin{pmatrix} A & U \\ U^T & -J^{-1} \end{pmatrix} = \text{In}(A) + \text{In}(-\Phi) = \text{In}(-J^{-1}) + \text{In}(A + U J U^T),$$

with $\Phi = J^{-1} + U^T A^{-1} U$, is sorry-free (**haynsworth.mixed**) [lean-verified]. The two-rank theorem is the case $J = I_2$.

Census building blocks. Three further finite objects are available to a specialist proof of (P1)–(P2). All three are typed CENSUS or THEOREM-GRADE as marked; none is an extrapolation.

- Wedge-sign stability [census, round 371]: the canonical CD two-form satisfies $\langle \omega, \wedge^2 A_0^{-1} \omega \rangle < 0$ on 45/45 of the nneg-1 branch (range $[-637.7, -1.32]$, zero flips). The second compound matrix therefore has source structure. The capture identity that would turn this into (P2) is algebraically identical to (P2) via the exact expansion of $\det K_2$; it is not independent arithmetic. The $\wedge^2$ -subspace share is *not* stable (majority ≥ 0.50 only on 27/45).
- Φ -monotonicity [theorem-grade, round 370]: the matrix-valued Weyl function $\Phi_N(z) = J^{-1} + U^T (A - zI)^{-1} U$ is a block resolvent (Fractions determinant lemma exact at $z = -2$), and $\Phi' \succeq 0$ (live minimum eigenvalue $8.949 \cdot 10^{-9}$ on 18/18 admissible grid nodes). The phase census is two-sided: MAIN 74/74 resolvable rows are classified positive-definite with exact signature balance; the six terminal-dead character-twisted rungs ($q_N > 1$) are classified negative (χ_3 5/5, χ_4 1/1).
- Prüfer dictionary [theorem-grade, round 372]: on the terminal surface, the $\lfloor \theta/\pi \rfloor$ -runs of the chain Prüfer phase coincide with the sign-runs of v_2 on 181/181 windows, with the XOR identity $\text{sign}(c_t) = \text{sign}(w \cdot x \cdot v_2)$ on the same 181. The dictionary is a theorem; the one-defect extraction that would have closed the terminal half from a single turning is [refuted].

Documented dead ends. Four internal routes are closed, each at a named place.

- No 3×3 Jacobi Casoratian [refuted, round 370]: dual p and dual-measure Cauchy q satisfy the three-term recurrence; the border Cauchy sequence r_n is not a homogeneous dual solution (residual $3.615 \cdot 10^{-2}$ on the running instance). There is no (p, q, r) Casoratian, so a matrix-Weyl index that would replace Haynsworth by an oriented winding does not linearize.
- Phase restatement [refuted, round 370]: the real-line census index $\Xi = \text{nneg } \Phi(0)$ is Haynsworth inertia in other coordinates. The phase language does not replace the inertia cut.

- $O(1)$ buffer of mass counting [refuted, round 366]: proportional Markov–Stieltjes counting in the pair gap predicts occupation $Z = 0$ ($nM_I/U = 2.40 \cdot 10^{-5}$ on the running instance); measured $Z = 1$ lives in the $O(1)$ sandwich buffer, which cannot separate 0 from 1 (0/74 scaled-band candidates).
- Edge pinning [refuted, round 363]: dual OP zeros are not exponentially node-pinned in the saturated block $\{1, 2, 3\}$ (Hurwitz maximum 0.462 against bar 0.15). The zone is dual-void / primal-saturated; a BKMM pinning argument at the right hard edge does not apply.

Lean landing pad (historical state through r459). At the r459 cut the sorry census was five (two arithmetic holes `lstar_canonical`, `terminal_q_canonical`, now the alternative route; the named dictionary `pair_terminal_dictionary`; the opacity bridge `mainWindow_iff_builtFromPrimeSource` the remaining Level-C arch statement `arch_elementwise_stabilization`). The entire $R^\dagger$ layer — Haynsworth, the dual-resolvent dictionary (A2)–(A5), (A7-min), the window bridge, the one-defect absorption, and the r412 graph-resolvent identities — consumes no `sorryAx`. A proof of (P1) and (P2) for all selected windows beyond the certified head lands, via the kernel-checked chain, at window-local master positivity. That is a landing pad, not a theorem of the family. The live mincut is the named Prop `selected_augDualResolvent_gt_half`.

9 Problem statement

Problem 1 (Minimal inertia of the rest block). *Let (μ, ν) run over the admissible family of [4], with dual kernel R as in Section 2, rest block $A_0 = R_{N-3} - \frac{1}{2}I$, and $K_2 = I + U^T A_0^{-1} U$ the two-rank Schur matrix of Section 8. Prove (P1) and (P2) uniformly for the unbounded family: at most one negative inertia direction in A_0 , and $\det K_2 \leq -c_0 < 0$ on the nneg-1 branch. Which theorem in the Riemann–Hilbert analysis of discrete orthogonal polynomials — in particular, which statement in the saturation regime of Baik–Kriecherbauer–McLaughlin–Miller [2, Thm 2.7, Remark 2.8] — supplies these two coarser facts? The target is an integer mode count and the sign of an $O(1)$ minor, not a rate.*

By Theorem 1 a proof of Problem 1 gives $R \succ \frac{1}{2}I$. The $R^\dagger$ layer then lifts to $L^\dagger$ once the terminal Schur is in hand (Proposition 2(A3),(A5), Lean-verified). The mixed identity (5) is the bordered analogue: the same Haynsworth additivity, with $J = \text{diag}(1, 1, 1/\text{den})$, applies to $M^\dagger$.

Problem 2 (Stronger saturation form). *In the notation of Definition 1, which local parametrix at a hard edge adjacent to a saturated band guarantees*

$$R^\dagger \succ \frac{1}{2}I \tag{6}$$

uniformly, equivalently that the critical 3×3 Schur block of $R^\dagger - \frac{1}{2}I$ (binding pair plus border) stays positive-definite? This is the pre-367 target: a margin asymptotic $S_N = N^{-\alpha}(M_0 + \dots)$ with relative error control. It implies Problem 1 and is strictly stronger.

The finite input to either theorem is assembled above: reciprocal weight (1); occupation duality (3); a census-grade saturated block $\{1, 2, 3\}$ with the binding pair astride the edge $3 \mid 4$; the dichotomy $\text{ind}_-(A_0) \leq 1$ on 74/74 with (P2) on the 45-row nneg-1 branch; mixed form (5); observed rate $N^{-3.332}$ (now optional); local-parametrix datum t_{geo} margin-locked in $[0.222, 0.287]$; error control in relative/projective form, with rest-block theorem only at $c = 1$. The AC/Bessel class is the wrong class [3]. A Gamma-class derivation at saturation adjacency remains a candidate for the *stronger* form and is open. Relative to that stronger form the cheaper target remains (P1)–(P2). The current externalisable architecture is the open program of Section 14.

10 The graph-resolvent form (rounds 407–412)

Rounds 407, 409 and 411 produced three equivalent finite-algebra dictionaries for the rest-block count (P1). Round 412 records them as sorry-free matrix identities in `rh/lean/RH/GraphResolvent.lean`. Nothing below asserts (P1) or $R^\dagger \succ \frac{1}{2}I$ on the family; the mincut remains the named `Prop selected_augDualResolvent_gt_half`. This is a coordinate consolidation, not a claim.

Dictionary. Let $C \succ 0$ be a real positive-definite Gram (on the campaign, the μ -only dual CD Gram on the hole set Y). The *graph resolvent* is

$$R = C(I + C)^{-1} = (I + C^{-1})^{-1} = I - (I + C)^{-1}. \quad (7)$$

The first form is the r407 intertwiner; the second is the r356 dual resolvent of the L-ensemble kernel $E = C^{-1}$ (r409 graph identity). Well-definedness, $0 \prec R \prec I$, and the spectral law $\text{eig}(R) = \lambda/(1 + \lambda)$ are finite algebra [lean-verified]. The half-shift identity

$$R - \frac{1}{2}I = \frac{1}{2}(I + C)^{-1}(C - I) \quad (8)$$

is an equality of matrices. Because $(I + C)^{-1} \succ 0$, Sylvester gives

$$\text{ind}_-(R - \frac{1}{2}I) = \text{ind}_-(C - I). \quad (9)$$

The factorization $I - C^{-1} = C^{-1}(C - I)$ is *not* a congruence. The true sandwich is $C^{-1}(C - I)C^{-1}$, equivalently the simultaneous eigenbasis, and yields the Möbius law

$$\text{ind}_-(C - I) = \text{ind}_-(I - C^{-1}). \quad (10)$$

If a rectangular factor satisfies $\mathfrak{T}^*\mathfrak{T} = C^{-1}$ (r409: the min-norm Birkhoff extension $\mathfrak{T}_0$), the energy split is

$$\|\mathfrak{T}\| = 1/\sqrt{\lambda_{\min}(C)}, \quad \|\mathfrak{T}\| \leq 1 \iff C \succeq I, \quad (11)$$

and “at most one singular value > 1 ” is $\text{ind}_-(I - \mathfrak{T}^*\mathfrak{T}) \leq 1$, which is (9)–(10) [lean-verified].

Four coordinates for (P1). The integer count (P1) is the same statement in four languages:

$$\begin{aligned} \text{ind}_-(R - \tfrac{1}{2}I) &\leq 1, & (\text{inertia}) \\ \lambda_2(C) &\geq 1, & (\text{sampling}) \\ \mathfrak{T}_0 &\text{ has at most one singular value } > 1, & (\text{contraction}) \\ C &\succeq I \text{ except in at most one direction.} & (\text{energy split}) \end{aligned}$$

The zero-defect face $C \succ I$ together with the A5 Schur $q^\dagger < 1$ lifts to $R^\dagger \succ \frac{1}{2}I$ [lean-verified, `augDualResolvent_gt_half_of_C_gt_one`]. The one-defect lift at the same depth is blocked by A7-min (a one-defect of R cannot be absorbed into $R^\dagger$ at the same size). Identifying the campaign Gram C with E^{-1} on a μ -ONB transcription is the named `Prop GraphResolventIsLEnsembleInv`, the same class as `P1EqCapInertia`; it is not a `sorry`.

Running-instance margins. On the frozen window $z = 16$ ($S = 367$, $N = 184$, $|Y| = 104$) [census, rounds 407–411]: $C_{\min} = 0.85712$, $\lambda_2(C) = 1.00018$, $r_{\min}(R) = 0.46153 = C_{\min}/(1 + C_{\min})$, $\|\mathfrak{T}_0\| = 1.08014 = 1/\sqrt{C_{\min}}$. Exact identity saturation $\|\mathfrak{T}_0\| = 1$ on already-PD windows is [refuted, round 411]: $C_{\min} - 1 = +4.57 \cdot 10^{-8}$ at $k_z = 42$, $+4.9 \cdot 10^{-9}$ at $k_z = 130$, core-42 PD in $[+1.33 \cdot 10^{-7}, +7.54 \cdot 10^{-5}]$. These numbers certify the dictionary; they do not prove (P1) off the certified head.

Table 1: Twenty-eight named closed routes (bulk and terminal), DCCXLVI–DCCXCVI. Verdicts are probe-local. No family theorem is claimed.

Language	Round	Verdict
L^∞/d_{arm} majorant	r381, DCCXLVI	too coarse
Fold-parity pairing as SATZ	r383, DCCXLIX	residual 0.512
Finite-head Chebyshev / family-uniform Z	r383/r386, DCCL–DCCLI	χ mutants
Gershgorin/Schur assist	r387, DCCLII	$k^* \sim 0.11 N$
OSC-Geronimus / Neumann R_2	r388, DCCLIII	$ \text{quad} / \text{lin} \sim 0.86$
Checkerboard / M -matrix	r394, DCCLIX	agree 0.504
Three-gap mask	r395, DCCLX	$n_{\text{uniq}} \in [8, 18]$
Isolation of the fold mask	r396, DCCLXI	pair count does not $\rightarrow 0$
Source Weyl energy decay	r399, DCCLXV	E^{bulk} grows; RH-near
Bulk FORM.T / FORM.P	r400, DCCLXVII	rank-1 / phase predictor
(P1) as construction class	r403, DCCLXVIII	REFUTED
Moments / Euler source Gram	r404, DCCLXXI	residual 209
Ones-mode / lift overflow	r404–r405, DCCLXXI–II	$\cos 0.007$; dead χ keep $\ c\ < 1$
ν -rank one / Loewner sandwich	r407, DCCLXXIII	$N_\nu = 104$
Nyquist density $\frac{1}{2}$ / $\text{diag } C \approx 1$	r408, DCCLXXV	density 0.283
Literal $\Theta = C^T \Phi_N C$	r409, DCCLXXVI	Haynsworth template, not the Schur
Unfitted Krein transport	r409, DCCLXXVI	not a (P1) proof
Testpoly / Fourier / Bernstein / SEQ	r410, DCCLXXVII	factor 90; off-fraction 0.968
Identity saturation $\ \mathfrak{T}_0\ = 1$	r411, DCCLXXVIII	$C_{\min} - 1 = +4.57 \cdot 10^{-8}$
Hole top mode v_{top}	r413, DCCLXXXII	leftover $\sigma = 1.024$
Top-mode as edge defect	r415, DCCLXXXI	$\cos 0.6766/0.058$; H PD 18/42
de Branges phase dominance $\kappa \leq 1$	r416, DCCLXXXIII	$\text{yy}_A = 3$; $\text{corr} = -0.17$
Uniform $\varphi_{bb} < 0$	r418, DCCLXXXV	vacuous overflow 6/14
Hoped $O(1)$ den-gap as SATZ	r420, DCCLXXXVII	reserve shrinks $0.105 \rightarrow 0.038$
Razor-overlap as global death coordinate	r428, DCCXCIV	vacuous-chart only
$C_{\min}$ Nyquist face	r428, DCCXCIV	$\cos = -0.007$ on VAC
Unsigned $ Z_{\text{loc}} $ triangle	r429, DCCXCVI	$[2.240, 19.245]$; slope $+0.719$
Chebyshev ψ / left–right Abel	r429, DCCXCVI	$\text{pair}_c = 1$; no k_0

Closed internal languages. Table 1 lists twenty-eight named routes closed between DC-CXLVI and DCCXCVI: the original seventeen languages for (P1), the two r409 graph-factor closures previously omitted from the count, and the r413–r429 bulk and terminal kills. Each line is a sealed probe record, not an extrapolation. Nothing in the table is a family theorem.

Lean layer. RH/GraphResolvent.lean is sorry-free. The axiom audit (RH/Audit.lean, section (m)) prints [propext, Classical.choice, Quot.sound] on the identities (7)–(11), on the four-coordinate bridge, and on the zero-defect composition with $R^\dagger$. The pilot sorry census stays five. The live remainder is still the CD identification plus the depth gap $N - 3$ versus the cap, i.e. the named mincut selected.augDualResolvent.gt.half.

11 The edge balance chain (rounds 417–425)

Rounds 417–424 produced eight finite identities for the edge Schur scalar $\text{sch} < 0$; round 425 reduced the last chain step $\gamma < 1$ to the already-named terminal ratio $q_N < 1$. Round 426 records the abstract halves in rh/lean/RH/EdgeBalance.lean. Nothing below asserts $\text{sch} < 0$ cofinally, nor $R^\dagger \succ \frac{1}{2}I$, on the unbounded family. This is a coordinate consolidation, not a claim.

Eight identities. Write $A_0 = R_{N-3} - \frac{1}{2}I$ on the hole nodes, U_{CD} the last two dual Christoffel–Darboux columns, s the Sherman–Morrison border, and $\text{den} > 0$ the border scale. The mixed 3×3 is

$$\Phi = \begin{pmatrix} K_2 & c_0 \\ c_0^T & \varphi_{bb} \end{pmatrix}, \quad K_2 = I + U_{\text{CD}}^T A_0^{-1} U_{\text{CD}}, \quad \varphi_{bb} = \text{den} - 2 + s^T A_0^{-1} s,$$

and sch is the Schur complement of K_2 in Φ . After the 2×2 Sylvester map of K_2 , write $(\tilde{a}, \tilde{b})^T = D_2^T c_0$ for the *unnormalized* couplings and $\tau^2 = \tilde{a}^2 + \tilde{b}^2$.

1. **Woodbury and the unnormalized chart** [exact, lean-verified].

$$\text{sch} = \text{den} - 2 + s^T (A_0 + U_{\text{CD}} U_{\text{CD}}^T)^{-1} s = \varphi_{bb} - c_0^T K_2^{-1} c_0.$$

The first form is the r406 discriminant Δ at $J = I$: $\text{sch} = (\text{den} - 1) - \Delta$. At $\text{den} = 1$ one has $\text{sch} < 0$ if and only if the one-defect update is positive definite. The chart of $J_2 = \text{diag}(\varepsilon_a, \varepsilon_b)$ with $\varepsilon \in \{\pm 1\}$ is the 3×3 Schur $\text{sch} = \varphi_{bb} - (\varepsilon_a \tilde{a}^2 + \varepsilon_b \tilde{b}^2)$, because $J_2^{-1} = J_2$:

$$\begin{aligned} \text{P1 } (\varepsilon = (-1, +1)) &: \text{sch} = \varphi_{bb} + \tilde{a}^2 - \tilde{b}^2, \\ \text{vacuous } (\varepsilon = (+1, +1)) &: \text{sch} = \varphi_{bb} - (\tilde{a}^2 + \tilde{b}^2), \\ \text{tot } (\varepsilon = (-1, -1)) &: \text{sch} = \varphi_{bb} + \tilde{a}^2 + \tilde{b}^2. \end{aligned}$$

Rational toys: $\text{sch} = -2/3$ (P1) and $-7/6$ (vacuous).

2. **One balance** [exact, lean-verified]. On both charts $\varphi_{bb} = c_J + \Sigma$ with $c_J = \text{den} - 2$ and $\Sigma = s^T A_0^{-1} s$. Flagship $z = 16$: $-0.0856 = -0.3989 + 0.3133$.
3. **Vacuous τ^2 -separator** [exact, lean-verified]. On the vacuous chart, $\text{sch} < 0 \iff \tau^2 > \varphi_{bb}$. In particular $\varphi_{bb} < 0$ forces $\text{sch} < 0$ for every coupling. On the $\varphi_{bb} > 0$ class this is the exact life/death cut (live overflow iff $\tau^2 > \varphi_{bb}$; dead vacuous χ_3 -23/33, χ_4 -20 iff $\tau^2 < \varphi_{bb}$).
4. **Branch τ -dynamics** [census]. Vacuous $|\tau|$ slope -1.24 (vanishes); P1 slope $+0.05$ (flat). The r417 decay was vacuous-driven. Six core overflow windows live now by $\tau^2 > \varphi_{bb}$ (margins 0.0015–0.046) and do not cross $\varphi_{bb} = 0$ (slope -0.003); they are finite exceptions.
5. **Stieltjes form of Σ** [exact]. Via the r407 dictionary $A_0^{-1} = 2(C + I)(C - I)^{-1}$,

$$\Sigma = 2\|s\|^2 + 4s^T(C - I)^{-1}s = 2 \sum_i \text{occ}_i g(\lambda_i),$$

the Stieltjes transform of the occupation measure at $z = 1$. Near-1 mass stays $O(0.01)$: n_{eff} growth is bulk broadening, not pole-creep. No elementary closed occupancy density (one-pole relative error 0.73; Cauchy two-moment 0.95).

6. **den formula** [exact, lean-verified].

$$\text{den} = 1 + \gamma - v_t \cdot s, \quad \gamma = \|b\|^2 / B_w, \quad B_w = S_{N-2} + \frac{5}{7},$$

and $\text{den} < 2 \iff \gamma < 1 + v_t \cdot s$. If $\gamma < 1$ and $v_t \cdot s \geq 0$ then $\text{den} < 2$ even at vanishing correction. Census: $\text{den} \in [1.460, 1.652]$ on all 42 (gap to 2 at least 0.348); $v_t \cdot s \in [0.061, 0.096]$.

7. **b -Parseval** [exact; identification named in Lean]. $b_k = \langle \sigma, \hat{\pi}_k^\mu \rangle$ with σ the signed border measure, so $\|b\|^2$ is the μ -side Parseval budget (w9 residual $4 \cdot 10^{-14}$; $\mathbb{Q}$ -toy $b = (3/5, 4/5)$, $\|b\|^2 = 1 < 16/9$). Termwise $w_k = b_k^2 / \rho_k$ is not uniform ($\sim 43\%$ of terms > 1).

8. **Floor and three saturation scales** [census]. Selected $a_k = 2^k$ prefers $R = -\varphi_{bb} = R_\infty + c k^{-\alpha}$ with $R_\infty \approx +0.030$ against decay-to-0 ($\Delta\text{AIC } 6.9$) and against $c/\log k$ (killed, $\Delta\text{AIC } 18.4$). EXT-vacuous sits independently in $[0.0285, 0.0373]$. Three scales, not one exponent: budget gap $2 - \text{den} \sim O(0.5)$ flat; razor $C_{\min} - 1 \sim 10^{-7}$ with $\|\mathfrak{T}_0\| - 1 = -(C_{\min} - 1)/2$ exactly (ratio 1.0000); edge reserve R of order 0.03.

Branch map. Exact separators, not a fitted partition:

branch	chart / K_2	live criterion
P1 (28/42 core)	$J_2 = \text{diag}(-1, +1)$	$\text{sch} = \varphi_{bb} + \tilde{a}^2 - \tilde{b}^2 < 0$ (hyperbola; danger axis large $ \tilde{a} $)
vacuous (14/42)	$J_2 = I_2$	$\varphi_{bb} < 0$, or $\tau^2 > \varphi_{bb}$ on the $\varphi_{bb} > 0$ overflow class
tot / dead χ	$J_2 = -I$ or $\text{sch} > 0$	vacuous death: $\varphi_{bb} > 0$ and $\tau^2 < \varphi_{bb}$; P1 death: τ -terms, $\varphi_{bb} < 0$ retained

Round 425: $\|b\|^2 \leq S$ is the kernel Loewner $K_\mu \preceq K_{\mu-\nu}$ on $P_{<N}$ [exact]; $\|b\|^2 < B_w$ is that identity plus $q_N < 1$. On the 42-rung $q_N < 1$ census this composes to $\gamma < 1$. Dead χ have $q_N > 1$ and need kernel slack; they keep $\gamma < 1$ nonetheless.

Two open limit questions. The identities above are finite. What remains, for the unbounded selected sequence $a_k = 2^k$, are two existence statements. They are the cheapest remaining targets; neither is a Riemann–Hilbert steepest-descent, and neither is a statement about the Riemann hypothesis.

Problem 3 (cofinal $q_N < 1$). *On the selected sequence, prove $q_N < 1$ (equivalently $S < B_w = S_{N-2} + \frac{5}{7}$) for all large k . Together with the Loewner identity $\|b\|^2 \leq S$ this yields $\gamma < 1$, hence $\text{den} < 2$ by the formula of item 6, even at $v_t \cdot s = 0$. The 42-rung census is already certified (v960); cofinality is the named mincut `selected_augDualResolvent_gt_half`.*

Problem 4 (constructed Σ_∞). *Construct $\text{den}_\infty < 2$ and Σ_∞ with*

$$\limsup_k \Sigma_k < 2 - \text{den}_\infty = |c_{J,\infty}|,$$

equivalently a strictly positive floor $R_\infty > 0$ for $R = -\varphi_{bb}$. The occupation Stieltjes form is exact; a closed occupancy density is refuted; near-1 mass is stable $O(0.01)$. The census prefers $R_\infty \approx 0.030$ (two independent lines: selected AIC, EXT-vacuous band). No k_0 is proved. Depth $k = 10$ border-fails computationally.

Lean layer. RH/EdgeBalance.lean is sorry-free. The axiom audit (RH/Audit.lean, section (n)) prints [propext, Classical.choice, Quot.sound] on the Woodbury-sch corollary, the chart trichotomy, the τ^2 -separator, the den formula and the γ -bridge $\|b\|^2 \leq S < B_w \Rightarrow \gamma < 1 \Rightarrow \text{den} < 2$. Named Props BorderIsMuParseval, BorderLoewnerLeS, QNLtOne are hypotheses, not holes: they name the window transcription of items 7–8 and Problem 3. The pilot sorry census stays five. Rounds 428–429 retype the cheapest remaining terminal statement on the selected sequence (Section 12); cofinality stays open.

12 The two cancellation cores (rounds 413–429)

The graph dictionary reduces the bulk hole to $\lambda_2(C) \geq 1$; the edge chain reduces the terminal hole to $q_N < 1$, equivalently $|Z_{\text{loc}}| \leq \frac{1}{2}$ on the selected sequence once COMPOSE[−] is applied. After r429 both remainders are the same kind of object: an unsigned cancellation of explicit

von Mangoldt data that has survived every majorant, structure, phase, sampling, kernel, and interpolation language of the campaign. The identities below are finite; the bounds are not. This is a consolidation, not a claim.

Bulk: window-emergent one-defect. On the flagship window $z = 16$, $C_{\min} = 0.85712$, $\lambda_2(C) = 1.00018$, $\|\mathfrak{T}_0\| = 1.08014$. Four candidate vectors for the single excess direction are named and closed:

Candidate	Round	Independent numbers
ones-mode $v \equiv 1$	r408	$ \langle v, 1 \rangle = 0.007$
centred mass difference of A_0	r400	cosine 0.0075 with the genuine negative mode
Fourier / hole-Nyquist face v_F	r410	overlap 0.386 with $C\text{-ev}_0$; $\ \mathfrak{T}_0 v_{\text{Nyq}}\ < \ \mathfrak{T}_0\ $
hole top OP v_{top}	r413	corr($v_{\text{top}}, C\text{-ev}_0$) = 0.6766; leftover $\sigma = 1.02354$; $\lambda_{\min}(Q(I - \mathfrak{T}_0^* \mathfrak{T}_0)Q) = -0.0476$

Quadrature dominance already fails for $p \equiv 1$: the mass asymmetry of a single full-carrier dual is $\sum_X u^\vee / \sum_Y u^\vee = 1.357130$ (r413; r427 route A = route B at $3.9 \cdot 10^{-12}$; a hypothetical subset- P' bug would have given 1.525 and is not live). The min-norm interpolant *does* save the constant direction, $\|\mathfrak{T}_0 v_1\| = 0.359 < 1$, and does *not* save the leftover excess. Round 415: v_{top} is not the bulk defect at depth either (cos = 0.6766 at $k_z = 9$, 0.058 at $k_z = 18$). Round 416: Hermite–Biehler interlacing holds, but phase dominance with Potapov index ≤ 1 fails ($\text{yy}_A = 3 \not\leq 1$, corr = -0.17 with the $\mathfrak{T}_0$ -index). The defect is window-emergent: no source-pure vector known before the Gram is built is the excess. Circularity verdict of r399: E^{bulk} grows as $N_w^{0.54}$ and sits at quadratic-mean; a bound forcing decay below that level is RH-near, hence circular as a strategy for $R^\dagger \succ \frac{1}{2}I$, and is in any case contradicted by the census.

Terminal: the Z_{loc} formula. On a canonical window the terminal drive splits [exact, Lean `canonical_split`]

$$Z = t_{N-2} + \text{chain}, \quad Z_{\text{loc}} = t_{\text{loc}} + \text{chain}, \quad (12)$$

with t_{loc} the hull-edge sum at geometric mask $F = 1/5$ (not the set of small prime powers) [7]. Reconstruction residual 10^{-14} . Live pins $k = 3, 4, 5$ reproduce 0.486846, 0.157211, 0.227187. Edge cardinalities on selected occupy 85–1691. The formula exhibits death of the χ mutants (1.021 and 0.892, both $> M$) and reproduces the old $k_z = 16$ razor 0.756. Unsigned triangle, Chebyshev ψ , and left–right Abel are all closed on selected: the triangle occupies [2.240, 19.245] with two-point slope +0.719; cancellation of the signed head against that triangle is a factor 0.013–0.145. Selected is not milder for majorants than the old 42-rung ladder. There is no explicit k_0 .

Selected location change. On the selected sequence $a_k = 2^k$ ($k = 3..9$, builder pin at $k = 8$) one has $q_N \in \{0.3828, 0.2143, 0.0228, 0.00176, 0.00151, [\text{pin}], 0.00070\}$, falling in k , with minimum margin to 1 equal to 0.617 against 0.0195 on the old 42-rung ladder ($k_z = 16$: $q_N = 0.9805$) [6]. The razor rungs of the old ladder are a quantifier artefact: the minimal quantifier of round 397 skips them. COMPOSE[−] premises on selected are comfortable ($R = S_F/D \leq 1.40 \ll 4$, $L_1 \leq 0.63 \ll 3$, $|Z_{\text{loc}}| \leq 0.487 < \frac{1}{2}$). The sharpest remaining sublemma is $|Z_{\text{loc}}(W_k)| \leq \frac{1}{2}$ on MAIN-selected (air 0.013 at $k = 3$, scramble-unstable at 0.524). Hoped $L_1 \leq 3/5$ fails at $k = 9$ (0.628); $2/3$ still holds as census. Honest terminal balance: identity (12) and COMPOSE[−] are theorems; $|Z_{\text{loc}}| \leq \frac{1}{2}$, $R \leq 3/2$, $L_1 \leq 2/3$ are finite censuses; $k = 8$ is a disclosed pin; $k \rightarrow \infty$ is open.

13 Audit (round 427)

Round 427 re-computed seven load-bearing campaign numbers of R397–R426 on independent code paths (r367 block $A_0 + \frac{1}{2}I$ versus dense Chebyshev–Hankel C ; full-carrier product dual; KKT $\|\mathfrak{T}_0\|$; truncated positive-definite Grams; AIC re-fit; Woodbury-sch). Search raster: truncation/depth mismatch, signed versus occupied chain mix-up, shared-builder bugs, pin circularity, float64 on 10^{-8} razors. Ausgang: **0 BUG / 2 SUSPECT / 5 CLEAN**. No round verdict of R397–R426 changes. This is an audit of recorded numbers, not a theorem of the wall [5].

Table 2: Round-427 verdict per target. Independent re-computation, then CLEAN / BUG / SUSPECT.

T	Verdict	Independent numbers
1	CLEAN	$R = C(I + C)^{-1}$ on $k_z = 9, 17, 12$; $\ R_f - R\ _{w9} = 3.38 \cdot 10^{-12}$; $\text{ind}_- = \#\{\lambda(C) < 1\} (1, 1, 0)$; mpmath dps-40 relative $\leq 1.5 \cdot 10^{-14}$.
2	CLEAN	$\sum_X u^\vee / \sum_Y u^\vee = 1.357130$ from one full-carrier dual; subset- P' would have given 1.525.
3	CLEAN	$\ \mathfrak{T}_0\ _{w9} = 1.080138437324$; $\kappa(C) = 2.086 \cdot 10^3$ so the $4.57 \cdot 10^{-8}$ razor is $56\times$ eigvalsh noise; r409 print 1.00000 is display rounding.
4	CLEAN	$K^{\mu-\nu} \succeq K^\mu$ on truncated PD Grams $n = 16$ (evmin $\sim -10^{-9}$). <i>Definiteness hypothesis</i> : inversion monotonicity holds only when the tilde form is positive definite; an indefinite $\mathbb{Q}$ -toy ($\mu = (3, 1)$ at $(0, 1)$, $\nu = (2)$ at 1, depth 2) has $\lambda_{\min}(G(\mu - \nu)) = -1.30$ and $\lambda_{\min}(K_t - K_\mu) = -2$. The campaign states the tilde chain is definite; the implication then holds. The hypothesis is part of the recorded theorem, not a silent extra.
5	SUSPECT	vs. k : M1 $R_\infty = +0.02983$ (r421 $+0.02982$); vs. N : winner M3 ($N = 5690$ at $k = 8$, $N = 1433$ at $k = 9$); $k = 8$ is a disclosed builder pin (r421→r425 circularity). Floor remains a census.
6	CLEAN	6/6 dead: $q_N > 1 \Leftrightarrow Z_{\text{loc}} _{\text{term}} \geq \sqrt{5/7} \Leftrightarrow \text{sch} > 0$; live χ_{3-9} does not fire. $C_{\min} - 1$ is <i>not</i> a death coordinate (three dead windows have $C_{\min} > 1$ and die at the edge).
7	SUSPECT	no n_{ref} /float32 in the 17 campaign probes; razor $C_{\min}$ in float64 eigvalsh; precision ward: a window with $\kappa(C) \gtrsim 10^8$ would make a 10^{-8} gap undecidable (has not occurred; $k_z = 42$ was not native-Hankel rebuilt).

The two historical artefact classes (r395 $n_{\text{ref}} = 60$; r381 signed versus occupied Fejér) are absent from the campaign probes. The r428 location change (selected margin 0.617) stands on audited numbers.

14 Current architecture (Potapov program)

After the symmetric closing line of Section 12 the remaining work is not a further lemma-first grind on $\lambda_2(C)$ or Z_{loc} . Both holes are restated as two steps of one generalised Schur process in the Krein–Langer class $\mathcal{S}_\kappa$ (κ = number of negative squares of the kernel K_Θ). Status of every statement in this section: [open program]. Nothing below is proved on the unbounded family, and nothing is a claim about the Riemann hypothesis.

Two quantifier weakenings. Elementwise extraction from a selected window does not need a strict spectral gap.

Problem 5 (Semidefinite extraction). *Prove that $R_k^\dagger \succeq \frac{1}{2}I$ (equivalently, the augmented dual form is positive semidefinite: the window pairing is nonnegative) already yields the window contribution of that k . A strict free-window margin is not required. Named Lean target: `Rdagger_ge_half_iff_augmented_posSemidef`.*

Problem 6 (Frequently selected). *Prove: $\forall K \exists k \geq K$ with $R_k^\dagger \succeq \frac{1}{2}I$ implies that the Weil quadratic form is nonnegative on every admissible test function (each test function instantiates only finitely many windows). In particular, a subsequence of positive density of good windows suffices. Named Lean targets: `weil_nonneg_of_frequently_selected`, `internal_weil_nonneg_of_frequently_selected`. The second statement is internal window-local positivity, not a claim about zeros of ζ .*

Fallback route, still [open program]: if a block average of Potapov indices satisfies

$$\frac{1}{K} \sum_{k=1}^K \kappa(\Theta_k^\dagger) < 1,$$

then some window in the block has index 0. Mean-value theorems for Dirichlet polynomials apply to this averaged index in a way the pointwise r399 majorant could not (that majorant was circular).

Bulk and edge as Schur steps.

Problem 7 (Bulk in $\mathcal{S}_{\leq 1}$). *On every selected window, construct a rational transfer matrix Θ_k with $\Theta_k \in \mathcal{S}_{\leq 1}$. The defect vector need not be known in advance: at index 1 one has $\Theta = B_\alpha^{-1}S$ with a wandering Blaschke location α_k . This is the language in which window-emergent, collective, birth-at- $k = 37$, and Nyquist-face-without-eigenvector are the same statement.*

Problem 8 (Edge in $\mathcal{S}_0$ via Redheffer). *Given Problem 7, prove $\Theta_k^\dagger \in \mathcal{S}_0$ by a Redheffer completion whose pivot is*

$$\delta_k = c\left(\frac{1}{4} - |Z_{\text{loc}}|^2\right) = c'(-\text{sch}). \quad (13)$$

The sought identity is the Schur-parameter law $\delta = 1 - |\alpha_{\text{edge}}|^2$ (energy conservation, not a triangle inequality; the 2.2–19× unsigned loss of r429 is the wrong currency).

Source Potapov completion (target form). A construction $\Theta_k = \Theta_\infty \star \prod_p \Theta_{p,k}$ (Redheffer product; elementary factors from p^r , $r \log p$, $\Lambda = \log p$; Θ_∞ the Gamma/pole/completion factor) is an open program if it satisfies all four:

- (A) Pick matrix of Θ isomorphic to $I - \mathfrak{T}_0^* \mathfrak{T}_0$;
- (B) source-pure constructor (forbidden inputs: eigenvalues of C , SVD of $\mathfrak{T}$, target Schur complement, Z_{loc} as a constructor);
- (C) $\kappa(\Theta) \leq 1$ read from the factorisation, before the target matrix is assembled;
- (D) edge Redheffer with the δ -identity (13) $\Rightarrow \kappa(\Theta^\dagger) = 0 \Rightarrow R^\dagger \succeq \frac{1}{2}I$.

The index is counted from the source factorisation, not after the target is built (the r416 de Branges route counted afterwards). Permutation is allowed to jump the index ($1 \rightarrow 20\text{--}100$) by breaking the $p^r \leftrightarrow r \log p \leftrightarrow \log p$ pairing; a χ -mutant may keep the elementary phases (bulk budget 1 survives, the edge factor dies). That pattern matches the sealed census. Round 399 is not reopened: the statement is a topological product, not an absolute-value majorant.

These four problems, together with Problems 3 and 4, are the externalisable remainder. They are not theorems, not landing pads, and not claims.

15 What a proof would give, and what it would not

A proof of Problem 1 for the unbounded family would give, in order: $R \succ \frac{1}{2}I$ by Theorem 1 [lean-verified finite algebra]; then $R^\dagger \succ \frac{1}{2}I$ once $q^\dagger < 1$, equivalently $L^\dagger$, by Proposition 2(A3),(A5) [theorem-grade and lean-verified]; $L^\dagger \Rightarrow L^* \wedge \text{Terminal}$ and $L^* \wedge \text{Terminal} \Rightarrow$ positivity of every bordered window matrix through the cap [Lean, sorry-free on the reconstruction layer, axiom audit RH/Audit.lean]. That is window-local master positivity. A proof of the stronger form (6) would give the same chain without a separate terminal input. A proof of the open program of Section 14 would give the same chain with the two quantifier weakenings (semidefinite; frequently) and without identifying the defect vector in advance. None of these is an asymptotic law, a bound on the Riemann ξ -function, or a statement about the Riemann hypothesis in either direction.

What is on the table now is a finite census (at most 85 windows for the hole kernel, 75 plus one deep row for the border, four sealed obstacle instances for the partition, 74 resolvable rows for the two-rank dichotomy, 181 windows for the Prüfer dictionary) together with exact finite-matrix identities and a sorry-free $R^\dagger$ Lean layer. No Riemann–Hilbert steepest-descent analysis has been carried out. The equilibrium-measure limit, the Gamma parametrix, and the error operator E are open, and they are no longer the cheapest target. The cheapest remaining targets are the four open-program statements of Section 14 together with Problems 3 and 4. Comparison inputs show that the identities survive source destruction while positivity and the saturated partition do not — so an existence proof for (P1)–(P2) must consume the construction, not merely the dual algebra.

Table 3 records the running instance. Every number is a sealed probe record. Numbers not in those records are omitted.

Table 3: Running instance $z = 16$ ($S = 367$, $N = 184$, pair folds $(2, 4)$). Dual quantities: round 356/359. Occupation and obstacle: round 360. Bordered quantities: round 362. Two-rank inertia: round 367.

quantity	value
$\lambda_{\min}(R)$	0.500041882
$\lambda_{\min}(R^\dagger)$	0.500041459
margin / margin †	$1.6752 \cdot 10^{-4}$ / $1.6582 \cdot 10^{-4}$
bind / bind †	1.0029 / 1.0028
cross share / share †	0.6973 / 0.6990
t_{geo}	0.2646
o_{dual} at folds 1, 2, 4	0.1497, 0.5118, 0.5137
QP dual-void block (KKT gap)	$\{1, 2, 3\}$ (+0.092)
$q^\dagger$ / border Schur	0.933044 / $1.7319 \cdot 10^{-2}$
$\text{ind}_-(A_0)$ / $\det K_2$	1 / -5.0389
$-\det K_2$ (nneg-1 family)	[1.157, 1126.389]

16 R463 Lean-fidelity repair

The r459 vitality statement is now qualified as *Python object, fidelity comparison completed only in r463*; before this repair the selected Lean object was a different raw-atom cosine fold. The selected Lean path now uses the production sequence

tent lags $\longrightarrow c_A + c_P \longrightarrow$ circulant spectral density $\longrightarrow$ grid fold and sign split, $L = 2(m+1) - 2$.

The exported $k=5,9,10$ pins reproduce the Python positive-measure Hankel entries and support/cap values. The remaining archimedean lag coefficients, v958 border, and r243 budget are

explicit typed interfaces rather than per-atom completion data.

The endpoint is now `internal_weil_nonneg_of_frequently_selected`. Its compound min-cut visibly includes `SelectedACapPsdImpliesPlainReads`. Three previously absent external links are named `sorrys`: dense extension to the full Weil class, identification with the standard explicit formula, and the standard Weil criterion against Mathlib’s `RiemannHypothesis`. Hence the honest census is $5 \rightarrow 8$. No statement in this section claims the Riemann hypothesis.

Claim boundary. Research documentation. Not a proof of L^* , of $L^\dagger$, of (P1)–(P2), of $R^\dagger \succ \frac{1}{2}I$, or of Problems 5–8 for the unbounded family. Not evidence for or against the Riemann hypothesis in either direction.

References

- [1] A. Borodin, *Duality of orthogonal polynomials on a finite set*, J. Statist. Phys. **109** (2002), 1109–1120.
- [2] J. Baik, T. Kriecherbauer, K. T.-R. McLaughlin and P. D. Miller, *Discrete Orthogonal Polynomials: Asymptotics and Applications*, Ann. of Math. Studies **164**, Princeton Univ. Press, 2007. Theorem 2.7 and Remark 2.8.
- [3] D. S. Lubinsky, *A new approach to universality at the edge of the spectrum*, Int. Math. Res. Papers (2008).
- [4] S. Hamann, *A strict polynomial subordination inequality for two explicit prime-generated discrete measures (open problem)*, TFPT research repository, `rh/problem/lstar_problem.tex`, August 2026.
- [5] S. Hamann, *Campaign audit R397–R426*, TFPT research repository, `rh/problem/campaign_audit.tex`, August 2026.
- [6] S. Hamann, *q_N reopened on the selected sequence*, TFPT research repository, `rh/problem/qn_reopened.tex`, August 2026.
- [7] S. Hamann, *The selected head bound*, TFPT research repository, `rh/problem/zloc_head.tex`, August 2026.