

TFPT — Architecture and the E_8 Compiler

The two axioms $\{c_3, g_{\text{car}}\}$, the derivation map, and the Coxeter–cyclotomic $D_5 \times A_3 \rightarrow E_8$ construction

Stefan Hamann

Alessandro Rizzo

August 16, 2026 · v5.4 (rev 642)

What this document is about

The architecture layer: how the two axioms $c_3 = \frac{1}{8\pi}$ and $g_{\text{car}} = 5$ build the Coxeter–cyclotomic compiler — the carrier $C^+ = D_5$, the family geometry $\mathbb{P}^1 \setminus \mu_4 = A_3$, the μ_4 glue $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$, the EM fixed point α^{-1} (with its ablation), and the whole number alphabet 16, 40, 41, 48, 240, 248 as carrier traces.

The TFPT document set — what is treated where

Plain language: TFPT (Topological Fixed-Point Theory) is a small discrete compiler. Two inputs — the seam constant $c_3 = \frac{1}{8\pi}$ and the carrier rank $g_{\text{car}} = 5$ — build $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ and read off the Standard-Model skeleton, the constants and the scale grammar. The series is **the introduction plus five numbered papers and five companions** (plus two standalone working notes), best read in order:

0. **introduction** — reading guide, compiler closure, predictions, the dependency DAG and the single proof ledger (`verification/status_ledger.csv`). (*entry point*)
1. **tfpt_1_architecture_e8** — the two axioms, the derivation map, the EM fixed point α^{-1} , the $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ construction, the QBL theorem chain.
2. **tfpt_2_standard_model** — the SM in one φ_0 -ladder formula, flavor from parabolic transport (sheet diamond, dual anchor, Q_+ cohomology), the worked closures.
3. **tfpt_3_e8_audit_bootstrap** — the seven E_8 slices as an audit raster, the cascade bridge, the Möbius bootstrap.
4. **tfpt_4_frontier** — honest status of η_B , m_p/m_e , Koide, dark matter, full QG.
5. **tfpt_5_redteam** — the adversarial audit: declared attacks, what survives, what each reduces to, and the kill tests.

Companions: **R.** **tfpt_research_contracts** — the open interfaces ($v_{\text{geo}}, G_{\text{net}}, F_{\text{transfer}}$) as numbered contracts; **H.** **tfpt_horizon_readouts** — Appendix H (unit-system reframe): one seam constant as the universal horizon thermal code, the Nariai/anchor surface, the resummed seam clock; **O.** **origin_theory** — why no free fundamental number remains (exact core + one honestly-typed cyclic interpretation); **S.** **tfpt_safeguards** — the verification discipline: every mechanism that defends a claim against coincidence and numerology (status calculus, no-free-pattern, the over-determination map, the fire-wall, frozen predictions + null model, the independent Wolfram and Lean paths, the red team); **P.** **tfpt_prime_front** — research documentation of the prime / zeta line: the seven load-bearing modules on the E_8 glue census (Hecke from geometry through the transport ledger), the sandbox localization program that follows them, and the kill list. Documentation, not a result: it makes no claim about the Riemann Hypothesis.

Working notes: **N1.** **note_e8_gaussian_code** — the Gaussian code bridge: E_8 over $\mathbb{Z}[i]$ via Construction A over the extended Hamming code $[8, 4, 4]$, the canonical four-bit quotient $L/(1+i)L \cong \mathbb{F}_2^4$, and the G_{31} quartic companion (exact algebra, machine-certified); **N2.** **note_hilbert_polya_truncations** — a computable, zeta-free truncation family for the Weil measure: measurements on a Hilbert–Pólya candidate (prediction-freeze methodology; no RH claim).

You are reading document 1 (Architecture).

TFPT status at a glance — read this card the same way in every document

Two inputs, one compiler. From the boundary constant $c_3 = \frac{1}{8\pi}$ (axiom P1) and the five-slot carrier $g_{\text{car}} = 5$ (axiom P2) — jointly the elementary symmetric data of one anchor $a = (1, 1, 2)$ — the discrete compiler $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ is *built*, and the Standard-Model skeleton (gauge group, three families, hypercharges, the flavor matrix), the dimensionless constants ($\alpha^{-1} = 137.0359992$) and the scale grammar are read off as fixed points. The algebraic core is machine-checked; physical readouts run through *named bridges* whose status is typed below.


What is closed, and what genuinely remains. The discrete/algebraic layer is closed ([E]); the everything-open residual reduces to *three named interface problems*:

Interface	Question	Status
v_{geo}	the one metrology unit ($= 1/\sqrt{G} = m/\mu$); No-Unit Thm: no compiler scale	primitive [O]
G_{net}	simple-current extension $(D_5)_1 \otimes (A_3)_1 \rtimes \langle (1, 1) \rangle \cong (E_8)_1$	[C]
F_{transfer}	one functor, four typed interfaces (Koide, η_B , axion, m_p/m_e)	[C]

Gravity is parameter-free. The classical metric-sector field equation is no longer only an $R+R^2$ readout: the fixed-volume entanglement equilibrium (Jacobson; Faulkner et al.), run with TFPT's atoms, gives the *full* covariant Einstein equation $G_{ab} + \Lambda g_{ab} = c_3^{-1} T_{ab}$ with *both* coefficients fixed — $c_3^{-1} = 8\pi$ (no free dimensionless Newton dial; the thermodynamic origin $2\pi/\eta$ *coincides* with the geometric one $|\mathbb{Z}_2|2\pi\chi$ via $|\mu_4| = |\mathbb{Z}_2|\chi(S^2) = 4$, so c_3 is triply over-determined — anchor, geometry, thermodynamics, v358) and Λ from α ($\rho_\Lambda = (3/4\pi^2)e^{-2\alpha^{-1}}$, v60); the Einstein tensor (not Ricci) is forced by Lovelock, so matter conservation is an output (v359). The residual here is only the equation-of-state status (an external candidate action — Bianconi's entropic action, arXiv:2408.14391 — is quantified in v473 without changing the typing) and the one unit v_{geo} ; the global measure (C7 = QG.AMB.01) is now a [C] redundancy (v369), a certification object rather than missing dynamics, conditional on SEAM.EQUIV.01 and Bisognano–Wichmann.

The seam-net structural theorem G_{net} (= SEAM.EQUIV.01: the raw seam *is* the holomorphic $(E_8)_1$ net) now carries two named routes (route split 2026-07-22, no status change in substance): the MMST route SEAM.EQUIV.MMST.01 is *closed modulo cited theorems*; the twistor route SEAM.EQUIV.TWISTOR.01 (Costello–Li, prepared by CELEST.SEAM.01) and the unconditional parent stay [O]. On the MMST route: the explicit lattice model (v367/v368) and the S_3 closure stack pin the target at every computable level — central charge $c = 8$ (v376), the $(E_8)_1$ character with 248 currents and one primary (v377), genus-one torus GSD = 1 (v378) and reflection positivity (v379) — and it is Lean-formalised as FORM.SEAM.MMST.01: the collar's MMST hypotheses are kernel-proved, the MMST scaling-limit and Adamo–Moriwaki–Tanimoto OS-reconstruction theorems enter as named cited axioms, and the #print axioms check is clean. The *only* residual of that route that stays [O] is the abstract continuum existence of the scaling limit (v336) — its 128-spinor extension leg since certified at net level by the peer-reviewed crossed-product package ($h_s=16/16=1 \in \mathbb{Z}$, the Longo–Rehren locality criterion; Böckenhauer; KLM $\mu=4/2^2=1$; AGT/AMT the independent second witness) with the realisation input reduced to invariant level and the parallel Lean route seamResidualClosed' (v469); so the MMST route is *closed modulo cited theorems*, not solved. QGEO.SYM.01 (the μ_4 deck acting geometrically) is its *corollary*: a conformal net's vacuum is rotation-invariant by axiom, so the deck follows with no extra premise (v335, Lean FORM.QGEO.BW.01). The nonperturbative quantum-gravity measure QG.AMB.01 is no longer carried as an open item: it is a [C] redundancy (v369), a certification object rather than missing dynamics, conditional on SEAM.EQUIV.01 and Bisognano–Wichmann (and TFPT is moreover rigorously gap-decoupled from the general Euclidean-QG conformal-factor problem, margin $1.648 > 0$, v76/v330). So the honest residual is the one unit v_{geo} plus the typed F_{transfer} interfaces, above the cited-theorem ceiling on the seam and with QG.AMB.01 a [C] redundancy. Everything carries a claim ID resolving to a row of verification/status_ledger.csv (the single source of truth); **if the text and the ledger ever disagree, the ledger wins.** The historical labels ($U_{\text{wall}}, G_{\text{metric}}, F_{\text{frontier}}$) are retained only for ledger continuity — the live residual is $v_{\text{geo}} \oplus F_{\text{transfer}}$, with the seam G_{net} closed modulo cited theorems on its MMST route (parent [O]) and QG.AMB.01 a [C] redundancy.

Marker key: [E] exact / machine-proven · [C] conditional (holds under named hypotheses) · [O] open / axiom · [X] falsifiable kill test. Per-claim fine type in verification/status_ledger.csv.



Contents

I	Architecture and the derivation map	6
1	How to read this document	7
2	Headline: the Pascal compiler on five carrier slots	22
2.1	$g_{\text{car}} = 5$ as the Pascal closure	23
2.2	One master moment generates 40, 41, 240, δ_2	24
2.3	Compact E_8 compression and the role of ν^c	24
3	The $D_5 \times A_3$ compiler: E_8 as a closure, not an input	24
3.1	Carrier = $D_5 = \mathfrak{so}_{10}$ (proved)	26
3.2	Family = $A_3 = \mathfrak{su}_4$ from $\mathbb{P}^1 \setminus \mu_4$ (proved, with the metric settled)	26
3.3	The μ_4 glue and the $5/4 + 3/4 = 2$ discriminant theorem (proved)	27
3.4	Glue-norm arithmetic, and three readings of 41, 60, 120	43
3.5	What the compiler outputs for the Standard Model and our reality	45
4	Grammar I — the carrier grammar (3 + 2)	46
4.1	Dual-root dictionary (everything from two charges)	46
4.2	Carrier arithmetic (proved identities)	47
5	Grammar II — the seed grammar (one decoder u)	48
6	Grammar III — the action grammar (exponentials of $\alpha_\star^{-1}$)	48
6.1	The EM closure: $\alpha_\star^{-1}$ as an explicit fixed point (made explicit here)	49
6.2	The decuple bridge (new, exact headline)	53
6.3	The full power tower: x^1, x^5, x^{10} on the carrier graph K_5	53
6.4	Hubble is the square root of Λ (corollary, exact)	53
6.5	The ~ 123 orders of the cosmological constant	54
6.6	The ladder is binomial; $2g_{\text{car}} = \binom{g_{\text{car}}}{2}$ singles out $g_{\text{car}} = 5$	54
6.7	Discriminant action theorem: the ladder reads the hypercharge discriminant	54
6.8	A two-dimensional action lattice	55

7	The Einstein normaliser as a corrected tree value	55
8	Λ-anchored metrology closure, and two anchors	55
9	The landscape: $g_{\text{car}} = 5$ is overdetermined	56
10	The E_8 carrier atlas — stated unambiguously	57
10.1	Level 1 — original E_8 data of the series ([E])	57
10.2	Level 2 — new exact counting identities ([E])	57
10.3	Level 2.5 — the even-flip transition atlas (a falsifiable program)	57
10.4	Level 2.7 — the carrier Coxeter theorem (the strongest new bridge)	58
10.5	Level 2.8 — positive-root moment: $ R^+(E_8) = \text{Tr}_{S^+} X^2 = 5!$	59
10.6	Level 2.9 — the 2·3·5 Coxeter cyclotomic compiler (meta-theorem)	59
10.7	Level 3 — conjectural geometric readings ([C])	59
11	$\mathbb{Z}_6$ monodromy, flavor and CP readouts	60
11.1	CP as triple seed variance; a closed observable quadrilateral	60
12	Mass hierarchy as a hexagon with one extra turn	61
12.1	Lepton flavor decouple: the charged-lepton determinant carries u^{10}	61
12.2	Why the top Yukawa is ≈ 1 : the $L = 0$ anchor (Froggatt–Nielsen ladder)	61
13	Inflation interface	62
14	Further simple bridges	62
15	Closures that the architecture rests on	63
16	Status summary and open audit points	63
17	Consolidated picture	65
II	The Coxeter–cyclotomic compiler and the explicit E_8 construction	67
	Theorem 1	67
	Theorem 2	68
	Theorem 3	69
	Theorem 4	69
	Theorem 5	69
	Theorem 6	70
	Theorem 7	71
	Theorem 8 — anchor microcode	75
	Master statement	75
	Appendix: explicit E_8 construction	76

III	Appendix: ablation of the EM fixed point	78
	Appendix: computational verification	79

Part I

Architecture and the derivation map

Abstract

This document develops the *architecture layer* of TFPT: the two axioms, the derivation map, and the explicit E_8 construction. The thesis is that the bridge layer of TFPT is governed by *three nested grammars*: a **carrier grammar** (from $3 + 2$ come charges, families, hypercharge, $\Omega_{\text{adm}} = 48$, $b_1 = 41/10$), a **seed grammar** (one seed $u = \varphi_0^{\text{ret}}$ generates the Cabibbo angle, birefringence, the baryon fraction and the reactor angle), and an **action grammar** (the large scale ratios are exponential actions of $\alpha_\star^{-1}$ with rungs $1/g_{\text{car}}$, 1 , 2). Two new exact results sharpen the action grammar: the *decuple bridge* $\rho_\Lambda/\bar{M}_{\text{Pl}}^4 = \frac{3}{4\pi^2} e^{2\delta_{\text{ph}}} [v_{\text{geo}}/(5\beta_{\text{rad}}^2 \bar{M}_{\text{Pl}})]^{10}$ (the cosmological constant is, in essence, the tenth power of the stripped electroweak scale, with $10 = 2g_{\text{car}}$), and the Hubble corollary $H_0/\bar{M}_{\text{Pl}} = e^{-\alpha_\star^{-1}}/(2\pi\sqrt{\Omega_\Lambda})$. We also give the exact corrected tree form of the Einstein normaliser $\xi = \frac{3}{4}/(1 + 9/128\pi^3)$, the dual-root dictionary that generates all hypercharges and transport cusps from $y_\pm = \{1/2, -1/3\}$, the Λ -anchored metrology closure, the overdetermination of $g_{\text{car}} = 5$, and the E_8 carrier atlas in three clearly separated levels. Section 3 adds the strongest structural result: the carrier code is the $D_5 = \mathfrak{so}_{10}$ half-spinor ($\text{Aut} = W(D_5)$, order 1920; $R_4 = \text{Clebsch graph}$), the family geometry $\mathbb{P}^1 \setminus \mu_4$ is an A_3 , and E_8 is the explicit unimodular glue closure of $D_5 \oplus A_3$ — with the discriminant-form norms $q(D_5) = 5/4$ and $q(A_3) = 3/4$ summing to the E_8 root norm 2 and coinciding with the TFPT constants $\delta_2/\delta_{\text{top}}^2$ and ξ_{tree} . Status markers used throughout: **[E]** exact identity, **[E]** Lie/lattice theorem, **[E]** formalised, **[E]** numerical fixed point, **[C]** physical/conditional, **[O]** axiom/open. (All markers are explicit; the legacy **[M]**/**[C]** compatibility aliases have been removed.)

1 How to read this document

Notation convention. We write $D_5 \oplus A_3$ for the *root-lattice* direct sum (used in the glue/lattice statements: discriminant forms, μ_4 glue, E_8 as an even unimodular lattice), and $D_5 \times A_3$ for the *group/representation* branching (used for subalgebra embeddings and E_8 -rep decompositions). The objects are the same atoms; the symbol marks whether the statement is lattice-theoretic ($\oplus$) or branching/representation-theoretic ($\times$).

Nothing here introduces a new primitive assumption; it reorganises existing outputs and proves the cross-links. The whole bridge layer rests on two inputs,

$$c_3 = \frac{1}{8\pi} \quad (\text{axiom P1}), \quad g_{\text{car}} = 5 \quad (\text{axiom P2}). \quad (1)$$

Foundational postulates — the two genuine axioms (hardenable, not derivable)

Everything downstream is a theorem *given* two structural inputs that are not reduced further and are honestly flagged as the theory’s axioms:

(P1) Boundary kernel.

A single oriented seam Σ carries a primitive, reflection-positive boundary kernel with unit winding $[u_\Sigma] = 1$; its self-linking normalisation fixes $c_3 = 1/(8\pi)$. *Status*: primitive postulate. *Organizing identity* (2026-08-06, markers unchanged): the closed normalization $I \cdot \beta_{\text{angle}} \cdot c_3 = 4 \cdot 2\pi \cdot \frac{1}{8\pi} = 1$ holds exactly with the Jones index $I = 4 = |\mu_4|$ a c_3 -free theorem and $\beta_{\text{angle}} = 2\pi$ independently measured — the identity *explains* the value, it does not derive it; the open bridge is the even modular-response normalization (P1.INDEX.KMS.01 [O], [verification/v813_p1_index_kms.py]). *Hardenable by*: the Calderón projector / determinant-character chain (the P1 hardening) and the Lean carrier-rigidity development (TfptCarrier.*, boundary-polarization and Calderón interfaces).

(P2) Carrier interface.

This splits into an *axiom* and a *theorem*, which must not be conflated:

- **P2_{interface} [O] (axiom)**: the seam is read out by a five-slot carrier $g_{\text{car}} = 5$ (3 colour + 2 weak) — a postulate (the readout interface).
- **P2_{algebra} [E] (theorem)**: *given* the carrier, hypercharge ($y_- = -\frac{1}{3}$, $y_+ = \frac{1}{2}$), rigidity and the family count follow from the idempotent/lattice axioms — machine-verified in Lean (Hypercharge.lean, LatticeRigidityGeneral.lean, Rigidity.lean).

So P2 is an *interface axiom* whose *algebraic consequences* are a *formalised theorem* — “axiom or theorem?” has the precise answer: interface axiom, algebra theorem.

These two are *inputs*: they can be formalised and stress-tested (more Lean, more interface theorems) but not computed away. Every other number in this note — $N_{\text{fam}} = 3$, $\Omega_{\text{adm}} = 48$, $b_1 = 41/10$, $\alpha_\star^{-1}$, the flavor matrix, the E_8 glue, A_s — is a consequence of (P1),(P2).

Anchor-first refinement. The two inputs are themselves the elementary symmetric polynomials of the single parabolic anchor $a = (1, 1, 2)$ (the exponents-at-infinity of the splitting type $\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$): $e_1(a) = 4 = |\mu_4|$, $e_2(a) = 5 = g_{\text{car}}$, $e_3(a) = 2 = |\mathbb{Z}_2|$, with $c_3 = 1/(2e_1(a)\pi) = 1/(8\pi)$. The power sums $p_n(a) = 2 + 2^n$ then give $|R(E_8)| = p_1 p_2 p_3 = 240$, $\dim E_8 = 240 + (p_4 - p_3) = 248$ and the binary ladder $p_{n+1} - p_n = 2^n$. So the input set reduces to $\{a = (1, 1, 2), \pi\}$. [verification/v23_anchor_generator.py] The whole ladder is one affine map: $p_{n+1} = 2p_n - 2$ identically, with unique fixed point $2 = |\mathbb{Z}_2|$ and the compiler quintet (4, 6, 10, 18, 34) as the T -orbit of 4 — the affine normal form of Theorem 8 ([verification/v832_anchor_flavor_checksum.py]).

The parabolic self-code of the anchor ([verification/v566_parabolic_anchor_selfcode.py]). The anchor is not only generated — it is *coded twice* by one operator structure that already lives in the compiler. The sheet-diamond direction operators $U = Q \text{diag}(1, 0, 0)$ and $V = Q \text{diag}(0, 1, 1)$ (v218; $\text{Spec } V = \{0, 1, 2\}$) generate the *full* $2|1$ parabolic (line-stabiliser) algebra $\mathcal{A}$ (dimension 7, word determinant -81 , multiplicatively closed; a perturbed Q breaks it) [E]. The Lagrange projector onto V ’s dominant eigenvalue gives $H = I + \frac{1}{2}V(V - I)$ with spectrum *exactly* $\{1, 1, 2\} = a$, and $\chi_H(x) = x^3 - 4x^2 + 5x - 2$ carries the double code $(4, 5, 2) = (|\mu_4|, g_{\text{car}}, |\mathbb{Z}_2|) = (\dim_{\mathbb{R}}(GL_3/P), \dim(\mathcal{A}/J), \dim J)$ — flag, Levi quotient and radical of $\mathcal{A}$ [E]. A uniqueness theorem closes the family: the line-stabiliser structure polynomial factorises as $(x - m)(x^2 - mx + 1)$, and all-positive-integer spectra force $m = 2$

— $a = (1, 1, 2)$ is the *only* anchor this parabolic family can code [E]. Honest typing: this is a *compression* of one operator system, not new independent evidence (no-free-pattern), and it is *not* a P^2 reduction — the route “ $g_{\text{car}} = \dim(\mathcal{A}/J)$ as a derived value” is conditional on the *No Carrier Reconstruction* audit [O]; the axioms’ markers are untouched. *Stage A of that audit is since executed* ([verification/v567_carrier_free_q_reconstruction.py]): with the v11 carrier budgets *forbidden* (rows (4, 5, 6), columns (9, 5, 1), χ_Q with its 5), the carrier-free inputs — sheet involution, winding normalisation, A_3 exponent spectrum, monodromy budget, seam-line demand — admit exactly 28 parabolic reconstructions, and **all 28 code the same anchor** $\chi_{H'} = x^3 - 4x^2 + 5x - 2$: the anchor self-code is carrier-free; the single family invariant $81 = N_{\text{fam}}^4$ then pins Q to a two-element residual family ($q_{32} \in \{2=|\mathbb{Z}_2|, 0\}$) on which every classical carrier-free certificate coincides — the one separating certificate is exactly the forbidden column budget. Verdict RECONSTRUCTS-TO-FAMILY; the audit relocates to GATE.QGEO.01 (derive the geometric inputs themselves) and to the one binary sheet-coupling selector [O] — which is since *located and given a canonical witness* ([verification/v568_bit_selector_ladder.py]): the twins generate different integral orders, and the *ladder self-consistency* (the dominant mode reproduces its own eigenvalue as the geometric ladder $(1, 2, 4) = (2^0, 2^1, 2^2)$, the binary AnchorLadder) selects the compiler Q uniquely on all 28 members; the split geometry ($q_{32} = 1$) is excluded by the demands already imposed; *why* the ladder self-consistency should hold physically stays open (the RP-selection slot) [O]. Three follow-ups since executed: the *kinematic RP form* on the twins ([verification/v572_rp_bit_form.py]) is spectrally blind (identical inertia for every declared corpus state; never PSD — no kinematic positivity selection exists) but *integrally anchor-visible*: the Smith towers of the anchor-state Gram separate the twins in one slot (18 vs 2, a factor $9 = N_{\text{fam}}^2$) — the bit shows to the RP form only through anchor-coupled states [E]; and *Stage B is narrowed* ([verification/v574_stageb_winding.py]): the involution *type* is forced by a rank theorem (a two-dim-positive involution cannot carry the binary trace code), and the winding normalisation $N_{\text{fam}} = 3$ is *forced* by the double self-consistency — the ladder selects one member for every $N \leq 3$ (higher N die on the exponent demand), and $\text{sat} = N^4$ holds only at $N = 3$ ($81 = 3^4$; $N = 1, 2$ give $32 \neq 1, 512 \neq 16$): two of the five declared geometric inputs relocate to consistency-forced; GATE.QGEO.01 is narrowed, not closed [E]/[C]. The involution-*existence* input is since *derived* ([verification/v590_involution_existence.py]): the integral V -commuting involutions are exactly the spectral sign group $(\mathbb{Z}/2)^3$ (rigidity, both twins), and the v574 type theorem plus two named demands (seam reversal, zero-mode fixing) single out the explicit S_0 with exact Σ -signature uniquely — existence is unconditional; the two demands’ continuum derivation stays the named open [E]/[C]. The *interacting* follow-up ([verification/v584_ext_source_response.py]) then separates the twins *directionally*: rank-one Ext-source coupling on the Gibbs pencil has exact rational flip thresholds $J_c(s) = -1/(v^T K(s)^{-1} v)$ whose numerators are exactly the twins’ pencil factors; in the crossed phase positive coupling $J_c \leq 15/16 < 1$ heals exactly the compiler twin (through either Ext arrow), while the alt twin cannot be flipped by any positive coupling in the open unit box — the healing direction is strictly positive, consistent with the v534 toy RP selection; the response pole sits on the golden polynomial $s^2 - s - 1$ (observed). Honest: a response-level selection, not an OS/RP positivity theorem — the alignment-bit contract stays open [E]/[C]. A *geometric candidate* for the whole GATE.QGEO realization has since emerged ([verification/v597_mu4_cover_model.py]): the μ_3 -cover $y^3 = x^4 - 1$ of $\mathbb{P}^1$ branched at μ_4 hits four compiler fingerprints at once — H_1 is a rank-3 module over the Eisenstein integers; the D_4 rotation closes projectively via the deck ($r^4 = \omega$); the cusp-class group is $(\mathbb{Z}/3)^4$ of order 81 (the saturation index, with the per-mark $\mathbb{Z}/3$ mechanism the self-code program asked for); and the unique bilinear hermitian form has Lorentz signature (1, 2) — so the RP-positivity route would become Riemann bilinear positivity if the identification completes. The dictionary since stands at *three* exact entries ([verification/v598_cover_dictionary.py]): the winding operator U is a single-cusp Gram, the μ_4 sheet coupling Q_- is the *difference of the opposite cusp Grams* (the diamond diagonal), and the binary AnchorLadder (1, 2, 4) is identity plus the diagonal sum; the grading half (Q_+ , hence V and Q) is absent from 7296 scanned canonical combinations — the named suspect for the missing generator is the anti-holomorphic D_4 reflection (the real structure); the gate does not move [E]/[C]. The suspect has since been *constructed* ([verification/v599_real_structure.py]: R unique up to scale with $R^2 = +1$; its grading functor closes the three gaps at char level, with the anchor spectrum (1, 1, 2) appearing; the invariant form has $\det J = 8$ and Lorentz signature (1, 2)), and the *joint embedding has landed* ([verification/v600_joint_embedding.py]): a Γ -even degree-2 pair (U^*, V^*) is $\mathbb{Z}[\omega]$ -integral, GL_3 -conjugate to the compiler pair (explicit g , exact), restricts to *integer* matrices on the saturated real fixed lattice, and its seven-word $\mathbb{Z}$ -order has saturation index $81 = 3^4$ with the exact v566 divisor pattern — the self-code algebra lives on the cover; canonicity stays open and the gate still does not move [E]/[C]. The canonicity question has since been answered ([verification/v601_equivariant_dual.py]):

the J-adapted cusp projectors give a presentation that is D_4 -equivariant by construction (mark choice = winding choice) and realizes the *dual* parabolic exactly (conjugate to the transposed compiler pair), and the v590 Klein four-group lives integrally on the curve $[E]/[C]$. *The cover program at a glance (v597–v605)*. Nine rounds close into one narrative: the μ_3 -cover $y^3 = x^4 - 1$ carries the fingerprints (v597), the dictionary (v598), the real structure with $R^2=1$ and the Lorentz form $\det J = 8$ (v599), the joint pair with saturation index $81 = 3^4$ (v600, Lean-certified), the D_4 -equivariant canonical presentation and the integral Klein group (v601), the J-adjoint duality, the verbatim anchor formula $H = 1 + V(V - 1)/2$ and exact Weil positivity (v602), the *real* seam line with the geometric seam-reversal demand and the first per-mark $\mathbb{Z}/3$ witness (v603), the anti-isometry $M^\dagger JM = J^T$ with the integer form J_{fix} ($\det 72$), the full four-frame order of index $27 = 3^3$ and the torsion localization (81 at the marks, 27 at infinity) (v604), and the RP translator $\theta_{\text{corpus}} = J_{\text{adj}} \circ \text{inner}(c)$ with $14c$ integral (v605). The gate does *not* move: the residual is the conformal seam identification (QGE0.SYM.01), the zero-mode selection (the zero-mode line is *not* R -real — honest negative), and the geometric identity of c $[E]/[C]/[O]$. The honest decomposition (spectrum ablation): freeing the exponent spectrum too, the one surviving double-selfconsistent family is exactly $\{1, 2, N\}$ with the compiler shape — the self-consistency forces the *sheet* spectrum $\{1, 2\}$ (the binary code) and the shape, and the A_3 exponent demand contributes exactly *one* datum, $N = 3$ $[E]$. And the redundancy ablation shrinks the input list further: relative to the self-code demands, the monodromy budget (M) and the seam-line demand (L) are *redundant* (both dropped, the class widens to 1274 members and the self-code still picks the compiler Q uniquely), and the maximal ablation drops (E) too: the winding column $3(1, 1, 1)$ plus integrality plus the self-code already pin the compiler alone. Even the winding *form* is now located: dropped entirely, the self-code leaves a four-column family of which exactly one is uniform — the compiler; the winding demand *is* the uniformity choice (1 of 4). What remains declared: the involution *existence*, the column *uniformity*, and $N_{\text{fam}} = 3$ (independently anchored in the E_8 glue, $240 = 16 \times 5 \times 3$) $[E]$. On the bit itself, the anchor-Hamiltonian *Gibbs pencil* delivers the first dynamical distinction: the compiler twin's RP kernel degenerates at a finite critical temperature $s^* = (8 - 4\sqrt{19})/15$ (determinant constant 81^2), the alternative twin's never does — and the critical mode is the seam-row, sheet-flow-fixed direction (the A_2 -Ext carrier): a rank-drop marker, not yet a positivity selection, but the resonance with the alignment bit now has a shared carrier $[E]$.

The Gaussian code bridge and the quartic half (2026-08-03) — the code section closes two loops

Two exact-arithmetic results tie the binary code layer directly to the E_8 lattice and its complex μ_4 structure. *The Gaussian code bridge* (`[verification/v689_gaussian_code_bridge.py]`, 26 checks, exact): the μ_4 complex structure J makes E_8 a rank-4 module over the Gaussian integers $\mathbb{Z}[i]$, and the quotient $E_8(\mathbb{Z}[i])/(1+i)$ is $\mathbb{F}_2^4$ — the information space of the Reed–Muller code $\text{RM}(1, 3)$ (Smith normal form exact; standard model $|\det| = 256 = 2^8$, SNF $[1^4, 2^4]$, presentation-independent). The 240 roots fall 15×16 over the fifteen nontrivial classes (the zero class provably empty), each class is 4 of the 60 G31 lines ($60 = 15 \cdot 4$), the compiler clock $\sigma = c^4$ acts as the $\text{RM}(1, 3)$ *family permutation* (3-cycle plus fixed anchor — the v638 info-bit action) with a residual gauge group of order 18, and the coordinate block is the sum-of-families label $F_1 + F_2 + F_3$. Three must-fail controls fire (naive placement, non- πJ placement, plain $\mathbb{Z}[i]^4$) $[E]$. *The quartic half* (`[verification/v690_quartic_half.py]`, 22 checks, no floats): the vanishing of the $W(E_8)$ invariants of degrees $\{2, 14, 18, 30\}$ under holomorphic μ_4 restriction is proven symbolically but *trivial* — $W(J\alpha) = -iW(\alpha)$ kills every J -orbit contribution for all $d \not\equiv 0 \pmod 4$ (honest typing); the substance is the surviving half: the restrictions $F_8|, F_{12}|, F_{20}|, F_{24}|$ are G31-invariant (60/60 mirrors), algebraically independent (exact Jacobian), with Molien-unique degrees of product $46080 = |G31|$ — by Chevalley (cited) a system of *basic* invariants, i.e. the holomorphic restriction of the $W(E_8)$ ring generates the full G31 invariant ring; the Jacobian carries the 60 mirror lines point-verified. Controls: a non-root-preserving complex structure breaks the informative half, a root-preserving one keeps it — the statement is about the compiler-compatible conjugacy class $[E]$. No marker moves; both are structure statements of the code layer, registered as E8.GAUSSCODE.01 and E8.QUARTICHALF.01. *Follow-up (2026-08-03, the Hecke tower)*: the same $\mathbb{Z}[i]$ -module structure carries a Hecke tower whose primitive degrees are *exactly* the Gaussian prime norms and whose σ -descended traces reproduce the atom layer of the Weil measure at deviation 0.00 (`[verification/v714_moonshot_hecke_groupoid.py]`, 19/19, MOONSHOT-GO; honestly fenced: the finite counting facts are E8-unspecific on that slice — the specificity question belonged to stage 2 of PRIME.Z1.MOONSHOT.01 and has since been *measured*: the archimedean place of the same tower glues with *one* normalization ($\kappa = (1, 1, 1)$ to $< 10^{-12}$; `[verification/v716_moonshot_arch_glue.py]`, 18/18, MOONSHOT-STAGE2-GLUED), and among the dimension-8 unimodular lattices $\{\mathbb{Z}^8, E_8\}$ (Mordell

1938) *only* the Gaussian E_8 glues ([verification/v719_moonshot_traceformula.py], 24/24) — E_8 becomes forced *at the gluing*, not at the finite places; the uniqueness *theorem* stays open, no marker moves). *The orbit packet lemma (2026-08-04)*: the code bridge’s number family closes as one lemma ([verification/v736_orbit_packet.py], 26/26, ORBIT-PACKET-EXACT; companion Lean module OrbitPacket.lean, 34 theorems, kernel-checked):

σ orbit census (Burnside)	$7 = (15 + 3 + 3)/3$
canonical packet split	$240 = 5 \times 48$ (1 fixed + 4 moved, 12 Gaussian lines each, label-free)
carrier reading	packets = $1 + 4 = 5 = g_{\text{car}}$, $240 = g_{\text{car}} \times 48$
anchor	leader weights $(0, 1, 1, 2) \rightarrow (1, 1, 2)$, pair e_6/e_7
$q = 16$ normal form	$240 = 16 \cdot 15$, $248 = 240 + 8$

The circularity fence is part of the lemma: P1 ($c_3 = 1/(8\pi)$) and P2 ($g_{\text{car}} = 5$) come back from the code as *exact internal reconstruction* — the packet law reconstructs the axioms, it does *not* derive them; and the $1 + 4$ law is σ -class-specific (the 160 free order-3 elements of G31 give Burnside 5, not 7). The sharpening to a canonical intertwiner with the 48 regular five-cycles is *dead* on its preregistered criterion ([verification/v737_seam48_intertwiner.py], 24/24 machinery, SEAM48-DEAD: incidence distribution 0:96/1:64/2:64/3:16 instead of $\equiv 1$) — $240 = 48 \cdot 5$ remains a factorization *without* coordinatization [E]. *The symplectic code lemma (2026-08-04)*: the quotient carries its own incidence geometry ([verification/v752_projective_hamming_incidence.py], 36/36, PROJ-HAMMING-EXACT; companion Lean module ProjectiveHamming.lean, 32 theorems): $h = H/4$ is $\mathbb{Z}[i]$ -valued with hermitian Gram determinant 1 — the *unimodular* hermitian E_8 lattice over $\mathbb{Z}[i]$ — and the reduction $\bar{h}$ makes $V = L/(1+i)L$ a symplectic $\mathbb{F}_2^4$ (doubly-even $\Rightarrow$ alternating; explicit Darboux basis); on the 15 nonzero classes the hyperplane incidence matrix satisfies $BB^T = 4I + 3J$ *exactly*, with design parameters 2/3/4/7/15/16 and eigenvalues $7^{(+2)^9}(-2)^5$ (the multiplicities (1, 9, 5) are the symplectic signature), the channel $K = B/7$ obeys $K^2 = (4/49)I + (45/49)\Pi_0$, and the NS/R parity is the point $y_0 = \text{FSUM}$ of the geometry ($\chi_{\text{NSR}} = \bar{h}(\cdot, y_0)$; $112 + 128 = 240$, $248 = 120 + 128$). The lemma also *resolves* the seam-48 kill: the distribution above is derived exactly ($16 \times$ the 3×5 label table), and the 5×48 coordinatization exists via exactly one $\bar{u}$ -invariant spread — the kill is explained (wrong blocks), not revised [E]. *The Arf/spinor compiler (2026-08-05)*: the code layer now carries its canonical quadratic refinement ([verification/v774_arf_spinor_compiler.py], 46/46, ARF-SPINOR-EXACT; companion Lean module ArfSpinorCompiler.lean, 23 kernel theorems): of the exactly 16 refinements of $\bar{h}$ the frozen selector (σ -invariance, $q(A)=1$, $q(F_\Sigma)=0$) picks a *unique* q^* , which is the parity-lift weight form under ι : $V \cong C_{\text{even}}(5)$ (crosslink $\beta = \bar{h}$ exact in all 256 cells); $\text{Stab}(q^*) \cong S_5$ makes $16 = 1 + \bar{5} + 10$ an *Arf geometry* ($Sp(4, 2) \cong S_6$; K_6 duads; $B = I + A_{KG(6,2)}$ with exact ovoid eigenvectors), and the hypercharge table $X = (-2, -2, -2, 3, 3)$ reproduces the v310/v14 dictionary with the master-moment chain 120/40/41/240/248; the same σ acts as the family 3-cycle A_3^{Fam} on message bits and as the 3+2 slot split A_3^{PS} on carrier legs — naming discipline, not a physics claim [E]. *The fired matter fence (2026-08-05)*: the preregistered kill test of the code $\rightarrow$ matter reading *fired* ([verification/v775_gaussian_class_d5_purity.py], 18/18, ROOTCLASS-MIXED): no $D_5 \oplus A_3/SU(5)$ convention makes even one full set of 15 classes pure (≥ 7 classes are adjoint-side for *every* embedding, by the 128-spinor-root counting bound); the code $\rightarrow$ matter table — including the “empty mailbox = right-handed neutrino” reading — must not enter any main claim [E]. *Boundary code uniqueness (2026-08-05)*: the non-circular selection chain exists ([verification/v776_boundary_hamming_uniqueness.py], 31/31, BOUNDARY-CODE-UNIQUE): among all doubly-even self-dual binary codes of length 8 (direct census 30, no Hamming/ E_8 input) the boundary data — four J -paired marks, $\mathbb{Z}_3$ deck pair-coherence, and *one* deck-invariant $\mathbb{Z}_2$ (the product of the four side orientations) — select exactly one code; Construction A of it *is* E_8 , and the deck packets output $g_{\text{car}} = 1 + 4 = 5$. The circularity fence above is thereby *relocated and shrunk*, not removed: the residuals named then were the doubly-even self-dual *universe premise* (R1), the semantic deck placement (R2) and the anchor naming (R3); all three moved in round 21 (see the Type-II and metaplectic entries below) — R1 is replaced by three physical axioms, R2 is placed, R3 is named; no P2-removal claim — the sharpened contract form is SEAM.CODE.TYPEII.01 [C]. *The $d = 4$ forcing census (2026-08-05)*: the glue pattern itself forces the four marks ([verification/v781_one_object_clock_census.py], 21/21 + 23/23, ONE-OBJECT-PARTIAL+CLOCK-FIBER-DEAD): the TFPT glue conditions (cyclic isotropic glue, evenness, unimodularity $|H|^2 = 4d$) admit an even unimodular *diagonal* glue of $A_{d-1} \oplus D_{d+1}$ iff $d = 4$ (exact census $d = 2..12$; at $d = 4$ exactly the two v92 Lagrangians with the explicit 240-root E_8 certificate; $d = 9$ dies at

isotropy; both relaxation controls fire) — so $g_{\text{car}} = d+1 = 5$ and $N_{\text{fam}} = d-1 = 3$ are the two functor readouts of *one* four-point boundary object. *Honesty gate*: $c_3 = 1/(2\pi d)|_{d=4} = 1/(8\pi)$ is a corpus-legitimate *rewriting* of the anchor-first refinement ($e_1(a) = 4$), not a new derivation of P1 — P1 stays a primitive postulate. The companion clock census is an honest kill: no $\mathbb{Z}_{12} \times_{\mathbb{Z}_6} \mathbb{Z}_{30}$ realization can act on the 60 Gaussian lines (full line-compatible group order 23040, maximal element order 12) [E]. *The transition-bus torsor (2026-08-05)*: the $240 = 16 \times 15$ reading is settled at its sharp level ([verification/v782_e8_transition_bus.py], 35/35, TRANSITION-BUS-TORSOR): the roots are 15 coordinate 8-frames; each class fiber is a canonical *torsor* under $v^\perp$ on $L/2L$ (no distinguished origin), the deck acts as translation by v , and the obstruction census is proof-grade — *no* deck-equivariant labeling $R(E_8) = V \times (V \setminus 0)$ exists; the frozen symplectic predicate reproduces all 57600 inner products, and the A_{15} comparison quantifies what a strict transition bus would need (a canonical start state, which E_8 provably lacks) [E]. *The two-qubit Clifford identification (2026-08-05)*: under one predeclared chart the 60 Gaussian lines are the 60 pure two-qubit stabilizer rays, entry-exactly over $\mathbb{Z}[i]$ ([verification/v783_two_qubit_clifford.py], 32/32, CLIFFORD-PARTIAL(H4)): $UG_{31}U^{-1}$ is an index-2 subgroup of the Clifford matrix group, $46080 = \mu_4 \times \text{Pauli}_{16} \times Sp(4, 2)$, classes = Pauli contexts and spreads = MUB pentads; the index bit is the Galois $\sqrt{2}$ /Hadamard bit (not $J \leftrightarrow -J$, the honest H4 partial; identified in round 21 as the $\tau = i$ metaplectic bit — see below). Carrier mathematics only; the fired matter fence above is unaffected [E]. *The curve/code 2-torsion decider (2026-08-05)*: the seam-curve realization of the code geometry is honestly *partial* ([verification/v784_curve_code_2torsion.py], 33/33, CURVE-CODE-PARTIAL): $(A[2], e_2) \cong (V, \hbar)$ as bare symplectic spaces, but the correspondence dies at two exact layers — the deck acts syntheme-side ($[3, 3]$ -free) while the code acts duad-side, and the curve's canonical theta refinement is even (Arf 0, $1+6+9$) while q^* is odd (Arf 1, $1+5+10$); joint census 0 with positive control 6 [E]. *The canonical outer bridge (2026-08-06)*: the two kill layers above are the two faces of one outer twist, and that twist is canonical ([verification/v796_duad_syntheme_bridge.py], 24/24, BRIDGE-CANONICAL): the Sylvester duality β (code odd forms $\rightarrow$ curve isotropic spreads) with $\beta\sigma = \tau\beta$ and $\beta(q^*) = S^*$ — the unique $\langle \tau, \rho \rangle$ -fixed spread — exists and is unique up to the order-6 code joint stabilizer (census $18 + 18 \rightarrow 6 + 6$, one orbit including the τ/τ^2 swap); it resolves the Arf mismatch canonically (q_Θ pulls back constantly to $q_{\text{even}}^* = q^* + \bar{h}(\cdot, A)$ — the anchor shift), transports $\bar{5}/10$ and $3+2$ exactly onto the S^* -line geometry, sends the anchor transvection t_A to the curve rotation $\rho = J$ on every census bridge, and types χ_{NSR} as an honest exact two-variant torsor. The inner identification stays dead; structure labels only [E]. *Type II from the seam axioms (2026-08-06)*: residual R1 of the boundary-code chain is replaced by three physical axioms ([verification/v799_seam_code_typeii.py], 19/19, TYPEII-FORCED): exhaustively over all 308,993 subspaces of $\mathbb{F}_2^8$ ($\dim \leq 4$), mutual locality of the Construction-A boundary vertex words $\iff$ self-orthogonal, integer conformal spin $\iff$ doubly even, and holomorphy/index one $\iff$ self-dual — the survivor set is exactly the 30 $\hat{e}_8$ copies, and no boundary-legal non-Type-II code exists; the universe premise reduces to A1 (bose statistics / trivial monodromy), A2 (modular T -alignment), A3 (holomorphic index one), each a standard physical demand; imported lattice-VOA/theta ingredients cited and machine-verified in every instance. No P2 marker move [E] + [C] (the seam satisfies A1–A3). *The metaplectic S-lift (2026-08-06)*: the missing Clifford coset is identified ([verification/v798_seam_clifford_modular_s.py], 27/27, MODULAR-S-PARTIAL): the per-binary-factor Hadamard is the metaplectic/Weil lift of modular S at its fixed point $\tau = i$ and lands exactly in $\zeta_8 G_{31}$ ($S^2 = I$, $T^2 = \text{Pauli parity}$, $(TK)^3 = \zeta_8 I$); the total Fourier $H \otimes H$ lies inside G_{31} with metaplectic anomaly $(T_3 K_3)^3 = iI = J$ — the deck is the anomaly of the total modular lift (R2 placed), and $\zeta_8 = (1+J)/\sqrt{2}$ is the real half-deck swapping the seam pair $\{L, \zeta_8 L\}$; the strict uniqueness census over all 46080 coset elements is *empty* (no Θ -real metaplectic involution fixes q^* — the Arf defect is unavoidable), one G_{31} -conjugacy class survives retyped, and the σ -orbit Arf defects XOR to the anchor A itself (R3 named, not fixed) [E]. *The torsor Fourier modes (2026-08-06)*: the origin-free fibers are read at character level ([verification/v800_e8_torsor_fourier.py], 30/30, TORSOR-FOURIER-PARTIAL): the 120 projective Fourier rays are canonical (an origin shift is a global sign), orthogonal, deck/Pauli-diagonal and G_{31} -monomial through $Sp(4, 2)$; the carrier block — the 8 R-fibers' 64 rays — factorizes as $(1+5+10) \boxtimes (1+3)$ under $W(D_5) \times W(A_3)$ with σ the family 3-cycle plus fixed anchor — the carrier Pascal at *character* level, provably impossible at point level; the honest cap: the weight and character bases are mutually unbiased ($1/8$ exactly), so the fired matter fence stands unmoved; the 45 deck-even modes are the 45 context Pauli observables under the certified chart [E]. *The jet-code normal form (2026-08-06)*: the no-origin torsor becomes a ring-theoretic theorem ([verification/v803_e8_jetcode_affine_nsr.py], 18/18+15/15, JETCODE-EXACT+AFFINE-NSR-EXACT): because 2 ramifies in $\mathbb{Z}[i]$, $\mathbb{Z}[i]/(2) \cong \mathbb{F}_2[\epsilon]/(\epsilon^2)$ (unique iso) and $W = L/2L$ is *free* of rank 4 over the dual numbers — the mod-2 root system is a first-order jet space, the deck is the jet unit $J = 1+\epsilon$,

and among all 65536 sections of $0 \rightarrow \epsilon W \rightarrow W \rightarrow V \rightarrow 0$ there are *zero* μ_4 -equivariant splittings: non-splitness *is* the no-canonical-origin obstruction; $240 = 15 \times 8 \times 2$ (address $\times$ error jet $\times$ orientation). The affine companion: $\text{Stab}(\chi_{\text{NSR}}) = \text{AGL}(3, 2)$ of order 1344 = $|\text{Aut}(H_8)|$ in the same natural 8-point action, and affine functions on the NS/R torsor reconstruct $\text{RM}(1, 3) = H_8$ — a typed internal reconstruction, not a P2 removal [E]. *The null selector (2026-08-06)*: a second, independent $d = 4$ mechanism ([verification/v807_lorentz_nullselector.py], 14/14 on this part, NULLSELECTOR-ALGEBRAIC): $|v_d|^2 = \det M_d = d(4-d)$ vanishes iff $d = 4$ on the RR/Serre functor vector, with anchor identities $i+j = N_{\text{fam}}$, $ij = |\mathbb{Z}_2|$, $i^2+j^2 = g_{\text{car}}$, $2ij = |\mu_4|$ and the bridge $g^2 - N^2 = 4d = |H|^2$ tying null to glue index on the same invariant $4d$ as the census above (the honest downgrade — no d -dependence in the seam family to vary — is typed) [E]. *The Petersen carrier machine (2026-08-06)*: the K_5 edge geometry is certified ([verification/v808_petersen_sixthroot.py], 19/19 + 16/16, PETERSEN-RADIAL-EXACT+SIXTHROOT-SPECTRAL-ONLY): edge charges $(-4, +1, +6)$ = the certified $SU(5)$ decuple, the edge graph is Petersen $\text{SRG}(10, 3, 0, 1)$, the flavor hexagon *is* its distance-2 shell edge-by-edge, and $\text{spec}(P^6) = \{1, 64/729, 1/729\}$ *is* the deployed transport spectrum exactly — a graph-geometric origin for the $\{1, (2/3)^6, (1/3)^6\}$ triple; the sixth root is spectral-only (the corpus freezes the spectrum, not the basis) [E]. [Dated correction, 2026-08-06 round 23: the sixth-root census had missed the deployed v_{221} evaluation — $T_{v_{221}}$, the exact spectral sum of v_{221} 's transfer $(1-w)^6$, equals B^6 bit-exactly, so the verdict upgrades to BITEXACT-vs- v_{221} ([verification/v814_k5_sixstep_transport.py]); the frozen SPECTRAL-ONLY stands as that probe's adjudication.] *The doily incidence identity (2026-08-06)*: the outer bridge sharpens ([verification/v809_curve_code_doily.py], 21/21, DOILY-EXACT): $NN^T = B + 2I$ on the 15×15 duad-syntheme doily, $N/3$ with singular values $\{1, 2/3, 0\}$ — the recovery base rate $2/3$ as the unique nontrivial singular value; every census bridge factors exactly $N_{\text{code}} = P_{\Delta}^T N_{\text{curve}}^T P_{\Gamma}$ (a transpose duality, not an isomorphism); the $(2/3)^6$ identification stays typed pending the six-step composition [E] [decided 2026-08-06 round 23: clock-only — see the six-step entry below]. *The cut code (2026-08-06)*: the Arf split is the cut classification of K_5 ([verification/v810_cutcode_moment.py], 17/17 + 16/16, CUTCODE-EXACT+MOMENT-INCIDENCE-PARTIAL): the cut map is injective-isometric on all 256 cells, $16 = 1 + \bar{5} + 10$ *is* $(0|5) + (1|4) + (2|3)$, Clebsch $40 = 5 + 20 + 15$ over the Arf shells with within-10 = Petersen; the $210 = \binom{10}{4}$ map is decorative per the frozen rule (the equivariant bijection killed exhaustively); Arf/cut grading, not matter semantics [E]. *The six-step composition and the T_{10} proposal (2026-08-06)*: the $(2/3)^6$ question of the doily entry is decided ([verification/v814_k5_sixstep_transport.py], 23/23, SIXSTEP-CLOCK-ONLY+TRANSPORT10-PROPOSED): the rate arises canonically on *three* spatial routes (doily alternation, hexagon compression, hexagon circulation) but each carries a typed obstruction (no unit mode; the bipartite sign $-2/3$; the wrong space), and sixth-power blindness (C_6 vs $2K_3$) makes per-step data mandatory — the deployed six is the *clock* exponent (v_{124}/v_{486} , fixed eigenbasis); the canonical proposal $T_{10} = P_{10}^6$ in the Λ^2 basis passes all gates with the equitable v_{774} quotient = $(Q_{\text{Pet}}/3)^6$ bit-exact — a proposal, no deploy claim [E]. *The Reed-Muller cascade (2026-08-06)*: the 15-label channel is a code ([verification/v819_prime_packet_rm14.py], 21/21, RM14-EXACT; Lean `PacketRM14.lean`): the row hull of B is the punctured $\text{RM}(1, 4)^* = [15, 5, 7]$ (enumerator $1 + 15z^7 + 15z^8 + z^{15}$ exact), C_0 the $[15, 4, 8]$ simplex, χ_{NSR} *one* codeword (the $7+8$ split its two sides, C_2 the affine bit), and the self-reproduction cycle $\text{RM}(1, 3) \rightarrow E_8 \rightarrow V \rightarrow \text{RM}(1, 4)^* \rightarrow \text{RM}(1, 3)$ closes with exactly 1344 equivalences; the CSS code $[[15, 1, 3]]$ with dual $[15, 10, 4]$, logical distances $(7, 3)$ and triorthogonality $8/4/2$; ζ_8 typed [H] [E]. *The canonical check rule (2026-08-06)*: the Kraus protocol set meets the dual code ([verification/v820_prime_kraus_rm24.py], 16/16 + 14/14, KRAUS-RM24-RULE-CANONICAL+PLANEFRAMES-EXACT): the literal leg-support claim is dead twice (weights $1/2/3 < \text{distance } 4$; $Sp(4, 2)$ orbits $15+90$ vs $45+60$), but the canonical ω -built rule identifies the rule-A protocol set (45 reversal pairs + the 15×4 fiber) with the 105 weight-4 checks of $[15, 10, 4]$ exactly, $Sp/\sigma/\text{deck}/\text{KMS}$ -covariant [E]. *The E_7 vacuum completion (2026-08-06)*: the check system completes ([verification/v821_prime_vacuum35.py], 22/22, VACUUM35-EXACT): $140 = 105 + 35$, Grams $22I + 6J / 6I + J / 28I + 7J$ entrywise, and the spectral completion $112 + 21 = 133 = \dim E_7$ is a *matrix identity* (the $E_7 \times A_1$ corpus reading a [C] fingerprint, anchor v_{170}); Weitzenböck $H_{105}^T H_{105} = 2B^2 + 14I$ as the canonical label-side left factor (the v_{758} continuum wall untouched); the $\pm B$ anisotropy cancels inside one Gram; $\text{RM}(2, 4)$ dual-containing [E]. *The $(1+i)$ ramification ladder (2026-08-07)*: the four documented roles of the ramified Gaussian prime $\pi_2 = 1+i$ are *one* machine-checked object with four rungs, each re-certified from scratch ([verification/v833_gaussian_ramification_ladder.py], 33/33, RAMIFICATION-LADDER-EXACT). *Rung 1, norm doubling*: $N(\pi_2) = 2$, $(\pi_2^2) = (2)$; on the rebuilt Gaussian lattice $(1+J)^T(1+J) = 2I_8$ exactly, $\min L = 4$ by census, so the zero class of $L/(1+i)L$ carries *no* root ($0/240$ directly), $240 = 15 \times 16$, each class 4 of the 60 Gaussian lines, elementary divisors $(1^4, 2^4)$, index $16 = N(\pi_2)^4$. *Rung 2*: $L/(1+i)L \cong \mathbb{F}_2^4$ with the deck acting *trivially* — the 4-bit address. *Rung 3*,

the non-split jet: $\mathbb{Z}[i]/(2) \cong \mathbb{F}_2[\epsilon]/(\epsilon^2)$ is the unique ring iso among all 24 bijections, $W = L/2L$ is free of rank 4, the deck is the jet unit $1+\epsilon$, and exactly 0 of the 65536 sections are deck-equivariant (the v803 obstruction re-proved); the 120 root pairs are exactly the $q = 1$ shell of the plus-type form (136/120, Arf 0). *Rung 4, the metaplectic lift*: $\zeta_8 = (1+i)/\sqrt{2}$, $(1+J)^2 = 2J$, $(SH)^3 = \zeta_8 I$ per binary factor and $(T_3 K_3)^3 = iI$ = the deck (v798); the exact $\mathbb{Z}[\zeta_8]$ BFS census gives $|C_2/\mu_8| = 11520 = 16 \times 720$ with zero Galois-mixed classes, hence $|C_2| = 92160$ and $C_2 = K \sqcup \zeta_8 K$ (the v783 chart theorem cited), with witness parities $\chi(H \otimes I) = 1$ (the RM-CSS ζ_8 phase bit) and $\chi(H \otimes H) = \chi(CZ) = 0$. Controls: the split ring $\mathbb{F}_2 \times \mathbb{F}_2$ admits 0 ring isos (2 *must* ramify) and the split self-extension admits 65536/65536 equivariant sections (non-splitness is contentful). One ladder — the same ramified prime read at (π_2) , (π_2^2) and $\pi_2/\sqrt{2}$; a compression of v689/v803/v783/v798/v819, no marker moves [E]. *The message ladder (2026-08-07)*: the budget lines of this paper are *one formula* ([verification/v844_message_doily_rank.py], 23/23, MESSAGE-LADDER-EXACT): $M_n = 15 \cdot \Delta p_n = 15 \cdot 2^n = (15, 30, 60, 120, 240)$ — the anchor's binary shift (v832) times the 15 nonzero classes of $V = L/(1+i)L$ (v833) — with each rung realized by a *named structure of the same rebuilt object*: $M_0 = 15$ the address space; $M_1 = 30 = h(E_8)$, derived here *from the rebuilt root system* (exactly 8 simple roots, connected simply-laced Cartan matrix with $\det = 1$ and the E_8 degree sequence, exact nonnegative integer simple-root coordinates, the highest root unique at height 29 with marks $\{2, 2, 3, 3, 4, 4, 5, 6\}$, $h = 1 + 29 = 30 = |R|/8$); $M_2 = 60$ the Gaussian root lines (μ_4 orbits, 4 per class); $M_3 = 120 = |R^+|$ (the sign pairs); $M_4 = 240 = |R|$ with the census $240 = 15 \times 16$ (zero class empty); and the coda $248 = 15 \cdot 16 + 8 = M_4 + \text{rank}$ against the actual census and the actual HNF rank. The structural doubling $60 \rightarrow 120 \rightarrow 240$ (each line exactly two sign pairs, each pair two roots) and the halving $60 \rightarrow 30 = |\mu_4|$ vs. rank are typed. Controls fire: the wrong anchor (1, 2, 2) gives (45, 105, 225, 465) — all four targets missed — and the wrong multipliers 14 (the weight-4 codeword count) and 16 (the class size) miss all four targets and break the coda [E]. *The Doily-Pascal rank theorem (2026-08-07)*: the doily incidence identity above sharpens into a *rank theorem* ([verification/v844_message_doily_rank.py], 16/16, DOILY-PASCAL-RANK-EXACT): under the K_6 duad model of the 16 quadratic refinements (Arf census $6+10$, rebuilt by brute force over 2^{16}), the duad-syntheme incidence N of the Cremona-Richmond doily carries $NN^T = B + 2I$ *entrywise* (all 225 cells) — duad collinearity *is* symplectic orthogonality, and the spectrum shift $7/2/-2 \rightarrow 9/4/0$ identifies the ovoid (-2) -eigenspace of v774 with the kernel: $\text{charpoly}(NN^T) = (x-9)(x-4)^9 x^5$ exactly, so $\text{rank } N = 10 = A_\Lambda = \binom{g_{\text{car}}}{2}$ (the decuple sector), $\dim \ker N = 5 = g_{\text{car}}$ (the five-slot carrier; the six ovoid indicators $3 \cdot \mathbf{1}_{O_q} - 1$ *span* it), multiplicity $9 = N_{\text{fam}}^2$, top singular value $3 = N_{\text{fam}}$ — *the P2 integers are the singular-value data of the Cremona-Richmond incidence*. The recovery value: $\text{sing}(N/3) = \{1, 2/3^{\times 9}, 0^{\times 5}\}$ exactly, $2/3 = (N_{\text{fam}} - 1)/N_{\text{fam}}$; the six-step exponentiation identity gives exactly $(2/3)^6 = 64/729$ as the deployed λ_2 , and the typed gap is re-affirmed with no upgrade: $1/729$ is *not* a root of the six-step charpoly — the deployed $(1/3)^6$ stays on the clock side (SIXSTEP-CLOCK-ONLY, cited). Both wrong-pairing controls (circulant, frozen-seed LCG; same row sums 3) fire with rank $13 \neq 10$ [E]. *The Arf-vacuum syndrome theorem and the code pencil (2026-08-07)*: the flat counts, orbit splits and Hecke matches of this code section compress into *one two-bit syndrome table* ([verification/v852_arf_vacuum_ovoid.py], 24/24, ARF-VACUUM-SYNDROME-EXACT). The Arf bit is a *position-independent direction bit*: for every affine flat $t+U$ of $(\mathbb{F}_2^4, h)$ the sum $a(t+U) = \sum_{x \in t+U} q^*(x) \bmod 2$ equals $h(u, v)$ of the underlying plane, for *all* 140 flats; the pair $(s_{\text{vac}}, s_{\text{Arf}}) = (\text{origin bit}, \text{direction bit})$ splits the 140 flats exactly into 15/20/45/60 — the doily lines, the non-isotropic lines, and the two off-origin plane classes, i.e. the v821 and v834 counts as rows of *one* table. The *local* Hecke theorem holds at every one of the 15 points: 3 isotropic + 4 non-isotropic directions through x , the isotropic ones in bijection with the points of $x^\perp/\langle x \rangle$, and the 28 off-origin flats through x split $12 + 16$ with $A_3 + B_3 = 28 = \sigma_3(3)$ and $A_3 - B_3 = -4 = a_3$ (the η -product Hecke eigenvalue, now a *local* statement; $28 - (-4) = 32 = 2^5$, the π_{cusp} denominator). The code side is a *pencil*: $H = [15, 11, 3]$ (weight-3 words = the 35 lines, weight-4 = the 105 off-origin flats, set equalities) $\supset \{D = [15, 10, 4], C_{\text{iso}}, C_{\text{non}} = [15, 10, 3]\} \supset K = [15, 9, 4]$, all pairwise intersections K , all pairwise sums H , syndrome fibers 4×512 ; the *canonicity lemma*: all 16 refinements differ from q^* by simplex $(= H^\perp)$ words, hence induce *one* Arf functional on H ; and puncturing/shortening $\text{RM}(2, 4) = [16, 11, 4]$ at the vacuum coordinate gives exactly H/D — *the vacuum bit is the parity bit*. Controls fire (dot form: no refinements, ill-defined direction bit; Arf-0: table invariant but labeling broken, zero set $10 \neq 6$; frozen-LCG column permutation: parameters preserved, line labeling destroyed) [E]. *The bent phase code and the two CSS codes (2026-08-07)*: the canonical q^* is *bent* ([verification/v853_bent_css_mangoldt.py], 22/22, ARF-BENT-CSS-EXACT): its phase vector satisfies $\hat{s}_q = -4s_q$ on all 16 characters ($W^2 = 16I$; perfect autocorrelation $16\delta_{a,b}$ on all 256 pairs), the zero set $\{0\} \cup \bar{5}$ is a $(16, 6, 2)$ Hadamard difference set (every nonzero shift meets it in exactly 2 points — the arithmetic face of the ovoid), and $A_q = C_{\text{iso}}^\perp = [15, 5, 6]$ carries the 31-word structural census (15 weight-8 = the characters, 10 weight-6

= the Arf-0, 6 weight-10 = the Arf-1 refinements). Two CSS codes follow: $A_q \subset C_{\text{iso}}$ self-orthogonal gives $[[15, 5, 3]]$ (distance words = the doily lines), and $A_{16} = \text{RM}(1, 4) + \langle q^* \rangle = [16, 6, 6]$ inside $C_{16} = A_{16}^\perp = [16, 10, 4]$ gives $[[16, 4, 4]]$ whose 60 minimal words are *exactly* the 60 isotropic affine planes — the $s_{\text{Arf}} = 0$ rows of the syndrome table; the *vacuum transformation* $[[15, 5, 3]] \rightarrow [[16, 4, 4]]$ is typed (one physical slot added, one logical qubit paid, distance $3 \rightarrow 4$ bought: shortening A_{16} at the vacuum slot = A_q , puncturing $C_{16} = C_{\text{iso}}$). *The -4 triple point*: the local Hecke count $12 - 16$, the bent Gauss sum $-2^{4/2}$ and the Walsh eigenvalue are the *same machine-checked integer* = a_3 ; and the 16 bent translates form a second Gram-16I frame *mutually unbiased* against the 16 characters (all 256 cross products $\pm 4 = \sqrt{16}$) — the unbiased partner of the v800 torsor Fourier rays, reported with no upgrade. Controls fire (Arf-0 flips the Walsh sign to $+4$ and kills the $(16, 6, 2)$ set; the dot form has no bent seed and its $[16, 6, 6]$ extension dies at dimension $5 \neq 6$) [E]. *The ovoid decoder (2026-08-07)*: the doily rank theorem above gains its *integral* layer ([verification/v852_arf_vacuum_ovoid.py], 23/23, OVOID-DECODER-EXACT): $N^T v_a = 0$ for all six ovoid vectors $v_a = 3 \cdot \mathbf{1}_{O_a} - 1$ (each syntheme covers each letter once — kernel facts from perfect matchings; any five of the six independent, $\sum_a v_a = 0$); the Gram is $V^T V = 36I - 6J$ entrywise; the Smith normal form is $\text{SNF}(N) = \text{diag}(1^{10}, 0^5)$ exactly over $\mathbb{Z}$ — rank 10, kernel 5, *no torsion*, with a $\det = \pm 1$ minor exhibited; the Moore–Penrose inverse has the closed form $N^+ = N^T/4 - J/36$ (all four Penrose axioms in exact $\mathbb{Q}$), the carrier projector is $P_5 = I/2 - B/4 + J/12$ with $\text{tr } P_5 = 5 = g_{\text{car}}$ and $NN^+ = I - P_5$, and the closed-form decoder $\hat{m} = (I - P_5)m$ is exact on the predeclared test set (kernel modes invisible, im N modes fully visible) — the recovery channel now has an explicit integral inverse, not just a spectrum. The kernel *is* the S_6 standard representation ($\text{tr}(P_5 \Pi(\pi)) = \text{fix}(\pi) - 1$ on all 11 conjugacy classes), and the mod-2 shadow $v_a \bmod 2 = q_a$ identifies $\ker_{\mathbb{F}_2}(N^T) = A_q = [15, 5, 6]$ — *three rings, one kernel*: the $\mathbb{R}$ ovoid span, the torsion-free $\mathbb{Z}$ kernel, and the $\mathbb{F}_2$ code of the bent layer. The wrong-pairing control fires (torsion appears, the $\mathbb{F}_2$ kernel collapses to 4) [E]. *The simplex-Fourier character theorem (2026-08-08)*: the census $240 = 15 \times 16$ (zero class empty) *is* a Fourier law ([verification/v857_simplex_fourier_winding.py], 26/26, SIMPLEX-FOURIER-EXACT + INTERTWINER-IDENTIFIED): the Walsh transform of the root multiplicity is $\hat{r} = (240, -16^{15})$, i.e. the uniform-nonzero channel $P = (J - I)/15$ with spectrum $\{1, (-1/15)^{15}\}$ and two-step contraction $1/225$ (exact in Fractions). Ramification (the zero class is empty at *every* odd theta level — one-line proof) and equidistribution (one value over the 15 nonzero classes at every level ≤ 16) force the character theorem, and the cross-machinery identity lands: the v817 prime-packet numerator *equals* the code Walsh transform at every level ≤ 16 (Eisenstein normal form $15 \Theta_{D_8} = 7 \Theta_L + 8 \Theta_0$), so $\hat{m}_2(n) = -1/15$ holds as an *integer identity at every odd $n \leq 16500$* (all 1911 odd primes covered; upgraded to *all* odd shells in round 36 — see the global shell theorem below) — the per-prime measurement was the prime restriction of a character theorem of this code. Typed honestly: the identification is *spectral* (the packet grading $(52, 60, 128)$ and the code grading 15×16 do not refine each other); controls fire, and the law is invariant under the 24 explicit isometries and the other invariant placement C^{**} [E]. *The G31 clock alphabet (2026-08-08)*: the compiler clock alphabet is a *normal form* of G_{31} ([verification/v858_g31_clock_alphabet.py], 19/19, CLOCK-ALPHABET-SELECTIVE): $\text{Deg}(G_{31})/4 = \{2, 3, 5, 6\} = \{|Z_2|, N_{\text{fam}}, g_{\text{car}}, |R^+(A_3)|\}$ with $\text{gcd} = 4 = |\mu_4|$, $\text{lcm} = 120 = |R^+(E_8)|$, $\text{lcm}/\text{gcd} = 30 = h(E_8)$, and the audit identity $(\sum d/4)/(\sum (d-1)/4) = 16 \times 15 = 240$ *realized* on the roots (the 60 reflections are the 60 Gaussian lines, 4 per class — the same 15×16 census as the code side); $(8, 12, 20, 24)$ is the *unique* integer quadruple with product 46080 and sum 64; the kill scan over all 607 rank-4 Shephard–Todd groups leaves G_{31} the sole full-battery passer (weak-triple impostors and raw-sum-64 coincidences typed by name; honesty line: degree-only audits cannot see the reducible torus), and the $W(D_5) \times W(A_3)$ order coincidence is re-killed by computed centers, $|Z(G_{31})| = 4 = \langle J \rangle$ vs 1 (the structural refutation cited) [E]. *The global Gaussian shell theorem (2026-08-08, round 36)*: the $-1/15$ census law above is now a *theorem for all shells* ([verification/v875_shell_global_lean.py], 15/15, SHELL-GLOBAL-THEOREM): on $V = L/(1+i)L$, for every $n \geq 1$, $\Theta_v(n) = (\Theta_L(n) - \Theta_0(n))/15$ for all $v \neq 0$ (the division exact), $\hat{\Theta}_u(n) = 16(16 \cdot \mathbf{1}_{2|n} \sigma_3(n/2) - \sigma_3(n))$, and at every *odd n* : $\hat{\Theta}_u(n)/\Theta_L(n) = -1/15$ *exactly* — no census cap. The three inputs, each warded: the G_{31} class action is the 60 symplectic transvections generating $\text{Sp}(4, 2)$ (order 720), *transitive* on the 15 nonzero classes and identical to the shell-module action; norm doubling by $(1+i)$ ($\Theta_0(2m) = \Theta_L(m)$, $\Theta_0(\text{odd}) = 0$, exact from $2 = -i(1+i)^2$); and $\Theta_L = 240 \sigma_3$ (classical Eisenstein E_4 ; the $n \leq 16500$ census retained as the exact ward). A non-transitive control subgroup admits an invariant *non-uniform* census (the law breaks — transitivity is load-bearing), and the even-shell formula differs correctly ($7/135$ at $n = 2$). The consequence chain is kernel-checked from the named hypothesis bundle in `TfptCarrier/GaussianShells.lean` (7 theorems, exact integer identities, standard axioms, no `sorry`) [E]. *The winding decoder of the clock alphabet (2026-08-08, round 36)*: the Paper-2 winding quadratic $q_{\text{wind}}(t) = t^2 - g_{\text{car}}t + |Z_2|$ *decodes* the clock alphabet

(`[verification/v874_feedback_decoder_lean.py]`, 11/11, WINDING-DECODER-EXACT): $q(\{2, 3, 5, 6\}) = (-|\mu_4|, -|\mu_4|, |Z_2|, h(D_5)) = (-4, -4, 2, 8)$, with $q(2) = q(3)$ forced by the palindrome axis $g_{\text{car}}/2$ and $2 + 3 = 5 = g_{\text{car}}$; the decoder polynomial $D(y) = (y+4)^2(y-2)(y-8) = y^4 - 2y^3 - 48y^2 - 32y + 256$ carries the budget coefficients ($|Z_2|$, $\Omega_{\text{adm}} = 48$, $2^{g_{\text{car}}} = 32$, $2^{\text{rank}(E_8)} = 256$); and the Fourier bridge lands: $|\mu_4| q(|Z_2|)/|R(E_8)| = 4 \times (-4)/240 = -1/15$ is *exactly* the nonzero-character Walsh eigenvalue of the census above ($-16 = |\mu_4| q(|Z_2|) = \hat{r}(u \neq 0)$). Sibling deformation directions collapse the bridge and alternative alphabets change the decoder; honest lines typed: the -32 coefficient alone is not selective (the selective object is the full vector), and the 6-vs-7 tension carries over verbatim. Type: exact *audit* theorem, not a physical functor — an explicit intertwiner would be needed to make it causal [E]. *The divisor lattice of 210, with its canonicity guard (2026-08-08)*: the label register $\mathbb{F}_2^4$ is the divisor lattice of $210 = 2 \cdot 3 \cdot 5 \cdot 7$ — the Walsh–Hadamard transform carries the lattice Möbius function ($4 \text{ WH}[1111, v] = \mu(d(v)) = (-1)^{\text{wt}(v)}$ on all 16 labels), and the vacuum column is the rank-one Redheffer closure on the register’s own lattice, with the matrix-determinant lemma giving the Euler products *exactly* (half-weighted $\det = \prod_p (1 - 1/p) = 8/35$; unweighted $= \prod_p (1 - p^{-1/2})$, exact radical maps) (`[verification/v863_redheffer_colligation.py]`, `sympy-exact`). The hard canonicity guard then quantifies what selects the primes (`[verification/v868_divisor210_audits.py]`, 36/36, DIVISOR210-GAUGE-FAMILY(2)): the Boolean/Walsh/ μ layer is *measured generic* (210/210 quadruples of primes < 30 pass it — layer one carries no selection content, exactly as warned), the two deployed Euler determinants pin $\{2, 3, 5, 7\}$ *uniquely* in the predeclared field (1 match each; non-matches separated by $\geq 5.2 \times 10^{-3}$ at 50 digits; the tautological $\prod p = 210$ test excluded from the verdict), the anchor prime 2 is *forced* by ramification (the unique $\mathbb{Z}[i]$ -ramified prime onto the unique σ -fixed $q^*=1$ class — the very prime at which $V = L/(1+i)L$ is cut), the μ_4 weld grade cuts the gauge $C_6 \rightarrow C_3$, and after the ramification crossref exactly *two* gauge classes remain — the family-cycle chirality $(3, 5, 7)$ vs $(3, 7, 5)$, which is then *proven gauge* by two exact no-go wards (15/15, CHIRALITY-DEGENERATE: the register orientation functional vanishes identically — the Möbius complement $d \mapsto 210/d$ pairs the orbits with reversed orientation — and quadratic readouts are provably chirality-blind by transposition). The Gram-only baseline is a simply transitive 720-torsor cut to 6 ordered bases by the deployed pins — the semantics, not the lattice shape, carry the selection; the 6-vs-7 honesty note stands untouched (the clock alphabet $\{2, 3, 5, 6\}$ and the quadruple $\{2, 3, 5, 7\}$ are different objects: $6 = 2 \cdot 3$ sits *inside* the lattice as the unique weight-2 divisor from two alphabet primes, 7 is prime and not in the alphabet); and the $17 = 12+3+1+1$ moving-sector identification is honestly *buried at space level* (12/12, MOVING12-DIMENSION-ONLY: the count is exact, the principal angles against the only deployed bridge are $\sim 90^\circ$) [E]. *The Redheffer/Mertens Smith echo (2026-08-08)*: the register’s vacuum-completion pattern has an exact arithmetic instantiation (`[verification/v861_krein_normalform_lean.py]`, 37/37, REDHEFFER-ECHO-EXACT): $\det R_n = M(n)$ (Bareiss integer determinants on 62 sizes; the matrix-determinant lemma extends the identity along all $n \leq 2 \times 10^6$), the witness minor is 1 exactly, so $\text{SNF}(R_n) = (1, \dots, 1, |M(n)|)$ — vacuum completion is a rank-one origin update whose *entire* Smith deviation is one new invariant factor; the mod-2 shadow is the squarefree-count parity; and the code-side instantiation is the vacuum transformation $[[15, 5, 3]] \rightarrow [[16, 4, 4]]$ above, typed at parameter grade (pattern-grade cross-domain identification — no functor claimed). The same module types the register wiring: the μ -sign and the deck are *two separate* C_2 s (the deployed reading character separates them; the identification is one faithful- μ_4 character away, three candidate constructions machine-verified — diagonal characters, the Galois bookkeeping $\mu_K(p) = \chi_4(p)$ on $609 + 619$ primes, the Hall chain parity), and the 2-torsion weld law explains the refusal structurally (`[verification/v862_defect_polar_weld.py]`, 22/22, WELD-WITHOUT-CONTRACTOR: the deck’s anticommutant is entirely faithful- μ_4 — the deck enters a Clifford weld only as the quarter turn $J = MD$, never as a second involution; all four weld candidates close exactly on their forced domains, the unwelded control refuses exactly) [E].

The finite compiler normal form $C_{\text{fin}} = (V, \hbar, q^*, \sigma, \iota)$ (2026-08-07) — one object, the kill built in [E] (`[verification/v845_cfin_normal_form.py]`)

The four-bit symplectic compiler that this paper assembles piecewise — quotient census, Arf refinements, selector, parity lift, S_5 geometry, Pascal counting — is now *one* machine-checked object, read off the Gaussian quotient and not postulated (28/28, NORMAL-FORM-ASSEMBLED, SPEC v2 declared): $V = L/(1+i)L$ (census 240 = 15×16 , zero class empty); $\hbar$ the descended symplectic form with a *found* family/anchor basis (F_1, F_2, F_3, A) of Gram $J - I$, equal to the bit form on all 256 label pairs; σ the descended order-3 deck ($\sigma^3 = \text{id}$, symplectic, exactly three fixed nonzero classes $\{A, F_\Sigma, F_{\Sigma+A}\}$ and four free 3-orbits — the family cycle); q^* the *unique* selector-picked quadratic refinement (exactly 16

refinements by brute force over 2^{16} , Arf census $6 + 10$; σ -invariant: $4 \rightarrow q(A)=1$: $2 \rightarrow q(F_\Sigma)=0$: 1; Arf type 1, σ -preserved); ι the parity lift, bijective onto the even five-slot code $C_{\text{even}}(5)$ with $\beta = \hbar$ on all 256 cells, $q_{\text{wt}} = q^*$, and σ acting on the five carrier slots as the 3+2 split — message register and carrier register intertwined. *Derived from the one object*: $|Sp(4, 2)| = 720$ by full census, $\text{Stab}(q^*) = 120$ acting *faithfully* as S_5 on the five, orbits $[1, 5, 10]$; $16 = 1 + 5 + 10$ and $15 = 5 + 10$; the Pascal reading (the ι -weight census $(1, 10, 5) = (\binom{5}{0}, \binom{5}{2}, \binom{5}{4})$); the projective reading (the $\bar{5}$ is an *ovoid* of the doily in $PG(3, 2)$: it meets every isotropic line exactly once); and the budget against the actual lattice objects: $240 = 16 \cdot 15$, 60 lines $= 4 \cdot 15$, $30 = 2 \cdot 15 = h(E_8)$ (rung 1 of the message ladder), rank 8, $248 = 16 \cdot 15 + 8$. *The kill built in* (mandatory negative, v775 re-anchored): the exact counting bound — every $D_5 \oplus A_3 \subset E_8$ has 128 spinor-side roots, so at most $\lfloor 128/16 \rfloor = 8$ all-spinor classes and at least 7 of the 15 classes *always* contain adjoint-side roots; the concrete doubled-model census *saturates* both bounds ($8 + 7$, zero mixed classes — the J -quotient respects the D_8 grading), and the weight-level kill at the canonical split shows each spinor-pure class restricting to 8 distinct half-spinor weight pairs, never one — the 15 mailboxes are *not* 15 matter multiplets; ROOTCLASS-MIXED re-verified, not overturned. *What the normal form does not remove*: all 30 placements are Type II $[8, 4]$ codes (the v799 echo), and C_{fin} does *not* derive the three seam axioms, the OPE-linearity scope note, or the v776 selection of one copy — P2 ($g_{\text{car}} = 5$, entering as the ι slot count) is *narrowed* to the v799 residual R1' ([C] class), not eliminated. Controls fire (the non-alternating dot form admits zero refinements; the trivial deck leaves four selector candidates; the wrong lift breaks $\beta = \hbar$). A normal form, not a new claim; no marker moves [E].

The uniqueness of C_{fin} and $\text{Aut}(C_{\text{fin}}) \cong C_6$ (2026-08-07) — one orbit, six automorphisms [E]
 ([verification/v849_cfin_unique_cofinal_lean.py])

The normal form above is not merely assembled — it is *unique* (22/22, CFIN-UNIQUE, exhaustive finite verification over $\mathbb{F}_2^4$). The WLOG ladder is exact: the 28 nondegenerate alternating forms are one $GL(4, 2)$ orbit; $Sp(4, 2)$ (order 720 by full enumeration) is transitive on the 6 Arf-1 refinements; per refinement there are exactly 20 admissible σ (one orbit under the order-120 stabilizer) and per (q^*, σ) exactly 120 admissible equivariant ι (one orbit). *The uniqueness theorem*: the full admissible tuple census at $\omega = \hbar$ has 14400 members and the $Sp(4, 2) \times S_5$ action has *exactly one* orbit (orbit-stabilizer verified: $14400 \times 6 = 86400 = |G|$) — every admissible compiler object is isomorphic to C_{fin} . *The automorphisms*: $|\text{Aut}(C_{\text{fin}})| = 6$ with element orders $[1, 2, 3, 3, 6, 6]$, abelian and cyclic — $\text{Aut}(C_{\text{fin}}) \cong C_6$, the projection to $Sp(4, 2)$ faithful and the S_5 slot permutation *determined* by the symplectic part. The Gaussian-reduction diagram is typed: the parent lattice fixes the carrier triple $(V, \hbar, \sigma)$ up to exactly the 18-element centralizer torsor, the selector pins q^* uniquely, the 120 ι -completions form a single orbit — and the *strict* initial/terminal reading fails honestly (120 completions against 6 automorphisms: unique up to *non-unique* isomorphism, a one-object groupoid, not a terminal object). Controls fire: dropping alternation kills the object (0 refinements); the wrong Arf class kills ι (the $C_{\text{even}}(5)$ weight census $(1, 10, 5)$ *forces* Arf 1); dropping σ kills *canonicity* (the completion census becomes $16 \rightarrow 8 \rightarrow 4 \neq 1$) while the bare (q, ι) orbit count survives — σ is load-bearing for the selector, typed either way. P2 sharpens accordingly: the interface object is unique up to isomorphism with C_6 automorphisms; what remains of P2 is unchanged — the v799 residual R1' (why *this* normal form at all). No marker moves [E].

The four finite anchor theorems (2026-08-09) — the census-free q^* , the exclusive anchor classification, the doily as a multiplication table, and the $g = 5$ exhaustion [E]
 ([verification/v880_finite_anchor_theorems.py])

Four theorems close the finite-anchor layer of this section at identity grade (26/26, FINITE-ANCHOR-CLOSED; the discovery probe embedded byte-exact). **(1) The census-free normal form.** The selector-picked refinement has the closed weight-only form

$$q^*(x) = \sum_{i < j} x_i x_j + \sum_i x_i = \binom{|x| + 1}{2} \bmod 2,$$

value table $(0, 1, 1, 0, 0)$ on weights 0–4 — and the v845 selector *forces* the linear part $(1, 1, 1, 1)$: the 2^{16} brute-force search of the normal-form keybox above is replaced by a theorem. **(2) The exclusive anchor classification.** Every line of $PG(3, 2)$ is classified by its q^* -weight multiset, and the classification is *exclusive*:

lines	count	weight multiset	anchor
doily (isotropic)	15	$\{2, 1, 1\}$	$a = (1, 1, 2)$
secants	10	$\{2, 2, 1\}$	the <i>wrong</i> anchor $(1, 2, 2)$
external	10	$\{1, 1, 1\}$	—

All 15 doily lines carry the anchor $a = (1, 1, 2)$ with per-line characteristic polynomial $(t - 2)(t - 1)^2 = t^3 - 4t^2 + 5t - 2$ and Vieta data $(e_1, e_2, e_3) = (4, 5, 2)$ — $e_2 = g_{\text{car}}$ and $e_3 = |Z_2|$ read off one line; the 20 non-isotropic lines split *exactly* as 10 secants carrying the wrong anchor $(1, 2, 2)$ (Vieta $(5, 8, 4)$ — the v844 control multiset) plus 10 external lines $\{1, 1, 1\}$. The incidence budget lands as counting: $15 = M_0$ and $30 = M_1 = h(E_8)$ (counting only, no new structure claimed). The anchor–Vieta report closes the loop: $p_2(a) = 6$, $\chi_L - \chi_R = -p_2(a)(t^2 - e_2t + e_3)$, and $M_n = 15(1, e_3, e_1, e_1e_3, e_1^2)$. **(3) The doily is a multiplication table.** The doily *is* the multiplication table $\Lambda^2 E \times \Lambda^2 E \rightarrow \Lambda^4 E$ of the five-slot carrier: each doily line has support partition $\{i\} \cup \{j, k\} \cup \{l, m\}$, the wedge $b_{jk} \wedge b_{lm} = \pm f_i$ holds exactly, the assignment is a bijection onto all $5 \times 3 = 15$ such configurations, and the wedge is nonzero *exactly* on the doily lines. **(4) The $g = 5$ exhaustion.** The quadratic one-step exhaustion $\Lambda^2 \wedge \Lambda^2 = \Lambda^{g-1}$ *forces* $g_{\text{car}} = 5$ uniquely among odd $g \in \{3, \dots, 11\}$. The honest split is explicit and load-bearing: the *theorem* (exhaustion $\Rightarrow g = 5$) is [E]; the *physical premise* (why the carrier must satisfy the quadratic one-step exhaustion — the Calderón reading) stays [O] — P2 is *narrowed*, not eliminated; no marker moves.

The finite Pfaffian/ K_6 signature (2026-08-09, rounds 44–49) — the $Gr(2, 5)$ Plücker reading, the canonical sign character, the self-hosting counting theorem, the flavor hexagon as the automorphism orbit, and the vacuum selection [E]
 ([verification/v888_finite_pfaffian_signature.py])

Rounds 44–49 close the finite Pfaffian/ K_6 layer of the compiler at identity grade $(27/27 + 19/19 + 42/42)$; verdicts S6PLUCKER-CLOSED, C6FLAVOR-MEASURED + INTERTWINED, K6PFAFFIAN-CLOSED; three discovery probes embedded byte-exact; exact integer / Fraction / sympy arithmetic in every decision — no floats, no RNG, no fits). **(1) The $Gr(2, 5)$ reading.** The 15 doily lines *are* the 15 monomials of the five Plücker quadrics of $Gr(2, 5)$: the bijection is 15/15 with *perfect* sign match — every quadric reads $(+, -, +)$, the v880 Hodge table (converse warned: secants/externals produce no monomial). The S_6 dual representation decomposes as $15 = 1 + 5 + 9$ (Specht $S^{(6)} + S^{(5,1)} + S^{(4,2)}$, characters from scratch by Murnaghan–Nakayama) and the normalized doily operator acts on the isotypics by the exact scalars $(1, 0, 2/3)$, with the anchor identities $2/3 = e_3/(e_2 - e_3)$ and $15 = 1 + e_2 + (e_2 - e_3)^2$ at $(e_1, e_2, e_3) = (4, 5, 2)$; and the q^* zero set is a $(16, 6, 2)$ Hadamard difference set with the anchor signature $(v, k, \lambda, v - k) = (2^{e_1}, p_2(a), e_3, \binom{e_2}{2}) = (16, 6, 2, 10)$. **(2) The K_6 motif theorem.** Under the dual labeling the 35 lines of $PG(3, 2)$ are *exactly* the 15 perfect matchings of K_6 (the doily) plus the 10 triangles through the vacuum vertex (the secants — the wrong-anchor class) plus the 10 triangles avoiding it (the external lines, 20 triangles exhausting $\binom{6}{3}$), and the edge law is $q^*(e) = 0$ iff e touches the vacuum vertex — q^* *is* vacuum incidence. **(3) The canonical sign character.** The fermionic Pfaffian signs of the symbolic 6×6 expansion (15 monomials, coefficients ± 1) are gauge-matched to the Hodge table via the canonical global character $\chi(i) = (-1)^{i+1}$ on the five vacuum edges — triple-identified as the vacuum-row Laplace sign of $\text{Pf}(A) = \sum_i (-1)^{i+1} a_{0i} \text{Pf}(A_{\hat{0}i})$, the Λ^4 -vs- Λ^5 volume-form conversion (verified on all 15), and the C_6 crossing character $(\text{Pf}(PAP^T) = \det(P) \text{Pf}(A)$ symbolically) — a global gauge, not a line-dependent fudge (the flipped- χ control produces exactly the 3 expected mismatches); each 4×4 minor Pfaffian *equals* the corresponding $Gr(2, 5)$ Plücker quadric. **(4) The self-hosting counting theorem.** The pairing count $\binom{g+1}{2} = g!!$ has exactly the solutions $\{1, 5\}$ over odd g ($g = 1$ trivial, stated; growth separation $g!! > \binom{g+1}{2}$ for all odd $g \geq 7$ by symbolic induction), so $g = 5$ is the *unique* nontrivial fixed point, with $(g - 2)!! = 3 = N_{\text{fam}}$. The honest split is explicit and load-bearing: the counting *theorem* is [E]; the *physical premise* (the boundary grammar is a self-hosting Wick pair compiler) stays [O] — P2 is narrowed to *one* premise, not eliminated; no marker moves. **(5) The flavor hexagon is the automorphism orbit.** $\text{Aut}(C_{\text{fin}}) \cong C_6$ is rebuilt exactly; the pinned generator g satisfies $g^2 = \sigma$ (the deployed family 3-cycle) *exactly* and g^3 is the W -edge sheet swap; its action on the ten pair-messages has cycle type $1 + 3 + 6$ with orbits equal to the charge shells, the g -orbit of $\{f_1, w_1\}$ traverses the deployed flavor hexagon edge by edge (D_6 -equivalent to the transport cycle), and the recovery chain reads $(|Z_2|/N_{\text{fam}})^{|\text{Aut}(C_{\text{fin}})|} = (2/3)^6 = 64/729$ — the frozen deployed transfer eigenvalue as an exact Fraction (typed honestly as a consistent *new reading*: the corpus derivation of the exponent via $p_2 = |R_+(A_3)|$ stands; 0 of 600 symplectic non-automorphisms factor, scrambled transport fails).

(6) The vacuum-selection theorem. $Sp(4, 2) \cong S_6$ acts transitively on the six Arf-1 refinements; $\text{Stab}(q^*)$ has order 120 and acts faithfully *and* surjectively as S_5 on the remaining five, with orbits on the 16 messages exactly the ι -weight classes $1 + 5 + 10 = (\Lambda^0, \Lambda^4, \Lambda^2)$ — the five-slot carrier is the residual symmetry after choosing one of six equivalent vacua. Two side reports: the anchor power-sum clock $p_n = 2 + 2^n$ has mod-30 orbit $4 \rightarrow 6 \rightarrow 10 \rightarrow 18$ of period exactly $4 = |\mu_4|$, typed CLOCK-DISTINGUISHED (the purely period-4 moduli are exactly $\{5, 10, 15, 30\}$, all dividing 30); the operative-Coxeter-30 reading is separately dead (sandbox, both sides). All controls fired (Arf-0 refinements, secant substitution, wrong scalar, flipped sign character, $g = 3/7$ counterexamples). No marker moves [E].

The Wick functor arc (2026-08-09, rounds 50–55) — the six compiler roles in the deployed seam, the proved scalar obstruction, the block resolution, the seam-diagonal demand, and the constructed channel-mixing candidate [E]/[O] ([verification/v896_wick_block_functor.py], [verification/v898_kms_schur_mixing.py])

Rounds 50–53 take the one remaining P2 premise of the keybox above — the physical Wick-compiler reading, [O] — and close its *structure* level ($27/27 + 33/33 + 26/26$; verdicts WICKFUNCTOR-PARTIAL, WICKMIX-MEASURED (WICKMIX-OBSTRUCTED), WICKBLOCK-MEASURED (BLOCKFUNCTOR-CONSTRUCTED / SEAM-DIAGONAL); three discovery probes embedded byte-exact; exact integer / rational / symbolic checks in every decision typed PROVED or SYMBOLIC). **(1) The six roles exist.** The compiler roles are present in the deployed seam one-particle space — five D_5 carrier slot pairs plus the A_3 boundary block (dims $2, 2, 2, 2, 2, 6$ from the v155/v156 split $16 = 10 + 6$); the $\mathfrak{so}(16)$ sector law holds (current-sector dims 12 on vacuum edges / 4 on pair edges = the q^* edge grading, ratio $3 = N_{\text{fam}}$); and the C_6 intertwining is *exact*: the $\text{Aut}(C_{\text{fin}})$ generator g ($g^2 = \sigma$) induces the channel permutation of cycle type (1)(2)(3) fixing channel 0. But the deployed *vacuum kernel* is channel-diagonal, so all 15 duad two-point functions vanish (DUADMAP-DEGENERATE-ON-VACUUM): the structure level of the functor exists, the value level needs a channel-mixing covariance — WICKFUNCTOR-PARTIAL, the honest typing (the candidate is constructed in round 55, item (5) below). **(2) The scalar obstruction is proved.** On the 6-dimensional scalar channel space *every* C_6 -covariant antisymmetric covariance vanishes on the transposed duad — the C_6 2-cycle $\{4, 5\}$ forces a duad zero, so the scalar route reaches at most 14/15 nonzero duads, never 15/15: a *theorem of the deployed action*, exhaustively verified over the isotypic decomposition (the multiplicity list $[3, 0, 1, 1, 1, 0]$ re-derived by hand and by exact root-of-unity sums; the frozen expected list carried a transcription error, caught fail-first); no equivariant scalarization rescues the scalar space, and both relaxation witnesses are constructed. **(3) The block resolution.** The full block-valued functor on the 16-dim one-particle space is BLOCKFUNCTOR-CONSTRUCTED — block commutant dimension 33; all 15/15 duad cross-blocks nonzero; all 15/15 block Wick monomials obey the canonical sign law $\chi(i) = (-1)^{i+1}$ (the round-49 gauge of the keybox above); the interleaving character +1 *symbolically*; the review’s five conditions hold at block level. **(4) The seam-diagonal demand.** The decisive fact: the *deployed* seam covariance (the v113 family rebuilt exactly) has *zero* cross-blocks everywhere (SEAM-DIAGONAL) — the value level of the reality bridge needs a *physical channel-mixing mechanism* (a boundary/transport-twisted state, not the vacuum); with it, the K_6 /Pfaffian grammar of v888 would be the measured correlation grammar of the seam. **(5) The constructed mixing candidate (round 55, [verification/v898_kms_schur_mixing.py], 30/30; verdict kmsmix-measured [kmsmix-found, schurmix-generates, rp-theta-open]).** The demanded state now *exists as a candidate*, by two routes. *The KMS route*: the Majorana-compatible C_6 -invariant Hamiltonians $K = ih$ are parametrized exactly (the hermitian O_{16} -commutant has dimension $112 = 62 + 50$; the antisymmetric slice splits as 33 cross — exactly the block space of (3) — plus 17 diagonal, with the two forced zeros of the transposed $\{4, 5\}$ duad), and already the *first* point of the frozen grid, $h = -(A_{16}^{\text{dep}} + \frac{1}{8}A_{\text{int}})$ at $\beta = 1$, passes *all* gates: strict CAR positivity, exact C_6 block covariance, all 15 duad cross-blocks at the frozen floor, the q^* /sheet grading carried exactly ($5/5$ vacuum + $10/10$ carrier), and all 15 block Wick monomials nonzero with the *canonical* Pf_4 sign pattern of (3); robust at $\beta = 2$. *The Schur route (exact, sympy)*: eliminating the boundary of the coupled diagonal state $A_{\text{full}} = [[\kappa A_{CC}, tV], [-tV^\top, mJ_3]]$ gives $A_{\text{eff}} = \kappa A_{CC} + \frac{t^2 m}{1-m^2} V J_3 V^\top$ *exactly*, and $S = V J_3 V^\top$ deposits nonzero correlation on *all ten* carrier duads (uniform $3J$ block, canonical Pf_4 signs) from the bare diagonal state — the same Schur/Feshbach elimination that creates the RH port creates the carrier mixing: on both fronts the real correlation lives in the *eliminated boundary*. The frozen 15-signature table of the winner is the registered prediction any physical seam state must reproduce; the OS-positivity under a *spatial* reflection is the named open check (RP-THETA-OPEN — no spatial θ is deployed on the 16-dim space; the Majorana real structure satisfies the v426/v440 dictionary automatically but is particle-hole, not spatial). The honest split: the

obstruction theorem, the block construction and the Schur generation identity [E]; the constructed states are *candidates* — whether the *actual* seam realizes the signature pattern is untouched, so the physical Wick-compiler premise (SEAM.CFIN.WICKFUNCTOR.01, SEAM.CFIN.KMSMIX.01) stays [O], unmoved — the physical realization is the named demand. *Dated addendum (2026-08-10, rounds 57–59)*: the open check RP-THETA-OPEN is now *measured* at family level and the answer is an exclusion — strict reflection positivity under both deployable spatial reflections forces $t = 0$ (OS positivity $\iff$ the mixing floor falls: mutually exclusive on this family), the complete C_6 -twisted census is exclusionary (0/6), and the parent-dilation route yields only a marginal cone-boundary witness (with the exact floor identity $t^2 \cdot 3m/(1 - m^2) = 1/200$) while the strict collar route is obstructed by an exact linear $2t$ law ([verification/v903_seam_rp_exclusion.py]; a measurement on the complete finite candidate set, not a universal no-go); the [O] premise stays unmoved. *Dated addendum (2026-08-10/11, rounds 60–63)*: the exclusion is now *anatomized* — strict 2-cycle RP is impossible on the *whole* C_6 -covariant class (the seat law reduces exactly to $\text{diag}(a_J, -a_J)$; the only escape is covariance-breaking at exact linear price 2ε , with the floor exchange $\text{Pf}_4(\varepsilon) = (\varepsilon - \frac{1}{200})(\varepsilon + \frac{1}{200})$), while the strict-collar obstruction is a two-seat linear law whose *equilibrium* kernel witnesses carry the full $1/200$ mixing at zero entropy production — no non-equilibrium state is needed, and whether the deployed PURE-I wiring is compiler-forced became the registered successor contract ([verification/v908_seam_equilibrium_wiring.py], SEAM.STATE.WIRING.SELECTOR.01); the [O] premise stays unmoved. *Dated addendum (2026-08-12, rounds 69–71)*: the successor contract is *closed*, and the answer is a *freedom theorem* ([verification/v911_wiring_freedom.py]): the exact Gröbner census of the orbit/edge rules leaves exactly *one* admissible three-dimensional wiring component modulo the rule gauge (the rotation cell); Z/X wirings are excluded gauge-invariantly by orientation propagation; the deployed PURE-I point is an *interior* point of that cell, connected to the pure- J ray by an integer rule-gauge element — PURE-I IS A DEPLOYMENT REPRESENTATIVE, NOT A COMPILER THEOREM. No compiler-side demand distinguishes the θ_S sheet-swap frame within the rule-gauge algebra (five demand classes, all silent on the angle), the strict-collar RP Grams transform *exactly covariantly* under the rule gauge, and the same deployed parent that fails strict collar RP at θ_S *passes it exactly* in the integer-conjugated frame: the rounds-58/59 obstruction is frame-relative (a statement at the fixed θ_S frame, not a wiring no-go), and a canonical wiring selection would require an additionally *derived* physical frame. The [O] premise of the Wick-compiler demand stays unmoved. No marker moves.

The current reduction state of P2 (2026-07-22) — text consolidation, markers unchanged

The axiom P2 stays *declared* (AX.P2.01); what has changed is how much of it is now carried by theorems. The honest ledger state: P1 ($c_3 = 1/(8\pi)$) and the *four marks* are [E]-anchored (Gauss–Bonnet $n = 2\chi = 4$, [verification/v216_marks_gauss_bonnet.py]); given the four marks, $g_{\text{car}} = 5$ follows from the partition $4 = 2+1+1$ plus the Deligne/Birkhoff–Grothendieck/stability typing ([verification/v491_p2_partition_corollary.py], [verification/v499_p2_weights_deligne_bg.py]); residual R1, the module identification, [C]; and the clock’s *order*, the $\mathbb{Z}_8$ tower, the deck’s *class* (a half-period translation) and the *position* half of the alignment bit are theorems ([verification/v506_seam_clock_rigidity.py], [verification/v507_seam_bit_origin.py], [verification/v510_seam_bit_freedom.py]). What remains as carrier input is *one discrete symmetry-lift bit*: flag transitivity of the four marks (marks *and* their two sides indistinguishable, $V_4 \rightarrow D_4$) $\iff \tau = i$ (equivalently: the clock / the CM-fixed half-period / the non-split extension class on the free seam circle) — on the free seam circle bare mark-transitivity is automatic for *every* configuration (the pair-exchanging V_4) and no local jet sees the side bit, so the flag form is the sharp discrete statement of the datum and its negation is concretely measurable (odd-doublet split $(2/\pi) \ln \cot(\delta/2) \neq 0$) ([verification/v512_seam_tau_flag.py], SEAM.TAU.FLAG.01). The bit has since survived *ten* side-blind derivation attacks on that scoreboard — free RP/ Θ existence (the eighth, v521), the exhausted Wick-computable free-plus-twist state class (the ninth, v525) and the first genuinely interacting seam toy (the tenth, v529) — and is now physically *defined* as the *twist-class choice*: facets #14/#15 extend the web to 15 exact equivalences (both directions each), the flip-axis fraction is a gauge-robust order parameter ($O = \frac{1}{2}$ at $m = 4$, 0 elsewhere), and its η holonomy is in principle interferometrically readable (v528, SEAM.BIT.TWISTCLASS.01; the bit remains formal input — a definition replaces no derivation, no marker moves); and it now carries the first *positive* selection datum: reflection positivity, imposed on the interacting mark family, keeps exactly *one* member alive — $\delta = \pi/2$ with positive coupling, the carrier of that order parameter ([verification/v534_seam_straddle_cone.py], SEAM.STRADDLE.CONE.01; toy-level evidence for a dynamical origin, not a derivation); plus the [C] identifications R1/R2 of v499. In one sentence: **the compiler’s dimensionless inputs reduce to four marks (derived from**

P1-side topology), one discrete symmetry-lift bit — the twist-class choice (flag transitivity $\iff \tau = i$) — and π . This is a reduction *state*, not an abolition of the axiom: AX.P2.01 remains the declared interface input, and its dynamical realisation (QGE0.REALIZE.01) stays [C]/[O]. [2026-08-07 addendum: the finite normal form of the interface is now assembled as *one* object — $C_{\text{fin}} = (V, \hbar, q^*, \sigma, \iota)$, the keybox above ([verification/v845_cfin_normal_form.py]) — with the code-to-matter kill built in as a checked negative; its honesty section types exactly what the assembly does *not* remove: P2's remaining content is the v799 residual R1' (the three seam axioms, the OPE-linearity scope, the v776 uniqueness selection). The assembly narrows; it does not eliminate. Markers unchanged.]

Seed normal form, the factorial spine, and the degree-2 pattern [E]/[C] ([verification/v106_review_validation.py])

Three review-validated sharpenings of the anchor picture. **(1) Seed normal form [E].** The seed formula is exactly

$$\varphi_0^{\text{ret}} = \frac{|\mu_4|}{N_{\text{fam}}} c_3 + \Omega_{\text{adm}} c_3^{|\mu_4|} \quad \left(\frac{4}{3} = \frac{|\mu_4|}{N_{\text{fam}}}, \quad 48 = \Omega_{\text{adm}} = N_{\text{fam}} \dim S^+ \right),$$

a *linear seam term* plus a *topological fourth-order correction* over all admissible fermion states — a structured anchor readout, not a fitted expression. **(2) Factorial spine [E].** With $X = 6Y$ on one 16-state generation, $\text{Tr } X = 0$ and $\text{Tr } X^2 = 120 = 5!$ (computed from the explicit charge table); the *same* factorial moment is $|R^+(E_8)| = \sum(E_8 \text{ exponents}) = 120$, doubles to $|R(E_8)| = 240$, and scales to $|W(D_5)| = 2^4 \cdot 5! = 1920$ (the Hawking-power fingerprint) — one moment, four projections (hypercharge, roots, Weyl group, horizon). **(3) The degree-2 inventory.** The compiler repeatedly selects *degree two*: the Pascal closure $2^{g-1} = \sum_{k \leq 2} \binom{g}{k}$ truncates at $K=2$ with the *unique* solution $g = 5$ (scan 1..40); the glue norms add to the root norm, $\frac{5}{4} + \frac{3}{4} = 2$; $c_3 = \frac{1}{2} \times \frac{1}{4\pi}$ (variational half $\times$ two-dimensional-slice Gauss–Bonnet); the pair sector $\binom{5}{2} = 10 = \mathcal{A}_\Lambda$; the hypercharge moment is a *second* moment; the seam is codimension *two*. *Named hypothesis [C] (recorded, not claimed): Quadratic Boundary Locality* — the seam supports only bilinear data, so the truncation $K = 2$ is *forced* rather than chosen; if proven, the Pascal *selection* (red-team Target B residual, typed [O]/[C]) upgrades to [E] and the chain locality $\Rightarrow K=2 \Rightarrow g=5 \Rightarrow 16 \Rightarrow D_5 \Rightarrow E_8$ closes the carrier choice. *Audit*: the even-code transition reading $240 = 16 \times 15$ is exact arithmetic but non-unique ($240 = 16 \times 5 \times 3$ is the established glue factorisation) and stays recorded, not promoted.

The Sheet-Pairing Lemma ([verification/v109_sheet_pairing.py]) — **the first exact anchor of the hypothesis.** Character-exact weight-multiset combinatorics of the D_5 spinor weights $(\pm \frac{1}{2})^5$ gives: (i) **no scalar within a sheet** — the zero-weight multiplicity of $S^+ \otimes S^+$ is 0, because the odd slot count $g = 5$ flips the sign parity of $-w$ (a parity theorem of the five-slot code); (ii) **the scalar lives across the sheets**:

$$S^+ \otimes S^- = \Lambda^0 \oplus \Lambda^2 \oplus \Lambda^4 \quad (256 = 1 + 45 + 210),$$

exact as weight multisets, with zero-mode grading $(1, 5, 10)$ = the carrier code; (iii) **sheet-diagonal = odd forms**: $(S^+ \otimes S^+) \cup (S^- \otimes S^-) = 2\Lambda^1 + 2\Lambda^3 + \Lambda^5$ exactly — within-sheet pairings are odd-form-valued, never scalar; (iv) the scalar-bearing tower **tops at Λ^{2K}** with $K = 2$ — bilinear seam data reach exactly the Pascal-ladder degree (the source-layer transport theorem; document 3). *Consequence [C]*: the seam's single scalar two-point kernel is necessarily a **sheet pairing** (chirality-off-diagonal — the same $\mathbb{Z}_2$ as the glue chiralities and the branch-kernel selection); the remaining QBL step is to show the Calderón kernel supplies exactly this pairing and nothing beyond.

The Calderón-sheet selection theorem ([verification/v110_calderon_sheet.py]) — **which involutions certify a scalar kernel.** On $H = S^+ \oplus S^-$ the certified bilinear data of a Calderón involution ε ($\varepsilon^2 = 1$) are the matrix elements $\langle f, \varepsilon g \rangle$. The lemma's singlet counts per sheet block, $(0, 0, 1, 1)$, give the exact selection statement [E]:

$$\text{a scalar two-point datum exists} \iff \varepsilon \text{ is sheet-odd}$$

— a sheet-odd involution (swapping $S^+ \leftrightarrow S^-$) certifies exactly $1 + 1 = 2 = |\mathbb{Z}_2|$ invariant scalar kernels (one per orientation), a sheet-even involution certifies *none*. The kernel count $2 = |\mathbb{Z}_2|$ matches exactly the *two Lagrangian glues* (swapped by single spinor conjugation): the Calderón scalar channel carries precisely the glue ambiguity, resolved by the same sheet choice. *Anti-overclaim control [E]*: the parity theorem is generic over the odd ladder ($g = 3, 5, 7$: within-sheet count 0, cross count 2^{g-1})

— sheet-oddness forces the *half-spinor relation* $K = (g - 1)/2$, **not** the value $g = 5$; the g -selection stays with the independent rank-8/integrality closure. *Consequence*: the QBL residue splits into two named parts [C] (recorded, not claimed): **(a) interface** — “the seam Calderón involution is sheet-odd” (structurally natural: the one-sided collar/double-cover deck is the *same* $|\mathbb{Z}_2|$ that halves $c_3 = \frac{1}{2}(4\pi)^{-1}$, and exchanging cover sides is sheet-odd by construction); **(b) analytic core** — “the two-point kernel certifies only slot-degrees ≤ 2 ”. Proving both closes $\text{QBL} \Rightarrow K=2 \Leftrightarrow g=5$.

The Quadratic-Transport Theorem ([verification/v111_quadratic_transport.py]) — **the transport half of the analytic core is a theorem**. Grade seam transport by Clifford degree in the ten field operators on the five-slot Fock space ($S = \Lambda^\bullet(\mathbb{C}^5)$, S^+ the 16-state code; integer Jordan–Wigner model, spanning claims by full-rank certificates mod p , valid over $\mathbb{Q}$). *Parity* [E]: all linear words are sheet-odd, all 45 quadratic words sheet-even — code-preserving transport has *even* Clifford degree (the same $\mathbb{Z}_2$ again). *Minimality* [E]: sheet-even words of degree ≤ 1 are the scalars alone. *Generation and completeness* [E]: the 45 quadratics ($= \mathfrak{so}(10)$, exactly the Λ^2 term of the certified tower $1 + 45 + 210$) generate the whole code from the vacuum *and* span the full operator algebra $\text{End}(S^+)$ ($\dim = 256$) at word length two — every code operation whatsoever is a word in pair transport, and all higher even-degree elements (e.g. the 210 quartics) are redundant. Hence

the transport degree is exactly 2

— degree ≤ 1 generates nothing, degree 2 generates everything: a theorem, not a hypothesis. *Anti-overclaim control* [E]: the $g = 3$ world behaves identically (15 quadratics generate $\text{End}(S^+) = 16$) — the theorem selects the *degree*, not the rank; the g -selection stays with Pascal closure + rank-8. *Residue after v109–v111* [C]: QBL’s remaining content is two *interface* statements — (a) the seam Calderón involution is sheet-odd; (b’) the certified *state* inventory is the ≤ 2 -slot tower. Nothing about transport remains hypothetical.

The Self-Counting Channel ([verification/v112_selfcounting_channel.py]) — **the Pascal closure is an identity of the certified channel**. For odd g , negation $w \mapsto -w$ maps the S^+ weights bijectively into the opposite sheet, so the Cartan-neutral kernels of the certified channel are exactly the pairs $(w, -w)$ — *one per code state*: $\#\{\text{neutral kernels}\} = \dim S^+ = 2^{g-1}$ exactly [E]. Grading the *same* neutral set by pair-degree m gives sizes $\binom{g}{m}$, $m = 0..K$ — for $g = 5$ the triple $(1, 5, 10)$ [E]. Two countings of one set:

$$2^{g-1} = \sum_{m \leq K} \binom{g}{m} \text{ is the channel counting itself}$$

(symbolically verified for all odd $g = 3..13$) — the Pascal *closure* (v2/v108) is not an extra requirement but an identity of the certified channel. *Hodge fold* [E]: the code’s minus-sign grading $\binom{g}{2j}$ equals the folded binomial $\binom{g}{\min(2j, g-2j)}$ with $\min(2j, g-2j) \leq K$ — every code sector appears at pair-degree $\leq K$. *Anti-overclaim*: the identity holds for every odd g and carries no g -selection; the selection stays with the rank-8 closure $g + N_{\text{fam}} = 8$ and ladder integrality. *Residue re-type* [C] (recorded, not claimed): clause (b’) loses its closure burden — given the sheet-odd kernel, the channel *automatically* contains one neutral kernel per code state at pair-degree $\leq K$; what remains physical is the single input “the seam certifies through one scalar two-point kernel” (for the established free $c = 8$ seam net, Wick extends the one kernel to the entire even tower by determinants). QBL is thereby the structural consistency frame of the carrier choice; the *selection* is carried by rank-8 + integrality.

One kernel is the whole net ([verification/v113_quasifree_kernel.py]) — **the QBL input merges with the holomorphy premise**. The remaining input is now anchored in the established Gate-A objects. *Bookkeeping* [E]: $c(D_5)_1 = \frac{45}{9} = 5 = g_{\text{car}}$, $c(A_3)_1 = \frac{15}{5} = 3 = N_{\text{fam}}$, and the carrier net $(D_5)_1 \times (A_3)_1 = SO(10)_1 \times SO(6)_1$ is $10 + 6 = 16$ *free Majorana fermions* ($c = \frac{16}{2} = 8$); the whole extension tower carrier $(\mu=16) \rightarrow SO(16)_1(\mu=4) \rightarrow (E_8)_1(\mu=1)$ carries the *same* sixteen fermions — only the certified glue grows (index $2 \times 2 = 4 = |\mu_4|$). *Wick/Pfaffian* [E]: in the exact ten-Majorana model *all* 210 vacuum four-point and all 210 six-point functions equal the Pfaffian of the single two-point kernel — one kernel determines all correlations. *The kernel is a Calderón involution of rank g* [E]: $M = \mathbf{1} + iA$ with $A^2 = -\mathbf{1}$, so $P = M/2$ is a projection with rank $P = 5 = g_{\text{car}}$ and $\varepsilon = iA$ an involution; at seam level (sixteen Majoranas) the rank is $8 = \text{rank } E_8$:

the central charge is the rank of the one kernel

($c = 5$ carrier block, $c = 8$ seam hull); and the polarization fixes the vacuum *uniquely* (joint annihilator kernel one-dimensional). *Premise merge* [C] (recorded, not claimed): “single scalar two-point kernel”

= “the seam state is quasi-free” = the defining property of the free-fermion net the holomorphy gate already posits — QBL adds *no* assumption beyond Gate A; when the gate closes, the carrier choice closes [E] along free net $\Rightarrow$ one kernel $\Rightarrow$ sheet-odd $\Rightarrow$ self-counting tower $\Rightarrow$ complete transport $\Rightarrow$ carrier. Gate typing [C]/[O] unchanged.

Hardening (P1): the Gauss–Bonnet origin of $c_3 = \frac{1}{8\pi}$

$c_3 = 1/(8\pi)$ is not an arbitrary number: it is the *one-sided Gauss–Bonnet normalisation of the seam*. The seam construction compactifies the oriented two-dimensional normal slice N_Σ to a sphere S^2 , so Gauss–Bonnet gives the seam curvature integral

$$\oint_{S^2} K dA = 2\pi \chi(S^2) = 4\pi \quad (\chi(S^2) = 2),$$

and the boundary datum is *one-sided* (the Calderón/double-cover construction uses a single collar), contributing a factor $1/|\mathbb{Z}_2| = \frac{1}{2}$. Hence

$$c_3 = \frac{1}{|\mathbb{Z}_2| \oint_{S^2} K dA} = \frac{1}{2 \cdot 4\pi} = \frac{1}{8\pi}$$

equivalently $c_3 = \frac{1}{2}(4\pi)^{-1}$, the one-sided 2D heat-kernel coefficient. This locates both factors precisely:

- the *discrete* $8 = |\mathbb{Z}_2| \cdot 2 \cdot \chi(S^2) = 2 \cdot 2 \cdot 2$ coincides with $h(D_5) = \text{rank } E_8 = \varphi(30)$ — the geometric and the bootstrap readings of the “8” agree;
- the *continuous* π is the irreducible Gauss–Bonnet / 2D-Gaussian primitive ($\int e^{-x^2} = \sqrt{\pi}$), the single primitive the self-consistency loop also isolates.

Status: this *hardens* (P1) — it gives c_3 a named geometric origin and reconciles the “8” with E_8 — but does *not* eliminate π ; π remains the one continuous primitive. [C]

The one-paragraph version. There is a boundary (the seam). It yields $c_3 = 1/(8\pi)$ and a five-slot carrier $g_{\text{car}} = 5$, split 3 colour + 2 weak. Two base charges $y_- = -1/3$ and $y_+ = 1/2$ generate the whole fermion family; the 16 even-occupation states are one generation, three families give $3 \times 16 = 48$. One seed $u = \varphi_0^{\text{ret}}$ drives four low-energy observables. Finally $\alpha_\star^{-1} \approx 137$ is the exponential engine: the electroweak scale takes one fifth of it ($e^{-\alpha_\star^{-1}/5}$), the cosmological constant takes it twice ($e^{-2\alpha_\star^{-1}}$), and the Hubble scale is the square root of Λ . A small grammar generates many sectors.

2 Headline: the Pascal compiler on five carrier slots

The strongest new statement is that three of the theory’s separate structures are the *same* truncated row of Pascal’s triangle on the five-slot carrier.

Family = action ladder = transition count (one Pascal row)

The even exterior algebra of E ($\dim E = g_{\text{car}} = 5$), the action ladder, and the E_8 root count are one Pascal row:

$$S^+ = \Lambda^0 E \oplus \Lambda^2 E \oplus \Lambda^4 E, \quad (\dim \Lambda^0, \dim \Lambda^2, \dim \Lambda^4) = (1, 10, 5), \quad \dim S^+ = 1 + 10 + 5 = 16,$$

$$A_{\text{EW}} : \mathcal{A}_H : \mathcal{A}_\Lambda = \binom{5}{0} : \binom{5}{1} : \binom{5}{2} = 1 : 5 : 10 = (\dim \Lambda^0 E, \dim \Lambda^4 E, \dim \Lambda^2 E),$$

$$|R(E_8)| = \dim S^+ (\dim S^+ - 1) = 16 \cdot 15 = 240.$$

Family (exterior degree), cosmology (action charge) and E_8 (root transitions) read the same $\{1, 5, 10\}$. [E]




Figure 1: Exterior–Action duality. The Standard-Model family is the valency side of the even Hamming scheme on the five-slot code; the cosmological action ladder is its spectral (multiplicity) side. The Λ (pair) sector $\Lambda^2 E$ has dimension $\binom{5}{2} = 10$, the origin of the decuple exponent.

Natural sector assignment:

$$EW \leftrightarrow \Lambda^0 E \text{ (singlet)}, \quad H \leftrightarrow \Lambda^4 E \cong E^\vee \text{ (dim 5)}, \quad \Lambda \leftrightarrow \Lambda^2 E \text{ (dim 10)}.$$

The cosmological constant lives on the *pair* sector $\Lambda^2 E$, $\dim = \binom{5}{2} = 10$; this is why the decuple bridge has exponent 10 — Λ is a product over all $\binom{5}{2}$ carrier pairs (K_5 edges), not over the full 16-state family.

2.1 $g_{\text{car}} = 5$ as the Pascal closure

A family is exactly the complete Pascal truncation to degree two:

$$\dim S^+ = 2^{g_{\text{car}}-1} = \binom{g_{\text{car}}}{0} + \binom{g_{\text{car}}}{1} + \binom{g_{\text{car}}}{2} \iff g_{\text{car}} = 5 \quad (16 = 1 + 5 + 10), \quad [\text{E}], \quad (2)$$

the unique positive-integer solution of $f(g) := 2^{g-1} - 1 - \frac{g(g+1)}{2} = 0$. *Proof of uniqueness.* Direct evaluation gives $f(1)=-1$, $f(2)=-2$, $f(3)=-3$, $f(4)=-3$, $f(5)=0$, $f(6)=+10$; so $g=5$ is the only root in $\{1, \dots, 6\}$. For $g \geq 6$ the forward difference is $f(g+1) - f(g) = 2^{g-1} - (g+1) \geq 2^5 - 7 = 25 > 0$ and increasing, so f is strictly increasing on $[6, \infty)$ from $f(6)=10 > 0$; hence $f(g) > 0$ for all $g \geq 6$ and there is *no* further root. Thus $g_{\text{car}} = 5$ is the unique solution (no finite-range caveat). [verification/v2_carrier_pascal.py] This is sharper than $2g_{\text{car}} = \binom{g_{\text{car}}}{2}$: the family code (even states, 2^{g-1}) equals the action code (sum of the first three Pascal entries) only at $g_{\text{car}} = 5$.

Carrier uniqueness theorem: $g_{\text{car}} = 5$ and the 3+2 split are forced [E]

The carrier rank and its colour/weak split are not chosen — they are the unique solutions of two integer conditions tied to the E_8 closure and the SM gauge algebra:

$$g_{\text{car}} = 5 \text{ is the unique } g \text{ with } N_{\text{fam}} = \frac{2^{g-1}-1}{g} \in \mathbb{Z}^+ \text{ and } \text{rank } E_8 = g + N_{\text{fam}} = 8 \quad (3)$$

$$(b, s) = (3, 2) \text{ is the unique split of } b + s = g_{\text{car}} = 5, \quad b^2 + s^2 = 13 = |R(A_3)| + 1 \quad (4)$$

The integer-family carriers are $g \in \{3, 5, 7, \dots\}$ (rank 4, 8, 16), so $\text{rank } E_8 = 8$ picks $g_{\text{car}} = 5$ uniquely; and $b^2 + s^2 = 13$ with $b+s = 5$ forces $(3, 2)$, reproducing $\dim \mathfrak{g}_{\text{SM}} = (b^2-1) + (s^2-1) + 1 = 12 = |R(A_3)|$. The induced hypercharge is anomaly-free ($\text{Tr } Y = \text{Tr } Y^3 = 0$ over the 16 states). This hardens P2: the *rank* and the *split* are forced; only the seam→carrier interface itself remains a declared input. [verification/v14_carrier_uniqueness.py]

The Pascal grammar: one $\Lambda^\bullet$ mechanism in four places (anti-numerology) [E]. The recurring small integers are *not* coincidences — they are the *same* exterior-algebra (Pascal)

mechanism read in four spaces, which is why the binomials of rows 4 and 5 keep reappearing:

space	Pascal reading
carrier (one generation)	$\dim S^+ = \Lambda^{\text{even}}(5) = \binom{5}{0} + \binom{5}{2} + \binom{5}{4} = 1 + 10 + 5 = 16$
scale ladder (cosmology)	$A_{\text{EW}} : \mathcal{A}_H : \mathcal{A}_\Lambda = \binom{5}{0} : \binom{5}{1} : \binom{5}{2} = 1 : 5 : 10$
flavor u/d readout	$\ \text{Pl}(K) \ _1 = \binom{4}{0} + \binom{4}{1} + \binom{4}{2} = 1 + 4 + 6 = 11 = 16 - g_{\text{car}}$
mass volume	$\det M_{\text{SM}} \sim (\varphi_0^{\text{ret}})^{6+9+10} = (\varphi_0^{\text{ret}})^{25} = (\varphi_0^{\text{ret}})^{g_{\text{car}}^2}$

So the carrier is the even row 5, the scale ladder is the lower row 5, the u/d leg is the cumulative row 4 up to $\deg u^c = 2$, and the mass volume closes on g_{car}^2 . *One* grammar (Λ^k of the 5-slot carrier and the $4 = |\mu_4|$ family fundamental), not four lucky hits. [\[verification/v44_carrier_exterior.py\]](#) [\[verification/v45_family_exterior.py\]](#) [\[verification/v46_grand_mass_volume.py\]](#)

2.2 One master moment generates 40, 41, 240, δ_2

With $X = 6Y$, the quadratic hypercharge trace on one family is $\text{Tr}_{S^+} X^2 = 120 = 5!$, and it generates the whole integer chain:

$$\sum_{f,j} L_{f,j} = \frac{\text{Tr}_{S^+} X^2}{3} = 40, \quad 10b_1 = \frac{\text{Tr}_{S^+} X^2}{3} + 1 = 41, \quad (5)$$

$$|R(E_8)| = 2 \text{Tr}_{S^+} X^2 = 240, \quad \delta_2 = 4! \text{Tr}_{S^+} X^2 c_3^8.$$

The flavor budget, the $U(1)$ budget, the E_8 count and the second seam defect all hang on the second hypercharge moment $5!$. [\[E\]](#) *Round-22 sharpening*: the moment itself is one symbolic identity, $\text{Tr}_{S^+} X^2 = 4|X|^2 = 4h(E_8) = 120$ — the Coxeter number carries the whole chain ([\[verification/v810_cutcode_moment.py\]](#)) [\[E\]](#).


2.3 Compact E_8 compression and the role of ν^c

$$\begin{aligned} |R(E_8)| &= \dim S^+ \cdot g_{\text{car}} \cdot N_{\text{fam}} = 16 \cdot 5 \cdot 3 = 240, \\ \dim E_8 &= \dim S^+(\mathcal{A}_H + \mathcal{A}_\Lambda) + (g_{\text{car}} + N_{\text{fam}}) = 240 + 8 = 248 \end{aligned} \quad (6)$$

with $N_{\text{fam}} = (2^{g_{\text{car}}-1} - 1)/g_{\text{car}} = 15/5 = 3$ and $\text{rank } E_8 = g_{\text{car}} + N_{\text{fam}} = 8$. [\[E\]](#). The right-handed neutrino ν^c is the Pascal singleton $\Lambda^0 E = (1, 1)_0$ that closes $16 = 1 + 5 + 10$; removing it breaks the truncation and shifts α^{-1} by 2.16×10^{-3} .

3 The $D_5 \times A_3$ compiler: E_8 as a closure, not an input

This section upgrades the E_8 atlas (Section 10) from a counting coincidence to an explicit lattice construction. The whole content is consolidated and verified in the standalone companion note “*The Even Carrier Code and the $\mathbb{Z}_{30}$ Coxeter Compiler*.” The thesis: the carrier code is the D_5 half-spinor, the family geometry is an A_3 , and E_8 is the standard unimodular glue closure of $D_5 \oplus A_3$ — with the four punctures μ_4 as the glue code.



$D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ — the unique even unimodular rank-8 glue [\[E\]](#) exact (lattice).

The compiler in one line:

five carrier slots $\Rightarrow D_5$, four family corners $\mu_4 \Rightarrow A_3$, $(D_5 \oplus A_3) + \mu_4\text{-glue} \Rightarrow E_8$.





Figure 2: The explanation hierarchy. The boundary fixes the seam seed c_3 ; the five-slot even carrier code is the D_5 half-spinor; the four punctures μ_4 give A_3 ; their common $\mathbb{Z}_4$ discriminant (μ_4) glues $D_5 \oplus A_3$ into the unimodular E_8 . E_8 is the *closure*, not the input; c_3 dresses the continuous observables.

The glue is *one* operation in two categories. The same order- $|\mu_4|=4$ discriminant glue closes $D_5 \oplus A_3$ into E_8 *twice over*: once on *root lattices* (the unimodular pushout $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$, v1) and once on *chiral conformal nets* (the simple-current / anyon-condensation extension $(D_5)_1 \otimes (A_3)_1 \rtimes \mu_4 \cong (E_8)_1$, v92/v281). These are not two coincidences but the two functorial faces of a single commuting square (v234/v277/v260); the boundary realisation on which the strict-TOE residual turns (SEAM.EQUIV.01) is exactly the statement that the raw seam realises this square — now *closed modulo cited theorems on its MMST route* SEAM.EQUIV.MMST.01 (the lattice/ S_3 stack pins it at every computable level, Lean FORM.SEAM.MMST.01; only the cited MMST/OS continuum-existence theorems stay [O], v367/v368/v376–v379/v336; the twistor route SEAM.EQUIV.TWISTOR.01 and the unconditional parent stay [O]). The extension leg of that citation package is itself re-founded at net level ([verification/v469_seam_crossedproduct_route.py]): the 128-spinor step is the *local* $\mathbb{Z}_2$ simple-current crossed product ($h_s(SO(16)_1) = 16/16 = 1 \in \mathbb{Z}$, the Longo–Rehren locality criterion; KLM $\mu = 4/2^2 = 1$), on 1995–2001 peer-reviewed subfactor theory, with the lattice-VOA preprint route demoted to an independent second witness, and the realisation input reduced to invariant level (R1'). Premise (i) of that package ($c_- = 8$) now also carries a *ground-state* witness ([verification/v489_seam_modular_commutator.py]): the Kim–Shehab–Kim modular commutator $J(A, B, C) = (\pi/3)c_-$, evaluated on the exact BdG ground state of the collar model, gives $c_- = 8 = g_{\text{car}} + N_{\text{fam}}$ for the 16-layer carrier (per layer 0.500000 at $L=48$; trivial and orientation-flip controls pass) — independent of both the Chern route (v367) and the KLM count (v378). SEAM.EQUIV.01 stays [O]. *Follow-up (2026-08-03, the G_{net} witnesses)*: the index-4 statement now carries two *exact* arithmetic/index witnesses. The NS/R grading is the parity character of $E_8(\mathbb{Z}[i])/(1+i) = \mathbb{F}_2^4 - (1+i)L$ lies in the even sublattice (proved on a $\mathbb{Z}$ -basis), all 15×16 root classes are parity-pure (7 NS + 8 R), so the Ramond projection is $(1+i)$ -adic at the one ramified edge of the $\mathbb{Z}[i]$ -E8 Hecke tower ([verification/v722_phys_ramified_ns_r.py], 9/9, RAMIFIED-SEES-NSR); and on the CAR ladder the μ_4 -average expectation has optimal Pimsner–Popa constant exactly $1/4$ and Watatani index exactly 4 via an explicit quasi-basis with *forced* weights, with μ_4 *derived* from the clock and the Ramond sector healed state-preservingly by the Klein zero-mode dressing ([verification/v726_phys_car_pp_index.py], 57/57, T2-SLICE-GO). The remaining half is the *identification* of the ladder Q-system with the constructed $\mathbb{C}[\mathbb{Z}_4]$ object (GNET.RAMIFIED.01, kills registered); GATE.METRIC.08/.10 typing unchanged. *Follow-up (2026-08-05, the continuum strand)*: the finite-level precursors of the GNS limit are measured *coherent* ([verification/v779_gnet_gns_arf.py], 27/27 + 27/27): the Haar scaling tower intertwines exactly, the index-4 expectation squares commute exactly (the finite half of the Q-system identification), the state converges geometrically, and the exact dim law $d_c(\ell) = 4^{\ell-1} + 2^{\ell-1} \cos(\pi c/2)$ types the

$\mathbb{C}[\mathbb{Z}_4]$ obstruction at exactly $2^{-\ell}$ (explaining the $\ell = 1$ three-sector anomaly); the honest negative: the $16 = 1+5+10$ counting echo with the Arf geometry is exact but *non-geometric*, and the two fours (index 4 vs channel 4/49) stay in their registers. The round-21 martingale probe measures the limit-existence hypotheses of the continuum strand as met ([[verification/v802_gnet_martingale.py](#)], 15/15, MARTINGALE-HYPOTHESES-MET; the analytic remainder stays typed). Typing unchanged.



One operation, two categories: the order- $|\mu_4|=4$ glue is the lattice pushout $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ and the simple-current / anyon-condensation extension $(D_5)_1 \otimes (A_3)_1 \rtimes \mu_4 \cong (E_8)_1$; the square commutes. [E] exact.

3.1 Carrier = $D_5 = \mathfrak{so}_{10}$ (proved)

The 16 even-occupation states $C^+ = \{x \in \mathbb{F}_2^{g_{\text{car}}} : |x| \text{ even}\}$ are exactly the weights of the D_5 half-spinor $\frac{1}{2}(\pm 1, \dots, \pm 1)$ (even number of minus signs). The flip-weight-4 relation is the **Clebsch graph** $\text{SRG}(16, 5, 0, 2)$ and the flip-weight-2 relation is its complement $\text{SRG}(16, 10, 6, 6)$; the *affine* automorphism group of the scheme (translations $\rtimes$ coordinate permutations) is the D_5 Weyl group,

$$\boxed{\text{Aut}_{\text{aff}}(C^+) = C^+ \rtimes S_5, \quad |\text{Aut}_{\text{aff}}(C^+)| = 2^{g_{\text{car}}-1} g_{\text{car}}! = 1920 = |W(D_5)| = r |R(E_8)|} \quad [\text{E}], \quad (7)$$

(the linear code automorphism group alone is only S_5 ; the order-1920 group is the affine/scheme one) and the Clebsch edge count is the flavor budget $|E(R_4)| = \frac{16 \cdot 5}{2} = 40 = |R(D_5)| = \sum_{f,j} L_{f,j}$. *Round-22 grading (2026-08-06)*: the Arf split $16 = 1 + \bar{5} + 10$ is the cut classification $(0|5) + (1|4) + (2|3)$ of K_5 (the cut map injective-isometric on all 256 cells), and the Clebsch edges grade as $40 = 5 + 20 + 15$ over the Arf shells with the within-10 block equal to the Petersen graph — an Arf/cut geometry, exact but carrying no matter semantics ([[verification/v810_cutcode_moment.py](#)]) [E].

3.2 Family = $A_3 = \mathfrak{su}_4$ from $\mathbb{P}^1 \setminus \mu_4$ (proved, with the metric settled)

The carrier geometry fixes $X_f^\circ \cong \mathbb{P}^1 \setminus \mu_4$, $H_1 = \mathbb{Z}^3$, rigid D_4 generated by $z \mapsto iz$, $z \mapsto 1/z$. The twelve puncture-difference cycles $\gamma_i - \gamma_j$ carry the residue pairing $\text{Res}(\omega_i, \gamma_j) = \delta_{ij}$ and form the A_3 root system:

$$\boxed{\#\{\gamma_i - \gamma_j\} = 12 = |R(A_3)| = \dim \mathfrak{g}_{\text{SM}}, \quad \text{rank} = 3 = N_{\text{fam}}, \quad \text{Gram} = \text{Cartan}(A_3), \quad \det = 4} \quad [\text{E}]. \quad (8)$$

The corner rotation $z \mapsto iz$ acts on $H^1 = \mathbb{C}^3$ as the A_3 Coxeter element (order $4 = h(A_3) = |\mu_4|$, spectrum $\{i, -1, -i\}$, $\chi = \Phi_4 \Phi_2$), an isometry of the Cartan form — the exact analogue of the E_8 Coxeter element reading Φ_{30} .

The μ_4 clock is the order-4 character of the E_8 cycle, and $(2, 3, 5)$ is the icosahedron ([[verification/v223_coxeter_totative_clock.py](#)], [[verification/v219_icosahedral_mckay.py](#)]). The two “Coxeter” clocks are one object read at two scales. The E_8 exponents are the totatives $(\mathbb{Z}/30)^\times = \{1, 7, 11, 13, 17, 19, 23, 29\}$, and the multiplicative order-4 element 7 ($7^2=19$, $7^4=1 \pmod{30}$) generates a literal μ_4 clock $\langle 7 \rangle = \{1, 7, 13, 19\}$ that permutes the four conjugate Coxeter 2-planes — the same μ_4 as the seam corner rotation. And the period $30 = 2 \cdot 3 \cdot 5$ itself is not a free factorisation: by the McKay correspondence E_8 is the exceptional *top* of the tower of finite $SU(2)$ subgroups ($2T \rightarrow \hat{E}_6$, $2O \rightarrow \hat{E}_7$, $2I \rightarrow \hat{E}_8$), so its branch data is the *icosahedron* and the three atoms $(|\mathbb{Z}_2|, N_{\text{fam}}, g_{\text{car}}) = (2, 3, 5)$

are its rotation-axis orders. The binary icosahedral group $2I$ (order $120 = |R^+(E_8)|$) has nine irreducible-representation degrees $\{1, 2, 2, 3, 3, 4, 4, 5, 6\}$ equal to the affine- E_8 Kac marks, with $\sum d_i = 30 = h(E_8)$ and $\sum d_i^2 = 120$. This is a *backward certificate* of the closed E_8 [E] (it explains the $(2, 3, 5)$ choice [C]; it does *not* replace the P2 input).

The metric, settled (former [C]). The transcendental geometric (Green) metric $G_{ij} = -\log |p_i - p_j|$ on μ_4 is only D_4 -symmetric, with a regularisation-independent anisotropy $-\log 2$ on the root space, so it is *not* proportional to the A_3 Cartan form. But it does not need to be: the E_8 -relevant A_3 is the canonical *residue* pairing (integral, S_4 -symmetric, $= \text{Cartan}(A_3)$), and the geometric metric is a D_4 -equivariant deformation preserving the root system, Weyl group and Coxeter element. So the lattice structure is exact [E]; the analytic metric is a characterised deformation, not an open gap.

3.3 The μ_4 glue and the $5/4 + 3/4 = 2$ discriminant theorem (proved)

The two factors have equal discriminant groups and E_8 is unimodular, so the glue index is exactly $|\mu_4|$:

$$\text{disc}(D_5) = \text{disc}(A_3) = \mathbb{Z}_4, \quad [E_8 : D_5 \oplus A_3] = \sqrt{\frac{\det D_5 \det A_3}{\det E_8}} = \sqrt{\frac{4 \cdot 4}{1}} = 4 = |\mu_4|. \quad (9)$$

The discriminant-form norms sum to the E_8 root norm, and — strikingly — they are pre-existing TFPT constants:

$$\boxed{\begin{aligned} q(D_5) &= \frac{5}{4} = \frac{\delta_2}{\delta_{\text{top}}^2}, & q(A_3) &= \frac{3}{4} = \xi_{\text{tree}}, \\ q(D_5) + q(A_3) &= 2 = |E_8 \text{ root}|^2 \end{aligned}} \quad [\text{E}] \quad [\text{verification/v1_e8_glue.py}] \quad (10)$$

This is literally the E_8 spinor-root split $\frac{1}{2}(\pm)^8 = \frac{1}{2}(\pm)^5 \oplus \frac{1}{2}(\pm)^3$, $5 \cdot \frac{1}{4} + 3 \cdot \frac{1}{4} = 2$.




Figure 3: The discriminant-norm glue. The E_8 spinor root $\frac{1}{2}(\pm)^8$ (norm 2) splits across $D_5 \times A_3$ as $5 \cdot \frac{1}{4} + 3 \cdot \frac{1}{4}$; the two halves are the pre-existing TFPT constants $\delta_2/\delta_{\text{top}}^2 = \frac{5}{4}$ and $\xi_{\text{tree}} = \frac{3}{4}$.

The standard branching is realised explicitly:

$$\mathfrak{e}_8 = (45, 1) \oplus (1, 15) \oplus (10, 6) \oplus (16, 4) \oplus (\overline{16}, \overline{4}), \quad 248 = 45 + 15 + 60 + 64 + 64, \quad 240 = 40 + 12 + 60 + 64 + 64, \quad (11)$$




Figure 4: The $E_8 \supset D_5 \times A_3$ branching as norm-2 blocks. The adjoint half is the hypercharge moment $\text{Tr}_{S^+} X^2 = 120 = \dim \mathfrak{so}_{16}$; the spinor half is two sheets over the four corners μ_4 .

and assembling $D_5 \oplus D_3 \subset D_8$ with the even spinor coset produces all 240 roots (norm 2, Cartan $\det = 1$, every root an integer combination of one simple-root basis): *the 240-metrisation is no longer blind — E_8 is built.* [E]

Hecke structure of the glue census is lattice-native (HECKE.GEOM.01). The μ_4 -glue class thetas that refine the 240-root branching ($240 = 52+64+60+64$ over $\mathbb{Z}/4$) now carry a machine-checked Hecke package ([[verification/v535_hecke_from_geometry.py](#)]): Kneser p -neighbours realise $\#iso_lines = \sigma_3(p) \# \mathbb{P}^3(\mathbb{F}_p)$ at $p = 2, 3, 5, 7$; the marked neighbour-sum is the affine element $\nu_p = a \text{Id} + b T_p$ with cusp output $a_3 = -4$, $a_5 = -2$; and the seven census q -series span exactly the 2-adic oldform hull of E_4 and f_8 (levels $\{1, 2, 4, 8, 16\}$ and $\{8, 16\}$). Classical scaffolding named classical; weight-4 $\rightarrow \text{GL}(1)$ stays [O]; no RH claim. [E] (in-suite mechanics)

Eichler trace layer at the E_8 lattice (HECKE.GEOM.EICHLER.01). The same census channels now carry a machine-checked Eichler/Witt package ([[verification/v536_eichler_trace_layer.py](#)]; 23 checks, ~ 30 s; builds on HECKE.GEOM.01 without duplicating Kneser/ ν_p /oldform checks). (A) The elementary Witt shadow $\lambda_{\text{Eis}} = \sigma_3(\# \mathbb{P}^3 - \sigma_3) = L - \sigma_3^2$ is closed in p (identically and for all odd $p \leq 100$). (B) The Eichler identity $\lambda_{\text{geom}} = \lambda_{\text{Eis}} + a_p^2$ is two-sided at $p = 3$ (live type-(i)/(ii) split: $352 = 364 - 12$) and at $p = 5$ via a T36-validated freeze ($3784 = 3906 - 122$); the assembler $R(p) = \sigma_3 - 1 - c(p^2)/8$ equals a_p^2 with Kneser filter = identity. (C) Closed Type-A/B local densities $N_A = \min(240(1 + p^3), p^7 + p^4 - p^3 - 1)$ (B empty iff $p^4 < 241$; $p = 7$ live 82560/743040; injectivity at $p = 11, 13$). (D) Signed extraction $a_p = -c(p)/8$ from geometric Shell(p) recovers $(-4, -2, +24)$ and $b \in \{24, 124, 368\}$. (E) $O(1)$ assembler for all odd $p \leq 31$ against the eta product f_8 . Classical Eichler/Siegel/Witt named classical; the claim is in-suite two-sidedness and the closed compiler densities. Honest fences: Shell(p^2) for $p \geq 7$ out of scope; fibre refinement scoped to $\{5, 7, 11\}$; weight-4 $\rightarrow \text{GL}(1)$ stays [O]; no RH claim; no marker moves. [E] (mechanics) / [O] (weight drop)

Half-integral bridge (HECKE.GEOM.HALFINT.01). The weight-4 cusp f_8 now has a machine-checked half-integral companion in the compiler theta-monoid ([[verification/v537_halfintegral_bridge.py](#)]; 20 checks, ~ 90 s; companion to HECKE.GEOM.01/HECKE.GEOM.EICHLER.01). (A) In the 70-element weight-5/2 theta-monoid (plain and q^2 -scaled $\vartheta_2, \vartheta_3, \vartheta_4$) there is exactly one object with the signed Shimura identity $\text{Sh}_{t=2, \psi=1}(g) = -8 f_8$ (witness $g = \vartheta_2(q^2)^2 \vartheta_3(q^2) \vartheta_4 \vartheta_4(q^2)$, verified coefficientwise through $n \leq 120$; three further monoid elements lift to scales $+8, -16, +16$). (B) g is an exact $T(p^2)$ -eigenform of weight 5/2, trivial nebentypus, with eigenvalues $a_p(f_8)$ for $p = 3, 5, 7, 11, 13$ $(-4, -2, 24, -44, 22)$; trial FE of the 5/2-side around centre 5/4 with $N = 512$, $\varepsilon = +1$. (C) $U_4(g) = 0$ exactly (mass mod 4 supported on $\{1, 2\}$); level $32 = 4 \cdot 8$ with $M = 8$ even lies outside Kohnen 1982 — the plus-route is structurally closed (negative result part of the claim). (D) With AFE-based twist L -values ($\varepsilon_d = \chi_d(8)$; $\varepsilon = -1 \Rightarrow L(2) = 0$), the Waldspurger quotient $R(d) = b(|d|)^2 / (|d|^{3/2} L(f_8 \times \chi_d, 2))$ is constant on ten fundamental discriminants $d \equiv 1 \pmod{8}$ ($R = 23.1873585645\dots$, relative spread $\leq 10^{-12}$, including $d = 89$); the class $d \equiv 5 \pmod{8}$ vanishes structurally. Classical Shimura/Kohnen/Waldspurger/Baruch–Mao named classical; the claim is the unique signed-scale monoid preimage, the in-suite chain, and the *measured* constant R . Honest fences: $\text{GL}(2)$ centre $s = 2$, not the ξ -line; no RH claim; ZETA.HP.CARRIER untouched; no marker moves. [E] (Shimura/Hecke/Kohnen fence) / [C] (measured R)

Compiler relative-trace identity (HECKE.GEOM.RTF.01). The three preceding packages are machine-checked as three *projections* of one finite relative-trace identity [E_8 /glue orbital counting] = [Newform/Waldspurger spectral sum] ([[verification/v538_relative_trace_identity.py](#)]; 18 checks, ~ 14 s; promotion of discovery T53; companion to HECKE.GEOM.01/HECKE.GEOM.EICHLER.01/HECKE.GEOM.HALFINT.01). (R1) $\text{Tr}_V(\nu_p \circ \pi)$ is computed *both sides independently*: geometric $a_p = -c(p)/8$ from the glue census (no modular input) equals spectral a_p from $\eta(2\tau)^4 \eta(4\tau)^4$ (no geometry) for all $(p, \pi) \in \{3, 5, 7\} \times \{\text{Id}, \pi_{\text{Eis}}, \pi_{\text{cusp}}, \pi_{E_4}, \pi_{f_8}\}$ (anchors e.g. $p=7$: full trace 727680, f_8 -channel 19840). (R2) The Eichler split $\lambda_{\text{geom}} = \lambda_{\text{Eis}} + a_p^2$ is the RTF orbit decomposition (principal orbit $\rightarrow \lambda_{\text{Eis}} = L - \sigma_3^2$; elliptic $\rightarrow a_p^2$; cross terms cancel; anchors $\lambda_{\text{Eis}} = 336/3780/19264$, $a_p^2 = 16/4/576$). (R3) The period side is lattice counting: g is the quaternary form $n = (x^2 + y^2)/2 + 2z^2 + u^2 + 2w^2$ (x, y odd; sign $(-1)^{|u|+|w|}$), with $b_{\text{rep}} = b$ through $n \leq 200$, and $[\text{signed lattice count}]^2 = R \cdot |d|^{3/2} \cdot L(f_8 \times \chi_d, 2)$ at $R = 23.1873585645$ (rel $\sim 10^{-12}$). Verdict **ONE-FORMULA**. Classical Jacquet-type RTF / Eichler / Witt / Waldspurger / theta correspondence named classical; NEW is the compiler-native realisation and bilateral independence. Honest fences: the identity is *finite* ($p \in \{3, 5, 7\}$, finite d -windows); the *infinite*

RTF (sum over all p and d + archimedean term) remains the named open step; no RH claim; not ZETA.HP.CARRIER-relevant; no marker moves. [E] (R1/R2 exact) / [C] (R3 measured R)

Weil structure of the compiler family (RTF.GNS.WEIL.01). The compiler family carries a fully identified Weil structure up to *two explicitly isolated obstructions* ([verification/v539_weil_structure_family.py]; 25 checks, ~ 6 s; promotion of discovery T55/T61/T62/T63; companion to HECKE.GEOM.RTF.01). (A) The canonical GNS pairing is the direct integral over the Gelfand spectrum of the character subalgebra: metric $\ell^2(d, b^2/|d|)$ diagonal; fibres $\sigma(d) = (\chi_d(3), \dots, \chi_d(13))$ partition the live family; Λ_{fam} constant on each fibre; twist-mixing certificate per fibre $\leq 5.037 \times 10^{-16} < 10^{-12}$. (B) The trivial Sato–Tate isotype is algebraically the $\text{GL}(1)$ core $G_0 = (1 + Y)/(1 - Y) = \zeta_p(w - 3)^2/\zeta_p(2w - 6)$, globally $\zeta(u)^2/\zeta(2u) = \sum 2^{\omega(n)} n^{-u}$ (Dirichlet coefficients exact through $n \leq 2000$; character expansions $\Phi_1 = \chi_0 + \chi_2$, $\Phi_2 = \chi_4 - \chi_2$, $\Phi_3 = 2\chi_0 - \chi_4 + \chi_6$, $\Phi_4 = \chi_8 - \chi_6$; core series $c_{k,0} = [1, 0, 2, 0, 2, 0]$). (C) The family form satisfies the exact linear relation $Q_{\text{fam}} = 2Q_\zeta(g) - 2Q_\zeta(g_b) + \text{Arch} + \text{Corr}$ ($\text{Prime}_F = 2P_\zeta(g) - 2P_\zeta(g_b)$) with max rel 9×10^{-16} ; $Q_{\text{fam}} = Q_F + \text{Corr}$ with rel 2×10^{-16} . **Obstruction 1:** the doubling term enters with a *minus* (family positivity does *not* imply $Q_\zeta \geq 0$). **Obstruction 2:** the Plancherel correction is demonstrably non-automorphic (no polar/arch absorption; no finite Euler factor; identified as $e^{-\sum p^{-u}}$ -type). Finite-class positivity $Q_{\text{fam}} \in [4.369, 11.486]$ is measured [C]; dense-class positivity stays open / RH-adjacent and is *not* claimed. Classical Weil 1952 / Gelfand / $\text{SU}(2)$ –Chebyshev / Sato–Tate / 2^ω -series / GNS named classical; NEW is the compiler-native identification and the machine-checked isolation of the two obstructions. Honest fences: *not* “almost RH” — the claim structurally shows why family positivity does not imply RH; ZETA.HP.CARRIER untouched; no marker moves. [E] (A/B/C/obstructions exact) / [C] (finite-class Q_{fam})

Amplitude route and positive linear carrier (RTF.GNS.AMP.01). The route out of the square plane behind the two RTF.GNS.WEIL.01 obstructions is now machine-checked ([verification/v540_amplitude_linear_carrier.py]; 34 checks, ~ 3 s; promotion of discovery T67–T72; companion to HECKE.GEOM.RTF.01/RTF.GNS.WEIL.01). (A) The amplitude Dirac $D = \begin{pmatrix} 0 & V \\ V^\top & 0 \end{pmatrix}$ with $V[d, m] = \sqrt{wd} b(d) \chi_d(m)$, $wd = |d|^{-1}$, exists exactly and is Hecke-equivariant: $D = D^\top$, $D^2 = \text{diag}(VV^\top, K)$ with $V^\top V = K$ (rel 2.7×10^{-16}), spectral $\pm$ -symmetry; m -side intertwining $VA_p^\top = \hat{A}_p V$ exact on the p -free locus ($p = 3, 5, 7$) and d -side Shimura $b(dp^2) = b(d)(a_p - \chi_d(p)p)$ integer-exact on 644 pairs. (B) The geometric polarisation $b = N_+ - N_-$ is exact (parity-split lattice count of the quinary form, plus the Θ -monomial identity coefficientwise); $\Theta = N_+ + N_-$ is a *pure* Siegel–Weil Eisenstein $T(p^2)$ -eigenform with eigenvalues $\sigma_3(p) = 1 + p^3$ ($p \leq 13$ exact; cusp component zero), carrying the Cohen seed identity $\Theta(d) = -48 L(-1, \chi_d)$ exact-rational on 159 live d (Cohen 1975: $H(2, d) = L(-1, \chi_d)$; generalised Bernoulli $B_{2, \chi}$). (C) Every coefficient bilinear form inherits the even- k deletion: the Cauchy–Littlewood window lemma with *free* Satake symbols has numerator $1 - stuv X^2$, hence $1 - p^6 X^2$ pair-independently at $st = uv = p^3$ (five channels checked); simultaneously the deletion *is* the square-class double counting of the towers, $\text{fam}[1, 0, 2, 0] + 2b[0, 2, 0, 2] + \delta_{k1} = [2, 2, 2, 2]$ (generating-function exact; Möbius square sieve). (D) The positive linear carrier $\ell^2(d, \mu)$ with $\mu = 48 |L(-1, \chi_d)| |d|^{-a}$ carries *full* weights $\lambda_k = 1 + p^{3k} - (\chi p)^k$ (layer $[1, 1, 1, 1]$; no even- k kill) and the plus balance $Q = Q_\zeta(g_-) + Q_\zeta(g_+)$, $g_\pm = e^{\pm 3u/2} g$ (rel 2×10^{-15}); the b /doubling kernel enters *with plus* — the sign of the v539 minus inverted. (E) The completed functional equation $\Lambda_\Theta(s) = 8^{1-s} \Lambda_{\Theta^\dagger}(5/2 - s)$ holds exactly (Fricke closed via the Jacobi inversions; rel $\sim 10^{-40}$ modular and in the strip; poles $5/2$ and 0 swapped; centre $5/4$, plus-balance locus off-centre by exactly $1/4$); the guaranteed cone and the gap functional are FE-self-dual/-covariant. **Open boundary as content:** the positivity is *Euler-region* positivity (edge L -values, not the central line); the residual distance to the Weil cone is the FE-covariant gap functional λ^* on $n \equiv 6 \pmod{8}$ — named, measured, Farkas-certified, and not removable by any finite signed theta library. Classical Cohen 1975 / Siegel–Weil / Jacobi–Fricke / Landen / Hecke split-Mellin / Cauchy–Littlewood / Shintani / Weil 1952 / Farkas named classical; NEW is the compiler-native consolidation. Honest fences: *not* “almost RH”; ZETA.HP.CARRIER untouched; no marker moves. [E] (A–E exact) / [C] (sampled cone facts; λ^* named open)

Matching lemma and transport ledger package (RTF.GNS.LEDGER.01). The discovery proof package T78–T85 is now load-bearing as one module in which every core check is *re-computed*, not cited ([[verification/v541_matching_lemma_ledger.py](#)]; 33 checks, ~ 10 s; companion to HECKE.GEOM.RTF.01/RTF.GNS.WEIL.01/RTF.GNS.AMP.01). (A) *Window proof.* The matching lemma of the hybrid certificate recipe is *proved exact-integer on* $[4, 10^6]$: the certificate $7S(j) < 40A(j)$ (i.e. $\rho_{\text{design}}F(j) < 1$ with $\rho = 21/20$) holds at *every* atom $j \leq 10^6$ (0 violations over 939 870 clash atoms; full enumeration, proven no-overflow, big-integer rechecks), with the exact Fraction margin $X = 22242575693/270725400000 = 0.082159$ at the exact argmax $j^* = 554400$ and $\rho_{\text{crit}} = 1.1440 > 21/20$; the maximal-greedy identity makes the certificate *uniform* in the target set M and the greedy weights. Structure laws at 0 tolerance: the 8-law $\Theta(4n) = 8\Theta(n)$ ($n \leq 250000$); the seed-tower divisor formula $\Theta(4^a s f^2) = 8^a \Theta(s) \alpha_s(f)$ and the ψ w -table (all $n \leq 50000$; Euler factor and w -recursion sympy-exact); the Cohen seeds $\Theta(s) = c L(-1, \chi_\Delta)$ with *one* exact-rational constant per residue class $\{-48, -80, -8, -8\}$ on all squarefree $s \leq 2000$; the bracket $\Theta \leq 2|\psi| \leq 3\Theta$ on the full window. (B) *Transport ledger.* The ledger $Q_{\text{Weil}} = Q_{\text{cert}} + \Delta_{\text{arch}} + \Delta_2$ closes *exactly*: $\Delta_{\text{pole}} \equiv 0$ and $\Delta_{\text{conv}} \equiv 0$ are *proven identities* (sympy kernel collapse; quadrature 2.8×10^{-16} ; exact scale invariance), the residual is 4.4×10^{-16} on 24 exact-autocorrelation battery rows, and the odd-prime side of the full Weil functional agrees *exactly* with the certified plus combination $P_\zeta^{\text{odd}}(h) = P_\zeta(g_-) + P_\zeta(g_+)$ (rel 3.9×10^{-16}): the prime side was never the wall. (C) *Character-exact signed envelope.* $-\psi = (\chi_{-4} + \frac{1}{4}\chi_8 + \frac{1}{4}\chi_{-8}) \cdot \Theta$ coefficient-exact on all odd $n \leq 50000$ on both independent builds (character system solved sympy-exact); the signed certificate $7S_{\text{net}} < 40A$ holds with 0 violations and margin factor $8.86\times$; the confinement is a *set equality* on 10^6 : $\{\text{zero-credit clash atoms}\} = \{\chi_{-4}\text{-coherent odd composites}\}$. (D) *The archimedean term is internal.* Via Legendre duplication $(2\pi)^{-s}\Gamma(s) = \frac{1}{2}\Gamma_{\mathbb{R}}(s)\Gamma_{\mathbb{R}}(s+1)$ (sympy-exact + 1.8×10^{-41}) the ledger kernel obeys $k_\zeta = K_{\text{fam}} - K_{\text{shift}}$ *pointwise* (7.9×10^{-31} on 51 points; Gauss digamma anchor at $t = 0$), and $\Delta_{\text{arch}}(h) = A_{\text{fam}}(h) - A_{\text{shift}}(h)$ on the battery subset at rel 9.9×10^{-16} — I5 becomes the self-consistency of *one* heat family. (E) *λ -equivariant channel.* The CM carrier $g_\lambda = \sum_{\mathfrak{a}} \lambda_1(\mathfrak{a}) q^{N_{\mathfrak{a}}}$ exists exactly in two independent routes ($\mathbb{Z}[i]$ lattice sum of z^4 vs Cornacchia reconstruction, 0 mismatches $n \leq 10^4$; CM laws exact; $\text{supp}(c_1) = \{\mathbb{Z}[i]\text{-norms}\}$ as a set equality with $c_1 \neq 0$ at every coherent atom; $\vartheta_3^2 \equiv r_2$ on 10^6); the canonical phase mean $\mu_1 = c_1/(c_0 d^2) \in [-1, 1]$ is exact, and the λ -window certificate over all $j \leq 10^6$ gives $\max 7|S_\lambda|/(40A) = 0.0653$ strictly below the unlifted 0.2365 (factor $3.6\times$; exact Fraction rechecks) — the sign-mixing of the Grossencharacter phases *is* the cancellation channel; coherent targets are reachable only by coherent rescalings (T81 reachability). (F) *Frontier facts [C]:* unlifted coherent-chain crossing $k^* = 14$ (exact Fractions, $\log_{10} N^* = 23.1$); absolute Robin tail factor 6.16 at $J = 10^6$ and divergent — the constant route closes nothing beyond the window. **Named limits as content:** (i) *one* classically-shaped open lemma remains — the correlated-cancellation lemma on credit-rich non-coherent tail atoms (uniform form); provably-shaped $\neq$ formal proof; (ii) I5 in one-family form ($Q_{\text{cert}} + \Delta_2 + A_{\text{fam}} - A_{\text{shift}} \geq 0$) is the single remaining TFPT-specific object — by the closed ledger *equivalent* to Weil positivity $\iff$ RH (Weil 1952): an equivalence *typing*, not a progress claim. Classical Gronwall 1913 / Robin 1983 *unconditional* / Cohen 1975 / Hecke 1918–1920 / Cornacchia / Fermat–Gauss / Dirichlet $L(1, \chi)$ / Mertens-AP / Landau 1908 / Legendre duplication / Weil 1952–Guinand–Bombieri / Alaoglu–Erdős named classical; NEW is the compiler-native consolidation and the machine-checked window proofs. Honest fences: no RH-evidence language; *not* “almost RH”; ZETA.HP.CARRIER untouched; no marker moves. [E] (A–E exact) / [C] (frontier facts; two named limits open)

The margin-chain identities of phase 2 (PRIME.MARGIN.IDENT.01). The identity/theorem block of the discovery parts T128–T131 is now load-bearing as one *narrow* module ([[verification/v542_margin_chain_identities.py](#)]; 44 checks, ~ 0.2 s; companion to RTF.GNS.LEDGER.01), recomputed on small frame-A windows: 24 seam instances ($n = 3 \dots 97$, $\rho = 1.25 \dots 4.00$, every inverted or diagonalised matrix ≤ 300), 43 graded/fine grid pairs and 400 randomised profiles, each identity a numerical residual against a preregistered tolerance *plus* a mutation control that must fail by $\geq 10^{-3}$ relative. *Border geometry.* $\sum_i (1 - g_i) = t_l + t_r$ exactly (5.5×10^{-13}), so the κ weights are a probability vector and the flat border vector gives

$\kappa_{\text{flat}} = 1$; the (T) four-term identity $\tau_{\text{dn}} = \|y\|^2 - m_{\text{prot}} + m_{\text{fill}} - V_{\text{bord}}$ holds at 2.2×10^{-16} , with τ recomputed independently from the full odd overlap matrix (1.1×10^{-13}). *Profile and curvature.* $p_{N-1} = 2 - p_0 + 2E$ at 2.3×10^{-16} , hence the equivalence $2E > p_0 \iff p_{N-1} > 2$ with 0 exceptions (both branches realised), and the Abel/Hölder chain $\kappa \leq 2 - p_0 + \frac{N-2}{N} \sum_j |\Delta p_j - \overline{\Delta p}| + (p_{\text{max}} - p_{N-1})$ holds per instance with 0 violations — while the same chain *without* the curvature term is violated on 88/400 profiles, which is what makes the curvature term load-bearing. *Assembly and compression.* The matrix-free two-scale assembly equals $J^T Q J$ (5.2×10^{-16}), the graded overlap operator equals J , and the C ea/Strang defect identity $S_{\text{graded}} - S_{\text{uniform}} = R^T X^{-1} R$ holds as a *matrix* identity on all 43 grids (2.1×10^{-14}) with a PSD defect — the compression error is one-sided *by* the identity. *Sandwich, sign, counting.* The secular sandwich $\varepsilon/(\|u\|^2 + \varepsilon/\mu_1) \leq \lambda_{\min}(Q) \leq \varepsilon/\|u\|^2$ holds on all 48 sections at which $A > 0$ and $\varepsilon > 0$ are *checked* (relative width ≤ 1.4448), with Albert’s sign equivalence verified in both directions; $S^{-1} > 0$ entrywise (measured 48/48) implies a sign-constant and simple ground vector (Perron–Frobenius), with a control matrix where hypothesis and conclusion fail together; and $n_{\text{run}} \leq n_{\text{cross}} + 1 \leq s_2 + 2$ together with $\sum_j |\Delta p_j - \overline{\Delta p}| = \text{TV}(P)$ hold without exception on a non-vacuous battery. **Named limits as content:** every statement is *per instance on small windows* — nothing is uniform in the zone index; the κ law is *not* claimed (T129 broke it), only the chain underneath it; positivity of the pole-free section at depth stays open and carries no floor value here; the Perron step has a *measured* hypothesis; and the defect identity says nothing about the *size* of the defect. Abel / H lder / Schur–Haynsworth 1968 / Albert 1969 / C ea 1964–Strang / Yserentant 1986 / Perron–Frobenius–Ostrowski 1937 named classical; Weil 1952 cited, never used as a criterion. [E] (per-instance identities; nothing uniform, no marker moves)

The lumped M -matrix pair (PRIME.MMATRIX.IDENT.01). The identity block of the discovery part T136 is load-bearing as a second narrow module ([`verification/v543_lumped_pair_identities.py`]; 35 checks, ~ 1 s; companion to PRIME.MARGIN.IDENT.01) on 20 frame-A seam windows ($n = 4 \dots 59$, $\rho = 1.25 \dots 4.00$, $h = 25 \dots 299$, every factorised matrix ≤ 300) giving 39 matrices with positive off-diagonals, 312 shifted ladder rungs and 8 whitened two-grid rungs. *The pair and the congruence.* With Δ the positive off-diagonal part and $L_\Delta = \text{diag}(\Delta \mathbf{1}) - \Delta$, the lumped pair $A_B = A + L_\Delta$ is Stieltjes and row-sum preserving (1.6×10^{-15}) with $A_B \succeq A$ — an *upper* bound on $\lambda_{\min}(A)$, never a floor; $A = A_B^{1/2}(I - W)A_B^{1/2}$ holds as a matrix identity (4.4×10^{-15}) with $\lambda_{\max}(W) = \rho(L_\Delta A_B^{-1})$, and the equivalence $\lambda_{\max}(W) < 1 \iff A \succ 0$ is checked in *both* directions, so the Loewner sandwich is circular by itself. *Anchor, ladder, Varga.* With $x = A_B^{-1} \mathbf{1} \geq 0$ and $A_B^{-1} \geq 0$ entrywise (Ostrowski 1937, checked), $\lambda_{\min}(A_B) \geq 1/\max_r x_r$ on all 39 rows (sharpest ratio 0.9070) from one solve and one sign check; lumping commutes with diagonal shifts and M -matrix inverses are entrywise monotone (Berman–Plemmons 1979), so the shifted ladder is *structurally* empty (0 exceptions of 312); and for the regular splitting $A_B = D_0 - N_0$ Varga’s $\rho(J) = \tau/(1 + \tau)$ is an identity (4.8×10^{-15}) whose Collatz–Wielandt bracket at the anchor vector gives an a-priori $\rho(J) \leq 0.9861 < 1$ with no eigenvalue anywhere. *One negative result [X].* The k -scanned M18 majorant is *valid* at every k (39 pairs, 0 violations) and still stays above its preregistered bar $1/2$ (best 3.080) while the measured bad mass is ≤ 0.0120 : the loss is the bound, the localisation term alone exceeds the bar, and the bar is not moved. **Named limits as content:** per instance on small windows, nothing uniform in the zone index, in D or in the gap; the T136 D -exponents are fits and stay in the sandbox; the anchor and Varga rows are implications with hypotheses checked at the instance. Stieltjes / Ostrowski 1937 / Fan 1958 / Berman–Plemmons 1979 / Varga 1962 / Collatz 1942–Wielandt 1950 named classical; Weil 1952 cited, never used as a criterion. [E] (per-instance identities) + [X] (one closed route; no marker moves)

The long-lag support structure (PRIME.LONGLAG.SUPP.01). The structure block of the discovery part T137 is load-bearing as the companion module ([`verification/v544_long_lag_support.py`]; 24 checks, ~ 0.2 s) on the same 20 windows (3–25 prime-power atoms, 20555 positive off-diagonal edges, 455 loaded stripes, 39 bracketed lumped comparisons). *Support and amplitude.* The kernel splits exactly as $c = c_{\text{arch}} - c_{\text{comb}}$ with $c_{\text{comb}} \geq 0$, and the archimedean section alone has *all* off-diagonals ≤ 0 on 20/20 windows, so every positive off-diagonal is comb-generated; all 20555 positive edges have their Hankel index $M - 1 - r - t$ inside the atom-hat index set (per-window fraction exactly

1.000000), so the support is a union of anti-diagonal stripes at prime-power atom positions, each stripe a perfect matching with *exactly* orthogonal edge vectors; and $\Delta_{rt} \leq c_{\text{comb}}[M - 1 - r - t]$ on all 20555 edges (sharpest ratio 0.99998532), with the Toeplitz lag in place of the Hankel index failing on 20415 of them. *The Green bracket.* For the lumped comparison, $q_J = 1 - \min_r 1/(d_r x_r) < 1$ is an identity (8.2×10^{-16}) needing no measurement, and the Neumann partial sum brackets the Green function entrywise from below *because the discarded terms are nonnegative*, the anchor-weighted tail from above, on 39 rows. *A death certificate* [X]. Gershgorin, every row-sum norm and every Collatz–Wielandt quotient at every *positive* weight are upper bounds for $\rho(|E|)$, $E = B^{-1}L_\Delta$; one rigorous lower bound therefore decides them all at once: $\rho(|E|) \geq 1.3141 \geq 1$ on 20/20 rows, so no member of the absolute-value envelope family can *ever* certify $\lambda_{\max}(W) < 1$ — while the *signed* radius $\rho(E) = \lambda_{\max}(W) \leq 0.99995781$ on the same rows. The absolute value alone destroys the cancellation. **Named limits as content:** per instance on small windows, no decay law and no amplitude law; the support statement is one inclusion; the amplitude bound does *not* bound $\lambda_{\max}(W)$, and the structure bound assembled from it does not beat the margin (T137: 0 of 35 windows) and is not promoted; the bracket is for the lumped comparison, not the target. Gershgorin 1931 / Neumann series / Haynsworth 1968 named classical; Demko–Moss–Smith 1984 cited as a model and *not* applied (the comparison is dense); Weil 1952 cited, never used as a criterion. [E] (per-instance structure) + [X] (one family closed; no marker moves)

The Hardy-core identities (PRIME.HARDY.IDENT.01). The identity blocks of the discovery parts T140 and T141 are load-bearing as the third companion module (36 checks, ~ 1 s) on 11 frame-A windows ($n = 8 \dots 139$, $h = 58 \dots 285 \leq 300$, 193–5102 positive off-diagonal edges, D spanning a factor 3.15). *Form lifting and finite core.* With $C = \text{diag}(\sqrt{\Delta})M$ and M the edge–interval incidence, $\text{Gram} = CHC^\top$ (3.0×10^{-15}), so $\text{rank}(\text{Gram}) \leq h - 1$; the covering kernel in closed geometric form $K_{rs} = W([r \wedge s, r \vee s])$ equals $M^\top \Delta M$ (1.3×10^{-15}); and $\rho(W) = \lambda_{\max}(K^{1/2}HK^{1/2})$ is an *exact* eigenvalue, agreeing with $1 - \lambda_{\min}(A, A_B)$ (5.4×10^{-14}) and with $\lambda_{\max}(\text{Gram})$ (7.8×10^{-16}) — three independent routes, no truncation — while $H = \text{diag}(s) + L_N$ (4.7×10^{-16}) reads the sign law as mass plus a long-range Dirichlet form. *The four Hardy identities.* $\rho(W) = u^\top Hu / u^\top K^+ u$ at $u = K^{1/2}v$ (6.6×10^{-13}); $L_N = D^\top BD$ with the signed crossing kernel $B_{kl} = \sum_{r \leq k \wedge l, k \vee l < s} N_{rs}$ (4.9×10^{-16}); $Q = B_+ \mathbf{1}$ (1.4×10^{-15}), so the Cauchy–Schwarz step is the Gershgorin diagonal step; and $DKD^\top = L_\Delta$ on the interior nodes (2.7×10^{-14}), whose diagonal is the endpoint edge mass. The crossing kernel of $|N|$ does *not* reproduce L_N : the signs are load-bearing. The closed Muckenhoupt quantity $A_{M0} = \max_k Q_k(\Delta \mathbf{1})_{k+1}$ follows from two routes agreeing to 8.3×10^{-15} ; its *size* is reported, not promoted. The checkerboard split $\text{Band}_b = D + A_{\text{even}} + A_{\text{odd}}$ is entrywise exact on 9 (window, b) pairs, with three Weyl steps dominating. *Two shapes, promoted as valid and not by size.* Every Loewner step is Cholesky-certified; the additive Hardy chain dominates $\rho(W)$ on 11/11 ($3.074\text{--}10.758\times$), its exact Weyl split dominates too ($1.691\text{--}2.503\times$) and floors the additive family, and the joint shape $\Lambda(H, Y) \cdot \Omega(Y)$ dominates on 11/11 for a positive definite Jacobi Y — the same Y without its mass term is singular and the certificate refuses. *A negative typing* [X]. Over the D range the exactly bounded object spreads $1.36\times$ while every absolute variant spreads $\geq 54.94\times$; the power laws ($D^{-0.138 \pm 0.094}$ signed against $D^{-0.776 \pm 0.352}$ and $D^{-2.748}$, $D^{-2.820}$, $D^{-2.918}$) are fits and labelled as fits. **Named limits as content:** per instance on small windows, nothing uniform in the zone index or in D ; the bound rows promote *validity* and not size — neither shape beats the target — and the finite-core row is an identity, so the zone-uniform discrete Hardy inequality stays open. Abel summation / Cauchy–Schwarz / Weyl 1912 / Gershgorin 1931 / Muckenhoupt 1972 / Bradley 1978 / Opic–Kufner 1990 / Gantmacher–Krein 1950–1960 named classical; Weil 1952 cited, never used as a criterion. Machine-checked: [verification/v545_hardy_core_identities.py]. [E] (per-instance identities) + [X] (one route typed; no marker moves)

The capacity-chain identities (PRIME.CAPCHAIN.IDENT.01). The identity spine of the capacity cycle T142–T145 is load-bearing as the fourth companion module (26 checks, ~ 0.5 s) on 12 frame-A windows ($n = 4 \dots 139$, $h = 26 \dots 285 \leq 300$, D spanning 7.6×10^{-3} – 2.8×10^{-2} ; the covering kernel positive definite on 11, where the first two items run). *The capacity decomposition.* $K^{-1} = D^\top J^{-1}D + xx^\top / \text{cap}$ with $J = DKD^\top$ (3.8×10^{-15} entrywise), $x = K^{-1}\mathbf{1}$, $\text{cap} = \mathbf{1}^\top K^{-1}\mathbf{1}$,

as a matrix identity (1.4×10^{-13}); under the congruence the Dirichlet half is an *orthogonal projection* — $\Omega = 1$ exactly (5.1×10^{-12}), where T141 had guessed 20.7–2724 — and its complement is the capacity rank-one. *The exact capacity–Rayleigh identity.* The assembled quotient for $1 - \rho(W)$ equals $v^\top(K^{-1} - H)v/v^\top K^{-1}v$ for every v (9.5×10^{-14}), reproduces $1 - \rho(W)$ at the exact top eigenvector under the declared 1/gap conditioning, and satisfies the inf property; on the constant vector the whole denominator *is* the capacity term. *Cholesky nestedness.* $\text{cap}_E([a, a+j])$ is a prefix sum of squared forward-substitution entries of one triangular solve (3404 interval capacities at 2.0×10^{-14} , exhaustive on the median window); the reversed ordering fails at $\geq 4.6 \times 10^{-1}$. *The whitening certificate.* $\lambda_{\max}(W) \leq \kappa_{\text{up}} \leq$ Gershgorin ceiling (1.2245–1.4236 against ≤ 2.1213), direction-correct, refusing at the shrunken shift. *The M_4 split and the certified chain.* The mass half of Maz’ya’s Markov step is a pointwise theorem, $\sigma_{\text{tot}} = 0.2197\text{--}0.4425 < 1$ on every window; licence 4 holds with $|R|$ and *not* R^+ (witness $[[1, -\frac{1}{2}], [-\frac{1}{2}, 1]]$ built in); and $S1'$ is *certified per window on both routes*, better route $c_0 = 2.5154\text{--}4.2265$ against the classical Dirichlet value 4, with Charikar’s cited density constant bracketed by exhaustive enumeration. **Named limits as content:** per instance on small windows, nothing uniform in the zone index or in D ; the c_0 is *measured* at the minimiser’s own layer cake and the a-priori level lemma (T145’s L1) stays open — Maz’ya’s strong-type half is not used as a theorem anywhere. Maz’ya 1985 / Muckenhoupt 1972 / Fukushima–Ōshima–Takeda 1994 / Miclo 1999 / Charikar 2000 / Goldberg 1984 / Gershgorin 1931 named classical; Weil 1952 cited, never used as a criterion. Machine-checked: `[verification/v546_capacity_chain_identities.py]`. **[E]** (per-instance identities and certified inequalities; no marker moves)

The level-lemma identities (PRIME.LEVEL.LEMMA.01). The a-priori core of T146 is load-bearing as the fifth companion module (20 checks, ~ 0.4 s) on 12 frame-A windows ($n = 4 \dots 139$, $m = 26 \dots 285 \leq 300$), 9 random positive definite forms for the form-independent items, and a stress ladder $m = 64 \dots 288$ — the companion to the capacity-chain module, whose certified chain carried a constant *measured* at the minimiser’s layer cake: here the same constant is a functional of the form alone. *The union identity for nested cakes.* $\sum_{j,l} c_j c_l |S_j \cup S_l| = 2T \|\psi_t\|_1 - \|\psi_t\|^2$ — form-independent and $O(m)$, the reason the cake base is movable; exact (0.0) against the $K \times K$ double sum and against union sizes computed with no nestedness used (controls 3.2×10^{-2} / 4.8×10^{-3}). *The counting lemma at a general base.* $G(b) \leq 2b^2 \|v\|_\infty \|v\|_1 / \|v\|^2 + \varepsilon$ for *any* vector at all 6 preregistered bases; the classical dyadic 8 recovered at $b = 2$ exactly, the base free (gain $4/b_*^2 = 3.7612$); the one-level-down cake breaks the domination on every draw, so the +1 is load-bearing. *The resolvent delocalisation bound.* $\psi = \lambda R \psi$ (9.0×10^{-14} , an identity — no Davis–Kahan step anywhere) $\Rightarrow \|\psi\|_\infty \leq \lambda_{\text{up}} \max_j \|Re_j\|$ with Rayleigh–Ritz λ_{up} (1.0592–1.2883 of the gap) and residual-paid certified column norms, within 1.061–1.337 of the truth; the lever: the columns sit 2.85–10.17 *below* the trivial ceiling $1/\lambda_{\min}$, giving $\Gamma = 1.8356\text{--}2.2797$ against the useless $\sqrt{m} = 5.1\text{--}16.9$. *The composite level lemma.* $c_0^{\text{ap}} = 2b^2 \Gamma \min(1, \Gamma_1) + \varepsilon = 3.9042\text{--}4.8488$ (slack 1.986–2.418 over the measured constant, under the ceiling $16 = 4 \times$ Maz’ya’s Dirichlet 4), the certified window inequality $\text{cert_lam_max}(R) \leq c_0^{\text{ap}} \Psi_{\text{abs}}$ on 12/12, chain loss 0.0423–0.1272 of the exact gap; the built-in no-go discriminator ($R = aa^\top + \varepsilon I$, $a_i = i^{-1/2}$: Γ grows $3.676 \rightarrow 6.798$, ratio $1.85 \geq 1.5$, $c_0^{\text{ap}} = 11.10$ exceeds every window value) fails exactly where it must, the theorem half surviving and the Dirichlet control flat (1.0063). **Named limits as content:** per instance on small windows — a finite list with an explicit maximum; nothing uniform in the zone index or in D , and explicitly *no* statement for all D (the asymptotic delocalisation of the Green columns stays open, typed open); the energy-route σ_{tot} stays measured. Maz’ya 1985 / Rayleigh 1877–Ritz 1909 / Cauchy–Schwarz / Miclo 1999 / Davis–Kahan 1970 (cited, deliberately not used) / Charikar 2000 / Goldberg 1984 / Gershgorin 1931 named classical; Weil 1952 cited, never used as a criterion. Machine-checked: `[verification/v547_level_lemma_identities.py]`. **[E]** (per-instance identities and certified inequalities; no marker moves)

The Green/Szegő identities (PRIME.GREEN.SZEGO.IDENT.01). The identity/certificate core of T147+T148 is load-bearing as the sixth companion module (21 checks, ~ 0.4 s) on 12 frame-A windows ($n = 4 \dots 139$, $m = 26 \dots 285 \leq 300$), 7 random positive definite forms for the form-independent items, and a weight-class model ladder $m = 64 \dots 288$ — the companion to the

level-lemma module: here the two scalars its constant is made of are certified. *The factorisation identity.* $\Gamma = \sqrt{m} \lambda_{\text{up}} \max_j \|Re_j\| = \sqrt{Q^*} \cdot S_w$ with $S_w = \lambda_{\text{up}} \|R\|_F$ purely spectral and $Q^* = m \max_j (R^2)_{jj} / \text{tr}(R^2)$ purely geometric — exact at 2.2×10^{-16} ; the certified factorisation dominates the true Γ everywhere ($S_w = 1.1186\text{--}1.5777$, $Q^* = 2.0879\text{--}2.8634$). *The bottom-block constant.* $\sum_j (\Pi_B)_{jj} = |B|$ exactly (3.3×10^{-16} ; B the preregistered $\theta = 10$ block, $|B| = 3\text{--}5$), so $Q_B \geq |B|$ is a theorem; measured $Q_B/|B| = 1.5091\text{--}1.7980$ inside the preregistered band $[1, 2.2]$ against the Szegő/Widom sharp reference $2|B|$ — the coordinate-projector control overshoots by ≥ 3.9 . *The second-difference ℓ^1 ceiling.* $|a_k| \leq \min(1, \|s_k\|_\infty \|L_0^p \psi\|_1 / \mu_k^p)$ from the exact Dirichlet eigenpair, with the m -free closed form $\|a\|_1 \leq 2\sqrt{\nu} + 1$ ($\nu = (m+1)^{3/2} \|L_0 \psi\|_1 / (2\sqrt{2})$) and the chain $m\|\psi\|_\infty^2 \leq 2\|a\|_1^2$; Dirichlet calibration $\|a\|_1 = 1$ exact, $\nu \rightarrow \pi$. *The layer-cake S_w certificate.* $\text{LDL}^\top$ inertia counts equal the direct spectral counts on all 96 (window, rung) pairs, and the layer cake certifies $S_w \leq 1.3288\text{--}1.8220$ (true $1.1186\text{--}1.5777$); dropping the bottom layer breaks it everywhere. *The Liouville transform.* $m\|\psi\|_\infty^2 \leq 2\kappa_\Lambda \text{ceil}(\hat{\varphi})^2$ with $\varphi = \Lambda^{-1/2}\psi$ and a-priori $\kappa_\Lambda = \max \Lambda / \min \Lambda = 1.3868\text{--}2.5805$, stepwise on every bottom-block mode; the weight-class discriminator at matched κ_Λ (mismatch 0.0013): the BV multiplier flat (1.003), the rough one (TV larger by 162.6) grows by 19.9 — the roughness, not the conditioning. **Named limits as content:** per instance on small windows — a finite list with an explicit maximum; nothing uniform in the zone index or in D , and explicitly *no* statement for all D (the one open lifting input, $\nu_L \leq F(\text{form})$ m -free / bounded TV($\log \Lambda$), stays open, typed open); the symbol-level order-2 hypothesis is refuted (T148: $f(0) < 0$ on every surface window) and enters nowhere as a theorem; σ_{tot} stays retired as a route. KMS 1953 / Widom 1958–Basor–Ehrhardt 2009–Böttcher–Silbermann (the mechanism’s address, never an authority) / Sylvester 1852–Bunch–Kaufman 1977 / Euler 1735 / Rayleigh 1877–Ritz 1909 / Davis–Kahan 1970 (cited, computed vacuous, not used) / Gershgorin 1931 named classical; Weil 1952 cited, never used as a criterion. Machine-checked: [\[verification/v548_green_szego_identities.py\]](#). **[E]** (per-instance identities and certified inequalities; no marker moves)

The gauge/parity identities (PRIME.GAUGE.PARITY.IDENT.01). The identity/certificate core of T149+T150 is load-bearing as the seventh companion module (21 checks, ~ 0.3 s) on 12 frame-A windows ($n = 4 \dots 139$, $m = 26 \dots 285 \leq 300$), 48 gauge assemblies (Jacobi, const, moving-geometric-mean, and the arithmetic-free archimedean diagonal), the full-section parity items on the 6 windows with $M \leq 300$, and a parity control model at $m = 2001$ — the companion to the Green/Szegő module: here the gauge *freedom* around that machinery and the parity *structure* underneath it are certified. *The gauge freedom.* For any positive diagonal $\tilde{\Lambda}$, $\lambda_{\min}(A) \geq \text{cert } \lambda_{\min}(\tilde{E}) \cdot \min_j \tilde{\Lambda}_j$ is a valid certified lower bound (an identity plus one Rayleigh step; family maximum valid); $\Gamma = \sqrt{Q^*} S_w$ holds gauge-invariantly at 2.4×10^{-16} ; the const gauge has TV($\log \tilde{\Lambda}$) = 0 *exactly* with the certified entrywise sandwich $\sigma = 1.3868\text{--}2.5805$; on the shifted indefinite form the completed-Cholesky floor refuses on 12/12. *The two-regime discriminator.* The flutter smoother removes TV by $\geq 2.88\times$ while the Liouville-transported ν_L moves ≤ 0.0088 — even the rough re-gauge moves it ≤ 0.0046 — while the untransported functional jumps 0.41–9.76 under the same rough gauge: the roughness is in the form, not in the multiplier (T149’s withdrawal of the TV hypothesis, wired per instance). *The explicit zone diagonal.* $\Lambda_r = c_0 - c_{M-1-2r} + \sum_{s \neq r} \max(c_{|r-s|} - c_{M-1-r-s}, 0)$ at residual 5.9×10^{-16} , the exact split $c = c^{\text{arch}} + c^{\text{atom}}$ with $c^{\text{atom}} \leq 0$, the closed atom budget $\|c^{\text{atom}}\|_1 \leq 4B\sqrt{N}$ ($B = 1.038821$ verified on the table); the archimedean section alone is indefinite on 12/12 (the atoms are co-responsible for the positivity — the additive perturbation reading is structurally empty). *The parity compression.* $U_-^\top T_M(c) U_- = T - H$ exactly (cross block 1.4×10^{-16}): the reflection-odd section is the compression of the full symmetric Toeplitz section onto its antisymmetric parity sector, and the negative inertia of the sign-changing symbol sits entirely in the *even* sector (odd sector zero, $\text{LDL}^\top$) — $f(0) < 0$ explained rather than worked around. *The parity-sine calibration.* The parity sines are exact eigenpairs of the parity Laplacian (corner 3 forced by antisymmetry), $\nu_k^P = \pi k^2$ on the exact model to 3.0×10^{-5} , and the per-window ladder $\nu_k^P \leq Ck^2$ holds with certified $C = 11.70\text{--}17.28$ and the derived per-mode ceiling $m\|\psi_k\|_\infty^2 \leq 2(2\sqrt{C}k + 1)^2$. **Named limits as content:** per instance on small windows — a

finite list with an explicit maximum; nothing uniform in the zone index or in D , and explicitly *no* statement for all D (the one open input, an m -free C in the odd-sector ladder for a sign-changing symbol, stays open, typed open); TV(log Λ) stays withdrawn as a hypothesis; σ_{tot} stays retired as a route. KMS 1953 (parity sector) / Widom 1958–Basor–Ehrhardt 2009–Böttcher–Silbermann (the parity sectors as the algebra’s natural objects; the sign-changing symbol is exactly what Basor–Ehrhardt does not cover) / Sylvester 1852–Bunch–Kaufman 1977 / Chebyshev 1852 (verified on the table) / Abel / Weyl 1912–Bauer–Fike 1960–Davis–Kahan 1970 (cited, computed dead, not used) / Rayleigh 1877–Ritz 1909 / Gershgorin 1931 named classical; Weil 1952 cited, never used as a criterion. Machine-checked: [verification/v549_gauge_parity_identities.py]. [E] (per-instance identities and certified inequalities; no marker moves)

The odd-sector identities (PRIME.ODD.SECTOR.IDENT.01). The identity/certificate core of T151 is load-bearing as the eighth companion module (19 checks, ~ 0.2 s) on 12 frame-A windows ($n = 4 \dots 139$, $m = 26 \dots 285 \leq 300$), parity control models at $m = 241$ and $m = 2001$, and a no-go size ladder $m = 64 \dots 256$ — the companion to the gauge/parity module: here the *reroute* off the quadratic ladder is certified — through the pencil against the parity Laplacian and one elementary Sobolev step, to a per-mode bound *linear* in k . *The grid step-over.* The parity sines live on the shifted grid $\theta_k = 2\pi k/N$ with $\mu_k^P = 2 - 2\cos\theta_k$ exactly; the symbol has $f(0) = -47.6 \dots -6.3 < 0$ with a nonempty negative window $[0, \theta_c)$, and $\theta_c/\theta_1 = 0.371\text{--}0.485$ on every window: the odd grid steps over the negative window — odd-sector positivity is a grid fact. The honest negative beside it, promoted as content: the symbol-only pencil floor is vacuous ($\rho \geq 1$ on 12/12) and the symbol moves by 0.45–3.27 across one mode spacing — the local model is a *matrix* statement, not a symbol statement. *The certified bottom ladder.* $\lambda_k(A) \leq S\mu_k^P$ for $k \leq 8$ by completed LDL^T counts (Sylvester 1852), $S = 1.1019\text{--}6.4725$ — order 2 in k , with $f(0) < 0$ absorbed by the Rayleigh floor $L_P \geq \mu_1^P I$. *The Sobolev step.* $\|\psi\|_\infty^2 \leq 2\|\psi\|_2\|\nabla\psi\|_2$, licensed by the odd sector’s virtual node $\psi_{-1} = 0$; on the exact parity model the pencil is (1, 1) (certified 1.1×10^{-12}) and the per-mode price is exactly π (5.1×10^{-8}). *The closed archimedean minimum.* $\min_r \Lambda_r^{\text{arch}} = c_0^{\text{arch}} - c_{M-1}^{\text{arch}}$ exactly (residual 0.0), attained at $r = 0$, from two per-window-checked shape properties of the smooth kernel. *The linear per-mode bound.* With the certified pencil floor $\kappa = 0.1939\text{--}0.3599$ and $K_{\text{bot}} = S$, $b_k = 2m\sqrt{K_{\text{bot}}\mu_k^P/\kappa} \leq C_S k$ with $C_S = 12.081\text{--}26.953 \leq 2\pi\sqrt{K_{\text{bot}}/\kappa}$ — linear where the quadratic ladder grew, nothing additive anywhere; on the T145 no-go the bottom pencil ratio explodes ($\times 15.8$ over $m = 64 \dots 256$) while the parity model holds 1. **Named limits as content:** per instance on small windows — a finite list with an explicit maximum; nothing uniform in the zone index or in D , and explicitly *no* statement for all D (the one open input, the m -freeness of the bottom pencil ratio $R = K_{\text{bot}}/\kappa = 3.7362\text{--}18.6607$, stays open, typed open); the symbol route stays certified vacuous. KMS 1953 / Widom 1958–Basor–Ehrhardt 2009–Böttcher–Silbermann (the symbol question’s address — the computed answer: the pointwise symbol language does not extend to the bottom mode) / Weyl 1912 / Parlett / Hardy–Littlewood–Pólya 1934–Agmon 1965 / Sylvester 1852–Bunch–Kaufman 1977 / Rayleigh 1877–Ritz 1909 / Wilkinson 1968–Higham 2002 / Chebyshev 1852 named classical; Weil 1952 cited, never used as a criterion. Machine-checked: [verification/v550_odd_sector_identities.py]. [E] (per-instance identities and certified inequalities; no marker moves)

The fixed-size Ritz ceiling certificate (PRIME.RITZ.CEIL.01). The certificate core of T154 is load-bearing as the ninth companion module (16 checks, ~ 0.4 s) on the 12 *deepest* prime-power zones admitting a frame-A window inside the cap ($n = 29 \dots 139$, $m = 96 \dots 291 \leq 300$ — declared, not tuned: on the shallowest zones $n \leq 11$ more than one bottom direction leaves the sine block and the fixed-depth certificate is not claimed there), a no-go size ladder $m = 48 \dots 288$, parity controls at $m = 64, 128, 256$ and 3 Dirichlet control windows — the companion to the odd-sector module: here the *fixed-size* replacement for the ceiling step is certified. *The direction lemma.* $\lambda_k(A) \leq \theta_k$ for every $k \leq \dim$ — Ritz values are upper bounds for the eigenvalues of the *same index* (Courant–Fischer 1920 / Cauchy 1829), verified against exact spectra with zero one-sided excess — so the ceiling needs *no* residual argument; the reversed reading is refuted on 12/12 (overshoot ≥ 18.6). *The sixteen-column certificate.* $\text{span}\{t_1..t_8\} + A^{-1}L_P \text{span}\{t_1..t_8\}$ (dimension 16 fixed,

independent of m ; 8 completed Choleskys of matrices $\leq 8 \times 8$ carries $K^F = 1.1019\text{--}1.5845$ attaining the inertia-certified K_{bot} to $\leq 9.9 \times 10^{-7}$ against the declared cap 10^{-4} on 12/12 — the 8 size- m LDL^T counts retired from the ceiling step (controls: the sines alone overshoot 41.6–739; corrupted Green columns break the attainment on 12/12). *The one-direction anatomy.* Median of the seven aligned angles $0.03\text{--}0.14^\circ$, the largest $83.79\text{--}89.70^\circ$ (measured): the single misaligned direction *is* the collapse price $\lambda_1(A)/(t\mu_1^P) = 4.408\text{--}7.042$, recovered in full per window by one Cholesky of $A - \gamma I$ (size m , certified, not fixed size). *The no-go discriminator.* On the T145 reconstruction $c(l) = 1/(1+l)$ the floor survives ($t = 0.05$) but K_{bot} explodes $\times 35.4$ and the ceiling explodes with it $\times 57.5$ — no manufactured closure; the Dirichlet control (the Hankel half removed) is certified non-positive on 3/3: the reflection half is load-bearing for positivity. **Named limits as content:** per instance on small windows — a finite list with an explicit maximum; nothing uniform in the zone index or in D , and explicitly *no* statement for all D ; the attainment is not promoted to a theorem (the no-go overshoots — it is a property of the real section); the two open inputs after T154 (R1', an m -free floor for $\lambda_{\min}(B_{HH})$ on the bulk parity block; R2', an m -free lower bound for the L_P floor on the Ritz complement, arithmetic-free and worth 91–100% of the collapse price) stay open, typed open. Courant–Fischer 1920–Cauchy 1829 / KMS 1953 / Schur 1917–Haynsworth 1968 / Sylvester 1852–Bunch–Kaufman 1977 / Weyl 1912 / Temple 1928–Kato 1949 (a floor device, not used here) / Kaniel 1966–Paige 1971 (quoted, never a bound) / Björck–Golub 1973 / Wilkinson 1968–Higham 2002 / Chebyshev 1852 named classical; Weil 1952 cited, never used as a criterion. Machine-checked: [verification/v551_ritz_ceiling_certificate.py]. [E] (per-instance theorems and certified inequalities; no marker moves)

The four fixed-size angle instruments (PRIME.ANGLE.INSTR.01). The instrument cores of T155 and T157 are load-bearing as the tenth companion module (18 checks, ~ 0.4 s) on the same declared surface as the Ritz module (the 12 deepest in-cap frame-A windows, $n = 29 \dots 139$, $m = 96 \dots 291 \leq 300$; a no-go size ladder $m = 48 \dots 288$; parity and Dirichlet controls at $m = 64, 128, 256$) — none of the four closes an angle, and none is promoted as anything but a per-instance instrument. *The complement-floor certificate (T155).* $\min_{v \perp W} v^T L_P v \geq \mu_{13}^P - \lambda_{\max}(M^{1/2}(I - GG^T)M^{1/2})$ with $G = T_{12}Q_W$ — an identity plus one certified Rayleigh bound, a theorem for every m and every subspace; never above the exact size- m complement bottom, agreement $\leq 1.01 \times 10^{-3}$ against the declared cap 2×10^{-3} on this small surface, both closed-form controls to 2.0×10^{-11} (including the *worthless* configuration $W = \text{span}\{t_{2..t_{13}}\} \rightarrow \mu_1^P$: the instrument reports its own worthlessness); a corrupted overlap row moves it by ≥ 0.549 . *The sine-block confinement (T157).* From the certified pencil floor and ladder ceiling alone, $\|\gamma_H\|^2 \leq \lambda_1/(t\mu_{17}^P) \leq (S/t)/\rho_{17} = 0.0153\text{--}0.0245 < 1$ on 12/12 — the bottom eigenvector lives 97.5–98.5% inside the first 16 parity sines by a per-instance theorem (measured tail $6.7 \times 10^{-6}\text{--}3.0 \times 10^{-5}$); with the pencil ceiling κ instead of the ladder S the bound is vacuous (17.6–477), and the inflated floor is refuted at $e_1(A)$ by $3.3 \times 10^3\text{--}1.0 \times 10^5$. *The Schur-entry identity (T157).* $s = \mu_1^P t_1^T A^{-1} t_1 = (B^{-1})_{11}$ exactly (1.1×10^{-11}), $(B^{-1})_{LL} S_L = I$ to 5.8×10^{-11} with S_L the fixed-size 16×16 Schur complement the chain already forms: $1/s = 4.0460\text{--}5.8962$ is an $O(1)$ number carried by *one* entry, while the inversion-free Cauchy–Schwarz ceiling misses by $250\text{--}7.27 \times 10^3$ (the cancellation is nearly complete); a one-index shift breaks the identity by 1.55–2.15. *The adaptive Lipschitz ceiling (T157).* Interval bisection with the local Lipschitz constant certifies $\inf f^{\text{arch}}/(4 \sin^2(\theta/2)) \geq 0.25$ on 12/12 windows in 519–1201 evaluations (a uniform grid at the global constant would need up to $61 \times$ more); fed a target above the true grid minimum it *refuses* on 12/12. *The no-go discriminator.* On $c(l) = 1/(1+l)$ the floor survives ($t = 0.05$) while the confinement goes vacuous ($\times 52.0$) and $(S_L)_{11}$ explodes ($\times 60.0$) against the flat 4.05–5.90 real band; parity control exact to 4.9×10^{-14} , the Dirichlet residual hits $2/\sqrt{N}$ to 7.4×10^{-5} . **Named limits as content:** per instance on small windows — nothing uniform in the zone index or in D , and explicitly *no* statement for all D ; nothing about the angles is closed — the two open terms after T157 (an m -free lower bound on $\hat{g}_1^2$, equivalently an m -free upper bound on the one Schur diagonal entry; an m -free R1'' domination with a non-shrinking margin) stay open, typed open. KMS 1953 / Schur 1917–Haynsworth 1968 / Courant–Fischer 1920–Cauchy 1829 / Cauchy–Schwarz (refuted as a ceiling, wired as a control) / Kantorovich 1948 / Temple 1928–Kato 1949 (floor devices, not used

here) / Sylvester 1852–Bunch–Kaufman 1977 / Wilkinson 1968–Higham 2002 / Chebyshev 1852 / Szegő 1915–Widom 1958 (a named address, never a licence) named classical; Weil 1952 cited, never used as a criterion. Machine-checked: [verification/v552_angle_instruments.py]. [E] (per-instance identities and certified inequalities; no marker moves)

The exact-form identities (PRIME.EXACT.FORM.IDENT.01). The theorem cores of T158 and T159 are load-bearing as the eleventh companion module (25 checks, ~ 0.2 s), recomputed on the same declared surface as the two modules above (12 deepest in-cap frame-A windows, $m = 96 \dots 291 \leq 300$; no-go ladders $m = 48 \dots 288$ on *both* T145 forms; parity and Dirichlet controls at $m = 64, 128, 256$) — the algebra of the sixteen-form itself, and it closes neither open term. *The Thomson dual form and the positive Cholesky ladder (T158).* $s = (B^{-1})_{11} = \max_x (2x_1 - x^T Bx)$ with the direction *executed* (8 random trials per window never exceed s); the ladder g_K from one Cholesky of the leading 16×16 block: terms strictly positive, partial sums monotone, $1/s \leq 1/g_{16} \leq 1/g_1 = B_{11}$ literally, tight to 1.0642–1.1522; a 5% corruption of one off-diagonal entry makes the Cholesky *refuse* on 12/12. *The two-dimensionality (T158).* $\text{span}\{t_1, A^{-1}L_{Pt1}\}$ attains s exactly (9.0×10^{-13}) because $L_{Pt1} = \mu_1^P t_1$; without the Green column the two-sine span misses by 1.42–4.8, the corrupted column by $255\text{--}7.7 \times 10^3$. *The gauge identity and the signature (T159).* The reflection-odd section annihilates constant lag vectors entrywise with *zero* tolerance, hence $\sum_d w_d = 0$ to 1.1×10^{-14} of $\sup |w|$: the form is blind to the lag mass of c — exactly where the archimedean and atom halves are h^2 -sized and cancel; the Toeplitz-minus-Hankel signature holds to 1.4×10^{-16} ; a one-index Hankel shift breaks the identity loudly, and the T145 form is not even a parity section (signature defect 7.4×10^{-2}). *The exact lag sum and the two closed scalars (T159).* $x^T B_{LL} x = \sum_d c_d w_d$ to 2.7×10^{-16} of the absolute scale, with the honest price measured (relative to the cancelled total only 1.4×10^{-11} on this small surface); $w_0 = \sum_k x_k^2 / \mu_k^P$ and $2w_0 - w_1 = \|x\|^2$ exact. *The Z-matrix law and the Collatz–Wielandt floor (T159).* B_{HH}^{arch} is a symmetric Z-matrix *raw* (no tolerance, 12/12); the closed sign-based floor $\min_k (B_{HH}^{\text{arch}} \mathbf{1})_k = 0.9093\text{--}1.1284 \geq 0.25$ with the theorem direction verified against the exact $\lambda_{\min} = 0.9138\text{--}1.1432$; the atom block obeys *no* sign law (a coin toss) and the full-block row-sum floor is negative — admissible and vacuous; the checkerboard similarity (an isometry) destroys the raw law: an entrywise fact, not a spectral one. *The no-go discriminator.* The T145 family breaks on three axes of this module’s structure, while on the $c(l) = 1/(1+l)$ reconstruction the ladder theorem survives and $1/g_{16}$ explodes ($1187 \rightarrow 7.11 \times 10^4$) — the instrument reports the collapse. **Named limits as content:** per instance on small windows — nothing uniform in the zone index or in D , and explicitly *no* statement for all D ; the two open terms after T159 (the one explicit pairing inequality for R2'', needing correlation instead of sizes; a third device for the atom off-band block for R1'') stay open, typed open. Maz’ya 1985 / Miclo 1999 / Schur 1917–Haynsworth 1968 / KMS 1953 / Dirichlet 1829 / Perron 1907–Frobenius 1912 / Collatz 1942–Wielandt 1950 / Gantmacher–Krein 1950 named classical; Weil 1952 cited, never used as a criterion. Machine-checked: [verification/v553_exact_form_identities.py]. [E] (per-instance identities and certified inequalities; no marker moves)

The sampling/harmonics identities (PRIME.SAMPLING.HARM.01). The closed theorem cores of T160 and T161 are load-bearing as the twelfth companion module (21 checks, ~ 0.5 s), recomputed on the same declared surface as the module above (12 deepest in-cap frame-A windows, $m = 96 \dots 291 \leq 300$) plus a declared Λ -battery of six log-spaced cut-offs $X = 10^3 \dots 3.2 \times 10^5$ (finite von-Mangoldt sums only, no matrix anywhere) — what the sixteen-form *pairs against*, and it closes no open term. *The sampling identity (T160).* The atom half of the pairing equals $-\frac{1}{2} \sum_{n \leq X} (2\Lambda(n)/\sqrt{n}) \hat{w}(\log n)$ ($\hat{w}$ the linear interpolant of the lag weights; exact to 5.6×10^{-15} of the absolute atom scale on 12/12): identically a finite combination of sums $\sum_n \Lambda(n) n^{-1/2} \cos(t \log n)$ at 32 explicit frequencies; a one-index grid shift breaks the identity loudly. *The moment laws and the TV theorem (T160).* $m_0 = 0$, $m_1 = -[S_0^2 + 2 \sum_j P_j^2] \leq 0$ and $m_2 = -[2S_1 - (M-1)S_0]^2 \leq 0$ (a perfect square), both strictly negative on every window; and U3 as a theorem with *checked* hypotheses: $\text{TV}(c^{\text{arch}}) = 3.98\text{--}4.90 \leq |c_0| + 2 \sup < 6$, closed and window-independent — the full atom-including kernel violates the monotone hypothesis on 12/12 (the wired mutation). *The harmonic-frequency theorem and the PNT main term (T161).* $t_j(2\alpha) = 2\pi j$ *exactly*: the 32 frequencies are the Fourier harmonics of

the log-window, and the closed parameter-free prediction $(\sqrt{X} - 1)/(\frac{1}{4} + t^2)$ (de la Vallée Poussin 1896) matches the finite sums to 0.0007–0.097 of the trivial mass against the declared cap 0.12; a wrong Mellin pole and a mass-matched uniform fake measure both fail loudly — the agreement is a property of the von-Mangoldt measure. *The head split and the scale-free Bernstein rate (T161).* $c_d^{\text{arch}} = \Psi_d + D\hat{G}(dD, D)$ with $\Psi_d = -\frac{1}{2}[(d+1)\log(d+1) - 2d\log d + (d-1)\log(d-1)]$ closed, D -free and m -free, exact to 3.2×10^{-16} at every lag; $\rho^* = (3 + \sqrt{5})/2 = 2.618034$ is the root of $x^2 - 3x + 1$, scale-free (interval ratio 5 only), with every window's $\rho(D) = 2.588$ – 2.608 below it. *The fraction bound and the sign refutation (T161).* $|\text{off-diagonal}| \leq \frac{1}{4}|Q^{\text{arch}}|$ certified per window (0.171–0.212 on 12/12), with the *refuted* sign law wired as the negative control (arch half negative, off-diagonal block positive, 176/240 pairs violate the per-pair form) — nothing single-signed is promoted. **Named limits as content:** per instance on small windows — nothing uniform in the zone index or in D ; finite Λ -sums allowed and used, *no* RH statement, and the δ triage of T161 (required depth against RH strength) is a documentation item, not a module claim; the open terms after T161 (R-A' the prime-free log-moment; R-B' the m -free 16×16 Gram fraction bound; R-C' one split of the pairing with $\delta_{\text{eff}} < \frac{1}{2}$; R-D a fifth R1'' device) stay open, typed open. Dirichlet 1829 / Abel 1826 / Fejér 1915–Schur 1917 / Bernstein 1912 / Chebyshev 1852–Mertens 1874 / de la Vallée Poussin 1896 / Clausen–Lerch / KMS 1953 named classical; Weil 1952 / Bombieri 2000 cited, never used as a criterion. Machine-checked: [verification/v554_sampling_harmonics_identities.py]. [E] (per-instance identities and certified inequalities; no marker moves)

The Pareto/total-variation identities (PRIME.PARETO.TV.01). The closed theorem cores of T162 and T163 are load-bearing as the thirteenth companion module (23 checks, ~ 0.4 s), recomputed on the same declared surface as the module above (12 deepest in-cap frame-A windows, $m = 96 \dots 291 \leq 300$) with the one arithmetic input $\kappa = 0.038821$ *verified* at every jump point of $\psi(x)/x$ on the von-Mangoldt table — the *price structure* of the pairing, and it closes no open term beyond the negative closure that the floor inequality itself is. *The exchange law (T163).* $\delta_{\text{bnd}}(x) = \frac{1}{2} + \log(2\kappa g_{16}\text{TV}(x)/P(x))/\log X$ is an identity at all 612 grid points of the declared Fejér knob (5.1×10^{-16}), with the crossing criterion $\delta_{\text{bnd}} < \frac{1}{2} \iff P > 2\kappa g_{16}\text{TV}$ an exact equivalence at every point: price and demand are two coordinates of one object; the wrong constant breaks the law loudly. *The total-variation floor (T163).* $w_0 = \|a\|^2$ (4.9×10^{-16}), $\text{TV} \geq |w_0|$ by telescoping, and $x_1 = 1$ forces $\mu_1^P \text{TV} = 5.00$ – $28.84 \geq 1$ on all 708 trial vectors built — the closed floor $1/\mu_1^P = 1/(4\sin^2(\pi/N)) \sim h^2$ — so every sub- $\frac{1}{2}$ point pays $P > 2\kappa g_{16}/\mu_1^P$: a flat-price sub- $\frac{1}{2}$ demand is impossible in the parity sector on this surface (closes R-C''' negatively); the inadmissible vector $0.01x^*$ undercuts the floor and is recognised (the wired negative control). *The chain-derived price axis (T163).* $\sigma = 1$ is exactly the $K = 1$ rung of the T158 Cholesky ladder (4.1×10^{-16}) and $\sigma = \infty$ the $K = 16$ top rung (1.2×10^{-12}); the knob is monotone in both coordinates (its curve is the Pareto front); a corrupted sixteen-block makes the Cholesky refuse. *The Mellin cell moments and the Abel step (T162).* The k -th Mellin rung $\sum_d w_d C_d(-(2k + \frac{1}{2}))$, closed, agrees with an independent quadrature to 2.8×10^{-13} ($k = 1 \dots 4$); the boundary-free Abel identities hold at levels 1–3 (8.3×10^{-17}), the optimal level is one for the closed reason $32\pi/\alpha = 40.6$ – $59.4 > 1$, and a gauge violation breaks by its closed prediction (1.2×10^{-15}). *The Lerch/Frullani log-moment (T162).* $L = \sum_d \Psi_d w_d$ on three routes (Wronskian form 2.8×10^{-14} , integral with the $d = 1$ peel 1.8×10^{-15} , $2\log 2$ to 2.2×10^{-16} , $\Psi_1 = -\log 2$ exact); plus the monotone no-go staircase ($1.00 \rightarrow 3.49 \rightarrow 10.81 \rightarrow 24.98$, exact null control). **Named limits as content:** per instance on small windows — nothing uniform in the zone index or in D ; the T163 h -exponents are sandbox fits, *not* consumed; finite Λ -sums allowed and used, *no* RH statement; the open terms after T163 (R-E the successor, both arms prime-free; R-B''' the h -uniform positivity margin; R-D a fifth R1'' device) stay open, typed open. Fejér 1915 / Mellin 1896 / Abel 1826 / Chebyshev 1852–Rosser–Schoenfeld 1962 / Legendre / Dirichlet 1829 / KMS 1953 / Frullani 1828–Lerch 1887–Clausen 1832 / Schur 1917–Haynsworth 1968 named classical; Weil 1952 / Bombieri 2000 cited, never used as a criterion. Machine-checked: [verification/v555_pareto_tv_identities.py]. [E] (per-instance identities and certified inequalities; no marker moves)

The gauge/ P_{pr} identities (PRIME.GAUGE.PPR.01). The closed theorem cores of T164 and T165

are load-bearing as the fourteenth companion module (23 checks, ~ 0.3 s), recomputed on the same declared surface as the module above (12 deepest in-cap frame-A windows, $m = 96 \dots 291 \leq 300$) with the one arithmetic input $\kappa = 0.038821$ *verified* at every jump point of $\psi(x)/x$ on the von-Mangoldt table — the *gauge structure* of the pairing, and it does not close the one remaining quantifier. *The gauge theorem (T164)*. $x_1 = 1$ fixes only the *scale* of a ; Q and TV are homogeneous of degree two; hence the demand ratio Q/TV of the same a -direction is the *same number in every sector* — unchanged to 2.9×10^{-11} on 2520 sector-by-trial-vector combinations (5 weight families $\times$ 7 shifts $\times$ 6 taper widths), with the pure shift scaling the certified value by exactly $\theta = \mu_1^P/(\mu_1^P + \kappa_s)$; the full space is settled strictly worse ($\mu_0 = 0$ exactly); the wrong transport map breaks loudly ($2.0 \times 10^2 - 5.4 \times 10^3$). *The h^2 conservation law (T164)*. floor $\times$ transfer $= 1/(\mu_1^P + \kappa_s) \cdot (1 + \kappa_s/\mu_1^P) = 1/\mu_1^P$ *identically* at every shift (2.2×10^{-16}): the h^2 is conserved, never removed — it lives in the ratio, not in the floor; the wrong transfer misses by 0.9995 at the largest shift. *The power-one spend (T164)*. The r -ceiling $r \leq U/L$ with $L = \lambda_1(A)/\mu_1^P$ is linear in the entry ceiling, so $d \log r / d \log U = +1$ exactly (1.1×10^{-14}); the quadratic consumption reads slope 2 and is recognised. *The P_{pr} identity and the predicate equivalence (T165)*. $P_{\text{pr}} = g_{16} \cdot R \cdot (\text{TV}/(t_1 v)^2)/\mu_1^P$ exact to 4.3×10^{-16} on all 192 battery vectors, with the predicates $R > 2\kappa$ and $P_{\text{pr}} > 2\kappa g_{16} \text{ tvf}/\mu_1^P$ agreeing as booleans everywhere and every crossing vector paying $P_{\text{pr}} > 2\kappa g_{16}/\mu_1^P = 15.4\text{--}142.8$ above the preregistered tolerance 10 on 12/12 — the successor question R-F closes negatively by an identity plus a floor; the price with $(t_1 v)$ unsquared breaks the gauge invariance loudly. *The Schur-cascade direction and the exponent closure (T164/T165)*. $s = \mu_1^P t_1^T A^{-1} t_1 \geq g_{16} \geq g_1$ window by window with $P_K = \hat{a} s \geq 1$ (Kantorovich 1948), and the exponent closure $\log(1/g_1) = \log(g_{16}/g_1) + \log(1/g_{16})$ is exact per instance and on the fitted lists (1.3×10^{-15}); plus the monotone no-go staircase and exact Dirichlet/parity controls. **Named limits as content:** per instance on small windows — nothing uniform in the zone index or in D ; the T164/T165 h -exponents and the measured anatomy are sandbox, *not* consumed; finite Λ -sums allowed and used, *no* RH statement; **the one open object after T165 stays open, typed open:** $\inf_m g_{16}(m) > 0$, equivalently a lower bound on the sixteen-step Schur-cascade gain g_{16}/g_1 uniform in m — a quantifier over a certified flat list. Schur 1917–Haynsworth 1968 / KMS 1953 / Abel 1826 / Chebyshev 1852–Rosser–Schoenfeld 1962 / Kantorovich 1948 / Fejér 1915 / Dirichlet 1829 named classical; Weil 1952 / Bombieri 2000 cited, never used as a criterion. Machine-checked: [verification/v556_gauge_ppr_identities.py]. [E] (per-instance identities and certified inequalities; no marker moves)

The cascade/vector identities (PRIME.CASCADE.VECT.01). The closed theorem cores of T166 and T167 are load-bearing as the fifteenth companion module (23 checks, ~ 0.1 s), recomputed on the same declared surface as the module above (12 deepest in-cap frame-A windows, $m = 96 \dots 291 \leq 300$) with the one arithmetic input $\kappa = 0.038821$ *verified* at every jump point of $\psi(x)/x$ on the von-Mangoldt table — the *anatomy of the cascade* and the *exactness of the closed vector at $K = 2$* , and it does not close the one remaining scalar. *The cascade identities (T166)*. The prefix property (one Cholesky of the sixteen-block yields all sixteen rungs; 192 comparisons, 1.1×10^{-12}), the minor form $1/g_K = \det G_{[1..K]}/(\mu_1^P \det G_{[2..K]})$ on the raw arithmetic Gram block (2.7×10^{-12} ; Schur 1917–Haynsworth 1968), and the regression form $g_K/g_1 = 1/(1 - R_K^2)$ (8.8×10^{-13}) — the open object *is* a near-collinearity statement; the wrong minor misses by 0.99–1.45. *The diagonal invariance (T166)*. The gain $g_K/g_1 = B_{11}(Q_K^{-1})_{11}$ is unchanged under $B \rightarrow DBD$ on 144 random positive diagonals (2.7×10^{-12}): the cascade does not see the KMS normalisation — the gain belongs to the arithmetic Gram block alone; a non-diagonal congruence moves it by 2.7–3.8. *The exact $K = 2$ identity (T167)*. The closed vector $u = (1, -Q_{21}/Q_{22})$ attains $1/g_2 = Q_{11}(1 - r_{12}^2)$ exactly (1.1×10^{-12}) and $B_{11}g_2 = 1/(1 - r_{12}^2)$ identically (1.2×10^{-12}): at $K = 2$ the fast-null-vector is free — the third dress collapses onto the second, one scalar built from three closed lag sums; the wrong-sign vector pays $4r_{12}^2/(1 - r_{12}^2) = 6.7 \times 10^2\text{--}1.5 \times 10^4$. *The pivot identity and the unification (T167)*. $u^T Q_K u = 1/g_K + \delta^T Q_K \delta$ for $\delta_1 = 0$ (2.0×10^{-12} on 252 evaluations) — every candidate excess is an exact number; the sign-aligned entry attainment $|x^{*T} E x^*| = \varepsilon S_K$ (3.8×10^{-16}) makes the entry threshold exact, and the unification $(1/g_2)/S_2 = (1 - r_{12}^2)/(1 + 3r_{12}^2)$ (1.2×10^{-12}) makes the three dresses *one* inequality; the $\delta_1 = 0.1$ mutation breaks by exactly the closed cross term

$2\delta_1$. *The trial theorem and the two must-breaks (T166/T167)*. All 288 random trials with $u_1 = 1$ bound g_K from below (min ratio 15.14; Schur 1917); the L_P no-go gives cascade gain exactly 1 (deviation 0.0); the scramble type-change makes B_{11} itself lose positivity on a majority of draws on 12/12 windows — at $K = 2$ the collapse under scrambling is a change of type, not a factor; plus exact Dirichlet/parity controls. **Named limits as content:** per instance on small windows — nothing uniform in the zone index or in D ; the T166/T167 h -exponents and the measured anatomy are sandbox, *not* consumed; finite Λ -sums allowed and used, *no* RH statement; **the one open object after T167 stays open, typed open:** R1, the single scalar — an m -free upper bound on $1 - r_{12}^2 = 1 - Q_{12}^2/(Q_{11}Q_{22}) \leq Ch^{-3+\varepsilon}$, three closed lag sums and nothing else. Schur 1917–Haynsworth 1968 / KMS 1953 / Rayleigh 1877–Courant–Fischer 1920 / Cauchy 1829 / Kato 1949 (cited as the closed-off route, never used) / Chebyshev 1852–Rosser–Schoenfeld 1962 / Dirichlet 1829 named classical; Weil 1952 / Bombieri 2000 cited, never used as a criterion. Machine-checked: [verification/v557_cascade_vector_identities.py]. [E] (per-instance identities and certified inequalities; no marker moves)

The bilinear/rank identities (PRIME.BILINEAR.RANK.01). The closed theorem cores of T168, T169 and T170 are load-bearing as the sixteenth companion module (27 checks, ~ 0.2 s), recomputed on the same declared surface as the module above (12 deepest in-cap frame-A windows, $m = 96 \dots 291 \leq 300$) with the one arithmetic input $\kappa = 0.038821$ *verified* at every jump point of $\psi(x)/x$ on the von-Mangoldt table — the *exact shape of the one remaining scalar*, and it does not close that scalar. *The Lagrange/Wronskian anatomy (T168)*. The lag-sum identity $\hat{a}_{ij} = t_i^T A_h t_j = \sum_d c_d W_d^{ij}$ with the closed correlation weights (1.6×10^{-15} ; the $1/\sqrt{\mu}$ factors cancel, T169-TH3); Euclidean orthogonality $t_1 \cdot t_2 = 0$ (4.4×10^{-16}) — the near-parallelism is created *entirely* by the arithmetic metric; the Wronskian telescope $(\mu_1 - \mu_2)u_r v_r = W_{r-1} - W_r$ with $W_{-1} = W_{h-1} = 0$ (1.9×10^{-17}) and $\frac{1}{2} \sum M^2 = 1/(\mu_1 \mu_2)$ exactly (4.6×10^{-16} ; Lagrange 1773); the wrong eigenvalue pair misses by 1.66–1.67 relative. *The exponent account and the Gershgorin certificate (T168/T169)*. $1 - r_{12}^2 = \nu_1 \nu_2 / (\hat{a}_{11} \hat{a}_{22})$ identically (2.5×10^{-13}); $\nu_1 \leq \max(\hat{a}_{11}, \hat{a}_{22}) + |\hat{a}_{12}|$ (Gershgorin 1931, unconditional) on 12/12 with loss 1.1885–1.1947; dropping $|\hat{a}_{12}|$ fails on every window. *The det-collapse theorem (T169-TH7)*. $R(K7) = 2\sqrt{\hat{a}_{11}\hat{a}_{22}} \det \hat{A} / [(\hat{a}_{11} + \hat{a}_{22})(\sqrt{\hat{a}_{11}\hat{a}_{22}} + \hat{a}_{12})]$ in closed form (8.7×10^{-13}) — the loop T167 scalar $\rightarrow$ T168 factor $\rightarrow$ T169 candidate closes as an identity; usable form $\nu_2 \leq 2 \det \hat{A} / (\hat{a}_{11} + \hat{a}_{22})$, tight to < 2 exactly; the closed constant $1/\sqrt{2}$ overshoots by $17\text{--}2.3 \times 10^2$. *The exact bilinear von Mangoldt form (T170)*. $\hat{A} = B - S$ against the direct block (1.7×10^{-14}); $\det \hat{A} = \det B - D(B, S) + \det S$ (1.0×10^{-10}); $\det S = \sum_{n,m} \Lambda(n)\Lambda(m)K(n, m)/\sqrt{nm}$ with $K = \frac{1}{2}D(X_n, X_m)$ through the *explicit* kernel matrix (7.5×10^{-16} ; Cauchy–Binet); the wedge reading to the measured rank-one defect 4.6×10^{-3} – 3.1×10^{-2} . *The rank-3 theorem and the Type II collapse (T170)*. The polarisation matrix has eigenvalues $\{-2, -1, +1\}$ exactly — rank 3, signature (1, 2) — so $\sigma_4/\sigma_1 \leq 1.6 \times 10^{-16}$ on every window: the bilinear form *is* the rank-3 polynomial $S_{11}S_{22} - S_{12}^2$ in three linear Λ -sums; Vaughan’s identity is coefficient-exact for every $U = V \in \{3, 4, 5, 6\}$ and the Type II kernel has 99.9%-energy rank $[2, 2, 2, 2]$ against generic $[20, 15, 11, 9]$. *The R_4 -free chain and the discriminators (T170/T169)*. $1 - r_{12}^2 = \det \hat{A} / (\hat{a}_{11}\hat{a}_{22})$ is an identity — no A -positivity, the Weil fence never approached; the L_P no-go gives $\hat{a}_{12} = 1.1 \times 10^{-17}$ identically (no decay); the scramble leaves the rank-3 collapse *untouched* on all 60 draws while the value moves $\times 834.8$ median — the arithmetic sits in the joint values of (S_{11}, S_{22}, S_{12}) against B ; plus exact Dirichlet/parity controls. **Named limits as content:** per instance on small windows — nothing uniform in the zone index or in D ; the T168–T170 h -exponents and route-table deltas are sandbox, *not* consumed; finite Λ -sums allowed and used, *no* RH statement — the one open object after T170 (R1, now finally *classified* but not closed: an m -free unconditional certificate that the two rows of $\hat{A}$ become collinear at rate $h^{-3+\varepsilon}$ — a joint near-degeneracy of three explicit linear Λ -sums, not a size) stays open, typed open. Lagrange 1773 / Cauchy–Binet 1812–1815 / Gershgorin 1931 / Rayleigh 1877 / Montgomery–Vaughan 1973–Vaughan 1977 (the priced route) / KMS 1953 / Chebyshev 1852–Rosser–Schoenfeld 1962 / Dirichlet 1829 named classical. Machine-checked: [verification/v558_bilinear_rank_identities.py]. [E] (per-instance identities and certified inequalities; no marker moves)

The phase-2 capstone (PRIME.PHASE2.CAPSTONE.01). The capstone *verifies the map and claims nothing new*: T126–T170 reduced the I5 uniformity question through sixteen exact reformulations to one object (R1), T171 (contract FINAL.MAP, verdict MAP-COMPLETE) reproduced every link of that reduction in one connected sandbox run, and the seventeenth companion module (35 checks, ~ 0.3 s) is the load-bearing version of that run on the same declared surface as the module above (12 deepest in-cap frame-A windows, $m = 96 \dots 291 \leq 300$; $\kappa = 0.038821$ *verified* at every jump point of $\psi(x)/x$; the 16×16 low block positive definite on 12/12 — a numerical fact, never routed through the Weil criterion). *The chain, sixteen links in one pass* (13 THEOREM / 3 CERT): the T152 Schur two-block criterion sharp on both sides (holds at $0.5\lambda_{\min}$, fails at $1.02\lambda_{\min}$; KMS 1.3×10^{-15} , Haynsworth 3.8×10^{-15}); the T151 Sobolev bound $b_k \leq Csk$ (worst ratio 0.4816); the T153 Ψ -collapse $\Psi = \sum_d c_d w_d = x^T G x$ (6.3×10^{-16} ; $\sum_d w_d = 0$ to 9.0×10^{-17}); the T154 Ritz ceiling from *below* ($g_{32}/s = 0.9520$ – 0.9849); the sharp T155 12×12 floor ($t = 0.2295$ – 0.3719); the T156 $F(P, r)$ chain ($L(16) = 6.6$ – 15.6); the T158 Thomson/ladder duality (3.1×10^{-12} ; 36/36 trials below); the T160 sampling identity (1.1×10^{-15} / 9.9×10^{-15}); the T163 Abel swap with the TV floor (overshoot $\times 5.2 \times 10^4$ – 1.1×10^6); the T164 gauge/quantifier collapse ($\Psi(x^*) = 1/g_{16}$ to 7.0×10^{-12} ; 48/48 perturbed vectors larger); the T165 P_{pr} identity ($= 1$ at x^*); the T166 cascade $1/(1 - R_{16}^2)$ (1.8×10^{-12} , diagonal-invariant); the T167 $K = 2$ exactness (milder $\times 1.62$ – 5.20); the T169 det collapse, R4-free ($\leq 4.2 \times 10^{-12}$, no positivity of A_h consumed — the Weil fence never approached); the T170 rank-3 theorem (J -eigenvalues $\{-2, -1, +1\}$ exact; $\sigma_4/\sigma_1 \leq 1.6 \times 10^{-16}$); and the T170 R1 shape via the 2×2 Lagrange identity ($\sin \angle = 1.169 \times 10^{-4}$ – $2.467 \times 10^{-3} \leq 10^{-2}$ per window — *rate and quantifier typed OPEN*). *The no-go battery*, all eight failing as classified: size budget $\times 2.7 \times 10^3$ – 1.6×10^4 ; symbol infimum < 0 on 12/12; sector change an invariance ($\leq 4.2 \times 10^{-12}$); perturbation $\times 9.8 \times 10^2$ – 2.0×10^4 off scale (Kato 1949); atom band dominates $\times 1.71$ – 2.83 ; sieve operator-norm $\times 8.8 \times 10^3$ – 7.5×10^4 over (Montgomery–Vaughan 1973 / Vaughan 1977 the priced route); scramble $\times 679$ median with rank-3 *untouched* on 60 draws; L_P gain = 1 exactly. *The precision ledger*: needed 2.752×10^{-4} at the deepest window — finer than the RH yardstick $h^{-0.5}$ by $215\times$ and the best unconditional $h^{-0.996}$ by $13\times$ on this surface; both self-similarity identities $\leq 6.8 \times 10^{-13}$; mutations fail loudly. **Named limits as content**: per instance on small windows — nothing uniform in the zone index or in D ; the T171 trend fits (pooled $h^{-2.33}$, per-frame $h^{-1.86}/h^{-2.87}$, $\delta = +1.044$ against 3.0, the full-surface numbers 2.284×10^{-7} / $1.2 \times 10^5 \times$ / $3.1 \times 10^3 \times$) are sandbox, *not* consumed; *zero* new uniform-in- m statements — which is exactly why the capstone is promotable; the one open phase-2 object (R1, an m -free unconditional certificate that the two rows of $\hat{A}$ become collinear at rate $h^{-3+\varepsilon}$) stays open, typed open; *no* RH statement. Schur 1917–Haynsworth 1968 / KMS 1953 / Cauchy 1829–Courant–Fischer 1920 / Rayleigh 1877 / Abel 1826 / Dirichlet 1829 / Lagrange 1773 / Cauchy–Binet 1812–1815 / Kato 1949 (the closed-off route) / Montgomery–Vaughan 1973–Vaughan 1977 / Chebyshev 1852–Rosser–Schoenfeld 1962 / Fejér 1915 named classical; Weil 1952 / Bombieri 2000 cited, never used as a criterion. Machine-checked: [verification/v559_phase2_capstone.py]. [E] (per-instance identities and certified inequalities; no marker moves)

The frame/deficit identities (PRIME.FRAME.DEFICIT.01). The gauge route of phase 2, closed as algebra: the eighteenth companion module (24 checks, ~ 0.5 s) promotes the theorem cores of T172 (contract FRAME.BEYOND), T173 (FRAME.RATE) and T174 (CANCEL.IDENTITY) — the companion to the capstone above, certifying what those three parts established *about* the map’s frame dependence, on a small declared surface (a 4×3 (anchor, h) rectangle of gap-blind frame-B cells, $n = 41 \dots 557 \times h = (160, 226, 320)$, comb complete, positive definite on 12/12; one non-prime-power instance $n = 210$; one truncated-comb stress instance $n = 1009$; $\kappa = 0.038821$ *verified* at every jump point of $\psi(x)/x$). *The identity transfer + sieve-horizon discriminator* (T172): four chain links — the KMS/Schur pair (3.2×10^{-16}), the lag-sum identity (1.4×10^{-16} of the larger half; Abel 1826), the R4-free det collapse (at 0.020 of the conditioned bar) and the additive split (7.6×10^{-16}) — re-derived *exactly* on a frame-B and a non-prime-power instance; the truncated-comb instance turns indefinite ($\lambda_{\min} = -9.6 \times 10^4$) against 12/12 positive definite complete-comb cells, with the identity links holding either way (none uses positivity): indefiniteness sits at the declared sieve

horizon, not at a frame or zone. *The $q = 1 - s$ identity + calibration* (T173): $hD = \alpha$ per cell forces the OLS slopes to obey $q = 1 - s$ exactly (2.9×10^{-14}); the gauge degrees come out as integers — $G2 = t/\lambda_{\max}(B_{LL})$ and the delivery $1 - r_{12}^2$ both $(0, 0)$, so $G2$ is the unique doubly gauge-invariant arithmetic member of the family; $G3 = \mu_1^P G2$ identically (the offset-2 explained); $(D/\alpha)^3 h^3 = 1$ a tautology. *The blindness theorems + the additive limit* (T174): the amplitude $c \rightarrow \kappa c$ is a gauge of both sides and of R over twelve decades (worst 0.026 of the per-cell conditioned bar); the scalar μ_1^P — *the entire h^{-2} channel of the demand* — cancels exactly on both sides; the delivery is invariant under the full positive diagonal group while a non-scalar diagonal moves the demand by 0.23–1.40 (the asymmetry is the anatomy); and neither the arch term nor the comb term alone has a positive Schur floor on a single cell ($0/12 + 0/12$ against $12/12$ for the sum): no factorisation $\hat{A} = (\text{frame}) \times (\text{arithmetic})$ exists — P6 is *not* a theorem, only its multiplicative half. *The KMS shape band* (unconditional, declared $\varepsilon = (K\pi/(2h+1))^2/6$): the k^2 -swap moves $\log \text{GAP}$ by ≤ 0.312 of the certified band while the delivery does not move at all. *The placement discriminator*: the value-multiset scramble destroys the floor on 24/24 draws. **Named limits as content**: per instance on a small declared surface — nothing uniform in the zone index or in D ; the frame-free deficit $+0.1111 \pm 0.0222$ (5.0σ , 151 anchor clusters) is a sandbox MEASURED number, *not* consumed; the one open phase-2 object (R1, now stated frame-free) stays open, typed open; *no* RH statement. Schur 1917 / KMS 1953 / Dirichlet 1829 / Rayleigh 1877 / Abel 1826 / Chebyshev 1852–Rosser–Schoenfeld 1962 / Wilkinson 1968–Higham 2002 named classical; Weil 1952 / Bombieri 2000 cited, never used as a criterion. Machine-checked: [verification/v560_frame_deficit_identities.py]. [E] (per-instance identities and certified inequalities; no marker moves)

The dense-limit identities (PRIME.DENSE.LIMIT.01). The exact cores of the phase-2 endgame (T176, contract DENSE.LIMIT, verdict SITS-AT-ZERO — the probe that raised the sieve twentyfold, extended the deficit(dens) curve to a densest bin consistent with zero, and *closed the phase-2 measurement programme as planned*), as one machine-checked module (16 checks, ~ 0.5 s, deterministic) on a small declared surface (18 prime-power anchors in $[40, 565] \times$ the declared $2^{1/4}$ ladder $h = (128 \dots 304)$, comb complete, positive definite on 108/108; $\kappa = 0.038821$ verified at every jump point; the big sieve is *not* rebuilt — the identities are sieve-size-free). *R1 closed* (T176 P2): the Feynman–Hellmann identity $d \log R / d\delta = v^\top d\hat{A}v / \lambda_{\min} - w^\top d\hat{A}w / \lambda_{\max} - d\text{DEL} / \text{DEL}$ with the exact lag derivative reproduces the full nonlinear rebuild on 6/6 instances at the minimum of a declared step ladder whose U-shape is exhibited (2.3×10^{-7} at $\delta = 10^{-6}$; a wrong formula has a floor, not a minimum) — T175’s measured intervention slope is retired as a separate claim, and the near-degeneracy channel $d\text{DEL} / \text{DEL}$ carries the largest share (0.54–0.98). *The phase pole* (T176 P3): positive definiteness survives only 2% of the phase period, $\delta_{\text{PD}} \cdot |d \log R / d\delta| = O(1)$, and the two lowest modes are already near-degenerate at $\delta = 0$ — the $\sim 300\times$ harmonic anomaly is retired (a first-harmonic regression of a function with a pole in the interval). *Sieve consistency at construction level* (T176 P4): complete-comb lag vectors agree bit for bit under a bigger table, R exactly, the bin deficit identically — what lets any later probe raise the cap without re-deriving the chain; the truncated must-break moves as it must. *The estimator + the ladder caveat as a check* (T176 P1): the cluster-robust bin statistic is deterministic and exactly reproducible — and *ladder-dependent* (fine vs coarse rung ladder shift 0.036 here, up to 0.20 on T176’s surface): any quoted bin deficit must cite its rung ladder, and the ledger row does. **Named limits as content**: T176’s SITS-AT-ZERO (pooled $+0.0496 \pm 0.0372$ over the two new bins; plateau window $3.6\times$ narrower) is a sandbox MEASURED number with its rung ladder, *not* consumed; the limit — true zero, power law, plateau — stays open, typed open, under *every* resource-reachable sieve; the closure is of the measurement programme, not of the question; *no* RH statement. Feynman–Hellmann (Hellmann 1937, Feynman 1939) / Schur 1917 / KMS 1953 / Dirichlet 1829 / Chebyshev 1852–Rosser–Schoenfeld 1962 named classical; Weil 1952 cited, never used as a criterion. Machine-checked: [verification/v562_dense_limit_identities.py]. [E] (per-instance identities and certified inequalities; no marker moves)

The Paper-II readout closure (PRIME.PAPER2.READOUT.01). The check closure of the classification paper (Paper II, parity_toeplitz_classification): every number that paper quoted from

an exploration probe without a recomputing module is recomputed on the identical declared surface (24 checks, ~ 0.5 s) — the T170 reference window bit for bit ($h = 540$, $X = 157^2$; $\det \hat{A} = 4.22 \times 10^{-3}$, a 4.7-order cancellation), the ℓ^1 -to-signed ratio $40.6\times$ (Cauchy–Binet at round-off), the tightness envelope $\mu_1^P \text{TV} \in [5.00, 23.20]$ with both ends attained, the density-curve endpoints on identical bin edges — and the one caveat Paper II declared without exhibiting it is exhibited as a structure fact: the rung-ladder shift of the binned exponent is *anchor selection* (fine vs coarse ladders select different anchor sets in the same bin, 13 vs 12; removing the extra anchors reproduces the whole shift, curvature transport moves it by only -0.0111). Pure readout/identity checks — no new claim, no rate, no uniformity; every measured quantity stays MEASURED with its rung ladder; no RH statement. Machine-checked: `[verification/v563_paper2_readouts.py]`. [E] (readout/identity checks with mutation controls; no marker moves)

3.4 Glue-norm arithmetic, and three readings of 41, 60, 120

The two glue norms generate the carrier integers through sum, difference and product. With $\dim S^+ = 16$:

$$\boxed{16(q_D + q_A) = 32 = 2^{g_{\text{car}}} = h(D_5)h(A_3), \quad 16(q_D - q_A) = 8 = r(E_8) = h(D_5), \quad 16 q_D q_A = 15 = \dim S^+ - 1} \quad [\text{E}], \quad (12)$$

so the glue already contains the full carrier exhaustion $2^{g_{\text{car}}}$, the E_8 rank, and the non-trivial transition count $\mathcal{A}_H + \mathcal{A}_\Lambda = 15$; and $|\mu_4| = (q_D - 1)^{-1} = (1 - q_A)^{-1} = 4$ is the inverse displacement of the two norms from the unit norm.

Spine Quotient Lemma: the recurring factors are adjacent quotients of $(2, 3, 4, 5)$ [E]
(`[verification/v74_compiler_micro_lemmas.py]`)

The integers of the spine are $(|\mathbb{Z}_2|, N_{\text{fam}}, |\mu_4|, g_{\text{car}}) = (2, 3, 4, 5)$. Every “mysterious” $O(1)$ factor in TFPT is an *adjacent quotient* of this chain, not a separate special number:

$$\frac{|\mathbb{Z}_2|}{N_{\text{fam}}} = \frac{2}{3}, \quad \frac{N_{\text{fam}}}{|\mu_4|} = \frac{3}{4} = q(A_3), \quad \frac{|\mu_4|}{g_{\text{car}}} = \frac{4}{5}, \quad \frac{g_{\text{car}}}{|\mu_4|} = \frac{5}{4} = q(D_5), \quad \frac{|\mu_4|}{N_{\text{fam}}} = \frac{4}{3}.$$

These are exactly the factors that recur across the theory: $\lambda_2 = (\frac{2}{3})^6$ (the transport gap = Page recovery rate, and the Koide target $Q_\star = \frac{2}{3}$); $\varepsilon = \frac{3}{4}\varphi_0 = q(A_3)\varphi_0$ (the solar misalignment); $\varphi_0 = \frac{4}{3}c_3 + 48c_3^4$ with leading $\frac{4}{3}c_3 = \frac{1}{6\pi}$; and $q(D_5) + q(A_3) = \frac{5}{4} + \frac{3}{4} = 2$ (the E_8 root norm). So $(\frac{2}{3}, \frac{3}{4}, \frac{4}{5}, \frac{5}{4}, \frac{4}{3})$ are five projections of one discrete chain — not scattered constants. The spine is moreover *one* finite object: as a tetrahedron on $T = \{e_3(a), p_0(a), e_1(a), e_2(a)\}$ its edge and face products and its volume $120 = |R^+(E_8)| = 5!$ reproduce the central integer grammar, with the honest negative control that the graph reading $240 = |\mu_4| \cdot |E(K_4)| \cdot |E(K_5)|$ breaks at K_6 (`[verification/v91_spine_tetrahedron.py]`).

The carrier adjoint $\dim D_5 = |R(D_5)| + g_{\text{car}} = 45$ then gives a second reading of the $U(1)$ integer,

$$\boxed{10b_1 = \dim D_5 - |\mu_4| = 45 - 4 = 41 = \underbrace{40}_{\sum L} + \underbrace{(g_{\text{car}} - |\mu_4|)}_{=N_{\Phi}=1}} \quad [\text{E}], \quad (13)$$

i.e. the single Higgs doublet is the carrier rank minus the glue corners, $5 - 4 = 1$.

A fourth, metric reading of 41 (`[verification/v222_cm_norm_duality.py]`). On the square seam ($j=1728$, complex multiplication by the Gaussian integers $\mathbb{Z}[i]$) the $U(1)$ index is the Gaussian norm of the carrier-glue vector,

$$10b_1 = N_{\mathbb{Z}[i]}(g_{\text{car}} + i|\mu_4|) = g_{\text{car}}^2 + |\mu_4|^2 = 5^2 + 4^2 = 41,$$

and the hexagonal partner ($j=0$, Eisenstein $\mathbb{Z}[\omega]$) reads the same $(3, 2)$ carrier split as the scalaron exponent $N_{\mathbb{Z}[\omega]}(3+2\omega) = 3^2 - 3 \cdot 2 + 2^2 = 7$. So 41 (EM) and 7 (scalaron) are the two exceptional CM

norms of the one carrier, square vs. hexagonal — not two unrelated $O(1)$ numbers [E]. The start defect 60 and the hypercharge moment 120 are “bilingual”:

$$D_{\text{start}} = \dim D_5 + \dim A_3 = 45 + 15 = 60 = \mathcal{A}_\Lambda \cdot |R^+(A_3)| = 10 \cdot 6,$$

$$\text{Tr}_{S^+} X^2 = 120 = \underbrace{60}_{\text{adjoint}} + \underbrace{60}_{\text{mixed (10,6)}}, \quad (14)$$

and $\dim E_8 = 248 = \text{Tr}_{S^+} X^2 + 2 \dim S^+ |\mu_4| = 120 + 2 \cdot 16 \cdot 4$. Finally the hypercharge norm and the EW exponent are A_3 -rational,

$$\boxed{\gamma_{\text{car}} = \frac{\mathcal{A}_\Lambda}{|R(A_3)|} = \frac{10}{12} = \frac{5}{6}, \quad I_1^{\text{EW}} = \frac{|E(K_5)|}{|R(A_3)|} + \frac{1}{|\mu_4| |R(A_3)|} = \frac{10}{12} + \frac{1}{48} = \frac{41}{48}} \quad [\text{E}]. \quad (15)$$

Transport reduction. The flavor matrix L has its mod-6 content fixed by a single combinatorial datum: $\{0, 1, 3\}$ is the *unique* (up to D_6) 3-subset of $C_6 = |R^+(A_3)|$ with all distinct cyclic distances $\{1, 2, 3\}$, and the three sectors are its D_6 orbit ($u \equiv \{0, 1, 3\}$, $d \equiv 2 - \{0, 1, 3\}$, $e \equiv 2 + \{0, 1, 3\}$). The total is fixed by the carrier, $\sum L = 40 = |R(D_5)|$, with winding unit $6 = |R^+(A_3)|$. Given the residue *set* of a sector, the full ordered packet is reproduced by two universal rules — the lightest generation carries the single +6 winding, and the entries are sorted in decreasing length (monotone hierarchy). The *only* remaining datum is then, per sector, which residue is the first-generation anchor that receives the winding:

$$\boxed{\text{anchors } (a_u, a_d, a_e) = (1, 1, 2) \Rightarrow L_u = (7, 3, 0), L_d = (7, 5, 2), L_e = (8, 5, 3)} \quad [\text{E}] \text{ (given anchors)}. \quad (16)$$

So the open step of a full “Clebsch walk” theorem is sharply isolated: it is exactly these three per-sector anchors. They are *not* free: the three transport cusps are exactly the D_5 dual-root combinations, and each sector is tied to its cusp by the right-handed hypercharge,

$$\boxed{\{1, \frac{2}{3}, \frac{1}{3}\} = \{2y_+, -2y_-, 2(y_+ + y_-)\}, \quad \text{lepton} \leftrightarrow 2y_+, \text{ up} \leftrightarrow -2y_-, \text{ down} \leftrightarrow 2(y_+ + y_-)} \quad [\text{E}]. \quad (17)$$

The canonical-transport-kernel theorem (document 3, Part II) derives the residue packets — and hence the anchors — as the *word-lengths of the family-connection holonomy* $\text{Hol}_{\mathcal{G}_{D_4}}(\nabla_F^*)$ along the D_4 -invariant geodesic spine of $\mathbb{P}^1 \setminus \mu_4$ with these cusps as boundary data. The anchors are therefore a genuine $D_5 \times A_3$ readout (cusps from the D_5 carrier, spine holonomy from the A_3 four-punctured sphere), not an extra parameter. A compact candidate closed form is the nearest-integer cusp map

$$\boxed{a_f = \text{round}(2 \text{ cusp}_f) : (a_u, a_d, a_e) = \text{round}(\frac{4}{3}, \frac{2}{3}, 2) = (1, 1, 2)} \quad [\text{C}], \quad (18)$$

which reproduces the matrix exactly. The remaining open step is now precise and purely geometric: an *independent* closed-form evaluation of the D_4 -spine holonomy word-lengths from the lattice data alone (the same $\mathbb{P}^1 \setminus \mu_4$ whose residue pairing is $\text{Cartan}(A_3)$ and whose corner rotation is the A_3 Coxeter element), rather than from the family-connection theorem (document 3).

Riemann–Hilbert check, and what it does (and does not) fix. Setting up the rigid $\mathbb{Z}_4$ -symmetric rank-3 $SU(3)$ local system explicitly — four puncture monodromies $M_k = R^k C R^{-k}$ with the order-3 cusp class $C \sim \text{diag}(1, \omega, \omega^2)$ ($\omega = e^{2\pi i/3}$, the three cusps as eigenvalue-exponents) and the rotation R ($R^4 = \mathbb{K}$) — the closure $M_0 M_1 M_2 M_3 = \mathbb{K}$ has a solution to machine precision, and the unique R has eigenvalues $\{i, -1, -i\}$: the A_3 Coxeter element. This *independently* confirms the rigidity of the family connection and the corner-rotation = Coxeter identity. However, the flat monodromy *alone* does not fix the integer word-lengths $\{0, 1, 3\}$: those are *parabolic degrees* (the cusp exponents as parabolic weights, together with the $\mathbb{Z}_2$ sheet parity), i.e. Hodge/metric data on the spine, not topological monodromy. *Status update (the computation has since been carried out)*: the exact residue normal form A_0^* with $\text{spec} = \{0, \frac{1}{3}, \frac{2}{3}\}$ is pinned algebraically

(`[verification/v115_anchor_residue.py]`, `[verification/v116_resonance_uniqueness.py]`), the unitarised holonomy is the exact 24-element $W(A_3) = S_4$ (`[verification/v117_monodromy_weyl_a3.py]`), and the parabolic weights are the μ_4 -character degrees of $H^1(\mathbb{P}^1 \setminus \mu_4)$ — one spectrum in four readouts (`[verification/v137_qplus_cohomology.py]`); the deck itself is *derived* — the integer model selects the sheet-twisted class, no discrete freedom left (`[verification/v141_deck_selection.py]`), and the full Möbius D_4 of the curve matches the integer model parity by parity (`[verification/v146_moebius_d4.py]`). What remains of the old residual is the much smaller selector establishment $R4'$ (research contracts): derive the *one* lift map of the torsion normal — the pairing values are atom identities (`[verification/v142_frame_integrality.py]`/`[verification/v145_pairing_atoms.py]`). [E]/[C]




Figure 5: Flavor transport on the hexagon $C_6 = Z_6 = |R^+(A_3)|$. The unique distinct-distance triple $\{0, 1, 3\}$ (cyclic distances $\{1, 2, 3\}$) and its D_6 orbit give the three sectors' residues mod 6; the total $\sum L = 40 = |R(D_5)|$ and the first-generation $+6$ winding come from the carrier.

3.5 What the compiler outputs for the Standard Model and our reality

Every entry below is an *output* of (c_3, g_{car}) through the $D_5 \times A_3$ compiler; none is an independent input. “Reads” names the lattice object the observable comes from.

SM / cosmology observable	Compiler output	Reads	St.
Number of generations	$N_{\text{fam}} = 3 = \text{rank } A_3 = \dim H^1(\mathbb{P}^1 \setminus \mu_4)$	A_3	[E]
Gauge algebra dimension	$\dim \mathfrak{g}_{\text{SM}} = 12 = R(A_3) = \gcd(6r, 2h)$	A_3	[E]
One generation (16, incl. ν^c)	$S^+ = D_5$ half-spinor $= \Lambda^{0,2,4} E$	D_5	[E]
Hypercharge spectrum	all Y from $y_{\pm} = \{\frac{1}{2}, -\frac{1}{3}\}$; $\text{Tr } X^2 = 120$	D_5 root data	[E]
Abelian index b_1	$10b_1 = 1 + \mu_4 E(K_5) = 1 + 4 \cdot 10 = 41$	$\mu_4 \times K_5$	[E]
Fermion family occupancy	$\Omega_{\text{adm}} = N_{\text{fam}} \dim S^+ = (N_{\text{fam}} + N_{\Phi}) R(A_3) = 48$	A_3	[E]
Fine structure $\alpha_{\star}^{-1} = 137.0359992168$	root of $F_{U(1)}(\alpha) = 0$ (explicit, unique)	EM closure	[E]
Cabibbo angle	$\lambda_C = \sqrt{u(1-u)}$, $u = \frac{4}{3}c_3 + 48c_3^4$	seed	[E]
CKM CP phase	$\delta_{\text{CKM}} = \frac{2\pi}{ R^+(A_3) } + N_{\text{fam}} u(1-u) = \frac{\pi}{3} + 3\lambda_C^2$	A_3^+ , seed	[E]/[C]
PMNS angles/phase	$\sin^2 \theta_{12} = \frac{1}{3} - \frac{u}{2}$, $\sin^2 \theta_{13} = e^{-5/6} u$, $\delta^\nu = \frac{4\pi}{3}$	A_3^+ , seed	[E]/[C]
Charged-lepton hierarchy	$\det M_\ell \propto u^{h/N_{\text{fam}}} = u^{10}$; $(5, 3, 2)$	decuple	[E]/[C]
Mass-hierarchy transport	$\sum L = 40 = R(D_5) $; entries on $C_6 = R^+(A_3) $	$D_5 \times A_3$	[E]
Strong CP	$\theta_{\text{eff}} = 0$ (determinant erasure, given RP; document 3, Part III)	—	[E]/[C]
Weak CP	holonomy: $\delta_{\text{CKM}} - \frac{\pi}{3} = N_{\text{fam}} u(1-u)$	A_3^+	[C]
Gravity tree normaliser	$\xi_{\text{tree}} = \frac{\dim \mathfrak{g}_{\text{SM}}}{\dim S^+} = \frac{3}{4} = q(A_3)$	A_3 glue	[E]
Cosmological constant	$\rho_\Lambda / \bar{M}_{\text{Pl}}^4 = \frac{3}{4\pi^2} e^{2\delta_{\text{ph}}} [v_{\text{geo}} / (5\beta_{\text{rad}}^2 \bar{M}_{\text{Pl}})]^{10}$	decuple	[E]
Hubble anchor	$H_0 \sqrt{\Omega_\Lambda} = \bar{M}_{\text{Pl}} e^{-\alpha_{\star}^{-1}} / (2\pi)$	action	[E]
Baryon density	$\omega_b = \lambda_C / \mathcal{A}_\Lambda = \lambda_C / 10$	decuple	[E]/[C]
Inflation (scalaron)	$M_{\text{scal}}^2 / \bar{M}_{\text{Pl}}^2 = c_3^{\Omega_{\text{adm}} - 10b_1} = c_3^7$	deficit $r - 1$	[E]
Inflation amplitude	$A_s = N_{\star}^2 c_3^7 / 24\pi^2 \approx 2.2 \times 10^{-9}$	deficit	[C]/[O]

Marker key: [E] exact / machine-proven · [C] conditional (holds under named hypotheses) · [O] open / axiom · [X] falsifiable kill test. Per-claim fine type in `verification/status_ledger.csv`.

Reading. The discrete sector — generations, gauge dimension, family content, hypercharge, b_1 , occupancy — is now *group-theoretic output* of $D_5 \times A_3$. The continuous sector — α^{-1} , λ_C , the mixings, ρ_Λ , H_0 , inflation — is the seam seed c_3 and the action exponentials read on the *same* carrier graphs (K_5 pairs, the C_6 hexagon $= |R^+(A_3)|$, the decuple $\mathcal{A}_\Lambda = \binom{g_{\text{car}}}{2} = 10$). What used to be a list of separate hits is, structurally, one machine: D_5 carries the family and the flavor budget, A_3 carries the generations and the gauge dimension, μ_4 glues them (and reappears as the 41 chain, the discriminant norms, and the CP phase step $2\pi/6$), and c_3 dresses everything.

4 Grammar I — the carrier grammar (3 + 2)

Machine checks for this section: [verification/v2_carrier_pascal.py]

From $g_{\text{car}} = 5$ and the split $(b, s) = (3, 2)$ the carrier data follow:

$$\begin{aligned} \gamma_{\text{car}} &= \frac{g_{\text{car}}}{g_{\text{car}} + 1} = \frac{5}{6}, & N_{\text{fam}} &= \frac{g_{\text{car}} + 1}{2} = 3, & \dim S^+ &= 2^{g_{\text{car}} - 1} = 16, \\ \Omega_{\text{adm}} &= N_{\text{fam}} 2^{g_{\text{car}} - 1} = 48, & b_1 &= \frac{g_{\text{car}} 2^{g_{\text{car}} - 2} + 1}{10} = \frac{41}{10}. \end{aligned} \tag{19}$$

4.1 Dual-root dictionary (everything from two charges)

Two carrier roots generate the family and the transport cusps [E]. With $y_+ = \frac{1}{2}$ and $y_- = -\frac{1}{3}$ (the determinant-normalised carrier eigenvalues), the full one-family hypercharge spectrum

and the flavor-transport cusps are sums of the two roots:

$$\begin{aligned} Y(Q_L) &= y_+ + y_- = \frac{1}{6}, & Y(u^c) &= 2y_- = -\frac{2}{3}, & Y(d^c) &= -y_- = \frac{1}{3}, \\ Y(L_L) &= -y_+ = -\frac{1}{2}, & Y(e^c) &= 2y_+ = 1, & Y(\nu^c) &= 0, \\ \{1, \frac{2}{3}, \frac{1}{3}\} &= \{2y_+, -2y_-, 2(y_+ + y_-)\}, & \gamma_{\text{car}} &= \text{Tr}_E Y^2 = 3y_-^2 + 2y_+^2 = \frac{5}{6}, \\ \varphi_{\text{base}} &= \frac{1 - \gamma_{\text{car}}}{\pi} = \frac{1}{6\pi}. \end{aligned}$$

All residuals are exactly zero. The Standard-Model charge table is not assumed; it is built by addition from $\{1/2, -1/3\}$.

The $X = 6Y$ quadratic and the hypercharge Pascal-sign duality [E]

Clearing denominators with $X = 6Y$ turns the two base charges into the integer roots $x_+ = 3$, $x_- = -2$ of

$$\boxed{x^2 - x - 6 = 0} : \quad x_+ + x_- = 1 = N_\Phi, \quad x_+ - x_- = 5 = g_{\text{car}},$$

$$-x_+x_- = 6 = |R^+(A_3)|, \quad \sqrt{\Delta} = 5,$$

so the Higgs singleton, the carrier rank and the A_3 hexagon all fall out of one quadratic. Counting the signs of X over a generation ($X = 1^{\times 6}, 2^{\times 3}, 6^{\times 1}, (-4)^{\times 3}, (-3)^{\times 2}, 0^{\times 1}$) gives the *Pascal triple* again:

$$\boxed{(\#X=0, \#X<0, \#X>0) = (1, 5, 10) = \binom{5}{0}, \binom{5}{1}, \binom{5}{2}},$$

$$10-5=g_{\text{car}}, \quad 5-1=|\mu_4|, \quad 10-1=\text{tr } R,$$

with $\nu^c \leftrightarrow \binom{5}{0}$ the neutral singleton — the right-handed neutrino is the Pascal zero with a job. The moment ladder adds $\text{Tr } X^2 = 120 = 5!$ and $\text{Tr } X^4 / \text{Tr } X^2 = 2280/120 = 19$ (an E_8 exponent; audit). [E]

4.2 Carrier arithmetic (proved identities)

The discrete coefficients are not isolated numbers; they are exact carrier traces:

$$10b_1 = 41 = g_{\text{car}}^2 + (g_{\text{car}} - 1)^2 = \Omega_{\text{adm}} \gamma_{\text{car}} + 1 = 1 + h_{E_8} \frac{\dim S^+}{\dim \mathfrak{g}_{\text{SM}}} = 1 + 30 \cdot \frac{16}{12} = 40 + 1, \quad [\text{E}]$$

$$I_1^{\text{EW}} = \frac{41}{48} = \text{Var}_{S^+}(Y) \cdot b_1 = \frac{5}{24} \cdot \frac{41}{10} = \gamma_{\text{car}} + \frac{1}{\Omega_{\text{adm}}}, \quad [\text{E}]$$

$$\text{Tr}_{S^+}(X^2) = 120 = 5! \quad (X = 6Y), \quad \delta_2 = \frac{5}{4}\delta_{\text{top}}^2 = 2880 c_3^8 = 4! \cdot 5! c_3^8 = 4! \text{Tr}_{S^+}(X^2) c_3^8, \quad [\text{E}]$$

$$\frac{5}{4} = \underbrace{\frac{3}{2} \cdot \frac{5}{6}}_{B\gamma} = \underbrace{\frac{D_{\text{start}}}{\Omega_{\text{adm}}}}_{60/48} = \underbrace{\frac{g_{\text{car}}}{g_{\text{car}}-1}}_{5/4} = \frac{\delta_2}{\delta_{\text{top}}^2}. \quad [\text{E}]$$

So: b_1 is a Pythagorean carrier identity ($5^2 + 4^2$); equivalently the anchor-product form $10b_1 = p_1p_3 + (g_{\text{car}} - p_1) = 4 \cdot 10 + (5 - 4) = 41$ reads “all four glue corners times all ten carrier pairs, plus the one leftover Higgs slot”; the electroweak exponent $41/48$ is the hypercharge variance times the $U(1)$ budget; the second seam defect $2880 c_3^8$ is a product of factorial traces $4! \cdot 5!$; and the universal compression $5/4 = g_{\text{car}}/(g_{\text{car}} - 1)$ recurs whenever the five-slot carrier projects onto four effective directions. As a minor curiosity, the $SU(5)$ -tree weak mixing angle rewrites as $\sin^2 \theta_W^{\text{tree}} = \frac{3}{8} = N_{\text{fam}}/(g_{\text{car}} + N_{\text{fam}}) = N_{\text{fam}}/\text{rank}(E_8)$ (the value is the standard one; the rewrite is a coincidence to note, not a claim). [C]

5 Grammar II — the seed grammar (one decoder u)

Machine checks for this section: [verification/v106_review_validation.py]

The seed is, exactly, a pure function of c_3 :

$$u = \varphi_0^{\text{ret}} = \frac{4}{3} c_3 + \Omega_{\text{adm}} c_3^4 = \underbrace{\frac{1}{6\pi}}_{\varphi_{\text{base}}} + \underbrace{\frac{3}{256\pi^4}}_{\delta_{\text{top}}} = 0.0531719521768 \quad [\text{E}]. \quad (24)$$

Its two coefficients are themselves carrier data: $\frac{4}{3} = \dim S^+ / \dim \mathfrak{g}_{\text{SM}} = 16/12 = 8/(g_{\text{car}} + 1)$ (so $\varphi_{\text{base}} = 1/((g_{\text{car}} + 1)\pi)$) and $48 = \Omega_{\text{adm}}$. The seed is thus a pure function of c_3 and the carrier counts.


One seed, four readouts (and their inverses) [E].

$$\begin{aligned} \lambda_C^2 &= u(1-u), \quad \beta_{\text{rad}} = \frac{u}{4\pi}, \quad \Omega_b = \left(1 - \frac{1}{4\pi}\right)u, \quad \sin^2 \theta_{13} = e^{-5/6} u, \\ u &= \frac{1 - \sqrt{1 - 4\lambda_C^2}}{2} = 4\pi \beta_{\text{rad}} = \frac{\Omega_b}{1 - 1/(4\pi)} = e^{5/6} \sin^2 \theta_{13} \\ &= \sum_{n \geq 0} C_n \lambda_C^{2n+2} = \lambda_C^2 + \lambda_C^4 + 2\lambda_C^6 + 5\lambda_C^8 + \dots \end{aligned}$$

One knob u turns four dials simultaneously: Cabibbo mixing (the Bernoulli width $\sqrt{u(1-u)}$), cosmic birefringence, the baryon fraction, and the reactor angle. Turning one forces the others; there is no free per-observable tuning. The exponent $5/6$ is the carrier trace $\gamma_{\text{car}} = \text{Tr}_E Y^2$, *not* the Euler–Mascheroni constant $0.5772\dots$ (audit [O]).

6 Grammar III — the action grammar (exponentials of $\alpha_\star^{-1}$)

The relation in plain words. The fine-structure constant is not just “a number near 137”: it is the *one exponential engine* behind three wildly separated scales. Take the same $\alpha_\star^{-1} = 137.036$ and divide the carrier into it: $\frac{1}{5}$ of it sets the *electroweak* scale, $1 \times$ it sets the *Hubble* scale, $2 \times$ it sets the *cosmological constant*. The three action charges are the Pascal row $1 : 5 : 10$ of the five-slot carrier, and the Hubble rung is just the square root of the vacuum energy. So one constant fixes the electroweak hierarchy, dark energy and the expansion rate at once.



action charges $1 : g_{\text{car}} : 2g_{\text{car}} = 1 : 5 : 10 = \binom{5}{0} : \binom{5}{1} : \binom{5}{2}$ (Pascal row of K_5)

Read every dimensionless ratio through its *action* and ask for the action charge:

$$x = R_x e^{-A_x}, \quad A_x = q_x \alpha_\star^{-1} + \Delta_x, \quad q_x \in \left\{ \frac{1}{g_{\text{car}}}, 1, 2 \right\}. \quad (25)$$

The robust ladder and its residues are

sector	action A_x	charge q_x	residue R_x	value of $-\ln x$
electroweak $v_{\text{geo}}/\bar{M}_{\text{Pl}}$	$(\alpha_\star^{-1} + \delta_{\text{ph}})/g_{\text{car}}$	$1/g_{\text{car}}$	$g_{\text{car}}\beta_{\text{rad}}^2$	36.85
Hubble $H_0/\bar{M}_{\text{Pl}}$	$\alpha_\star^{-1}$	1	$1/(2\pi\sqrt{\Omega_\Lambda})$	138.68
cosm. const. $\rho_\Lambda/\bar{M}_{\text{Pl}}^4$	$2\alpha_\star^{-1}$	2	$3/(4\pi^2)$	276.65

$$A_{\text{EW}} = \frac{\alpha_\star^{-1} + \delta_{\text{ph}}}{g_{\text{car}}} = 27.5259, \quad \mathcal{A}_\Lambda = 2\alpha_\star^{-1} = 274.072, \quad \boxed{A_{\text{EW}}:\mathcal{A}_H:\mathcal{A}_\Lambda = 1:5:10}. \quad (26)$$

The leading ratio is exact: $\mathcal{A}_\Lambda/(\alpha_\star^{-1}/g_{\text{car}}) = 2g_{\text{car}} = 10$ (with the transport correction $\mathcal{A}_\Lambda/A_{\text{EW}} = 9.957$). The claim is that the *action charges*, not the raw logarithms, form the ladder.




Figure 6: Action tower (data plot, `verification/make_figures.py`). The three action charges $A_x = -\ln x$ for the electroweak, Hubble and cosmological-constant ratios fall on the line $A = (\alpha_\star^{-1}/5)\binom{5}{k}$: one engine $\alpha_\star^{-1}$ generates all three scales with the Pascal exponents 1 : 5 : 10.

6.1 The EM closure: $\alpha_\star^{-1}$ as an explicit fixed point (made explicit here)

The engine $\alpha_\star^{-1}$ is not an input — it is the unique root of an explicit closure equation (the exact electromagnetic closure theorem). With $c_3 = 1/(8\pi)$, $\varphi_{\text{base}} = 1/(6\pi)$, $q(\alpha) = \delta_{\text{top}}e^{-2\alpha}$ and the seam response $\varphi_{\text{seam}}(\alpha) = \varphi_{\text{base}} + q(\alpha)(1 - q(\alpha))^{-5/4}$,

$$\boxed{F_{U(1)}(\alpha) = \alpha^3 - 2c_3^3\alpha^2 - \frac{4}{5}c_3^6\left(\sum_{f,j} L_{f,j} + N_\Phi\right) \log \frac{1}{\varphi_{\text{seam}}(\alpha)} = 0} \quad [\mathbf{E}]/[\mathbf{C}], \quad (27)$$

where the coefficient is $\frac{4}{5}(\sum L + N_\Phi)c_3^6 = \frac{4}{5} \cdot 41 c_3^6 = 8b_1c_3^6$, so the fine-structure constant reads the *flavor transport budget* $\sum L = 40$ and the $U(1)$ index $b_1 = 41/10$.

Lemma 1 (Existence and uniqueness of the EM fixed point). $F_{U(1)}$ has *exactly one* positive root, and it is simple.

Proof. Write $F_{U(1)}(\alpha) = h(\alpha) - Cg(\alpha)$ with $h(\alpha) = \alpha^3 - 2c_3^3\alpha^2$, $C = \frac{4}{5}c_3^6M > 0$ ($M = 41$), and $g(\alpha) = -\log \varphi_{\text{seam}}(\alpha)$. Since $q(\alpha) = \delta_{\text{top}}e^{-2\alpha} \in (0, 1)$ is strictly decreasing, $\varphi_{\text{seam}} = \varphi_{\text{base}} + q(1 - q)^{-5/4}$ is C^1 and decreasing, so g is C^1 , bounded, and slowly increasing with $g'(\alpha) = O(q)$ tiny; numerically $g \in [2.9342, 2.9343]$ on the admissible window. The cubic h has its interior critical point at $\alpha_c = \frac{4}{3}c_3^3 = 8.399 \times 10^{-5}$, with $h(0) = 0$ and $h < 0$ on $(0, \alpha_c)$. *Monotonicity (corrected endpoint).* Since $F'_{U(1)}(\alpha) = 3\alpha^2 - 4c_3^3\alpha - Cg'(\alpha)$ and at α_c the first two terms *cancel* ($3\alpha_c = 4c_3^3$), one has $F'_{U(1)}(\alpha_c) = -Cg'(\alpha_c) \approx -5.89 \times 10^{-10} < 0$: the derivative is *not* yet positive at α_c . Its first zero lies slightly above, at $\alpha_0 = 8.6264 \times 10^{-5}$ (where $3\alpha_0^2 - 4c_3^3\alpha_0 = Cg'(\alpha_0)$), with $F'_{U(1)} < 0$ on (α_c, α_0)

and $F'_{U(1)} > 0$ on (α_0, ∞) . Now $F_{U(1)}(0) = -C g(0) < 0$, and $F_{U(1)}$ decreases on $(0, \alpha_0]$ (both h and $-Cg$ decrease on $(0, \alpha_c]$, then $F'_{U(1)} < 0$ on $[\alpha_c, \alpha_0]$), so $F_{U(1)}(\alpha_0) \approx -3.82 \times 10^{-7} < 0$; thereafter $F_{U(1)}$ is strictly increasing to $+\infty$. Hence $F_{U(1)} < 0$ on $(0, \alpha_0]$ and strictly increasing on $[\alpha_0, \infty)$, so it crosses zero *exactly once*. The crossing has $F'_{U(1)}(\alpha_*) = 1.58 \times 10^{-4} > 0$, so the root is simple. A sign scan on $(0, 0.05)$ confirms a single change. *Rigorous version:* an **interval-arithmetic** enclosure (`mpmath.iv`) brackets the root in $[0.00729735256220970, 0.00729735256221000]$ with $F_{U(1)} < 0$ on the lower and > 0 on the upper endpoint (definite signs), and a 200-cell interval partition of $(\alpha_c, 0.05)$ contains exactly *one* indeterminate (root-bearing) band — a verified uniqueness, not merely a sampled one. `[verification/v3_em_alpha.py]` $\square$

Solving:

$$\boxed{\alpha_* = 0.00729735256220985\dots, \quad \alpha_*^{-1} = 137.0359992168407\dots} \quad [\text{E}] \quad (28)$$

`[verification/v3_em_alpha.py]` versus CODATA-2022 $\alpha_*^{-1} = 137.035999177(21)$ — a relative deviation 2.9×10^{-10} ($\approx 1.9\sigma$ on the tiny experimental error; the older 2018 value 137.035999084 gave 9.7×10^{-10} , so the agreement *improved* in relative terms). The exponent $-5/4$ in φ_{seam} is the carrier discriminant norm $q(D_5)$, and $\delta_{\text{top}} = 48c_3^4$, so the closure is built only from c_3 , the carrier, and the flavor budget. This discharges the former open audit point: the EM closure is a stated, existence-/uniqueness-proven, numerically reproduced equation, not a fit.

Why *this* function: the boundary $U(1)$ Ward reading (Theorem C) `[E]/[C]`. $F_{U(1)}$ is not a chosen ansatz but the reduced Ward identity of the boundary $U(1)$ partition function $Z_\partial = Z_{\text{Maxwell}} Z_{\text{Calderón}} Z_{\text{transport}}$, with three terms whose coefficients are compiler atoms, not fits:

$$\underbrace{\alpha^3}_{\text{Maxwell moment}} \quad \underbrace{-2c_3^3\alpha^2}_{\text{Calderón sheet } (2=|\mathbb{Z}_2|)} \quad \underbrace{-8b_1c_3^6 \log \frac{1}{\varphi_{\text{seam}}}}_{C_6 \text{ transport det.}} = 0, \quad 8b_1 = \frac{4}{5}M = \frac{4}{5}(\sum L + N_\Phi) = \frac{164}{5}.$$

The Calderón factor $2 = |\mathbb{Z}_2|$ (sheet), the transport power c_3^6 is the C_6 hexagon, and the seam exponent $-\frac{5}{4} = -q(D_5)$ is the carrier discriminant norm. **Typing:** the three-term decomposition and the coefficient identities ($8b_1 = \frac{4}{5}M$, $-\frac{5}{4} = q(D_5)$) are exact `[E]`; the *physical origin* as a boundary Ward closure (the lemmas $\partial_\tau \log Z_{\text{Maxwell}} = \alpha^3$, $\partial_\tau \log Z_{\text{Calderón}} = -2c_3^3\alpha^2$, $\partial_\tau \log \det_\zeta(1 - \mathcal{T}_{C_6, U(1)}) = -8b_1c_3^6 \log \frac{1}{\varphi_{\text{seam}}}$) is a conjectured `[C]` interpretation, not yet machine-proven. `[verification/v48_em_ward.py]`

The determinant-line (Quillen) form. (`[verification/v341_alpha_quillen.py]`) The three terms assemble into a single *stationarity* statement for the boundary $U(1)$ determinant line,

$$\underbrace{[\alpha^3 - 2c_3^3\alpha^2]}_{\delta \log \det \Delta_{U(1)}} = \underbrace{[8b_1c_3^6]}_{\text{ctr.}} \underbrace{[\log \frac{1}{\varphi_{\text{seam}}}]_{\text{seam resp.}}}$$

so $F_{U(1)} = 0$ is $\delta \log \det \Delta_{U(1)} + \text{counterterm} + \text{seam response} = 0$. Every factor is a *named* object, not a knob: the counterterm coefficient is $8b_1c_3^6 = (\text{rank } E_8) b_1 c_3^6$ (index $\times$ heat-kernel), $c_3 = 1/(8\pi)$ is the Gauss–Bonnet boundary coefficient, the exponent $-\frac{5}{4} = -q(D_5)$ is one carrier discriminant norm and the seam base $\varphi_{\text{base}} = 1/(6\pi) = \frac{4}{3}c_3$ carries the *other*, $c_3/\varphi_{\text{base}} = \frac{3}{4} = q(A_3)$ — both glue forms appear, and the split is verified at the root. **Typing (and: no new gate).** The determinant-line assembly and all coefficient identities are exact `[E]`; α_*^{-1} itself stays closed `[E]` (unique root + ablation). This adds *no* open item — it *sharpens* the pre-existing `[O]`/physical-origin obligation **EM.WARD.01** (the “why *this* function” residual, already conjectural) by naming every coefficient. The one still-open part *is* that residual, now *named* as its own tracked target **ALPHA.QUILLEN.EXACT.01** (`[verification/v382_alpha_quillen_exact.py]`): the from-first-principles AQFT/ ζ proof that $\delta_\tau(\log \det_\zeta \Delta_{U(1)} + 8b_1c_3^6 \log \varphi_{\text{seam}}) = 0$ — i.e. that $F_{U(1)}$ is the *exact* Quillen functional of the seam $U(1)$, the variational principle *derived*, not assembled. It is a *face* of **SEAM.EQUIV.01 (v336)** — the seam $U(1)$ determinant line lives on the same holomorphic net, so once the net is $(E_8)_1$ its $U(1)$ sub-determinant is fixed — hence no second independent gate, just the sharpest remaining EM-Ward obligation, now citable on its own. *Honest progress towards it*

(`[verification/v391_alpha_quillen_progress.py]`), without closing it: a solvable model $\det_\zeta(-\partial_\tau^2 + m^2)$ on a circle $= (2\sinh(\beta m/2))^2$ has the closed m -variation $\beta \coth(\beta m/2) = 2/m + (\beta^2/6)m - (\beta^4/360)m^3 + \dots$, a heat-kernel/Laurent series — so a ζ -determinant's variation is *structured* heat-kernel data, and the counterterm $8b_1c_3^6$ (a Gauss–Bonnet heat-kernel power) is the right *kind* of object. This *reduces* ALPHA.QUILLEN.EXACT.01 to the named sub-target “the seam $U(1)$ Seeley–DeWitt coefficient $= 8b_1c_3^6$ ”, but the target itself stays [O] (an external math proof, v384); the value $\alpha_\star^{-1}$ stays [E]. A *second honest step* (`[verification/v433_alpha_quillen_heatkernel.py]`) bridges this reduction to the heat-kernel origin below, without closing it: v391's circle reached only the low orders a_0, a_1 , whereas a solvable $4D$ model — the flat $\Delta = -\partial^2 + m^2$ on T^4 , $\text{Tr } e^{-t\Delta} = V/(4\pi)^2 t^{-2} e^{-tm^2}$ — has $a_4 = Vm^4/(2(4\pi)^2) \neq 0$, reaching the t^0 (conformal-anomaly) order the circle could not, the very order in which the quadratic term is grounded below (v342, Gilkey a_4). So a ζ -determinant's variation is heat-kernel data *at the a_4 order*, and the measure itself is c_3 -built, $(4\pi)^{-2} = (2c_3)^2 = 4c_3^2$; the *actual* seam $a_4 = 8b_1c_3^6$ and the cubic α^3 Chern moment stay [O] (v384), the seam F -normalisation [C] (retyped by the fifth step below, v470, as the embedding index $k_Y=5/3$).

The heat-kernel origin of the terms (`[verification/v342_em_ward_heatkernel.py]`). The determinant-line form is more than bookkeeping: its terms are textbook heat-kernel coefficients. [E] $c_3 = 1/(8\pi)$ is the boundary Gauss–Bonnet coefficient ($\oint_{S^2} K = 2\pi\chi = 4\pi$, $c_3 = 1/(\mathbb{Z}_2 \oint K)$, the same $\chi=2$ as the marks, v216). [E] the α^2 (Calderón) term is the Gilkey gauge-curvature coefficient: the $U(1)$ Laplacian's Seeley–DeWitt a_4 contains $30\Omega_{ij}\Omega^{ij}/360 = \frac{1}{12}\Omega^2$ (Gilkey, *Invariance Theory*, Thm 4.8.16), and with $\Omega = i\alpha F$ this forces an $-(\alpha^2/12)F^2$ contribution to $\log \det \Delta_{U(1)}$ — so the Calderón $-2c_3^3\alpha^2$ is the gauge-curvature (Quillen-curvature) piece, not an ansatz. [E] the c_3 -orders $\{0, 3, 6\}$ are an arithmetic ladder (step 3 = the three channels; $6 = |R^+(A_3)|$, the C_6 hexagon), and the π -power test confirms it: $2c_3^3 = 1/(256\pi^3)$ carries π^3 , i.e. *three* boundary c_3 -insertions, not a single bulk coefficient. **Typing:** the boundary structure and the gauge-curvature origin of the quadratic term are heat-kernel-forced [E]; the exact coefficient value is [C] (the seam F -normalisation — retyped as the embedding index $k_Y=5/3$ by the fifth step below, v470); the cubic α^3 Maxwell moment (a topological/Chern level) and the full “this is the seam $U(1)$ ζ -determinant” claim stay [O] — exactly the EM.WARD.01 residual, unchanged. So the quadratic/curvature half of the EM-Ward origin is now derived; the cubic-moment half remains the honest open part. A *third honest step* (`[verification/v434_alpha_quillen_betafunction.py]`) settles those residuals and shows they are not three independent problems. The counterterm's matter factor is heat-kernel-forced too: by the $\beta = a_4$ theorem the one-loop $U(1)_Y$ β is the a_4 coefficient, and the carrier hypercharge content gives $\sum \frac{2}{3}Y_f^2 + \frac{1}{3}Y_s^2 = 41/6$ (SM) $\Rightarrow b_1 = 41/10$ (GUT, v159/v246), so $8b_1c_3^6 = b_1(\text{rank } E_8)c_3^6$ is an a_4 content coefficient times a boundary geometry — the same *kind* of object as the Calderón a_4 . The exact value then turns only on the seam F -normalisation, so the “actual seam a_4 ” residual is the F -normalisation [C] (one obligation, not two); and the cubic factors $a^3 - 2c_3^3a^2 = a^2(a - 2c_3^3)$, exhibiting a^3 as the first moment of the Chern density a^2 — a level whose from-first-principles origin stays [O]. The EM-Ward residual is thus *one* [C] + *one* [O], not three. A *fourth honest step* (`[verification/v435_alpha_quillen_chernlevel.py]`) sharpens that single remaining [O] — residual (2), the cubic α^3 Chern level — without closing it. A π -power / metric-independence test partitions the closure: the three coefficients carry π -powers exactly $\{0, 3, 6\}$ ($k_0 = 1$; $-2c_3^3 = -1/(256\pi^3)$; $-8b_1c_3^6 = -41/(327680\pi^6)$), so the Maxwell a^3 is the *unique* π^0/c_3^0 (metric-independent) term. A heat-kernel boundary coefficient always carries the $(4\pi)^{-d/2}$ measure; a metric-independent term carries none — the signature of a *topological* term. So the test confirms, now from a test rather than a claim, that a^3 is a level and not a Seeley–DeWitt curvature integral (the other two terms are heat-kernel boundary data). [E] *Conditional* on the Chern–Simons/ η reading of the Maxwell moment (the EM1 interpretation, v48), large-gauge invariance quantises the seam $U(1)$ level to $\mathbb{Z}$, so $k_0 = 1$ is the *unit* (minimal) level — an integer, not a tunable real. [C] What stays [O] is the EM1 lemma itself — the from-first-principles Chern–Simons-boundary proof that $\partial_\tau \log Z_{\text{Maxwell}} = a^3$ — external math (type A, v384). A *fifth honest step* (`[verification/v470_alpha_inflow_level.py]`) upgrades both leftovers, still without closing the target. [C] The level is computed, not minimal: under the same EM1 reading, the

a^3 coefficient $k_0 = 1$ equals the FHS bulk Chern invariant $|C| = 1$ of the very p+ip collar that realises S3 of SEAM.EQUIV.01 (v367/v461; control $C(M=3) = 0$) — “why exactly 1” is answered by a computation plus the cited inflow identification (TKNN / Avron–Seiler–Simon quantisation; Callan–Harvey inflow; the APS/Witten η =CS reading of $\delta \log \det$; Quillen/Bismut–Freed determinant-line integrality), replacing the minimality assumption of the fourth step. [E] *The F -normalisation is the embedding index:* $k_Y = \text{tr}_5(Y^2)/\text{tr}(T_3^2) = (5/6)/(1/2) = 5/3$ (Ginsparg 1987), so $(3/5) \times \frac{41}{6} = \frac{41}{10} = b_1$ exactly — the “GUT 3/5” is the inverse embedding index, and once the seam net is $(E_8)_1$ at level 1 the $U(1)_Y$ normalisation $\langle J(z)J(w) \rangle = k/(z-w)^2$ is fixed by it (level-1 current-algebra rigidity, no continuous dial): the seam F -normalisation [C] has zero independent content — a face of SEAM.EQUIV.01, per the v382 typing. [C] *One phase, two responses:* the collar phase carries $c_- = 8$ (gravitational, feeding SEAM.EQUIV.01/S3, v456) and $C = 1$ ($U(1)$, feeding EM1) — so discharging the one realisation input feeds both named targets at once. *A sixth honest step* ([verification/v472_quillen_detline_moduli.py]) exhibits that bridge lemma at the finite level. ALPHA.QUILLEN.EXACT.01 names the level as the curvature of the determinant line over the $U(1)$ moduli (Quillen/Bismut–Freed curvature, Dai–Freed section) — and the fifth step computed the Chern invariant only over the Bloch Brillouin zone, the translation-invariant shortcut. The sixth step computes the Quillen-shaped object itself: the same collar Hamiltonian in real space with twisted boundary conditions — the twist torus (θ_x, θ_y) is the moduli space of flat $U(1)$ connections — and the Fukui–Hatsugai–Suzuki curvature of the determinant line of the occupied frame over that torus integrates to $2\pi \cdot 1$ exactly (10^{-9}), size-independently ($L = 4, 6$), with clean controls (trivial collar $\rightarrow 0$, orientation flip $\rightarrow -1$), the twist-moduli integer equal to the Bloch-BZ integer for all three phases (Niu–Thouless–Wu), and the Fermi gap open over the whole torus (the Dai–Freed section has no zero). So “ $\delta \log \det(\text{seam model}) = \text{the inflow response}$ ” holds verbatim at the finite/model level, where $\log \det$ of the occupied frame replaces $\log \det_\zeta$. [E] (computation) / [C] (the EM1 reading). What stays [O] is the continuum identification “ $\delta \log \det_\zeta(\text{seam}) = \text{the inflow response}$ ” on the abstract seam (the SEAM.EQUIV.01 face); ALPHA.QUILLEN.EXACT.01 stays [O], and $\alpha_\star^{-1}$ stays [E] regardless. *A seventh step unifies this target with the φ_0^{ret} -puncture target* ([verification/v484_seam_contact_unit.py]). The $\{0, 3, 6\}$ c_3 -ladder above and the puncture’s per-mark weight c_3 (v483) are one rule — the seam contact unit — and it is not a new unknown: “ c_3 per insertion” is the KMS seam unit $2\pi = 1/(4c_3)$ with $\frac{1}{4} = 1/|\mu_4|$ (v239), i.e. one bare boundary propagator ($\frac{1}{2\pi} = 4c_3$) orbit-averaged over the four marks. [E] derived on the seam circle for the finite cycle sector (the bare Green function takes integer multiples of $c_3 \ln 2$ at the μ_4 separations; the log-det contact expansion is exactly graded in c_3 per insertion), with the grammar already suite-wide (the Λ prefactor $\frac{3}{4\pi^2} = 48c_3^2$, v60, carries the same $\Omega_{\text{adm}} = 48$ at two insertions). [C]/[O] Consequence: ALPHA.QUILLEN.EXACT.01 and HYP.PHIO.PUNCTURE.01 merge into one remaining analytic step — the diagonal zeta-renormalisation at the marks plus the state-multiplicity matching ($41 = 10b_1$ here, $48 = \Omega_{\text{adm}}$ there). *An eighth step settles that remaining step at the computable level* ([verification/v485_contact_diagonal_closed.py]). [E] The renormalised coincident-point Green function on a circle of circumference ℓ is $G_{\text{reg}}(0; \ell) = \frac{1}{\pi} \ln(\ell/2\pi)$ exactly — it vanishes precisely at $\ell = 2\pi = 1/(4c_3)$, the KMS seam circumference (v239): the seam scale is the unique zero-self-energy scheme point, and the log is the pure scale channel (matching the fact that $F_{U(1)}$ carries exactly one log term while the φ_0^{ret} split is log-free). [E] With the diagonal zero the mark contact calculus resums in closed form (the BFK/gluing route, v151): $\det(I - uC) = (1 - 4u)(1 + 2u)^2$ with $u = \varepsilon c_3 \ln 2$ — all orders, the linear term absent because $\text{Tr } C = 0$. [E] And the two multiplicities are one state set under two response weights: the same $48 = 3 \times 16$ admissible states counted flat give the φ_0^{ret} puncture count $48 = \Omega_{\text{adm}}$, and $Y^2/\text{loop}/\text{GUT}$ -weighted give the alpha budget $41 = 10b_1 = 40 + 1$ ($\text{Tr}_{16} Y^2 = \frac{10}{3}$ exact, $\nu^c = 0$; Ginsparg, v470). [C]/[O] Net: every finite/computable piece of the merged target is proven; the single remaining [O] is the identification of the abstract seam ζ -det with this glued mark determinant — exactly the v382 typing (a face of SEAM.EQUIV.01), now substantiated computationally rather than asserted.

The full ablation — which inputs are load-bearing and which only set the last digit — is in the Ablation appendix below.

Inversion test (this round). Running the closure backwards — inserting the *measured* α and solving for the integer budget $M = \sum L + N_\Phi$ — returns $M = 41.0000001\dots$, i.e. the external measurement reconstructs the internal discrete budget $41 = 40 + 1$ to 10^{-7} . This is stronger than “we hit α ”: the equation reads the discrete 41 out of nature. [E]

6.2 The decuple bridge (new, exact headline)

Eliminating $\alpha_\star^{-1} = g_{\text{car}} A_{\text{EW}} - \delta_{\text{ph}}$ between the electroweak and the cosmological-constant lines gives a direct relation between the two scales:

Decuple bridge: Λ is the tenth power of the stripped EW scale [E]

$$\frac{\rho_\Lambda}{\bar{M}_{\text{Pl}}^4} = \frac{3}{4\pi^2} e^{2\delta_{\text{ph}}} \left[\frac{v_{\text{geo}}}{5\beta_{\text{rad}}^2 \bar{M}_{\text{Pl}}} \right]^{10} \quad (29)$$

where the exponent $10 = 2g_{\text{car}}$ and $5 = g_{\text{car}}$. Verified to a relative residual $\sim 10^{-159}$.

Proof. From $v_{\text{geo}}/\bar{M}_{\text{Pl}} = g_{\text{car}}\beta_{\text{rad}}^2 e^{-A_{\text{EW}}}$, $e^{-A_{\text{EW}}} = (v_{\text{geo}}/\bar{M}_{\text{Pl}})/(g_{\text{car}}\beta_{\text{rad}}^2)$. From $\rho_\Lambda/\bar{M}_{\text{Pl}}^4 = \frac{3}{4\pi^2} e^{-2\alpha_\star^{-1}}$ and $\alpha_\star^{-1} = g_{\text{car}} A_{\text{EW}} - \delta_{\text{ph}}$, $e^{-2\alpha_\star^{-1}} = e^{2\delta_{\text{ph}}} e^{-2g_{\text{car}} A_{\text{EW}}} = e^{2\delta_{\text{ph}}} (e^{-A_{\text{EW}}})^{2g_{\text{car}}}$. Substituting and using $2g_{\text{car}} = 10$, $g_{\text{car}} = 5$ gives the boxed form. $\square$

In words: the electroweak scale is not directly the Λ scale, but once the electroweak ratio is stripped of its small β_{rad} prefactor, Λ is its tenth power, and the “ten” is $2g_{\text{car}}$, not a fitted exponent. This is the direct algebraic compression of the action ladder.

6.3 The full power tower: x^1, x^5, x^{10} on the carrier graph K_5

The decuple bridge is one rung of a tower. Writing $x := e^{-A_{\text{EW}}} = v_{\text{geo}}/(5\beta_{\text{rad}}^2 \bar{M}_{\text{Pl}})$ for the stripped electroweak unit, all three cosmological scales are powers of the *same* x with binomial exponents (verified exactly):

$$\frac{v_{\text{geo}}}{5\beta_{\text{rad}}^2 \bar{M}_{\text{Pl}}} = x^{\binom{5}{0}} = x, \quad \frac{H_0}{\bar{M}_{\text{Pl}}} = x^{\binom{5}{1}} R_H = x^5 R_H, \quad \frac{\rho_\Lambda}{\bar{M}_{\text{Pl}}^4} = x^{\binom{5}{2}} R_\Lambda = x^{10} R_\Lambda \quad (30)$$

with residues $R_H = e^{\delta_{\text{ph}}}/(2\pi\sqrt{\Omega_\Lambda})$ and $R_\Lambda = \frac{3}{4\pi^2} e^{2\delta_{\text{ph}}}$. [E]. The exponents are the complete-graph counts of the five-slot carrier K_5 :

$$H \leftrightarrow \text{vertices of } K_5 \text{ (}\# = \binom{5}{1} = 5\text{)}, \quad \Lambda \leftrightarrow \text{edges of } K_5 \text{ (}\# = \binom{5}{2} = 10\text{)}.$$

So the cosmological constant is the product of one stripped-electroweak unit over every carrier *pair* (K_5 edge product), and the Hubble scale the product over every carrier *slot* (K_5 vertex product). This is the cleanest single-variable statement of the action ladder.

6.4 Hubble is the square root of Λ (corollary, exact)

On the comparison layer $\rho_\Lambda/\bar{M}_{\text{Pl}}^4 = 3\Omega_\Lambda(H_0/\bar{M}_{\text{Pl}})^2$; equating to the TFPT line gives

$$\frac{H_0}{\bar{M}_{\text{Pl}}} = \frac{e^{-\alpha_\star^{-1}}}{2\pi\sqrt{\Omega_\Lambda}}, \quad \mathcal{A}_H = -\ln \frac{H_0}{\bar{M}_{\text{Pl}}} = \alpha_\star^{-1} + \ln(2\pi\sqrt{\Omega_\Lambda}) = 138.68. \quad (31)$$

So the Hubble rung is *not* an independent third hit: it is the square root of the vacuum energy plus a geometric cosmology factor. The genuine TFPT content is the single $e^{-\alpha_\star^{-1}}$. (The numerical coincidence $\mathcal{A}_H - \alpha_\star^{-1} \approx \pi^2/6$ is just $\ln(2\pi\sqrt{\Omega_\Lambda})$ and is Ω_Λ -dependent; not a claim. [O])

6.5 The ~ 123 orders of the cosmological constant

$$\log_{10} \frac{M_{\text{Pl}}^4}{\Lambda_{\text{IR}}} = \underbrace{\frac{2\alpha_\star^{-1}}{\ln 10}}_{119.028} + \underbrace{\log_{10} \frac{1}{\delta_{\text{top}}}}_{3.920} = 122.948 \approx 123. \quad (32)$$

The “119” is the action $2\alpha_\star^{-1}$; the “ ≈ 4 ” is the seam defect $\delta_{\text{top}} = \Omega_{\text{adm}} c_3^4$.

High-precision cosmological constant: a $\sim 10^{-13}$ prediction [E]

Because $\alpha_\star^{-1}$ is a *derived* number (the EM fixed point, known to 13 digits) and the residue $3/(4\pi^2)$ is exact, the dimensionless vacuum energy is predicted essentially exactly:

$$\frac{\rho_\Lambda}{\bar{M}_{\text{Pl}}^4} = \frac{3}{4\pi^2} e^{-2\alpha_\star^{-1}} = 7.12533 \times 10^{-121} \quad [\text{E}] \text{ [verification/v7_gravity_cosmo.py]}, \quad (33)$$

i.e. a dark-energy scale $\rho_\Lambda^{1/4} = (\frac{3}{4\pi^2})^{1/4} \bar{M}_{\text{Pl}} e^{-\alpha_\star^{-1}/2} = 2.23747 \text{ meV}$ (reduced Planck mass $\bar{M}_{\text{Pl}} = 2.435323 \times 10^{18} \text{ GeV}$). The *dimensionless* ratio carries no observational input — its uncertainty is that of $\alpha_\star^{-1}$, $\sim 10^{-13}$ relative; the absolute meV value inherits only the Planck-mass uncertainty ($\sim 2 \times 10^{-5}$). Both are *far* sharper than the measured vacuum energy (Planck $\rho_\Lambda^{1/4} \simeq 2.24 \text{ meV}$, known to $\sim 1\%$ and entangled with the H_0 tension). So TFPT pins the cosmological constant several orders of magnitude more precisely than it is currently measured — a genuine, falsifiable forward prediction: any future sub-percent ρ_Λ must land on 2.2375 meV.

6.6 The ladder is binomial; $2g_{\text{car}} = \binom{g_{\text{car}}}{2}$ singles out $g_{\text{car}} = 5$

In units of the electroweak action $\alpha_\star^{-1}/g_{\text{car}}$, the three rungs are the first three binomial coefficients of the five-slot carrier:

$$\frac{A_{\text{EW}}}{\alpha_\star^{-1}/g_{\text{car}}} : \frac{\mathcal{A}_H}{\alpha_\star^{-1}/g_{\text{car}}} : \frac{\mathcal{A}_\Lambda}{\alpha_\star^{-1}/g_{\text{car}}} = 1 : g_{\text{car}} : 2g_{\text{car}} = \binom{g_{\text{car}}}{0} : \binom{g_{\text{car}}}{1} : \binom{g_{\text{car}}}{2} = 1 : 5 : 10. \quad (34)$$

The first two are automatic ($\binom{g_{\text{car}}}{0} = 1$, $\binom{g_{\text{car}}}{1} = g_{\text{car}}$). The non-trivial content is the cosmological-constant charge:

$$2g_{\text{car}} = \binom{g_{\text{car}}}{2} \iff g_{\text{car}} = 5 \quad [\text{E}] \text{ [verification/v2_carrier_pascal.py]}, \quad (35)$$

since $\binom{g_{\text{car}}}{2} = \frac{g_{\text{car}}(g_{\text{car}}-1)}{2} = 2g_{\text{car}} \iff g_{\text{car}} = 5$. So the “double overlap” charge $2g_{\text{car}}$ equals the number of carrier-slot *pairs* $\binom{g_{\text{car}}}{2}$ *only* at $g_{\text{car}} = 5$: the cosmological constant is suppressed by one action unit per carrier pair, and this combinatorial reading is exclusive to the physical rank. It is an independent overdetermination of $g_{\text{car}} = 5$ from the action grammar (cf. the landscape section).

6.7 Discriminant action theorem: the ladder reads the hypercharge discriminant

The g_{car} in the action ladder is not merely a rank: it is the square root of the discriminant of the carrier (hypercharge) polynomial. For a general split $b + s = g$ the carrier polynomial is $bsY^2 + (s-b)Y - 1 = 0$ with discriminant

$$\Delta_Y = (s-b)^2 + 4bs = (b+s)^2 = g_{\text{car}}^2, \quad \text{so for } 6Y^2 - Y - 1 = 0 : \Delta_Y = 25, \sqrt{\Delta_Y} = 5 = g_{\text{car}}. \quad (36)$$

Hence the action ladder is, exactly,

$$A_{\text{EW}} = \frac{\alpha_\star^{-1}}{\sqrt{\Delta_Y}}, \quad \mathcal{A}_H = \sqrt{\Delta_Y} A_{\text{EW}}, \quad \mathcal{A}_\Lambda = \binom{\sqrt{\Delta_Y}}{2} A_{\text{EW}} \quad [\text{E}]. \quad (37)$$

The electroweak hierarchy therefore reads the *discriminant of the hypercharge polynomial*, tying the Standard-Model hypercharge derivation directly to the EW-hierarchy and Λ problems.

6.8 A two-dimensional action lattice

All scales sit on the integer lattice spanned by two generators — the gauge unit $\alpha_\star^{-1}/g_{\text{car}}$ and the flavor unit $L_0 = \ln(1/u)$ — as $S_x = a_x(\alpha_\star^{-1}/g_{\text{car}}) + b_x L_0 + (\text{residue})$:

scale	a_x	b_x	axis
$v_{\text{geo}}/\bar{M}_{\text{Pl}}$	$1 = \binom{5}{0}$	0	gauge
$H_0/\bar{M}_{\text{Pl}}$	$5 = \binom{5}{1}$	0	gauge
$\rho_\Lambda/\bar{M}_{\text{Pl}}^4$	$10 = \binom{5}{2}$	0	gauge
$m_e/\bar{M}_{\text{Pl}}$	1	5	flavor
$m_\mu/\bar{M}_{\text{Pl}}$	1	3	flavor
$m_\tau/\bar{M}_{\text{Pl}}$	1	2	flavor

The cosmological/gauge sector runs along $b = 0$ with binomial charges; the charged-fermion sector runs along $a = 1$ with the integer flavor ladder $n_f = (5, 3, 2)$. This is the compact “periodic table” of TFPT scales: two integer labels per scale. **[E]** (charges) / **[C]** (residues).

A third (seam/gravity) generator completes the lattice: $L_c = \ln(1/c_3) = \ln(8\pi)$. The scalaron lives purely on it, $M_{\text{scal}}/\bar{M}_{\text{Pl}} = c_3^{7/2} = e^{-\frac{7}{2}L_c}$, and the seam defect is $\delta_{\text{top}} = e^{-(4L_c - \ln 48)}$. So the three log-generators are $\{\alpha_\star^{-1}$ (hierarchy), $L_0 = \ln(1/u)$ (flavor), $L_c = \ln(1/c_3)$ (seam/gravity) $\}$. **[E]**

7 The Einstein normaliser as a corrected tree value

Machine checks for this section: `[verification/v1_e8_glue.py]`

The gravitational normaliser $\xi = c_3/u$ has a transparent exact form. Writing $u = \frac{1}{6\pi} [1 + \frac{9}{128\pi^3}]$,

$$\xi = \frac{c_3}{u} = \frac{3}{4} \cdot \frac{1}{1 + \frac{9}{128\pi^3}} = 0.748303 \quad \textbf{[E]}. \quad (38)$$

So $\xi_{\text{tree}} = \frac{3}{4} = \dim \mathfrak{g}_{\text{SM}} / \dim S^+ = 12/16$ is the exact leading value (the inverse of the $\frac{4}{3} = \dim S^+ / \dim \mathfrak{g}_{\text{SM}}$ family/gauge compression that sits in the seed $u_{\text{base}} = \frac{4}{3}c_3$, so $\xi_{\text{tree}} u_{\text{base}} = c_3$), and the deviation is precisely the small topological seed correction $9/(128\pi^3)$, i.e. 0.226%. Gravity does not pick up a mysterious factor: first comes $3/4$, then the seam defect shifts it minimally.

8 Λ -anchored metrology closure, and two anchors

Machine checks for this section: `[verification/v60_lambda_metrology_branch.py]`

The absolute SI scale is a calibration, not a compiler output. The dimensionless vacuum quotient lets one anchor it to the observed geometric curvature without circularity.

Theorem 1 (Λ -anchored metrology closure). With $\lambda_\Lambda := \rho_\Lambda/\bar{M}_{\text{Pl}}^4 = \frac{3}{4\pi^2} e^{-2\alpha_\star^{-1}}$ and the observed geometric curvature $\Lambda_{\text{geom}}^{\text{obs}} > 0$,

$$\bar{M}_{\text{Pl}}^2 = \frac{\Lambda_{\text{geom}}^{\text{obs}}}{\lambda_\Lambda}, \quad G_N^{\text{TFPT}} = \frac{c^3}{8\pi\hbar} \frac{\lambda_\Lambda}{\Lambda_{\text{geom}}^{\text{obs}}} = \frac{3c^3}{32\pi^3\hbar \Lambda_{\text{geom}}^{\text{obs}}} e^{-2\alpha_\star^{-1}}.$$

Proof. $\bar{M}_{\text{Pl}}^2 = 1/(8\pi G_N)$ and $\Lambda_{\text{geom}} = \rho_\Lambda/\bar{M}_{\text{Pl}}^2$; the branch sets $\rho_\Lambda = \lambda_\Lambda \bar{M}_{\text{Pl}}^4$, so $\Lambda_{\text{geom}} = \lambda_\Lambda \bar{M}_{\text{Pl}}^2$, giving $\bar{M}_{\text{Pl}}^2$ and then $G_N = \lambda_\Lambda/(8\pi\Lambda_{\text{geom}}^{\text{obs}})$; restoring $c^3/\hbar$ closes it. $\square$

Non-circularity ([O], essential). The anchor must be the curvature $\Lambda_{\text{geom}}^{\text{obs}}$ (units m^{-2}), *not* the SI vacuum energy density $\rho_\Lambda^{\text{obs}} = \Lambda_{\text{geom}} c^4/(8\pi G_N)$, which already contains G_N . The curvature follows from $\Lambda_{\text{geom}}^{\text{obs}} = 3\Omega_\Lambda H_0^2/c^2$.

Two independent anchors (crosscheck with teeth). The metrology can be pinned in two ways that should agree:

1. *Electroweak anchor*: the dimensionless $G_N v_{\text{phys}}^2 = 4.06717 \times 10^{-34}$ fixes G_N once v_{phys} is calibrated non-gravitationally.
2. *Λ -curvature anchor* (this note): $G_N = \frac{3c^3}{32\pi^3 \hbar \Lambda_{\text{geom}}^{\text{obs}}} e^{-2\alpha_\star^{-1}}$.

With the example anchor $H_0 = 67.36 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_\Lambda = 0.6847$ ($\Lambda_{\text{geom}}^{\text{obs}} = 1.08914 \times 10^{-52} \text{ m}^{-2}$) the Λ anchor gives

$$G_N^{\text{TFPT}} = 6.65071 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}, \quad G_N^{\text{TFPT}} / G_N^{\text{CODATA}} - 1 = -3.5 \times 10^{-3}. \quad (39)$$

Closure algebra [E]; SI anchor and numerical match [C] (the cosmological anchor is model dependent; the Hubble tension applies). EW crosscheck: the Λ -pinned $\bar{M}_{\text{Pl}} = 2.43964 \times 10^{18} \text{ GeV}$ gives $v_{\text{geo}} = 242.6 \text{ GeV}$, needing $Z_{\text{EW}} = (246.22/242.61)^2 = 1.030$.

Prefactor branch ([O]). This note uses $\Lambda_{\text{IR}}/M_{\text{Pl}}^4 = \delta_{\text{top}} e^{-2\alpha_\star^{-1}}$. A competing prefactor $2c_3 e^{-2\alpha_\star^{-1}}$ would differ by $2c_3/\delta_{\text{top}} = 661.5$ and misscale G_N by the same factor; the branch must be versioned.

Mandatory Λ -layer typing ([O]). Three distinct Λ objects must never be conflated:

$$\underbrace{\frac{\rho_\Lambda}{M_{\text{Pl}}^4} = \frac{3}{4\pi^2} e^{-2\alpha_\star^{-1}}}_{\text{vacuum density quotient (dimensionless)}} \neq \underbrace{\frac{\Lambda_{\text{IR}}}{M_{\text{Pl}}^4} = \delta_{\text{top}} e^{-2\alpha_\star^{-1}}}_{\text{seam-determinant / IR scale (dimensionless)}} \neq \underbrace{\Lambda_{\text{geom}}^{\text{obs}} = \frac{3\Omega_\Lambda H_0^2}{c^2}}_{\text{observed curvature [m}^{-2}\text{]}}.$$

Only $\Lambda_{\text{geom}}^{\text{obs}}$ carries a dimension and is the non-circular anchor. Keeping the three typed protects the decuple bridge and the metrology closure from a spurious prefactor objection.

Measurement-near (physical) decuple bridge. Since $v_{\text{phys}} = v_{\text{geo}} \sqrt{Z_{\text{EW}}}$ (charged-current residue, not a fit), the decuple bridge in the observable vev reads

$$\frac{\rho_\Lambda}{M_{\text{Pl}}^4} = \frac{3}{4\pi^2} e^{2\delta_{\text{ph}}} \left[\frac{v_{\text{phys}}}{5\beta_{\text{rad}}^2 \sqrt{Z_{\text{EW}}} \bar{M}_{\text{Pl}}} \right]^{10}. \quad (40)$$

This is the same identity, written in the low-energy charged-current scale rather than the geometric source scale; both forms should appear side by side. [E]/[C]

9 The landscape: $g_{\text{car}} = 5$ is overdetermined

Machine checks for this section: [verification/v6_bootstrap.py] [verification/v14_carrier_uniqueness.py]

Varying the carrier rank and rebuilding every constant from the same machinery (c_3 fixed), the EM-closure value $\alpha^{-1}(g_{\text{car}})$ is monotone and meets the observed value only at $g_{\text{car}} = 5$:

g_{car}	N_{fam}	$\dim S^+$	Ω_{adm}	$10b_1$	$\alpha^{-1}(g_{\text{car}})$	verdict
3	2	4	8	7	258.15	wrong α , wrong family count
4	2.5	8	20	17	187.31	non-integer families
5	3	16	48	41	137.036	SM, 3 families, observed α
6	3.5	32	112	97	101.30	non-integer families
7	4	64	256	225	75.61	wrong α

Several distinct requirements converge on $g_{\text{car}} = 5$: integer families (g_{car} odd); three families $(g_{\text{car}}+1)/2 = 3$; the observed α^{-1} on the monotone curve; the SM structure $(b, s) = (3, 2)$, $\dim S^+ = 16$; and $\Omega_{\text{adm}} = 48$, $10b_1 = 41$. A clean algebraic shadow: $\frac{g(g+1)}{2} = 2^{g-1} - 1$ has $g = 5$ as its non-trivial solution ([C]).

10 The E_8 carrier atlas — stated unambiguously

Machine checks for this section: [\[verification/v5_e8_cascade.py\]](#)

In the series E_8 is a *downstream scale grammar*, not a primitive cause: the boundary kernel fixes α , the carrier rank, families and admissibility without E_8 upstream. Three levels.

10.1 Level 1 — original E_8 data of the series ([E])

$$|R(E_8)| = 248 - 8 = 240 = 5 \cdot 48 = 5 \Omega_{\text{adm}} = \text{lcm}(\Omega_{\text{adm}}, D_{\text{start}}) = \text{lcm}(48, 60), \quad (41)$$

$$D_{\text{start}} = g_{\text{car}} \cdot \dim \mathfrak{g}_{\text{SM}} = 5 \cdot 12 = 60, \quad \gcd(\Omega_{\text{adm}}, D_{\text{start}}) = 12 = \dim \mathfrak{g}_{\text{SM}}, \quad (42)$$

$$\frac{|R(E_8)|}{D_{\text{start}}} = \frac{240}{60} = 4 = \frac{\Omega_{\text{adm}}}{\dim \mathfrak{g}_{\text{SM}}}, \quad 2880 = D_{\text{start}} \Omega_{\text{adm}} = |R(E_8)| \cdot \dim \mathfrak{g}_{\text{SM}}, \quad (43)$$

$$\kappa_{E_8} = \frac{\gamma_{\text{car}}}{\ln(248/60)} = 0.58723, \quad X(n, r, I_1) = \frac{\bar{M}_{\text{Pl}}}{8} \sin^2 \theta_{13} \left(\frac{60 - 2n}{60} \right)^{\kappa_{E_8}} e^{-(8-r)/64} e^{-12\pi I_1}. \quad (44)$$

Coxeter/gauge typing of the lcm–gcd pair. Writing $D_{\text{start}} = 60 = 2h_{E_8}$ and $\Omega_{\text{adm}} = 48 = 6 \text{rank}(E_8)$, the relations become $\gcd(2h_{E_8}, 6r_{E_8}) = \dim \mathfrak{g}_{\text{SM}} = 12$ and $\text{lcm}(2h_{E_8}, 6r_{E_8}) = |R(E_8)| = 240$: the local SM algebra is the gcd of the Coxeter double-period and the $\mathbb{Z}_6$ rank-occupancy, and the E_8 root count is their lcm. Likewise the EW exponent reads $I_1^{\text{EW}} = \frac{41}{48} = \frac{h_{E_8}}{36} + \frac{1}{6r_{E_8}} = \frac{5}{6} + \frac{1}{48}$ (Coxeter principal part plus a rank singlet). [E] (arithmetic) / [C] (typing).

10.2 Level 2 — new exact counting identities ([E])

$$|R(E_8)| = 240 = 16 \cdot 15 = \dim S^+ (\dim S^+ - 1), \quad (45)$$

$$\dim E_8 = 248 = 8 \cdot 31 = (g_{\text{car}} + N_{\text{fam}})(2^{g_{\text{car}}} - 1), \quad (46)$$

$$|R(E_8)| = 8(31 - 1) = \text{rank}(E_8)(2^{g_{\text{car}}} - 2), \quad \text{rank}(E_8) = 8 = g_{\text{car}} + N_{\text{fam}} = 5 + 3, \quad (47)$$

$$10b_1 = 41 = \frac{|R(E_8)|}{6} + 1, \quad I_1^{\text{EW}} = \frac{41}{48} = \gamma_{\text{car}} + \frac{1}{\Omega_{\text{adm}}}, \quad (48)$$

with $31 = 2^{g_{\text{car}}} - 1$ the number of non-empty occupation words of the $[5, 4, 2]$ carrier code.

10.3 Level 2.5 — the even-flip transition atlas (a falsifiable program)

Identify S^+ with the even-weight binary code on five bits. A transition $x \rightarrow y \neq x$ has flip $F = x \oplus y$, even and nonempty, so $|F| \in \{2, 4\}$; per state there are $\binom{5}{2} + \binom{5}{4} = 10 + 5 = 15$ flips, giving

$$240 = 16 \cdot (10 + 5) = \underbrace{160}_{|F|=2} + \underbrace{80}_{|F|=4}, \quad (49)$$

and the 120 undirected flips match the 120 E_8 root *pairs* ([E], counting). This makes the atlas conjecture *testable*: a genuine root map must send each flip to a norm-2 vector in $\mathbb{R}^8$ with inner products in $\{-2, -1, 0, 1, 2\}$ reproducing the E_8 profile (per root: 1 self, 56 at +1, 126 orthogonal, 56 at -1, 1 anti).

Honest obstruction ([O]). The flip atlas splits $160 + 80$ by flip size; the standard E_8 (D_8) split is $112 + 128$. Since $160 + 80 \neq 112 + 128$, no flip-size-preserving map is an E_8 root system; a metrization, if it exists, must mix flip sizes. Until a norm-2, E_8 -profile embedding is exhibited, E_8 stays a *counting* atlas, not a derived root origin.

Sharper two-stage audit (Coxeter before Weyl). A full root-system isomorphism is hard; a much earlier, cheaper test is the Coxeter spectrum. *Stage A*: find a natural order-30 operator C_{TFPT} on the even-flip atlas (or its 8-dimensional primitive-character space) with characteristic

polynomial $\chi_{C_{\text{TFPT}}}(t) = \Phi_{30}(t)$. *Stage B (only if A holds)*: the norm-2, E_8 -profile, Weyl-closure metrization. Stage A can fail or carry far sooner than B, so it is the right first falsifier.

Stage A is closed on the primitive-character space (`[verification/v531_coxeter_tensor_stage_a.py]`).

The explicit integer operator

$$T_{30} = \text{Comp}(\Phi_5) \otimes \text{Comp}(\Phi_6)$$

(the 4×4 carrier clock of `v418` tensor the 2×2 family/CP hexagon) has *exact* order 30 ($T_{30}^{30} = I$, $T_{30}^k \neq I$ for $1 \leq k < 30$) and $\chi_{T_{30}} = \Phi_{30} = t^8 + t^7 - t^5 - t^4 - t^3 + t + 1$ exactly: by CRT ($\gcd(5, 6) = 1$) its eigenvalues are the products (primitive 5th) $\times$ (primitive 6th) = exactly the eight primitive 30th roots — the carrier clock times the family clock *is* the Coxeter clock, as one integer operator. Its traces are the Ramanujan sums, $\text{tr}(T_{30}^k) = c_{30}(k) = c_5(k) c_6(k)$ for all k , with the signed divisor table $(-1, 1, 2, 4, -2, -4, -8, 8)$ at $k = (1, 2, 3, 5, 6, 10, 15, 30)$ and the atom readout $(|\text{tr } T^3|, |\text{tr } T^5|, |\text{tr } T^{15}|) = (2, 4, 8) = (|\mathbb{Z}_2|, |\mu_4|, \text{rank } E_8)$ — eight integer traces type the clock *without* diagonalisation, a cheap kill test for any flip-operator candidate (honest scope: Φ_{15} passes the unsigned triple since $\Phi_{30}(t) = \Phi_{15}(-t)$; the *signed* table is the discriminator, and $\Phi_{16}, \Phi_{20}, \Phi_{24}$ die on the triple already). **[E]** *What stays open ([O])*: a canonical intertwiner from the concrete even-flip atlas onto this primitive space (`COX.FLIP.INTERTWINER`), and Stage B.

Audit only — non-load-bearing speculation: the dim 496 heterotic count [O]. *This box feeds no derivation; it is an arithmetic coincidence kept for completeness only, under the no-free-pattern rule.* The even half S^+ and odd half S^- of the exterior algebra each have dimension $2^{g_{\text{car}}-1} = 16$ and each carry $16 \cdot 15 = 240$ transitions. Together the full carrier ($2^{g_{\text{car}}} = 32$ states) gives

$$\dim S^+(2^{g_{\text{car}}} - 1) = 16 \cdot 31 = 496 = \dim(E_8 \times E_8) = \dim SO(32) = \binom{32}{2} \quad (\text{count}),$$

i.e. the two halves read as two E_8 root atlases, all $\binom{32}{2} = 496$ pairs as $SO(32)$ — the two anomaly-free heterotic gauge dimensions. The *dimension* equality is exact; that the carrier algebra *is* either gauge group is unproven and not used anywhere. **[O]** (audit-only). The same fingerprint recurs at checksum level: the dual products of the E_8 invariant degrees sum to $744 = 3 \cdot 248$ (the j -function constant), and the identical construction for D_{16} gives $1488 = 3 \cdot 496 = 2 \cdot 744$ (`[verification/v532_e8_degree_modular_checksum.py]`) — the checksum measures the common heterotic rank-16/Coxeter-30 structure, *not* E_8 uniqueness. **[E]** (audit-only).

10.4 Level 2.7 — the carrier Coxeter theorem (the strongest new bridge)

Use the universal root-system identity $|R| = \text{rank} \cdot h$ (Coxeter number h). With the carrier values $\text{rank}(E_8) = g_{\text{car}} + N_{\text{fam}} = 8$ and $|R(E_8)| = \dim S^+(\dim S^+ - 1) = 240$,

$$h_{E_8} = \frac{|R(E_8)|}{\text{rank}(E_8)} = \frac{240}{8} = 30 = N_{\text{fam}} \binom{g_{\text{car}}}{2} = N_{\text{fam}} A_{\Lambda} = 2g_{\text{car}} N_{\text{fam}} = 2^{g_{\text{car}}} - 2 \quad \text{[E]}. \quad (50)$$

The Coxeter number of E_8 is exactly three families times the cosmological pair action: each family contributes one Λ -decuplet ($A_{\Lambda} = \binom{5}{2} = 10$) to the Coxeter period. Equivalently the action ladder is a Coxeter ladder,

$$A_H = \frac{h_{E_8}}{2N_{\text{fam}}} = 5, \quad A_{\Lambda} = \frac{h_{E_8}}{N_{\text{fam}}} = 10, \quad (51)$$

so Hubble reads half a Coxeter period per family and Λ a full Coxeter period per family. **[C]**

Rank–Coxeter coupling. Equivalently $\dim S^+ = 2 \text{rank}(E_8) = 16$ and $h_{E_8} = 2(\dim S^+ - 1) = 2 \cdot 15 = 30$, so $\dim E_8 = (g_{\text{car}} + N_{\text{fam}})(h + 1) = 8 \cdot 31$. The two counting forms $16 \cdot 15 = 8 \cdot 30 = 240$ coincide because $\dim S^+ = 2r$. Dropping ν^c sends $\dim S^+ \rightarrow 15$, which breaks *both* $\dim S^+ = 2r$ and $\dim S^+(\dim S^+ - 1) = 240$: ν^c is the state that closes the rank–Coxeter coupling, not an optional singlet. **[E]/[C]**

10.5 Level 2.8 — positive-root moment: $|R^+(E_8)| = \text{Tr}_{S^+} X^2 = 5!$

The number of positive E_8 roots equals the master hypercharge moment:

$$\boxed{|R^+(E_8)| = 120 = \text{Tr}_{S^+} X^2 = 5!} \quad [\mathbf{E}], \quad (52)$$

so the second hypercharge moment of one family *is* the positive-root half-system. The whole integer chain then reads off $|R^+|$:

$$\sum_{f,j} L_{f,j} = \frac{|R^+(E_8)|}{3} = 40, \quad 10b_1 = 1 + \frac{|R^+(E_8)|}{3} = 41, \quad \delta_2 = 4! |R^+(E_8)| c_3^8. \quad (53)$$

The $\mathbb{Z}_6$ slice is then $|R(E_8)|/6 = \frac{|R^+|}{3} = 40 = \Omega_{\text{adm}}\gamma$: one $\mathbb{Z}_6$ phase slice of the E_8 root atlas is exactly the full flavor transport budget. $[\mathbf{E}]$ (arithmetic) / $[\mathbf{C}]$ (interpretation).

10.6 Level 2.9 — the 2·3·5 Coxeter cyclotomic compiler (meta-theorem)

The three hard carrier atoms — weak rank 2, color rank / family count 3, carrier rank $g_{\text{car}} = 5$ — generate the E_8 data *cyclotomically*, without taking E_8 as input. Set

$$\boxed{h := 2 \cdot 3 \cdot g_{\text{car}} = 30, \quad r := \varphi(h) = \varphi(30) = 8 = g_{\text{car}} + N_{\text{fam}},} \quad (54)$$

where h is the Coxeter number and $r = \varphi(2 \cdot 3 \cdot 5) = 8$ the rank. Then the standard $|R| = rh$, $\dim = r(h+1)$ give

$$|R(E_8)| = rh = 8 \cdot 30 = 240, \quad \dim E_8 = r(h+1) = 8 \cdot 31 = 248, \quad h+1 = 2^{g_{\text{car}}} - 1 = 31. \quad (55)$$

Moreover the E_8 exponents are exactly the primitive residues modulo $h = 30$:

$$\text{Exp}(E_8) = (\mathbb{Z}/30\mathbb{Z})^\times = \{1, 7, 11, 13, 17, 19, 23, 29\}, \quad \sum_{m \in (\mathbb{Z}/30\mathbb{Z})^\times} m = 120 = \text{Tr}_{S^+} X^2 = |R^+(E_8)|, \quad (56)$$

so the second hypercharge moment equals the sum of the E_8 exponents and the positive-root count. The $\mathbb{Z}_6 \times \mathbb{Z}_5 \simeq \mathbb{Z}_{30}$ tact has $\varphi(30) = 8$ primitive characters, whose phases $e^{2\pi i m/30}$ are the Coxeter eigenphases of E_8 ; its characteristic polynomial is the 30th cyclotomic polynomial

$$\chi_C(t) = \Phi_{30}(t) = t^8 + t^7 - t^5 - t^4 - t^3 + t + 1. \quad (57)$$

Reading: E_8 here is not primarily a root count but the *Coxeter cyclotomic readout of the 2, 3, 5 carrier*. $[\mathbf{E}]$ (arithmetic) / $[\mathbf{C}]$ (interpretation).

10.7 Level 3 — conjectural geometric readings ($[\mathbf{C}]$)

Conjecture 1 (Carrier transition atlas). $R(E_8) \cong \{(a, b) \in S^+ \times S^+ : a \neq b\}$ as a structure-bearing transition set, since $16 \cdot 15 = 240$. *Open*: a structure-preserving map (roots, lengths, Weyl data), not only the count.

Conjecture 2 (Least common refinement / rank-8 over the exhausted carrier). $240 = \text{lcm}(\Omega_{\text{adm}}, D_{\text{start}})$ reads E_8 as the least common refinement of the occupancy lattice and the $\mathbb{Z}_6$ phase lattice; $248 = (g_{\text{car}} + N_{\text{fam}})(2^{g_{\text{car}}} - 1)$ reads it as a rank-8 structure over the 31 non-empty carrier words.

Two routes to E_8 — keep their status separate. Levels 1–2 are exact arithmetic. There are now *two* distinct routes to E_8 , and they must not be conflated:

- **Glue route (closed, $[\mathbf{E}]$):** $E_8 = (D_5 \oplus A_3) + \mu_4\text{-glue}$ is built explicitly in Section 3 (disc(D_5) = disc(A_3) = $\mathbb{Z}_4$, glue index 4, 240 roots of norm 2, simple-root Gram det 1). This closes the former “blind 240-metrisation”.

- **Direct flip-atlas route (open, [O], optional bonus):** the even-flip transition set $240 = 160 + 80$ is *not* a flip-size-preserving E_8 map, since $160 + 80 \neq 112 + 128$. A direct structure-preserving metrisation of the flip graph remains open — a nice-to-have, not load-bearing, because the glue route already delivers E_8 .

So E_8 is closed via the glue; only the direct flip atlas is open.

11 $\mathbb{Z}_6$ monodromy, flavor and CP readouts

Machine checks for this section: `[verification/v88_cp_phase_audit.py]`

A single $\mathbb{Z}_6$ motif recurs in the quotient $G_{\text{phys}} = (SU(3) \times SU(2) \times U(1)_Y) / \mathbb{Z}_6$, the transport cusps $\{1, (2/3)^6, (1/3)^6\}$, $\delta_{\text{ph}} = z_\star^{1/6}$, and the CP phases

$$\delta_{\text{CKM}}^{(0)} = \frac{\pi}{3}, \quad \delta_{\text{CKM}} = \frac{\pi}{3} + 3\lambda_C^2 = 1.19823 \text{ rad}, \quad \delta_{\text{CP}}^{\text{PMNS}} = \frac{4\pi}{3} = 240^\circ. \quad (58)$$

From the documented inputs $s_{12} = \lambda_C$, $s_{23} = u/(1 + \lambda_C)$, $s_{13} = \lambda_C^3/3$ one gets the new derived readouts $|V_{td}| = 0.00908$, $|V_{ts}| = 0.04264$, $J_{\text{CKM}} = 3.33 \times 10^{-5}$. On the neutrino side, with $\sin^2 \theta_{13} = e^{-5/6}u = 0.02311$, $\sin^2 \theta_{23} = 0.4557$, $\delta_{\text{CP}} = 4\pi/3$ and the solar candidate $\sin^2 \theta_{12} = \frac{1}{3} - \frac{u}{2} = 0.30675$, a near-maximal $|J_{\text{CP}}^\nu| \approx 0.030$ (sharp readout for DUNE/Hyper-Kamiokande). [C]

11.1 CP as triple seed variance; a closed observable quadrilateral

The Cabibbo angle is the Bernoulli width of the seed. The *exact* CKM phase is the family transport holonomy (document 3), $\delta_{\text{CKM}} = \arg \text{Tr} \Pi_\Delta(\nabla_F^*)$; on the small-area branch this has the compact readout form (with an $O(\lambda_C^4)$ remainder)

$$\lambda_C = \sqrt{u(1-u)}, \quad \boxed{\delta_{\text{CKM}} - \frac{\pi}{3} = N_{\text{fam}} u(1-u) = 3\lambda_C^2} \quad [\text{E}]/[\text{C}] \text{ (readout; exact = holonomy)}. \quad (59)$$

The coefficient 3 is the family number N_{fam} : weak CP is the $\mathbb{Z}_6$ base angle $\pi/3$ plus N_{fam} times the seed (Bernoulli) variance. The $\mathbb{Z}_6$ base angle now has an A_3 root: since $\mathbb{Z}_6 = |R^+(A_3)| = 6$ (the positive-root hexagon of the family algebra), the CP phases are integer steps of the A_3 holonomy quantum $2\pi/|R^+(A_3)|$,

$$\boxed{\delta_{\text{CKM}} = \frac{2\pi}{|R^+(A_3)|} + N_{\text{fam}} u(1-u), \quad \delta_{\text{CP}}^{\text{PMNS}} = 4 \cdot \frac{2\pi}{|R^+(A_3)|} = \frac{4\pi}{3}} \quad [\text{E}]/[\text{C}]. \quad (60)$$

So weak CP = (A_3 positive-root phase step) + (family count $\times$ seed variance), while strong CP is the *determinant* sector, erased. The solar angle reads as a carrier root plus a half-seed, $\sin^2 \theta_{12} = \frac{1}{3} - \frac{u}{2} = -y_- - \frac{u}{2}$ (since $-y_- = \frac{1}{3}$), so PMNS = carrier root + seed + A_3 holonomy phase. [C] Strong CP is killed by the determinant selector ($\arg \det M_u = \arg \det M_d = 0$, $\bar{\theta} = 0$), while weak CP survives as the seed-deformed $\mathbb{Z}_6$ holonomy: CP is not forbidden, only *determinant* CP is. [C] — the strong-CP closure lives on the admissible QFT branch and is *conditional* on the analytic hypotheses of the QFT closure (document 3, Part III: reflection positivity, temperedness, clustering, mass gap, OS reconstruction); its status is therefore weaker than the discrete lattice identities, even though the semantics “determinant CP forbidden, holonomy CP allowed” is robust. Since $u = \Omega_b + \beta_{\text{rad}}$ exactly, eliminating the seed closes the four UV-shadow observables into one cycle:

$$\lambda_C^2 = (\Omega_b + \beta_{\text{rad}})(1 - \Omega_b - \beta_{\text{rad}}), \quad \sin^2 \theta_{13} = e^{-5/6}(\Omega_b + \beta_{\text{rad}}), \quad \Omega_b = \left(1 - \frac{1}{4\pi}\right) e^{5/6} \sin^2 \theta_{13}. \quad (61)$$

One measured member of $\{\lambda_C, \theta_{13}, \Omega_b, \beta_{\text{rad}}\}$ fixes the other three. [E] (the four rows remain operationally distinct readout classes).

12 Mass hierarchy as a hexagon with one extra turn

Machine checks for this section: `[verification/v18_quark_yukawa.py]` `[verification/v20_lepton_c_derivation.py]`

The transport length packets are

$$L_u = (7, 3, 0), \quad L_d = (7, 5, 2), \quad L_e = (8, 5, 3), \quad \sum_{f,j} L_{f,j} = 40 = \Omega_{\text{adm}} \gamma_{\text{car}} = 10b_1 - 1. \quad (62)$$

Modulo 6 they lie on related triples of the C_6 hexagon; the first generation carries one extra full turn ($u, d : 1 + 6; e : 2 + 6$), costing $\lambda_Y^6 = (u(1-u))^3 \approx 1.28 \times 10^{-4}$ and explaining why the first generation is so light — “one turn more”, not a free fit. **[C]** *Flagged proximity ([O], not exact)*: the full-turn cost is close to the topological seam defect, $\lambda_C^6 / \delta_{\text{top}} = 1.061$ ($\sim 6\%$), hinting that one flavor winding $\approx$ one seam-defect unit $\delta_{\text{top}} = \Omega_{\text{adm}} c_3^4$ — a structural smell, not a theorem. Writing the length matrix $L = R + W$ with $R = L \bmod 6$ the residue shadow and W the full-winding part,

$$L = \begin{pmatrix} 7 & 3 & 0 \\ 7 & 5 & 2 \\ 8 & 5 & 3 \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 3 & 0 \\ 1 & 5 & 2 \\ 2 & 5 & 3 \end{pmatrix}}_R + 6 \underbrace{\begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}}_W, \quad \sum R = 22, \quad \sum W = 18 = 3 \cdot 6, \quad \sum L = 40,$$

and the column sums are $(22, 13, 5)$ with $S_1 = 22 = \sum R$ and $S_2 + S_3 = 18 = \sum W$. So the first generation is light because it stores the *entire* residue budget plus the three full C_6 windings; generations 2 and 3 are the winding complement. **[E]** (arithmetic).

12.1 Lepton flavor decouple: the charged-lepton determinant carries u^{10}

The source masses on the v_{geo} branch are $(\hat{m}_e, \hat{m}_\mu, \hat{m}_\tau) = \frac{v_{\text{geo}}\pi}{\sqrt{2}} (\frac{16}{7}u^5, \frac{4}{3}u^3, \frac{7}{6}u^2)$. Their product is a pure decouple in the seed:

$$\boxed{\frac{\hat{m}_e \hat{m}_\mu \hat{m}_\tau}{(v_{\text{geo}}\pi/\sqrt{2})^3} = \left(\frac{16}{7} \cdot \frac{4}{3} \cdot \frac{7}{6}\right) u^{5+3+2} = \frac{32}{9} u^{10}} \quad \text{[E]}, \quad (63)$$

with $5 + 3 + 2 = g_{\text{car}} + b + s = 10 = \binom{5}{2}$. So the charged-lepton Yukawa powers $(n_e, n_\mu, n_\tau) = (5, 3, 2)$ are the three carrier numbers (g_{car}, b, s) , and the lepton determinant carries the same decouple (u^{10}) that the cosmological constant carries in the gauge sector ($A_\Lambda = 10$): Λ and the charged-lepton determinant are two realisations of the same degree- $\binom{5}{2}$ sector.

Sum–product reading. The lepton powers $(5, 3, 2)$ encode *both* decouple numbers at once: their sum is the Λ action, their product is the E_8 Coxeter number,

$$5 + 3 + 2 = 10 = A_\Lambda, \quad 5 \cdot 3 \cdot 2 = 30 = h_{E_8} = 2 \cdot 3 \cdot 5, \quad (64)$$

and $(5, 3, 2)$ is the prime factorisation of $h_{E_8} = 30$. So $\det M_\ell$, ρ_Λ and the E_8 Coxeter period read the same carrier core additively ($= 10$) and multiplicatively ($= 30$). **[E]/[C]**

12.2 Why the top Yukawa is ≈ 1 : the $L = 0$ anchor (Froggatt–Nielsen ladder)

The transport length L_f is a Froggatt–Nielsen-type charge with expansion parameter λ_C :

$$\boxed{y_f \simeq \lambda_C^{L_f} \times \Lambda_f \quad (\Lambda_f = \mathcal{O}(1)), \quad y_t \simeq 1 \text{ because } L_t = 0} \quad \text{[E]/[C]}. \quad (65)$$

The top sits at $L_t = 0$ (the *unwound* carrier direction), so $y_t = \Lambda_t \simeq 1$ ($\sqrt{2} m_t / v = 0.991$); every other charged fermion is suppressed by integer powers of $\lambda_C = 0.2244$. Numerically (with $\mathcal{O}(1)$ sector prefactors $\Lambda_f \in [0.43, 1.07]$):

f	t	b	τ	c	μ	s	u, d	e
L_f	0	2	3	3	5	5	7	8
$y_f/\lambda_C^{L_f}$	0.99	0.48	0.90	0.65	1.07	0.94	$\sim 0.4\text{--}0.9$	0.46

So the entire charged-fermion hierarchy is one λ_C -ladder anchored by the top at $L = 0$; the lightness of the first generation ($L = 7, 8$) and the heaviness of the top ($L = 0$) are the two ends of the same integer winding spectrum. This is the standard Froggatt–Nielsen mechanism with λ_C as the flavon ratio and L_f as the FN charge — a clean bridge between TFPT transport and an established flavor mechanism.

13 Inflation interface

Machine checks for this section: [\[verification/v7_gravity_cosmo.py\]](#) [\[verification/v86_nstar_reheating.py\]](#) [\[verification/v494_cosmo_running_mu_record.py\]](#)

The documented scalaron is Starobinsky R^2 ; with the pivot $N_\star = 57.1$:

$$n_s = 1 - \frac{2}{N_\star} = 0.96497, \quad r = \frac{12}{N_\star^2} = 0.00368, \quad (66)$$

$$\frac{M_{\text{scal}}}{\bar{M}_{\text{Pl}}} = \sqrt{8\pi} c_3^4 = c_3^{7/2} = (8\pi)^{-7/2} \Rightarrow M_{\text{scal}} \approx 3.06 \times 10^{13} \text{ GeV}.$$

n_s matches Planck; $r = 0.0037$ is the falsifiable target for LiteBIRD/CMB-S4.

Amplitude (status-clean). The scalaron mass is a pure seam power,

$$\boxed{\frac{M_{\text{scal}}^2}{\bar{M}_{\text{Pl}}^2} = c_3^{r-1} = c_3^{\Omega_{\text{adm}} - 10b_1} = c_3^7} \quad [\mathbf{E}], \quad (67)$$

an *exact* identity (the exponent $7 = 48 - 41$ is the post-flavor, post-Higgs occupancy deficit). The cosmology interface reduces the geometric gravity action to an FRW R^2 /scalaron branch in which M_{scal} is the Starobinsky mass M ; the identification is therefore derived in that sector, not posited. With the standard large- N amplitude $A_s = \frac{N_\star^2}{24\pi^2} (M/\bar{M}_{\text{Pl}})^2$ this gives

$$A_s = \frac{N_\star^2 c_3^7}{24\pi^2} = \frac{N_\star^2 \delta_2}{8640\pi} = 2.17 \times 10^{-9} \quad [\mathbf{C}]/[\mathbf{O}], \quad (68)$$

matching Planck ($\simeq 2.1 \times 10^{-9}$); $\delta_2 = 2880 c_3^8$ is the second seam defect. The seam-power form is exact [\[E\]](#); the amplitude is then determined *within* the R^2 sector, the only residual being the geometric-gravity (Hodge) closure on which that sector rests [\[C\]](#). *Convention note:* $A_s \propto N_\star^2$, so the quoted 2.17×10^{-9} uses the pivot $N_\star = 57.1$; the alternative seed pivot $N_\star = 3/u - 1 = 55.4$ gives 2.05×10^{-9} . A number should only be quoted with $N_\star$ fixed.

Running (consistency relation). The same attractor also fixes the *running* of the tilt parameter-free: $\alpha_s = \text{d}n_s/\text{d}\ln k = -2/N_\star^2 = -r/6$ and $\beta_s = -4/N_\star^3$ at leading order; the exact slow roll on the Starobinsky potential gives $\alpha_s = -7.09 \times 10^{-4}$, $\beta_s = -2.70 \times 10^{-5}$ at the reheating point $N_\star = 51.4$ (leading-order band $[-8.0, -5.6] \times 10^{-4}$ over $N_\star \in [50, 60]$). Current data are consistent (Planck 2018 -0.0045 ± 0.0067 : $+0.57\sigma$; P-ACT-LB $+0.0062 \pm 0.0052$: -1.33σ), and since plateau potentials give $\alpha_s < 0$, any robust *positive* running would kill the branch. [\[C\]](#) [\[verification/v494_cosmo_running_mu_record.py\]](#)

14 Further simple bridges

Machine checks for this section: [\[verification/v74_compiler_micro_lemmas.py\]](#)

- **Dark-energy equation of state $w = -1$ (exact).** The TFPT Λ is a genuine vacuum constant $\rho_\Lambda = \frac{3}{4\pi^2} e^{-2\alpha_\star^{-1}} \bar{M}_{\text{Pl}}^4$, not a rolling field, so $w = -1$ identically (no quintessence). A robust measurement $w \neq -1$ would falsify the Λ -branch. [\[E\]](#) (structural).

- **Effective neutrino number** $N_{\text{eff}} \simeq 3$. Three families ($N_{\text{fam}} = 3$) give three light neutrinos and the standard $N_{\text{eff}} \simeq 3.04$. [E].
- **Top Yukawa** $\simeq 1$ as the $L = 0$ anchor (above): a bridge to the Froggatt–Nielsen flavor mechanism.
- *Flagged (not clean)*: $m_H/v = 0.509$ is near $\frac{1}{2}$ (1.7%); $m_W/m_Z = 0.881$ matches the running $\sqrt{1 - \sin^2 \theta_W(M_Z)}$ (not the tree $\sqrt{5/8} = 0.79$). No clean closed bridge for the Higgs/ W/Z masses yet. [O]

15 Closures that the architecture rests on

Machine checks for this section: [verification/v36_spectral_action_g2.py] [verification/v4_flavor_matrix.py]

Five results close or sharpen the discrete core. They are *proved in this document set* — the φ_0^{ret} -ladder, the H2 realisation and the strong-CP/dynamics gap in document 2 (tfpt_2_standard_model), the $E_6 \times A_2$ shadow in document 3 (tfpt_3_e8_audit_bootstrap) — and are collected here because the architecture above depends on them.

1. **The φ_0^{ret} -ladder (whole spectrum, one formula).** Every fermion mass is $\hat{m}_{f,j} = \frac{v_{\text{geo}}}{\sqrt{2}} \lambda_Y^{L_{f,j}} \Lambda_{f,j}$ with $\lambda_Y = \sqrt{\varphi_0^{\text{ret}}(1 - \varphi_0^{\text{ret}})}$, and the closed evaluation gives the universal form $\lambda_Y^L \Lambda = \pi c(\varphi_0^{\text{ret}})^k$ — the nine Λ are hexagonal-resolvent outputs (not free numbers); the *charged-lepton* ones are exact, the *quark* ones are reduced to the (U_{wall}) selector. The φ_0^{ret} -order ladder is $t(0) > \{b, c, \tau\}(2) > \{s, \mu\}(3) > \{u, d\}(4) > e(5)$. [E] (leptons exact, quark ratios closed v49/v71) / [O] (absolute quark scale)
2. **H2 reduced via the Mehta–Seshadri framework.** Mehta–Seshadri supplies the equivalence (stable parabolic $\leftrightarrow$ irreducible unitary); the TFPT-specific step — “stable = transport-minimal” and the geodesic C_6 word length — is conditional on the unitarity (U) of ∇_F^* (document 3) and the parabolic-degree $\rightarrow C_6$ dictionary. H2 is the sharp realisation statement: the D_4 -equivariant stable structure on $\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$ realises the branch with $\det R = h(D_5) = 8$, principal 2-minors (2, 3, 5). [E] (given (U))
3. **Strong CP / dynamics via the explicit gap.** $\theta_{\text{eff}} = 0$ needs only γ_5 -Hermiticity + polar mass + sheet involution + reflection positivity (no mass gap). The full dynamics closes on the physical sector because the C_6 transport transfer operator has the *explicit* spectrum $\{(k/3)^6\}$, giving the mass gap $\Delta = 6 \log \frac{3}{2} = 2.43$ in closed form $\Rightarrow$ OS reconstruction $\Rightarrow$ measure on the admissible sector. [E] (phys. sector)
4. **$E_6 \times A_2$ flavor shadow.** The residue matrix sits inside an E_8 secondary branching: $248 = \|R\|_F^2 + \det R + 2(1^\top R a) N_{\text{fam}} = 78 + 8 + 2 \cdot 27 \cdot 3$, with $\|R\|_F^2 = 78 = \dim E_6$, $\det R = 8 = \dim A_2$. The seven E_8 slices form an audit raster (atlas note). [O]
5. **SM completeness.** The discrete structure and dimensionless core are closed; the dimensionful EW/QCD masses ($m_W, m_Z, m_H, \sin^2 \theta_W, \alpha_s$) are schema-level via $\mathcal{R}_{\text{SM}}$, with m_H resting on the closed UV quartic $\lambda_\Phi = 1/(16\pi^2)$ (near-criticality). The *exact* solar angle θ_{12} (leading $1/N_{\text{fam}} = \frac{1}{3}$) is *conditionally derived*, pending the seam-misalignment lemma — the sharpest remaining SM tension point. [C]

16 Status summary and open audit points

Machine checks for this section: [verification/v22_open_gates_audit.py]

Statement	Status	Note
Dual-root dictionary (Y , cusps, γ_{car} , φ_{base})	[E]	exact, from $\{1/2, -1/3\}$

$10b_1 = 5^2 + 4^2 = \Omega_{\text{adm}}\gamma + 1$	[E]	Pythagorean carrier identity
$I_1^{\text{EW}} = 41/48 = \text{Var}(Y) b_1$	[E]	exact
$\delta_2 = 4! \cdot 5! c_3^8, \text{Tr } X^2 = 5!$	[E]	factorial traces
$5/4 = g_{\text{car}}/(g_{\text{car}} - 1)$ universal compression	[E]	exact
$u = \frac{4}{3}c_3 + 48c_3^4$; one-decoder identities	[E]	exact
$\xi = \frac{3}{4}/(1 + 9/128\pi^3)$	[E]	corrected tree value
Ladder $A_{\text{EW}} : \mathcal{A}_H : \mathcal{A}_\Lambda = 1 : g_{\text{car}} : 2g_{\text{car}}$	[E]/[C]	Hubble rung is a corollary
Ladder $= \binom{g_{\text{car}}}{0} : \binom{g_{\text{car}}}{1} : \binom{g_{\text{car}}}{2}$; $2g_{\text{car}} = \binom{g_{\text{car}}}{2} \Leftrightarrow g_{\text{car}} = 5$	[E]	new overdetermination of $g_{\text{car}} = 5$
2D action lattice $S_x = a(\alpha_\star^{-1}/g_{\text{car}}) + bL_0$	[E]/[C]	two integer labels per scale
Decuple bridge $\rho_\Lambda/\bar{M}_{\text{Pl}}^4 = \frac{3}{4\pi^2}e^{2\delta_{\text{ph}}}[v/(5\beta_{\text{rad}}^2\bar{M}_{\text{Pl}})]^{10}$	[E]	exact, $10 = 2g_{\text{car}} = \dim \Lambda^2 E$
Family = ladder = Pascal row $\{1, 5, 10\}$; $16 = 1 + 5 + 10$	[E]	central new identification
$g_{\text{car}} = 5$ via $2^{g_{\text{car}}-1} = \binom{g_{\text{car}}}{0} + \binom{g_{\text{car}}}{1} + \binom{g_{\text{car}}}{2}$	[E]	Pascal closure (unique)
$\text{Tr}_{S^+} X^2 = 5!$ generates 40, 41, 240, δ_2	[E]	master moment
$ R(E_8) = 16 \cdot 5 \cdot 3$; $\dim E_8 = 240 + 8$	[E]	E_8 compression
Even-flip atlas $240 = 160 + 80$	[E]/[O]	metrization open ($\neq 112 + 128$); Coxeter Stage A closed on the character space (v531)
$\delta_{\text{CKM}} - \pi/3 = 3u(1 - u)$	[E]	CP = triple seed variance
Observable quadrilateral ($u = \Omega_b + \beta_{\text{rad}}$)	[E]	closed UV-shadow cycle
Power tower $v=x^1, H=x^5 R_H, \Lambda=x^{10} R_\Lambda$	[E]	K_5 vertices/edges
$496 = 16 \cdot 31 = \dim(E_8 \times E_8) = \dim SO(32)$	[E]/[C]	heterotic-dimension count
$A_{\text{EW}} = \alpha_\star^{-1}/\sqrt{\Delta_Y}, \Delta_Y = (b+s)^2 = 25$	[E]	discriminant action theorem
$\hat{m}_e \hat{m}_\mu \hat{m}_\tau \propto \frac{32}{9}u^{10}$; $(5, 3, 2) = (g_{\text{car}}, b, s)$	[E]	lepton flavor decuple
$M_{\text{scal}}/\bar{M}_{\text{Pl}} = c_3^{7/2}$	[E]	scalaron seam power
$\delta_{\text{CKM}} - \pi/3 = N_{\text{fam}} u(1 - u)$	[E]	weak CP = $N_{\text{fam}} \times$ seed variance
$L = R + W$: $\sum R = 22, \sum W = 18$	[E]	first generation = residue store
$ R(E_8) /6 = \sum L = 40$ ($\mathbb{Z}_6$ slice)	[E]	one $\mathbb{Z}_6$ slice = flavor budget
$h_{E_8} = R /\text{rank} = 30 = N_{\text{fam}} \binom{5}{2} = N_{\text{fam}} A_\Lambda$	[E]/[C]	carrier Coxeter theorem
$ R^+(E_8) = \text{Tr}_{S^+} X^2 = 120 = 5!$	[E]	positive-root moment
common decuple $\binom{5}{2} = A_\Lambda = u^{5+3+2} = h_{E_8}/N_{\text{fam}} = 10$	[E]/[C]	one carrier pair-decuple
$\lambda_C^6/\delta_{\text{top}} = 1.061$ (winding $\approx$ seam defect)	[O]	flagged proximity (not exact)
$y_f \simeq \lambda_C^{L_f}$, top $L_t = 0 \Rightarrow y_t \simeq 1$	[E]/[C]	Froggatt–Nielsen ladder
$w = -1$ (true Λ); $N_{\text{eff}} \simeq 3$	[E]	dark energy / neutrino species
3-generator lattice $\{\alpha_\star^{-1}, L_0, L_c\}$; $M_{\text{scal}} = e^{-\frac{7}{2}L_c}$	[E]	seam axis $L_c = \ln(1/c_3)$
$\dim S^+ = 2 \text{rank}(E_8), h_{E_8} = 2(\dim S^+ - 1) = 2^g - 2$	[E]	rank–Coxeter coupling (ν^c closes it)
$\gcd(2h, 6r) = \dim \mathfrak{g}_{\text{SM}} = 12, \text{lcm} = 240$	[E]/[C]	SM algebra = gcd; E_8 count = lcm
$I_1^{\text{EW}} = h_{E_8}/36 + 1/(6r) = 41/48$	[E]/[C]	Coxeter part + rank singlet
$M_{\text{scal}}^2/\bar{M}_{\text{Pl}}^2 = c_3^{r-1} = c_3^7$ exact; $A_s = N_\star^2 c_3^7/24\pi^2$	[E]/[C]/[O]	seam power exact; A_s conditional (Starobinsky)
$h_{E_8} = 2 \cdot 3 \cdot 5 = 30, r = \varphi(30) = 8$	[E]/[C]	2, 3, 5 Coxeter cyclotomic compiler
$\text{Exp}(E_8) = (\mathbb{Z}/30)^\times, \sum = 120, \chi_C = \Phi_{30}$	[E]/[C]	exponents = primitive residues
lepton $(5, 3, 2)$: $\sum = 10 = A_\Lambda, \prod = 30 = h_{E_8}$	[E]/[C]	sum=action, product=Coxeter
Hubble $H_0/\bar{M}_{\text{Pl}} = e^{-\alpha_\star^{-1}}/(2\pi\sqrt{\Omega_\Lambda})$	[E]	exact corollary
Λ -anchored closure $G_N = \frac{3c_3^3}{32\pi^3 \hbar \Lambda_{\text{geom}}} e^{-2\alpha_\star^{-1}}$	[E]/[C]	G_N to -0.35%
$g_{\text{car}} = 5$ overdetermined	[C]	monotone $\alpha^{-1}(g_{\text{car}})$
$ R(E_8) = 240 = 16 \cdot 15, 248 = 8 \cdot 31$	[E]	counting identities
$E_8 \leftrightarrow$ carrier-transition map	[C]	needs structure map
$\delta_{\text{CP}} = 4\pi/3, \theta_{12} = \frac{1}{3} - \frac{u}{2}$	[C]	PMNS readouts
$ V_{td} , V_{ts} , J_{\text{CKM}}, J_{\text{CP}}^\nu; n_s, r$	[C]	derived readouts

Open audit points: (1) *discharged* — the EM closure $F_{U(1)}(\alpha) = 0$ is now stated explicitly (Grammar III), with existence/uniqueness and root $\alpha_\star^{-1} = 137.0359992168$ (2.9×10^{-10} , 1.9σ vs CODATA-2022); (2) keep $\gamma_{\text{car}} = 5/6$ vs γ_E distinct; (3) type M_{Pl} vs $\bar{M}_{\text{Pl}}$ and “pure” ($\mathcal{A}_\Lambda = 274.07$) vs “total” ($-\ln(\delta_{\text{top}} e^{-2\alpha_\star^{-1}}) = 283.10$, $-\ln(\rho_\Lambda/\bar{M}_{\text{Pl}}^4) = 276.65$) actions consistently; (4) version the Λ prefactor branch; (5) the scalaron-mass $\rightarrow$ Starobinsky- M identification is the *only* condition for $A_s = N_\star^2 c_3^7/(24\pi^2)$ (the seam power $M_{\text{scal}}^2/\bar{M}_{\text{Pl}}^2 = c_3^7$ itself is exact); (6) the E_8 *lattice* is now built explicitly as $(D_5 \oplus A_3) + \mu_4$ -glue (Section 3), so the former “blind 240-metrisation” is closed. The flavor status (companion parabolic-weight note) now splits cleanly: the *anchor multiset* $\{1, 1, 2\}$ is derived (Schur–Horn + stability) [E]; the *sector assignment* $(1, 1, 2)$ follows from the D_5 cusp ordering [E]/[C]; the *first-generation* +6 *winding* is the hierarchy convention [C]; the six *non-anchor residues* are then forced (residue sets = D_6 orbit of $\{0, 1, 3\}$, minus anchor, monotone) [E]. The remaining geometric input at the time of writing was (H2), the specific D_6 cusp labelling on $\mathbb{P}^1 \setminus \mu_4$ — sharply characterised by the residue-matrix spectral selector ($\det R = h(D_5)$, principal minors $(2, 3, 5)$), with a naive cusp-position shortcut explicitly ruled out [C]. The explicit Riemann–Hilbert construction confirms the rigid $SU(3)$ system and its A_3 Coxeter rotation. *Status update:* the H2 dictionary has since been made *exact* (hexagon = sign-twisted family spectrum, addresses pinned, margins forced, v118–v122), the deck is *derived* (v141), and the residue is the $R4'$ selector establishment — two line pairings (v139/v142); what stays geometric is only the parabolic realisation premise (GATE.QGEO, research contracts).

Honest negatives (reported to keep the discipline visible). The TFPT lepton ladder gives a Koide ratio $Q = (\sum m_\ell)/(\sum \sqrt{m_\ell})^2 = 0.6645$, close to but *not* the empirical $2/3 = 0.66667$ (-0.33%); TFPT is Koide-*like* but does not reproduce the precise relation. The baryon asymmetry $\eta_B \approx 6 \times 10^{-10}$ and the proton/electron ratio $m_p/m_e = 1836$ do *not* fall on a clean action charge or L_0 ladder; they remain outside the present grammar. These are stated so the strong items are not read as a uniform success.

17 Consolidated picture

The three nested grammars.

Carrier: $3 + 2 \Rightarrow \{y_-, y_+\} = \{-\frac{1}{3}, \frac{1}{2}\} \Rightarrow Y$, $\dim S^+ = 16$, $N_{\text{fam}} = 3$, $\Omega_{\text{adm}} = 48$, $b_1 = \frac{41}{10}$,

Seed: $u = \frac{4}{3}c_3 + 48c_3^4 \Rightarrow \{\lambda_C, \beta_{\text{rad}}, \Omega_b, \sin^2 \theta_{13}\}$,

Action: $\alpha_\star^{-1} \Rightarrow A_{\text{EW}} = \frac{\alpha_\star^{-1}}{g_{\text{car}}}$, $\mathcal{A}_\Lambda = 2\alpha_\star^{-1}$, $\mathcal{A}_H = \alpha_\star^{-1} \Rightarrow \frac{\rho_\Lambda}{\bar{M}_{\text{Pl}}^4} = \frac{3}{4\pi^2} e^{2\delta_{\text{ph}}} \left[\frac{v_{\text{geo}}}{5\beta_{\text{rad}}^2 \bar{M}_{\text{Pl}}} \right]^{10}$.

A small grammar generates many sectors: the Standard-Model packet, flavor, the electroweak hierarchy, the cosmological constant, the metrology, and the inflation/CMB interface. The strongest new statements are exact: the decouple bridge ties Λ to the tenth power of the stripped electroweak scale with $10 = 2g_{\text{car}}$; the Hubble scale is the square root of Λ ; and the Einstein normaliser is the corrected tree value $\frac{3}{4}/(1 + 9/128\pi^3)$. None of this is claimed as a proof of a final theory; it is offered as a tightening of the existing architecture for your assessment.

The compiler in one line: TFPT = even carrier code + seed decoder + 2·3·5 Coxeter compiler. From the three atoms $(2, 3, g_{\text{car}}=5)$:

$$h_{E_8} = 2 \cdot 3 \cdot 5 = 30, \quad r_{E_8} = \varphi(30) = 8 = g_{\text{car}} + N_{\text{fam}},$$

$$|R(E_8)| = rh = 240, \quad \dim E_8 = r(h+1) = 8 \cdot 31 = 248,$$

$$\text{Exp}(E_8) = (\mathbb{Z}/30)^\times, \quad \sum \text{Exp} = 120 = \text{Tr}_{S^+} X^2 = |R^+|, \quad \chi_C = \Phi_{30},$$

$$\dim S^+ = 1 + 10 + 5 = 16 = 2r, \quad A_{\text{EW}} : \mathcal{A}_H : \mathcal{A}_\Lambda = 1:5:10, \quad \mathcal{A}_H = \frac{h}{6}, \quad \mathcal{A}_\Lambda = \frac{h}{3}, \quad N_{\text{fam}} = \frac{h}{A_\Lambda} = 3,$$

$$\sum L = \frac{|R^+|}{3} = 40, \quad 10b_1 = 1 + \frac{|R^+|}{3} = 41 = 1 + h \frac{\dim S^+}{\dim \mathfrak{g}_{\text{SM}}},$$

$$\det M_\ell \propto u^{10}, \quad \frac{\rho_\Lambda}{M_{\text{Pl}}^4} \propto \left[\frac{v_{\text{geo}}}{5\beta_{\text{rad}}^2 M_{\text{Pl}}} \right]^{10}, \quad A_{\text{EW}} = \frac{\alpha_\star^{-1}}{\sqrt{\Delta_Y}}, \quad \sqrt{\Delta_Y} = g_{\text{car}} = 5.$$

The common decuple (= 10): $\binom{g_{\text{car}}}{2}$ (carrier pairs) = A_Λ (cosmological action) = u^{5+3+2} (lepton determinant) = h_{E_8}/N_{fam} (Coxeter per family). Λ , the charged-lepton determinant, the E_8 Coxeter structure and the action ladder all read the *same* carrier pair-decuple. The open core is one question: can the even-flip atlas be metrised as a genuine E_8 root system (the 160+80 vs 112+128 obstruction)? Its Coxeter half (Stage A) is closed on the primitive-character space by the tensor clock $T_{30} = \text{Comp}(\Phi_5) \otimes \text{Comp}(\Phi_6)$ (**v531**); the atlas intertwiner and the metrization (Stage B) remain.

Part II

The Coxeter–cyclotomic compiler and the explicit E_8 construction

Abstract

This short companion compresses the TFPT bridge layer to a single discrete object. The claim is that the Standard-Model family, the cosmological action ladder, the E_8 counting data, the flavor budget, inflation and the cosmological constant are *not* independent coincidences but projections of one structure: the even-weight code C^+ on $g_{\text{car}} = 5$ slots, together with its $\mathbb{Z}_{30}$ Coxeter spectrum and the single seam seed $c_3 = 1/(8\pi)$. The decisive new result is that C^+ is precisely the half-spin weight orbit of $D_5 = \mathfrak{so}_{10}$ ($\text{Aut}(C^+) = W(D_5)$, order 1920), that the family geometry $\mathbb{P}^1 \setminus \mu_4$ is an $A_3 = \mathfrak{su}_4$, and that E_8 is then the standard *closure algebra* of $D_5 \times A_3$ — not an input. The fusion is exact: $\text{disc}(D_5) = \text{disc}(A_3) = \mathbb{Z}_4$, so the four family corners μ_4 are precisely the glue code that closes $D_5 \oplus A_3$ into the unimodular E_8 lattice. The glue is metric: the discriminant-form norms $q(D_5) = \frac{5}{4}$ and $q(A_3) = \frac{3}{4}$ sum to the E_8 root norm 2 — and these are exactly the pre-existing TFPT constants $\delta_2/\delta_{\text{top}}^2 = \frac{5}{4}$ and $\xi_{\text{tree}} = \frac{3}{4}$. The A_3 side is explicit: the residue pairing on $H^1(\mathbb{P}^1 \setminus \mu_4)$ is the A_3 Cartan form, and the carrier corner rotation $z \mapsto iz$ acts as the A_3 Coxeter element (order 4, spectrum $\{i, -1, -i\}$). The lift is no longer a count: the 240 E_8 roots are *explicitly* built from $D_5 \oplus A_3$ plus the spinor (μ_4) glue, all of norm 2, Cartan determinant 1. Seven statements carry it; every lattice identity below is checked exactly, and the analytic metric is computed and characterised (a D_4 -equivariant deformation), so no *lattice* structural step is left open in the $D_5 \times A_3$ glue route (the upstream boundary kernel, EM closure and flavor Hecke saturation remain separate open problems). Master statement: *five carrier slots give D_5 , four family corners give A_3 , and μ_4 glues $D_5 \times A_3$ into the E_8 atlas.*

The compression.

$$\begin{aligned} \text{TFPT} &= \underbrace{D_5 \text{ half-spin carrier}}_{C^+} + \underbrace{A_3 \text{ four-point family}}_{\mathbb{P}^1 \setminus \mu_4} + c_3 \text{ seam seed}, \\ E_8 &= \text{closure of } D_5 \times A_3 \end{aligned}$$

with $c_3 = \frac{1}{8\pi}$, $g_{\text{car}} = 5$, $h := 2 \cdot 3 \cdot g_{\text{car}} = 30$, $r := \varphi(h) = \varphi(30) = 8$, $N_{\text{fam}} = 3$, $N_{\Phi} = 1$, $\dim \mathfrak{g}_{\text{SM}} = 12$. Equivalently $C^+ + \mathbb{Z}_{30}$ Coxeter compiler + c_3 seed; the $D_5 \times A_3$ reading is the structural lift of the same object.

Theorem 1 — the even carrier code

Let E be the rank- $g_{\text{car}} = 5$ carrier, split $E = E_3 \oplus E_2$ (color $b = 3$, weak $s = 2$). The one-generation spinor packet is the even exterior algebra, equivalently the even-weight binary code

$$S^+ = \Lambda^0 E \oplus \Lambda^2 E \oplus \Lambda^4 E \cong C^+ = \{x \in \mathbb{F}_2^5 : |x| \text{ even}\}, \quad \dim S^+ = |C^+| = 2^{g_{\text{car}}-1} = 16. \quad (69)$$

The two carrier roots $y_+ = \frac{1}{2}$, $y_- = -\frac{1}{3}$ generate all hypercharges ($Y(Q_L) = y_+ + y_- = \frac{1}{6}$, $Y(u^c) = 2y_-$, $Y(e^c) = 2y_+$, $Y(\nu^c) = 0$) and the transport cusps $\{1, \frac{2}{3}, \frac{1}{3}\} = \{2y_+, -2y_-, 2(y_+ + y_-)\}$, with $\gamma_{\text{car}} = \text{Tr}_E Y^2 = 3y_-^2 + 2y_+^2 = \frac{5}{6}$. [E]

Gauge-rank factorisation ([[verification/v79_review_identities.py](#)]). The carrier polynomial factors into the two gauge ranks directly:

$$\boxed{6Y^2 - Y - 1 = (2Y - 1)(3Y + 1)}, \quad Y = \frac{1}{2} \text{ (} s=2, SU(2) \text{)}, \quad Y = -\frac{1}{3} \text{ (} b=3, SU(3) \text{)},$$

i.e. $(sY - 1)(bY + 1)$ — the polynomial splits into one linear representative of each Standard-Model non-abelian factor; the 5-slot carrier *forces* the split into $SU(3) \times SU(2)$.

Hypercharge [verification/v79_review_identities.py]	Lucas–Binet resonances	[E]	(mode reading	[C],
The integer carrier charges $(q_+, q_-) = (3, -2)$ (the primitive trace-free pair, $X^2 - X - 6 = (X-3)(X+2)$, Lean <code>unique_carrier_pair</code>) generate the Lucas U -sequence				
$D_n = \frac{3^n - (-2)^n}{5} : \quad D_1, \dots, D_6 = 1, 1, 7, 13, 55, 133,$				
which is, <i>in order</i> , $(N_\Phi, \cdot, \text{scalaron exponent } 7, \Delta_Q=13, \text{quark numerator } 55, \dim E_7=133)$. In particular the quark ratio reads $c_u/c_d = D_5/(N_{\text{fam}}^2 D_4) = \frac{55}{117}$ — a single recurrence of the hypercharge generator, not a free integer. [E] for the arithmetic; the per-mode physical identification is a [C] cross-domain reading (only $n \leq 6$, the carrier-slot range, is meaningful).				

Theorem 2 — Exterior–Action Duality

On C^+ the flip $F = x \oplus y$ is even, so $|F| \in \{0, 2, 4\}$. The three relations R_0, R_2, R_4 form a symmetric association scheme (the even Hamming scheme on five bits) with

$$\begin{array}{l} \text{valencies } (1, 10, 5) = (\dim \Lambda^0 E, \dim \Lambda^2 E, \dim \Lambda^4 E), \\ \text{spectral multiplicities } (1, 5, 10) = A_{\text{EW}} : \mathcal{A}_H : \mathcal{A}_\Lambda \end{array} \quad [\text{E}]. \quad (70)$$

The eigenvalues (computed directly) are $A_2 : \{10^{(1)}, 2^{(5)}, (-2)^{(10)}\}$ and $A_4 : \{5^{(1)}, (-3)^{(5)}, 1^{(10)}\}$, with $A_2 + A_4 = K_{16} - \mathbf{1}$. The two graphs are named objects:

$$\begin{array}{l} R_2(C^+) = \text{folded/complement, SRG}(16, 10, 6, 6) \text{ (the Clebsch complement);} \\ R_4(C^+) = \text{Clebsch graph, SRG}(16, 5, 0, 2) \end{array} \quad [\text{E}]. \quad (71)$$

Remark. The Standard-Model family is the *valency* side $(1, 10, 5)$ of one scheme; the cosmological action ladder is its *spectral* side $(1, 5, 10)$. The action ladder is therefore not merely “also binomial” — it is the spectral dual of the even exterior family. [C]

This duality is the MacWilliams transform ([verification/v79_review_identities.py]). C^+ is the even-weight code $[5, 4, 2]$; its dual $(C^+)^{\perp}$ is the repetition code $[5, 1, 5] = \{00000, 11111\}$ (the uniform/vacuum word). The MacWilliams identity sends the weight enumerator of the dual to that of C^+ , namely $(1, 10, 5)$ — so the valency $\leftrightarrow$ spectral duality *is* the coding-theoretic UV/IR duality, and the carrier grading $\Lambda^{\text{even}}(5) = 1+10+5$ is the MacWilliams image of the repetition vacuum. [E] (code identity) / [C] (holographic reading).

The full Bose–Mesner algebra (machine-checkable) fixes the duality beyond eigenvalues:

$$A_2^2 = 10 \mathbf{1} + 6A_2 + 6A_4, \quad A_2 A_4 = 3A_2 + 4A_4, \quad A_4^2 = 5 \mathbf{1} + 2A_2, \quad (72)$$

with eigenmatrix and multiplicity orthogonality

$$P = \begin{pmatrix} 1 & 10 & 5 \\ 1 & 2 & -3 \\ 1 & -2 & 1 \end{pmatrix}, \quad \det P = -64, \quad \boxed{P^{\top} \text{diag}(1, 5, 10) P = \text{diag}(16, 160, 80)} \quad [\text{E}], \quad (73)$$

so the flip split $240 = 160+80$ ($|F| = 2, 4$) appears directly in the scheme’s eigenmatrix orthogonality.

Carrier = D_5 (**verified**). The 16 vertices of C^+ are exactly the weights of the $D_5 = \mathfrak{so}_{10}$ half-spinor $\frac{1}{2}(\pm 1, \dots, \pm 1)$ with an even number of minus signs. The flip-2 and flip-4 graphs are translation- and permutation-invariant, so the *affine* group $C^+ \rtimes S_5$ (translations $\rtimes$ coordinate permutations) acts by graph/scheme automorphisms; the group closure gives the automorphism group of the even Hamming scheme (equivalently the Clebsch graph),

$$\boxed{\text{Aut}_{\text{aff}}(C^+) = C^+ \rtimes S_5, \quad |\text{Aut}_{\text{aff}}(C^+)| = 2^{g_{\text{car}}-1} g_{\text{car}}! = 16 \cdot 120 = 1920 = |W(D_5)|} \quad [\text{E}], \quad (74)$$

and $1920 = r |R(E_8)| = 8 \cdot 240$. (The *linear* code automorphism group alone is only the coordinate group S_5 of order 120; the factor $2^{g_{\text{car}}-1} = 16$ is the translation/scheme part. It is the affine/scheme group that equals $W(D_5)$.) Edges of the Clebsch graph: $|E(R_4)| = \frac{16 \cdot 5}{2} = 40 = |R(D_5)| = \sum_{f,j} L_{f,j}$ (the entire flavor transport budget). So the five-slot even code is the half-spin weight structure of $\mathfrak{so}_{10}$, the classical home of one SM family with ν^c .

Theorem 3 — the Coxeter closure equation selects $g_{\text{car}} = 5$

Requiring the even family to equal the action truncation through pair degree, $\dim S^+ = 2^{g-1} = \binom{g}{0} + \binom{g}{1} + \binom{g}{2}$, i.e.

$$\boxed{2^g - 2 = g(g+1) \iff g = 5} \quad [\mathbf{E}], \quad (75)$$

the unique positive-integer solution. At $g = 5$ the two sides equal 30, so *carrier exhaustion* $2^g - 2$ and the *Coxeter period* $g(g+1)$ coincide only at five slots: $2^{g_{\text{car}}} - 2 = 30 = h$ and $g_{\text{car}}(g_{\text{car}} + 1) = 30$.

Theorem 4 — the $\mathbb{Z}_{30}$ cyclotomic Coxeter compiler

From the three atoms $(2, 3, g_{\text{car}})$, with $h = 2 \cdot 3 \cdot g_{\text{car}} = 30$ and $r = \varphi(h) = 8$:

$$\boxed{|R(E_8)| = rh = 240, \quad \dim E_8 = r(h+1) = 8 \cdot 31 = 248, \quad h+1 = 2^{g_{\text{car}}} - 1 = 31} \quad [\mathbf{E}]. \quad (76)$$

The E_8 exponents are the primitive residues modulo 30,

$$\text{Exp}(E_8) = (\mathbb{Z}/30\mathbb{Z})^\times = \{1, 7, 11, 13, 17, 19, 23, 29\}, \quad \chi_C(t) = \Phi_{30}(t) = t^8 + t^7 - t^5 - t^4 - t^3 + t + 1, \quad (77)$$

and $|(\mathbb{Z}/30)^\times| = \varphi(30) = 8 = r$. The local SM algebra and the root atlas are the two arithmetic projections of $(r, h) = (8, 30)$:

$$\Omega_{\text{adm}} = 6r = 48, \quad D_{\text{start}} = 2h = 60, \quad \gcd(6r, 2h) = \dim \mathfrak{g}_{\text{SM}} = 12, \quad \text{lcm}(6r, 2h) = rh = 240. \quad (78)$$

Theorem 5 — the root transition atlas and the positive-edge moment

The directed even-flips of C^+ are the complete graph K_{16} :

$$\boxed{\begin{aligned} |R^+(E_8)| &= \binom{\dim S^+}{2} = \binom{16}{2} = 120 = \text{Tr}_{S^+} X^2 = 5! = \sum_{m \in (\mathbb{Z}/30)^\times} m, \\ |R(E_8)| &= 2 \binom{16}{2} = 240 \end{aligned}} \quad [\mathbf{E}]. \quad (79)$$

($X = 6Y$.) So the positive E_8 roots are the unordered transitions of one family; the second hypercharge moment, the positive-root count and the sum of the E_8 exponents coincide at 120. The CP pairing $m \leftrightarrow 30 - m$ of the exponents gives four pairs, each of sum $h = 30$:

$$\boxed{\text{Tr}_{S^+} X^2 = (N_{\text{fam}} + N_{\Phi}) h = 4 \cdot 30 = 120 = \dim \mathfrak{so}_{16} = \underbrace{45}_{D_5} + \underbrace{15}_{A_3} + \underbrace{60}_{(10,6)}} \quad [\mathbf{E}], \quad (80)$$

so the hypercharge moment is simultaneously four CP-paired Coxeter periods *and* the adjoint half $\mathfrak{so}_{16} = (45, 1) \oplus (1, 15) \oplus (10, 6)$ of the E_8 closure.

How the root *origin* is now obtained — see Theorem 7. A naive flip-size-preserving map $R_{\text{flip}} \rightarrow \mathbb{R}^8$ cannot be E_8 : the flip split $240 = 160 + 80$ ($|F| = 2, 4$) differs from the D_8

split $112 + 128$. The correct route is *not* a direct 240-metrization but the standard branching $E_8 \supset D_5 \times A_3$ (Theorem 7), which builds the root spaces from known D_5 and A_3 data. Until the A_3 lattice is derived from the four punctures (Theorem 7, [O]), E_8 remains a Coxeter atlas with a concrete, falsifiable lift.

Theorem 6 — downstream readouts from (h, r, c_3)

All discrete budgets and the seam seed are functions of $(h, r, c_3) = (30, 8, 1/(8\pi))$:

$$\dim S^+ = 2r = 16, \quad h = 2(\dim S^+ - 1) = 30, \quad (81)$$

$$10b_1 = 5r + 1 = 41 = 1 + h \frac{\dim S^+}{\dim \mathfrak{g}_{\text{SM}}}, \quad I_1^{\text{EW}} = \frac{5r + 1}{6r} = \frac{41}{48}, \quad (82)$$

$$u = \frac{r}{6}c_3 + 6r c_3^4 = \frac{4}{3}c_3 + 48c_3^4, \quad \delta_2 = 4! \binom{16}{2} c_3^8 = 2880 c_3^8, \quad (83)$$

$$\xi_{\text{tree}} = \frac{6}{r} = \frac{\dim \mathfrak{g}_{\text{SM}}}{\dim S^+} = \frac{3}{4}, \quad \frac{M_{\text{scal}}^2}{M_{\text{Pl}}^2} = c_3^{\Omega_{\text{adm}} - 10b_1} = c_3^{r-1} = c_3^7. \quad (84)$$

Two-graph product table. With the carrier-pair graph K_5 ($|E(K_5)| = \binom{5}{2} = 10 = \mathcal{A}_\Lambda$) and the family-corner graph K_4 on μ_4 ($|E(K_4)| = \binom{4}{2} = 6$, $|R(A_3)| = 2|E(K_4)| = 12$), every large budget is a product of the two graphs:

$$\boxed{40 = 10 \cdot 4, \quad 60 = 10 \cdot 6, \quad 120 = 10 \cdot 12, \quad 240 = 10 \cdot 24} \quad [\text{E}], \quad (85)$$

i.e. $(\sum L, D_{\text{start}}, |R^+(E_8)|, |R(E_8)|) = 10 \cdot (|\mu_4|, |E(K_4)|, |R(A_3)|, |S_4|)$. Two occupancy identities follow,

$$\boxed{10b_1 = 1 + |\mu_4| |E(K_5)| = 1 + 4 \cdot 10 = 41}, \quad (86)$$

$$\boxed{\Omega_{\text{adm}} = N_{\text{fam}} \dim S^+ = (N_{\text{fam}} + N_\Phi) |R(A_3)| = 3 \cdot 16 = 4 \cdot 12 = 48} \quad [\text{E}],$$

so the Higgs singleton is the fourth (μ_4) direction that closes $N_{\text{fam}} = 3$ to **4** of A_3 . **Occupancy deficit ladder.** The admissible occupancy splits cleanly,

$$\boxed{\begin{aligned} \Omega_{\text{adm}} &= \underbrace{40}_{\sum L (\text{flavor})} + \underbrace{1}_{\text{Higgs}} + \underbrace{7}_{\text{scalaron}} = 48, & \Omega_{\text{adm}} - \sum L &= 8 = h(D_5) = r(E_8), \\ \Omega_{\text{adm}} - 10b_1 &= 7 = h(D_5) - 1 \end{aligned}} \quad [\text{E}], \quad (87)$$

so the scalaron exponent $r - 1 = 7$ is exactly the post-flavor, post-Higgs occupancy remainder, $M_{\text{scal}}^2/\bar{M}_{\text{Pl}}^2 = c_3^{\Omega_{\text{adm}} - 10b_1} = c_3^7$. With $h(D_5) = 8$ and $|R^+(A_3)| = 6$ the seed itself is a D_5/A_3 ratio-product, $u = \frac{h(D_5)}{|R^+(A_3)|} c_3 + h(D_5) |R^+(A_3)| c_3^4$, and $\xi = \frac{|R^+(A_3)|}{h(D_5)} (1 + |R^+(A_3)|^2 c_3^3)^{-1} = \frac{3}{4} (1 + 36c_3^3)^{-1}$. The seed readouts and the two decuples are

$$\lambda_C^2 = u(1-u), \quad \beta_{\text{rad}} = 2c_3 u, \quad \sin^2 \theta_{13} = e^{-5/6} u, \quad \boxed{\omega_b = \frac{\lambda_C}{\mathcal{A}_\Lambda} = \frac{\lambda_C}{10} = 0.02244} \quad (\text{Planck } 0.02237), \quad (88)$$

$$\boxed{\begin{aligned} \det M_\ell &\propto u^{h/N_{\text{fam}}} = u^{10}, \\ \frac{\rho_\Lambda}{\bar{M}_{\text{Pl}}^4} &= \frac{3}{4\pi^2} e^{2\delta_{\text{ph}}} \left[\frac{v_{\text{geo}}}{5\beta_{\text{rad}}^2 M_{\text{Pl}}} \right]^{h/N_{\text{fam}}}, \end{aligned}} \quad (89)$$

with lepton powers $(5, 3, 2) = (g_{\text{car}}, b, s)$: $\sum = 10 = \mathcal{A}_\Lambda$ (the Λ decuple), $\prod = 30 = h$ (the Coxeter period). The first-generation transport sums read the non-primitive $\mathbb{Z}_{30}$ load,

$$(G_1, G_2, G_3) = (22, 13, 5) = (h - r, \dim \mathfrak{g}_{\text{SM}} + 1, g_{\text{car}}), \quad h - r = 30 - \varphi(30) = 22, \quad (90)$$

Flavor transport is a $D_5 \times A_3$ object. The transport packets $L_u=(7, 3, 0)$, $L_d=(7, 5, 2)$, $L_e=(8, 5, 3)$ split into a D_5 budget and an A_3 geometry:

$$\boxed{\sum_{f,j} L_{f,j} = 40 = |R(D_5)| \text{ (Clebsch edges)}, \quad L_{f,j} \bmod 6 \text{ lives on } C_6 = Z_6 = |R^+(A_3)|} \quad [\mathbf{E}], \quad (91)$$

the residues being the D_6 -orbit of the perfect set $\{0, 1, 3\} \subset C_6$ (cyclic distances $\{1, 2, 3\}$): $u \equiv \{0, 1, 3\}$, $d \equiv 2 - \{0, 1, 3\}$, $e \equiv 2 + \{0, 1, 3\}$, with the universal first-generation $+6$ winding. So the carrier D_5 sets the total budget and the family A_3 (its 6-element positive-root hexagon) sets the per-entry distances. The scalaron mass is a pure seam power $[\mathbf{E}]$; the amplitude is a *conditional* readout, kept $[\mathbf{C}]/[\mathbf{O}]$ until the scalaron variable is formally identified with the Starobinsky M parameter. That identification is now *heat-kernel grounded* (G2): the spectral-action Seeley–DeWitt coefficient $a_4|_{R^2} \sim R^2/72$ delivers exactly the Starobinsky R^2 term, with $M^2/\bar{M}_{\text{Pl}}^2 = 6(4\pi)^2/f_0$ and the TFPT closure c_3^7 fixing the cutoff moment f_0 (`[verification/v36_spectral_action_g2.py]`):

$$\underbrace{\frac{M_{\text{scal}}^2}{\bar{M}_{\text{Pl}}^2} = c_3^{r-1} = c_3^7}_{[\mathbf{E}]}, \quad \underbrace{A_s = \frac{N_\star^2}{24\pi^2} \frac{M_{\text{scal}}^2}{\bar{M}_{\text{Pl}}^2} = \frac{N_\star^2 c_3^7}{24\pi^2} = 2.17 \times 10^{-9}}_{[\mathbf{C}]/[\mathbf{O}] \text{ (if } M_{\text{scal}} = M_{\text{Star}})}. \quad (92)$$

Theorem 7 — the $D_5 \times A_3$ lift of the E_8 atlas

Since $A_3 = \mathfrak{su}_4 \cong \mathfrak{so}_6 = D_3$, one has the standard chain $E_8 \supset \mathfrak{so}_{16} \supset \mathfrak{so}_{10} \oplus \mathfrak{so}_6 = D_5 \times A_3$, with the exact branching (verified dimensions and root/weight counts):

$$\boxed{\mathfrak{e}_8 = (45, 1) \oplus (1, 15) \oplus (10, 6) \oplus (16, 4) \oplus (\overline{16}, \bar{4})}, \quad (93)$$

$$\underbrace{45+15+60+64+64}_{=248} = \dim E_8, \quad \underbrace{40+12+60+64+64}_{=240} = |R(E_8)|.$$

The adjoint **120** of $\mathfrak{so}_{16}$ is $(45, 1) \oplus (1, 15) \oplus (10, 6)$ and its spinor **128** is $(16, 4) \oplus (\overline{16}, \bar{4})$. The TFPT dictionary makes each summand a known object:

piece	value	TFPT reading
(45, 1)	$45 = 40 + 5$	D_5 carrier adjoint: Clebsch edges $40 = R(D_5) $ plus carrier rank $g_{\text{car}} = 5$
(1, 15)	$15 = 12 + 3$	A_3 family+Higgs adjoint: $\dim \mathfrak{g}_{\text{SM}} = 12$ plus family rank $N_{\text{fam}} = 3$
(10, 6)	$10 \cdot 6$	Λ decuple $\mathcal{A}_\Lambda = \binom{5}{2} = 10$ times Z_6 transport $\binom{4}{2} = 6$
$(16, 4) \oplus (\overline{16}, \bar{4})$	$2 \cdot 64$	matter half-spinor S^+ times the four corners μ_4 , plus conjugate sheet

Proof route (replaces the old blind 240-metrization). *Stage 1* (D_5 , $[\mathbf{E}]$): C^+ is the D_5 half-spin orbit, $R_4 = \text{Clebsch}$, $\text{Aut} = W(D_5)$, $|E(R_4)| = 40 = |R(D_5)|$ — Theorem 2. *Stage 2* (A_3 , $[\mathbf{E}]$): the carrier geometry fixes $X_f^\circ = \mathbb{P}^1 \setminus \mu_4$ with $H_1(X_f^\circ, \mathbb{Z}) = \mathbb{Z}^3$ and a rigid $D_4 \subset PGL_2$ generated by $z \mapsto iz$ and $z \mapsto 1/z$. The logarithmic forms $\omega_i = dz/(z - p_i)$ regular at infinity span H^1 by their differences, dual under the residue pairing to the puncture loops γ_i with $\text{Res}(\omega_i, \gamma_j) = \delta_{ij}$; hence the canonical (residue) metric on the puncture classes is I_4 , and the twelve puncture-difference cycles $\gamma_i - \gamma_j$ are exactly the A_3 roots:

$$\boxed{\{\gamma_i - \gamma_j\}_{i \neq j} : \# = 12 = |R(A_3)| = \dim \mathfrak{g}_{\text{SM}}, \quad \text{rank} = 3 = N_{\text{fam}}, \quad \text{Gram} = \text{Cartan}(A_3), \quad \det = 4} \quad [\mathbf{E}], \quad (94)$$

with $\dim A_3 = 15 = \dim S^+ - 1$, and the A_3 positive roots equal the exponent sum $1+2+3 = 6 = Z_6 = |E(K_4)|$. *The corner rotation realises the A_3 Coxeter element.* The carrier generator $z \mapsto iz$ permutes μ_4 as the 4-cycle $(p_1 p_2 p_3 p_4) \in S_4 = W(A_3)$ and acts on $H^1 = \mathbb{C}^3$ as an isometry of the Cartan form with

$$\boxed{\text{ord} = 4 = h(A_3) = |\mu_4|, \quad \text{spec} = \{i, -1, -i\}, \quad \chi(t) = \frac{t^4 - 1}{t - 1} = \Phi_4(t)\Phi_2(t)} \quad [\mathbf{E}], \quad (95)$$

i.e. the eigenvalues are the A_3 exponents $\{1, 2, 3\}$ as fourth roots of unity — the exact analogue of the E_8 Coxeter element reading Φ_{30} . The full $D_4 = \langle z \mapsto iz, z \mapsto 1/z \rangle$ embeds in $W(A_3) = S_4$ (order 8, index 3) by isometries. *Stage 3 (standard, [E]):* assemble E_8 from the $D_5 \times A_3$ branching — no 240-map required.

The μ_4 glue theorem (why the four corners appear everywhere). The two factors have equal discriminant groups, and E_8 is unimodular:

$$\boxed{\text{disc}(D_5) = \text{disc}(A_3) = \mathbb{Z}_4, \quad [E_8 : D_5 \oplus A_3] = \sqrt{\frac{\det D_5 \cdot \det A_3}{\det E_8}} = \sqrt{\frac{4 \cdot 4}{1}} = 4 = |\mu_4|} \quad [\text{E}]. \quad (96)$$

So $E_8 = (D_5 \oplus A_3) + \mu_4\text{-glue}$: the four family corners $\mu_4 = \mathbb{Z}_4$ are precisely the glue code that fuses the carrier and the family into the unimodular E_8 lattice. The glue pairs the D_5 spinor class ($= S^+$, the family) with the A_3 class — exactly the $(16, 4) \oplus (\overline{16}, \overline{4})$ spinor block. Hence μ_4 plays three roles at once: family corners, A_3 origin, and E_8 glue.

Discriminant-norm glue lemma (and the TFPT 5/4, 3/4 identification). The two glue generators carry the discriminant quadratic form $q \in \mathbb{Q}/2\mathbb{Z}$:

$$\boxed{\begin{aligned} q(D_5) &= \frac{n}{4}|_{n=5} = \frac{5}{4} \text{ (spinor class)}, & q(A_3) &= \frac{n}{n+1}|_{n=3} = \frac{3}{4} \text{ (corner class)}, \\ q(D_5) + q(A_3) &= 2 \equiv 0 \pmod{2}, & q(D_5) - q(A_3) &= \frac{1}{2} \end{aligned}} \quad [\text{E}]. \quad (97)$$

The diagonal $\langle (s, g) \rangle \cong \mathbb{Z}_4$ is isotropic ($k^2 \cdot 2 \equiv 0$ for all k), so the glue closes to an *even unimodular* lattice — this is E_8 .

The two glue norms are just $(g_{\text{car}}, N_{\text{fam}})$ over the glue index. Because $\text{rank } D_5 = g_{\text{car}} = 5$ (so $q(D_5) = \frac{n}{4}|_{n=5} = g_{\text{car}}/|\mu_4|$) and $\text{rank } A_3 = N_{\text{fam}} = 3$ (so $q(A_3) = \frac{n}{n+1}|_{n=3} = N_{\text{fam}}/|\mu_4|$, with $n+1 = 4 = |\mu_4|$),

$$\boxed{q(D_5) = \frac{g_{\text{car}}}{|\mu_4|} = \frac{5}{4}, \quad q(A_3) = \frac{N_{\text{fam}}}{|\mu_4|} = \frac{3}{4}}$$

and *all four arithmetic operations* on the pair are then forced compiler atoms:

$$\begin{aligned} \underbrace{q(D_5) + q(A_3) = \frac{g_{\text{car}} + N_{\text{fam}}}{|\mu_4|} = 2}_{E_8 \text{ root norm}}, & \quad \underbrace{q(D_5) - q(A_3) = \frac{|\mathbb{Z}_2|}{|\mu_4|} = \frac{1}{2}}_{\delta \text{ (lepton transport)}}, \\ \underbrace{q(D_5)q(A_3) = \frac{g_{\text{car}}N_{\text{fam}}}{|\mu_4|^2} = \frac{15}{16}}_{\dim \mathfrak{su}(4) / \dim S^+}, & \quad \underbrace{\frac{q(D_5)}{q(A_3)} = \frac{g_{\text{car}}}{N_{\text{fam}}} = \frac{5}{3}}. \end{aligned}$$

So the distinguished lepton transport value $\delta = \frac{1}{2} = (g_{\text{car}} - N_{\text{fam}})/|\mu_4| = |\mathbb{Z}_2|/|\mu_4|$ is the carrier-minus-family count over the glue index — a fully atomic origin, not a chosen rational (it also equals the harmonic-frame holonomy diagonal modulus, [\[verification/v41_leg_assignment.py\]](#)); and (sum, ratio) invert uniquely back to $(\frac{5}{4}, \frac{3}{4})$. [\[verification/v51_boundary_half_step.py\]](#) The sum $\frac{5}{4} + \frac{3}{4} = 2$ is literally the E_8 spinor-root norm $\frac{1}{2}(\pm)^8$ split across $D_5 \times D_3$: $5 \cdot \frac{1}{4} + 3 \cdot \frac{1}{4} = 2$. Strikingly, both norms are pre-existing TFPT constants:

$$\boxed{\frac{\delta_2}{\delta_{\text{top}}^2} = \frac{2880 c_3^8}{(48 c_3^4)^2} = \frac{5}{4} = q(D_5), \quad \xi_{\text{tree}} = \frac{\dim \mathfrak{g}_{\text{SM}}}{\dim S^+} = \frac{12}{16} = \frac{3}{4} = q(A_3)} \quad [\text{E}], \quad (98)$$

so $\delta_2/\delta_{\text{top}}^2 + \xi_{\text{tree}} = 2$: the two-loop/defect ratio is the D_5 spinor norm and the gravitational tree normalization is the A_3 corner norm, and their sum is the E_8 root norm.

The whole integer skeleton from one pair $(g_{\text{car}}, N_{\text{fam}}) = (5, 3)$ **[E]**
`([verification/v53_compiler_core.py])`

Every load-bearing integer of the compiler flows from the single boundary pair $(g_{\text{car}}, N_{\text{fam}}) = (5, 3)$ by sum, difference and squares:

$$\text{rank } E_8 = g_{\text{car}} + N_{\text{fam}} = 8, \quad |\mathbb{Z}_2| = g_{\text{car}} - N_{\text{fam}} = 2, \quad |\mu_4| = \frac{g_{\text{car}} + N_{\text{fam}}}{2} = 4,$$

and $(N_{\text{fam}}, |\mu_4|, g_{\text{car}}) = (3, 4, 5)$ is the Pythagorean triple. This is not decorative: the grand-mass-volume exponent $\Delta_Y = g_{\text{car}}^2 = 25$ ($\det M_{\text{SM}} \sim (\varphi_0^{\text{ret}})^{25}$; document 3) splits over the sectors as a *difference of squares*,

$$\Delta_Y = g_{\text{car}}^2 = \underbrace{N_{\text{fam}}^2}_{\text{down}=9} + \underbrace{(g_{\text{car}} - N_{\text{fam}})(g_{\text{car}} + N_{\text{fam}})}_{=|\mathbb{Z}_2| \cdot \text{rank } E_8} = 9 + 16 = \dim S^+$$

i.e. the down sector carries N_{fam}^2 and up+lepton carries $|\mathbb{Z}_2| \cdot \text{rank } E_8 = \dim S^+$ (the K -row sums $(6, 9, 10)$). **The anchor is the Dirac square root of this skeleton:** $a = (1, 1, 2)$ is the *unique* 3-multiset whose elementary symmetric functions are $(|\mu_4|, g_{\text{car}}, |\mathbb{Z}_2|) = (4, 5, 2)$, equivalently its characteristic polynomial is $\chi_a(t) = (t-1)^2(t-2) = t^3 - |\mu_4|t^2 + g_{\text{car}}t - |\mathbb{Z}_2|$ — the compiler atoms *are* its char-poly coefficients. Sharper still (`[verification/v491_p2_partition_corollary.py]`): the partition needs only $e_1 = 4$ — three *positive integers* summing to 4 already force $\{1, 1, 2\}$, so $g_{\text{car}} = e_2 = 5$ and $|\mathbb{Z}_2| = e_3 = 2$ are *corollaries* of the four marks given the weight typing (integrality = Birkhoff–Grothendieck splitting exponents, positivity = $h^0=0$, sum = par-deg-0 “given (U)”; every dropped assumption is negative-controlled, and **AX.P2.01** stays a declared axiom — the residual is the typing, not the integer 5. That typing is itself now hardened (`[verification/v499_p2_weights_deligne_bg.py]`, **P2.TYPING.01**): given the four marks, rank 3, the cusp class and (U), Deligne + Birkhoff–Grothendieck + Mehta–Seshadri stability force $E = \mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$ outright ($h^0=0$ is the positivity), leaving only the module identification and (U) as **[C]** residuals — and the module-identification rest is now *one* residual with the QGEO modulus rest, both reducing to the same order-4 clock = Coxeter/cusp-class carrier datum (**v503**, **QGEO.EMERGE.LIGHT.01**; markers unchanged); the clock-rigidity theorems then reduce that common residual to one alignment bit, with the clock’s order fermionically forced, $U^2 = (-1)^F$ exactly (`[verification/v506_seam_clock_rigidity.py]`, **SEAM.CLOCK.RIGIDITY.01**); and the bit-origin theorems derive the deck’s *class* (a half-period translation of the seam double) — only its position stays the **[C]** carrier input (`[verification/v507_seam_bit_origin.py]`, **SEAM.BIT.ORIGIN.01**); the freedom theorems then type that position half as *topology* (the covering deck is free on the seam circle, the edge class is excluded), so the residual is the square-modulus datum $\tau = i$ alone (`[verification/v510_seam_bit_freedom.py]`, **SEAM.BIT.FREEDOM.01**); and the flag-transitivity equivalence web restates that residual in discrete form — flag transitivity of the four marks ($V_4 \rightarrow D_4$) $\iff \tau = i$, with bare mark-transitivity automatic for every configuration and no local jet seeing the side bit (`[verification/v512_seam_tau_flag.py]`, **SEAM.TAU.FLAG.01**) — a residual that has since survived ten side-blind derivation attacks (**v521/v525/v529**) and carries a physical definition as the twist-class choice with a gauge-robust order parameter, in principle interferometrically readable (**v528**; it remains formal input) — and, since **v534**, the first *dynamical* selection datum: reflection positivity on the interacting mark family keeps exactly the $\delta = \pi/2$ member alive (toy-fenced). So the theory’s only transcendental is π (through $c_3 = \frac{1}{8\pi}$, $8 = \text{rank } E_8$, and the seed $\frac{1}{6\pi}$, $6 = |R^+(A_3)|$); everything else is this $(5, 3)$ -generated integer skeleton.

Boundary Carrier Selection Theorem (Theorem A) **[E]/[C]**

E_8 is not assumed: it is *selected* as the unique even-unimodular audit hull of the admissible boundary-carrier data. **Boundary side:** $\mu_4 = \{1, i, -1, -i\}$ gives $X_f = \mathbb{P}^1 \setminus \mu_4$ with $H_1 = \mathbb{Z}^3$, hence A_3 and $N_{\text{fam}} = 3$. **Carrier side:** a minimal D -type carrier with a 16-dim half-spinor forces D_5 (only $n = 5$ gives $2^{n-1} = 16$; $S^+ = \Lambda^{\text{even}}\mathbb{C}^5 = 1+10+5$). **Glue:** $\text{disc}(D_5) = \text{disc}(A_3) = \mathbb{Z}_4$ (isomorphic, cyclic, anti-isometric), $q(D_5) + q(A_3) = 2$ (the E_8 root norm), glue index $4 = |\mu_4|$, and E_8 is the unique even unimodular rank-8 lattice. **Exclusion:** the four conditions

$$(i) \dim S^+ = 16, \quad (ii) N_{\text{fam}} = 3, \quad (iii) \text{cyclic } \mathbb{Z}_4 \text{ glue}, \quad (iv) q_D + q_A = 2$$

are met *only* by $D_5 \oplus A_3$; D_8 (non-cyclic $\mathbb{Z}_2 \times \mathbb{Z}_2$, no family geometry), $E_7 \oplus A_1$ (1 family, no 16-spinor), $E_6 \oplus A_2$ (2 families, $\mathbb{Z}_3$), $A_4 \oplus A_4$ (no D -spinor, $\mathbb{Z}_5$) and A_8 (single factor) each fail at least one. **Typing:** the lattice/classification facts are **[E]**; the physical selection (reflection-positive seam $c_3 = 1/8\pi +$

minimal chiral carrier $\Rightarrow$ exactly these inputs) is the **[C]/[O]** selection axiom (P1 analytic interface + P2 carrier interface), not a zero-input derivation. `[verification/v47_selection_theorem.py]`

Group-level closure (μ_4 as the common centre). The lattice glue lifts to a clean group statement, because both factors share the *same* centre and μ_4 *is* the diagonal:

$$Z(\text{Spin}(10)) = \mathbb{Z}_4, \quad Z(SU(4)) = \mathbb{Z}_4, \quad |\mu_4| = 4, \quad G_{\text{closure}} = \frac{\text{Spin}(10) \times SU(4)}{\Delta \mathbb{Z}_4} \subset E_8 \quad [\text{E}], \quad (99)$$

with $\text{rank} = 5 + 3 = 8 = \text{rank } E_8$. Every adjoint block is diagonal- $\mathbb{Z}_4$ neutral (generator $\omega = i$; charges mod 4): $(45, 1), (1, 15) : 0$; $(10, 6) : 2+2=0$; $(16, 4) : 1+3=0$; $(\overline{16}, \overline{4}) : 3+1=0$, so $45+15+60+64+64 = 248$ are genuine reps of the quotient. This promotes μ_4 from a lattice glue index to the *common centre quotient* — the physically cleanest language for the gluing: the D_5 half-spinor 16 pairs with the $SU(4)$ fundamental 4 so the spinor charges ± 1 cancel the fundamental charges ∓ 1 .

Root-length completion. Every E_8 block reaches norm 2:

block	norm split	reading
$(45, 1), (1, 15)$	2, 2	D_5, A_3 root adjoints
$(10, 6)$	$1 + 1 = 2$	$SO(10)$ vector $\oplus$ $SO(6)$ vector
$(16, 4) \oplus (\overline{16}, \overline{4})$	$\frac{5}{4} + \frac{3}{4} = 2$	D_5 spinor norm + A_3 corner norm

The old question “can the 240 flip atlas be metrized to E_8 ?” is now the sharp question “are the TFPT D_5 and A_3 metrics in standard normalization?” — and the $5/4, 3/4$ match says yes.

Explicit construction (the lift is now built, not counted). Embedding $D_5 \oplus A_3 = D_5 \oplus D_3 \subset D_8$ and adjoining the even spinor coset (the μ_4 glue), the 240 roots are produced explicitly and verified:

$$\{\pm e_i \pm e_j\}_{i < j}^8 (112) \cup \{\frac{1}{2}(\pm)^8 : \text{even}\} (128) = 240 \text{ roots, all norm 2, Cartan det} = 1 \quad [\text{E}]. \quad (100)$$

Splitting each root into its D_5 (first five) and A_3 (last three) parts gives exactly the branching blocks — $(2, 0) : 40$, $(0, 2) : 12$, $(1, 1) : 60$, and $(\frac{5}{4}, \frac{3}{4}) : 128$ — and the 128 spinor roots realise $\frac{5}{4} + \frac{3}{4} = 2$ on the nose. All 240 vectors are integer combinations of one set of 8 simple roots, so the assembled lattice *is* E_8 . The blind 240-metrization is therefore eliminated: E_8 is explicitly $(D_5 \oplus A_3) + \mu_4$ -glue.

Remark (the metric, computed). This reframes E_8 from origin to closure: E_8 is the closure algebra of the D_5 carrier and the A_3 family geometry, glued by μ_4 . The root system, Cartan form, $W(A_3) = S_4$ and the corner-rotation = A_3 Coxeter element are exact (residue pairing + explicit carrier symmetry). The transcendental geometric metric $G_{ij} = -\log |p_i - p_j|$ on μ_4 has been computed: it is only D_4 -symmetric, with a regularisation-independent anisotropy $-\log 2$ on the root space, hence *not* proportional to $\text{Cartan}(A_3)$. It need not be: the E_8 -relevant A_3 is the canonical *residue* pairing, and the geometric metric is a D_4 -equivariant deformation that preserves the root system, Weyl group and Coxeter element. So the lattice data is exact **[E]**; the analytic metric is a *characterised* deformation, not an open gap.

Uniform Coxeter law. All three factors obey $|R| = h \cdot \text{rank}$, and the two TFPT Coxeter numbers are themselves carrier data:

$$\begin{aligned} h(D_5) &= 8 = r(E_8) = \varphi(30), & h(A_3) &= 4 = |\mu_4|, \\ h(D_5) h(A_3) &= 2^{g_{\text{car}}} = 32, & 2^{g_{\text{car}}} - 2 &= 30 = h(E_8) \end{aligned} \quad [\text{E}], \quad (101)$$

so the Coxeter closure of Theorem 3 reappears as $h(E_8) = h(D_5)h(A_3) - 2$, and $h(D_5) + h(A_3) = 12 = \dim \mathfrak{g}_{\text{SM}} = |R(A_3)|$.

Theorem 8 — the $(1, 1, 2)$ anchor microcode and the φ_0 -ladder

The parabolic flavor bundle $E_\bullet = \mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$ on $\mathbb{P}^1 \setminus \mu_4$ has splitting type $a = (1, 1, 2)$, whose power sums $p_n(a) = 2 + 2^n$ generate the whole compiler budget:

The anchor power compiler.

$$(p_1, \dots, p_5) = (4, 6, 10, 18, 34) = (|\mu_4|, |R^+(A_3)|, \mathcal{A}_\Lambda, N_{\text{fam}}|R^+(A_3)|, \mathbb{Z}_6 \text{ lift}),$$

$$q(A_3) = \frac{p_2}{2p_1} = \frac{3}{4}, \quad q(D_5) = \frac{p_3}{2p_1} = \frac{5}{4}, \quad \sum L = p_1 p_3 = 40,$$

$$\Omega_{\text{adm}} = 2p_1 p_2 = 48, \quad D_{\text{start}} = p_2 p_3 = 60, \quad |R(E_8)| = 4p_2 p_3 = 240.$$

The elementary symmetric data $\chi_a(t) = (t-1)^2(t-2)$ give $(e_1, e_2, e_3) = (4, 5, 2) = (|\mu_4|, g_{\text{car}}, |\mathbb{Z}_2|)$, so the anchor reconstructs the carrier rank, and $10b_1 = e_1^2 + e_2^2 = 4^2 + 5^2 = 41$. The seed itself is the two-term anchor form $\varphi_0^{\text{ret}} = \frac{2p_1}{p_2} c_3 + 2p_1 p_2 c_3^4 = \frac{4}{3} c_3 + 48c_3^4$.

The anchor affine normal form (2026-08-07) — all budgets from ONE recursion [E]
 ([verification/v832_anchor_flavor_checksum.py])

The anchor power compiler above is one affine map. The power sums $p_n(a) = 2 + 2^n$ satisfy

$$p_{n+1} = T(p_n), \quad T(x) = 2x - 2 \quad \text{identically in } n \text{ (sympy),}$$

with the shift $q_n = p_n - 2 = 2^n$ pure binary doubling. T has the *unique* fixed point $2 = |\mathbb{Z}_2|$ (the sheet), and the T -orbit of 4 is exactly the compiler quintet $(4, 6, 10, 18, 34) = (|\mu_4|, |R^+(A_3)|, \mathcal{A}_\Lambda, N_{\text{fam}}|R^+(A_3)|, \mathbb{Z}_6 \text{ lift})$, with $p_0 = 3 = N_{\text{fam}}$ prepended by $T(3) = 4$. Every budget line is then a word in the orbit and the fixed point: $|R(E_8)| = p_1 p_2 p_3 = 240$; $\dim E_8 = 240 + q_3 = 248$, where the ladder identity $p_4 - p_3 = p_3 - 2 = 8 = h(D_5) = \text{rank } E_8$ makes the corpus form $240 + (p_4 - p_3)$ the *same* formula; $h(E_8) = p_2 p_3 / 2 = 30$ (so $D_{\text{start}} = p_2 p_3 = 60$); $|R(D_5)| = p_1 p_3 = 40$; $\Omega_{\text{adm}} = 2p_1 p_2 = 48$; and the elementary layer $(e_1, e_2, e_3) = (4, 5, 2)$ has the fixed point of T as e_3 , with $10b_1 = e_1^2 + e_2^2 = 41$. Controls: the wrong map $T'(x) = 2x - 1$ (fixed point 1, orbit misses the quintet) and the wrong anchor $a' = (1, 2, 2)$ (recursion offset -1 , budget misses $(240, 48)$) both fire. A normal form of the existing compiler tables — a compression, not a new claim; 18/18 exact, verdict ANCHOR-AFFINE-EXACT.

The φ_0 -ladder. With $\lambda_Y = \sqrt{\varphi_0^{\text{ret}}(1 - \varphi_0^{\text{ret}})}$, every fermion mass is the single closed form $\hat{m}_{f,j} = \frac{v_{\text{geo}}}{\sqrt{2}} \lambda_Y^{L_{f,j}} \Lambda_{f,j}$ with $\lambda_Y^L \Lambda = \pi c (\varphi_0^{\text{ret}})^k$, so all nine holonomies are resolvent outputs and the masses form the φ_0^{ret} -order ladder $t(0) > \{b, c, \tau\}(2) > \{s, \mu\}(3) > \{u, d\}(4) > e(5)$.

Secondary E_8 atlas. Beyond the main $D_5 \times A_3$ split, the residue matrix appears in the $E_6 \times A_2$ branching, $248 = \|R\|_F^2 + \det R + 2(1^\top R a) N_{\text{fam}} = 78 + 8 + 2 \cdot 27 \cdot 3$, and the scalaron exponent in $E_7 \times A_1$, $112 = 7 \cdot 16$. The seven E_8 slices are charted in the companion note `e8_secondary_branching_atlas` as a falsifiable audit raster.

The $E_6 \times A_2$ block is the $\tau=\omega$ face ([verification/v407_dn_pairings_omega.py], SEAM.EQUIV.02). A_2 is the Eisenstein lattice of the family CM point $\tau=\omega$ (Cartan $\det = 3 = N_{\text{fam}}$), and the (d, n) selector's three pairings $\{n \cdot \mathbf{1}, n \cdot a, n \cdot \sigma\} = \{2, 8, 121\}$ are exactly its family-slice atoms $\{|\mathbb{Z}_2|, \dim A_2 = \det R = \text{rank } E_8, \|Pl(K)\|_1^2\}$, so deriving (d, n) (the $R4'$ residue, `v139`) folds into the dual keystone (`v405`), leaving only “why $11 = \|Pl(K)\|_1$ ”.

Master statement and status

Five carrier slots give D_5 , four family corners give A_3 , and $D_5 \times A_3$ gives the E_8 atlas.

Equivalently: $\mathcal{A}_\Lambda = \frac{h}{N_{\text{fam}}} = 10 = \binom{5}{2}$, $h = 30 = 2 \cdot 3 \cdot 5$, $(5, 3, 2) : \sum = 10, \prod = 30$; $\Lambda, \det M_\ell$, the E_8 Coxeter structure and the action ladder read the same five-slot pair-decuple. The Standard model, flavor, Λ , inflation and E_8 are projections of the even carrier code C^+ ($= D_5$ half-spinor) on five slots together with the four-point family A_3 .

Statement	Status	Note
Even code C^+ , $\dim S^+ = 16$, dual-root Y	[E]	Theorem 1
Exterior–Action duality $(1, 10, 5) \leftrightarrow (1, 5, 10)$	[E]	scheme eigenvalues verified
$R_4 = \text{Clebsch } (16, 5, 0, 2)$, $R_2 = (16, 10, 6, 6)$; $\text{Aut}_{\text{aff}} = W(D_5) = 1920$	[E]	carrier = D_5 half-spinor
$2^g - 2 = g(g+1) \Leftrightarrow g = 5$	[E]	Coxeter closure (unique)
$h=30=2\cdot 3\cdot 5$, $r=\varphi(30)=8$, $\chi_C=\Phi_{30}$	[E]/[C]	cyclotomic compiler
$ R^+ = \binom{16}{2} = \text{Tr } X^2 = 4h = 120$	[E]	positive-edge moment
$40, 60, 120, 240 = 10 \cdot (4, 6, 12, 24)$; $41=1+4\cdot 10$	[E]	$K_5 \times K_4$ product table
$10b_1=5r+1$, u totient, $M_{\text{scal}}^2=c_3^{r-1}$, $\omega_b=\lambda_C/10$	[E]/[C]	downstream readouts
$A_s = N_*^2 c_3^7 / 24\pi^2$	[C]/[O]	if $M_{\text{scal}} = M_{\text{Star}}$
$\mathfrak{e}_8 = (45, 1) \oplus (1, 15) \oplus (10, 6) \oplus (16, 4) \oplus (\overline{16}, \overline{4})$	[E]/[C]	$D_5 \times A_3$ lift
240 roots built from $D_5 \oplus A_3 + \mu_4$ glue, $\det 1$	[E]	explicit E_8 construction
flavor $\sum L=40= R(D_5) $, entries on $C_6= R^+(A_3) $	[E]	transport = $D_5 \times A_3$
A_3 from $\mathbb{P}^1 \setminus \mu_4$: residue pairing = Cartan, $\det 4$	[E]	Stage 2 (Theorem 7)
corner $z \mapsto iz = A_3$ Coxeter el.: $\{i, -1, -i\}$, $\Phi_4 \Phi_2$	[E]	order $4 = h(A_3) = \mu_4 $
$\text{disc}(D_5) = \text{disc}(A_3) = \mathbb{Z}_4$, glue $= \mu_4$	[E]	$E_8 = (D_5 \oplus A_3) + \mu_4$
$q(D_5)=\frac{5}{4}=\delta_2/\delta_{\text{top}}^2$, $q(A_3)=\frac{3}{4}=\xi_{\text{tree}}$, sum 2	[E]	discriminant-norm glue
$120 = \dim \mathfrak{so}_{16}$; $\Omega_{\text{adm}}=40+1+7$	[E]	adjoint half; deficit ladder
$h(D_5)=8=r(E_8)$, $h(A_3)=4= \mu_4 $, $ R =h \cdot r$	[E]	uniform Coxeter law
$\text{Exp}(E_8)$ as 41/48 budget deficits	[O]	exponent ledger (fingerprint)
geometric metric D_4 -deform (anisotropy $-\log 2$), residue $= A_3$	[E]	metric computed/characterised

Scope. This note compresses; it does not re-derive the upstream boundary kernel (the P1 hardening chain) or the carrier rigidity theorem (Lean), and it makes no claim of a final theory. The arithmetic and the $D_5/\text{Clebsch}/W(D_5)$, $A_3/\text{residue-Cartan}/\text{Coxeter}$, and μ_4 -glue identifications are exact, and the E_8 lattice is now built explicitly from $D_5 \oplus A_3$ plus the μ_4 glue (240 roots, $\det 1$). The only residual [C] is the analytic identification of the carrier geometry's L^2 cusp Hodge metric with the canonical residue metric up to scale, which does not affect the explicit root/lattice/Weyl construction. Honest non-fits (Koide $0.6645 \neq 2/3$, η_B , m_p/m_e) stay outside the compiler.

Appendix: machine-checkable $E_8 = (D_5 \oplus A_3) + \mu_4$ construction

The lift is reproducible by a short script (all entries below verified exactly).

Cartan matrices and discriminant groups.

$$C(D_5) = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & 0 \\ 0 & 0 & -1 & 0 & 2 \end{pmatrix}, \quad C(A_3) = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}; \quad \det C(D_5) = \det C(A_3) = 4.$$

Smith normal forms give $\text{disc}(D_5) = \text{disc}(A_3) = \mathbb{Z}_4$; the glue index is $[E_8 : D_5 \oplus A_3] = \sqrt{4 \cdot 4/1} = 4 = |\mu_4|$, with discriminant-form norms $q(D_5) = \frac{5}{4}$, $q(A_3) = \frac{3}{4}$, $q(D_5)+q(A_3) = 2 \equiv 0 \pmod{2}$ (even-glue). Writing $\mu := |\mu_4| = 4$ the norms are $q(D_5) = \frac{\mu+1}{\mu}$, $q(A_3) = \frac{\mu-1}{\mu}$, so $q(D_5)q(A_3) = 1 - \frac{1}{\mu^2} = \frac{15}{16}$ and

$$\dim S^+(1 - q(D_5)q(A_3)) = 16 \cdot \frac{1}{16} = 1 \quad [\text{E}], \quad (102)$$

i.e. the glue product fills all $15 = \dim S^+ - 1$ non-trivial half-spin states and the defect is exactly the singleton (the ν^c/Higgs closure). And E_8 has a tidy μ_4 -cardinal form, $\dim E_8 = \binom{|\mu_4|^2}{2} + 2|\mu_4|^3 = \binom{16}{2} + 2 \cdot 64 = 120 + 128 = 248$ [C].

Explicit roots. Split $\mathbb{R}^8 = \mathbb{R}_{(D_5)}^5 \oplus \mathbb{R}_{(A_3)}^3$. The 240 roots are

$$\{\pm e_i \pm e_j : 1 \leq i < j \leq 8\} \text{ (112)} \cup \left\{ \frac{1}{2}(\pm 1)^8 : \text{even } \# \text{ of } - \right\} \text{ (128),}$$

all of norm 2. Splitting each by $(\|\cdot\|_{D_5}^2, \|\cdot\|_{A_3}^2)$ gives the branching blocks

$$(2, 0): 40 [D_5], \quad (0, 2): 12 [A_3], \quad (1, 1): 60 [(10, 6)], \quad \left(\frac{5}{4}, \frac{3}{4}\right): 128 [(16, 4) \oplus (\overline{16}, \overline{4})],$$

so $\mathfrak{e}_8 = (45, 1) \oplus (1, 15) \oplus (10, 6) \oplus (16, 4) \oplus (\overline{16}, \overline{4})$, $248 = 45+15+60+64+64$, $240 = 40+12+60+64+64$.

Simple-root Gram (unimodular check). With the Bourbaki simple roots $\alpha_1 = \frac{1}{2}(e_1 - e_2 - \dots - e_7 + e_8)$, $\alpha_2 = e_1 + e_2$, and $\alpha_k = e_{k-1} - e_{k-2}$ for $k = 3, \dots, 8$ (so $\alpha_3 = e_2 - e_1$, $\alpha_4 = e_3 - e_2$, $\dots$, $\alpha_8 = e_7 - e_6$), the Gram matrix $G_{ij} = \langle \alpha_i, \alpha_j \rangle$ is the E_8 Cartan matrix with $\det G = 1$ and all $G_{ii} = 2$; every one of the 240 roots is an integer combination of the α_i . Hence the assembled lattice is E_8 , not merely a 240-count. [E] (The shifted index $e_{k-2} - e_{k-3}$ would duplicate α_3 and make G singular — use $e_{k-1} - e_{k-2}$.)

Part III

Appendix: ablation of the EM fixed point

The single sharpest reviewer test is: *was the closure $F_{U(1)}(\alpha) = 0$ engineered so that α comes out?* The honest answer is an ablation — vary each input and see which actually move the root. Three one-parameter scans (all others held at their compiler values) separate the *load-bearing* integers from the *fine* last-digit selector.

EM fixed point: only $M = 41$ and $N = 8$ land on α^{-1}




Figure 7: EM-closure ablation (data plot, `verification/make_figures.py`). (a) α_*^{-1} vs the integer budget M : only $M = 41$ lands on CODATA-2022. (b) α_*^{-1} vs the seam denominator N in $c_3 = 1/(N\pi)$: only $N = 8$ lands. Both integers are fixed elsewhere (flavor budget; rank/ $h(D_5)$), so the equation is not freely tunable to 137.036.

(A1) The transport budget $M = \sum L + N_\Phi$ (exponent held at $-\frac{5}{4}$):

M	38	39	40	41	42	43	44
α_*^{-1}	140.530	139.326	138.162	137.0359992	135.946	134.890	133.866

Each unit of M shifts α_*^{-1} by ≈ 1.09 , so *only* $M = 41$ lands on 137.036. This is a hard lock — and $41 = 10b_1 = \sum L + N_\Phi = 40 + 1$ is fixed by the compiler (flavor budget 40 + one Higgs). **Load-bearing.**

(A2) The seam “8” in $c_3 = 1/(N\pi)$ (here $\delta_{\text{top}} = 48c_3^4$ tracks c_3):

N	6	7	8	9	10
α_*^{-1}	76.97	104.85	137.036	173.53	214.34

Only $N = 8$ — i.e. $c_3 = 1/(8\pi) = 1/(\text{rank } E_8 \cdot \pi)$ — reproduces 137.036. The “8” is the same one fixed by the bootstrap ($h(D_5) = \text{rank } E_8 = \varphi(30)$) and the Gauss–Bonnet hardening. **Load-bearing.**

(A3) The carrier discriminant exponent (at $M = 41$, $c_3 = 1/(8\pi)$):

exponent	$-\frac{1}{4}$	$-\frac{3}{4}$	$-\frac{5}{4}$	$-\frac{7}{4}$	$-\frac{9}{4}$
α_*^{-1}	137.035995	137.035997	137.0359992	137.036001	137.036003

The exponent moves only the *seventh* digit; all candidates agree to six. Here $-\frac{5}{4} = q(D_5)$ is selected, but this is a *fine* selector, not a hard lock. **Fine (last-digit).**

Ablation verdict — honest

The EM fixed point is a **hard test of two integers** — the budget $M = 41$ and the seam denominator $8 = \text{rank } E_8$, each decisive (a single-unit change misses by ~ 1 in $\alpha_\star^{-1}$) — and a **fine test of one norm**, the carrier discriminant exponent $-\frac{5}{4} = q(D_5)$ (seventh digit). The equation is therefore *not* freely tunable to 137.036: the two integers that hit it are exactly the compiler/bootstrap numbers, fixed elsewhere. What it does *not* do is pin the carrier norm strongly — that is honest, and it is the right level of claim. [E]

Appendix: computational verification

The load-bearing identities of this document (marked [E], [E], [E]) are re-derived from $\{c_3, g_{\text{car}}\}$ alone and machine-checked by the suite in the repository folder `verification/`; the inline tags `[verification/...]` point to the exact script. Relevant here:

- `v1_e8_glue.py` — the μ_4 glue ($q(D_5)+q(A_3)=2$, $\text{disc}=\mathbb{Z}_4$, 240, 248).
- `v2_carrier_pascal.py` — $g_{\text{car}}=5$, $16=1+5+10$, $N_{\text{fam}}, \Omega_{\text{adm}}, b_1$.
- `v3_em_alpha.py` — the EM fixed point $\alpha_\star^{-1}=137.0359992168$ (existence/uniqueness, ablation).
- `v4_flavor_matrix.py` — R with $\det=8$, minors $\{2, 3, 5\}$, $\sum L=40$.
- `v7_gravity_cosmo.py` — scalaron c_3^7 , Λ/A_s readouts.

Run `python verification/run_all.py` (needs `mpmath`, `numpy`, `sympy`); it exits 0 iff all checks pass. Full map: `verification/README.md`.

External works built upon

The TFPT discrete core (the two axioms, the E_8 glue, the integer skeleton) is self-contained and machine-checked. Its *bridges* to continuum/AQFT physics, to gravity and to the lattice/singularity picture build on established mathematics, listed here by topic; the body refers to these by author name.

Modular theory and algebraic quantum field theory.

- M. Takesaki, *Tomita's Theory of Modular Hilbert Algebras and its Applications*, Lecture Notes in Math. **128**, Springer (1970).
- J. J. Bisognano, E. H. Wichmann, *On the duality condition for a Hermitian scalar field / for quantum fields*, J. Math. Phys. **16** (1975) 985–1007; **17** (1976) 303–321.
- K. Osterwalder, R. Schrader, *Axioms for Euclidean Green's functions* I, II, Comm. Math. Phys. **31** (1973) 83–112; **42** (1975) 281–305.
- S. Doplicher, R. Haag, J. E. Roberts, *Local observables and particle statistics* I, II, Comm. Math. Phys. **23** (1971) 199–230; **35** (1974) 49–85.
- R. Haag, *Quantum field theories with composite particles and asymptotic conditions*, Phys. Rev. **112** (1958) 669–673; D. Ruelle, *On the asymptotic condition in quantum field theory*, Helv. Phys. Acta **35** (1962) 147–163.
- R. Longo, *Index of subfactors and statistics of quantum fields* I, II, Comm. Math. Phys. **126** (1989) 217–247; **130** (1990) 285–309.
- Y. Kawahigashi, R. Longo, M. Müger, *Multi-interval subfactors and modularity of representations in conformal field theory*, Comm. Math. Phys. **219** (2001) 631–669.

Conformal nets, vertex operator algebras and lattice constructions.

- I. B. Frenkel, V. G. Kac, *Basic representations of affine Lie algebras and dual resonance models*, Invent. Math. **62** (1980) 23–66; G. Segal, *Unitary representations of some infinite-dimensional groups*, Comm. Math. Phys. **80** (1981) 301–342.
- T. J. Osborne, A. Stottmeister, *Conformal field theory from lattice fermions*, Comm. Math. Phys. **398** (2023) 219–289, [arXiv:2107.13834](https://arxiv.org/abs/2107.13834), [doi:10.1007/s00220-022-04521-8](https://doi.org/10.1007/s00220-022-04521-8). (This is the seam scaling-limit theorem the TFPT continuum leg cites — the Koo–Saleur $\rightarrow$ Virasoro convergence, $c = \frac{1}{2}$, and the smeared WZW-current Theorem 5.1. The internal TFPT code-token “MMST” and the legacy claim-ids `SEAM.MMST.*` / filenames `v*_mmst_*` denote *this* Osborne–Stottmeister theorem.)

- V. Morinelli, G. Morsella, A. Stottmeister, Y. Tanimoto, *Scaling limits of lattice quantum fields by wavelets*, Comm. Math. Phys. **387** (2021) 299–360, [arXiv:2010.11121](#), [doi:10.1007/s00220-021-04152-5](#); and *Operator-algebraic renormalization and wavelets*, Phys. Rev. Lett. **127** (2021) 230601. (The operator-algebraic (wavelet/Wilson–Kadanoff) renormalization *framework* the Osborne–Stottmeister theorem above builds on.)
- M. S. Adamo, Y. Moriwaki, Y. Tanimoto, *Osterwalder–Schrader axioms for unitary full vertex operator algebras*, [arXiv:2407.18222](#) (2024).
- M. S. Adamo, L. Giorgetti, Y. Tanimoto, *Rational and non-rational two-dimensional conformal field theories arising from lattices*, [arXiv:2506.01008](#) (2025); *Representations of conformal nets associated with infinite-dimensional groups*, [arXiv:2508.07109](#) (2025).
- P. Naaijken, D. Penneys, B. Wallick, *(commuting-projector reflection positivity / LTO-RP)*, [arXiv:2605.10693](#).

Lattices and quadratic forms.

- J. H. Conway, N. J. A. Sloane, *Sphere Packings, Lattices and Groups*, 3rd ed., Grundlehren der math. Wiss. **290**, Springer (1999).
- C. L. Siegel, *Über die analytische Theorie der quadratischen Formen* (the Minkowski–Siegel mass formula), Ann. of Math. **36** (1935) 527–606.

Singularities and geometry.

- E. Brieskorn, *Beispiele zur Differentialtopologie von Singularitäten*, Invent. Math. **2** (1966) 1–14.
- J. Milnor, *Singular Points of Complex Hypersurfaces*, Ann. of Math. Studies **61**, Princeton Univ. Press (1968).

Spectral geometry, gravity and thermal time.

- A. H. Chamseddine, A. Connes, *The spectral action principle*, Comm. Math. Phys. **186** (1997) 731–750.
- A. Connes, C. Rovelli, *Von Neumann algebra automorphisms and time–thermodynamics relation in general covariant quantum theories*, Class. Quantum Grav. **11** (1994) 2899–2918.
- R. M. Wald, *Black hole entropy is the Noether charge*, Phys. Rev. D **48** (1993) R3427–R3431.
- H. Casini, M. Huerta, R. C. Myers, *Towards a derivation of holographic entanglement entropy*, JHEP **05** (2011) 036, [arXiv:1102.0440](#).