

TFPT — The Prime Front

Research documentation of the prime / zeta line: seven verified modules, one localized residual, and the complete kill list

Stefan Hamann

Alessandro Rizzo

July 27, 2026 · v5.4 (rev 415)

The TFPT document set — what is treated where

Plain language: TFPT (Topological Fixed-Point Theory) is a small discrete compiler. Two inputs — the seam constant $c_3 = \frac{1}{8\pi}$ and the carrier rank $g_{\text{car}} = 5$ — build $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ and read off the Standard-Model skeleton, the constants and the scale grammar. The series is **the introduction plus five numbered papers and five companions**, best read in order:

0. **introduction** — reading guide, compiler closure, predictions, the dependency DAG and the single proof ledger (`verification/status_ledger.csv`). (*entry point*)
1. **tfpt_1_architecture_e8** — the two axioms, the derivation map, the EM fixed point α^{-1} , the $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ construction, the QBL theorem chain.
2. **tfpt_2_standard_model** — the SM in one φ_0 -ladder formula, flavor from parabolic transport (sheet diamond, dual anchor, Q_+ cohomology), the worked closures.
3. **tfpt_3_e8_audit_bootstrap** — the seven E_8 slices as an audit raster, the cascade bridge, the Möbius bootstrap.
4. **tfpt_4_frontier** — honest status of η_B , m_p/m_e , Koide, dark matter, full QG.
5. **tfpt_5_redteam** — the adversarial audit: declared attacks, what survives, what each reduces to, and the kill tests.

Companions: **R. tfpt_research_contracts** — the open interfaces ($v_{\text{geo}}, G_{\text{net}}, F_{\text{transfer}}$) as numbered contracts; **H. tfpt_horizon_readouts** — Appendix H (unit-system reframe): one seam constant as the universal horizon thermal code, the Nariai/anchor surface, the resummed seam clock; **O. origin_theory** — why no free fundamental number remains (exact core + one honestly-typed cyclic interpretation); **S. tfpt_safeguards** — the verification discipline: every mechanism that defends a claim against coincidence and numerology (status calculus, no-free-pattern, the over-determination map, the firewall, frozen predictions + null model, the independent Wolfram and Lean paths, the red team); **P. tfpt_prime_front** — research documentation of the prime / zeta line: the seven load-bearing modules on the E_8 glue census (Hecke from geometry through the transport ledger), the sandbox localization program that follows them, and the kill list. Documentation, not a result: it makes no claim about the Riemann Hypothesis.

*You are reading the **Prime Front** companion (P): the complete internal record of the prime / zeta line — how a discrete compiler’s bookkeeping kept producing classical number-theoretic objects, which of those links became load-bearing modules, and how the remaining question was compressed, over a hundred exploratory probes, down to one scalar ratio. It is a **research documentation**, not a results paper. Its purpose is that a reader (or a reviewer) can see the whole route, including everything that died on the way.*

Abstract

A two-axiom discrete compiler ($c_3 = 1/(8\pi)$, $g_{\text{car}} = 5$, $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$) has no business meeting the prime numbers. It does anyway: the μ_4 glue completion carries a quartic character, the signed glue census is a theta series, and the shell counts of that census turn out to be *Hecke* data. This companion documents that line end to end. Seven modules are load-bearing and ledger-carried (v535–v541): Hecke from geometry on the E_8 census, the Eichler trace layer, the half-integral bridge, one finite relative-trace identity of which the first three are projections, the Weil structure of the resulting family with two explicitly isolated obstructions, the amplitude route with a positive linear carrier, and the matching-lemma / transport ledger package. Everything beyond those seven is *sandbox*: an exploratory program, run as 108

preregistered probes with 2701 machine checks, which localized the residual object. That localization is the actual content of this document. The transport ledger closes exactly, leaving one coupled inequality (called **I5**) which is — *by identity, as a typing and not as progress* — equivalent to Weil positivity and hence to the Riemann Hypothesis. The subsequent arc maps I5 geographically (Connes–Consani dictionary, tightness curves, the Sonin crossing, the one-atom band), turns the positivity question into a per-zone handover induction, and then certifies its ingredients one by one until the entire remaining hardness sits in a single Loewner statement on the J -odd half of a window, reducible by Sherman–Morrison/Woodbury to one scalar ratio $r = \kappa/\varepsilon \leq 1$ (measured 0.0053–0.1814 on 16 zones) and then, once ε is identified as an exact energy, to two scalars whose open part is a boundary-decay statement about one explicit vector. **No claim of progress on the Riemann Hypothesis is made anywhere in this document**, and none of the sandbox findings moves any ledger marker. The document is deliberately symmetric about failure: a dedicated section lists the readings and routes that were refuted — an essential-singularity reading, a C/g proxy law, a density chain, an accumulated invariant with self-propagation, the Landau–Widom anchor, the symbol route for the odd channel, three certificate candidates for the avoidance norm, and the naive margin route — because in a program of this shape the kill list is the more reliable output.

[E] exact / machine-proven

[C] conditional

[O] open / axiom

[X] kill test

1 How to read this document

Honesty statement — read this before anything else

- 1. No RH claim.** Nothing in this document claims progress towards, or evidence for, the Riemann Hypothesis. Where an equivalence to RH appears (Section 4) it is an *equivalence typing*: a statement about which object carries the difficulty, produced by an exact identity, and deliberately *not* a statement that the difficulty has been reduced.
- 2. Two layers, never mixed.** Claims marked [ledger] are load-bearing: they have a row in `verification/status_ledger.csv`, a module in `verification/`, and a script citation here (the blue [verification/...] tag). Everything marked [sandbox] is exploratory work in `experiments/tfpt-discovery/`; it carries no ledger row, no marker, and no authority. Section 3 is the only section that is entirely [ledger].
- 3. Status markers are quoted, never upgraded.** The markers [E] [C] [O] [X] shown in Section 3 are read verbatim from the ledger rows of v535–v541. This document introduces no new markers and moves none.
- 4. Fits are declared as fits.** Every scaling law, slope, exponent and extrapolation quoted from the sandbox is a fit to finitely many measured windows and is labelled as such at the point of use. Sixteen resolved zones are sixteen zones, not all zones.
- 5. Classical results are named as classical.** Kneser neighbours, Eichler/Witt, Shimura, Waldspurger, Cohen, Weil/Guinand, Bombieri, Yoshida, Connes–Consani, Gronwall/Robin, Douglas, Slepian–Landau–Pollak, Sherman–Morrison/Woodbury, Grenander–Szegő and the rest are classical and are used as such; the compiler-native part is the wiring, not the theorems.
- 6. Numbers are literal.** Every numeric value in this document is transcribed from the research diary `experiments/next.txt` or from a ledger row. Figures that could not be drawn from literal data are marked *schematic* in their caption, and the caption states exactly which quantities in the figure are literal.

The diary parts are cited as **T n** (part n of the research diary, chronological, $n = 11 \dots 108$). The probe file, the contract name, the verdict and the check count of every part are listed in Appendix A. Section 2 explains why a physics compiler meets the primes at all; Section 3 is the verified layer; Sections 4, 5 and 6 are the sandbox program in three acts; Section 7 is the kill list; Section 8 is the honest status.

2 Motivation: why a compiler meets the primes

TFPT is a discrete compiler with two inputs: the seam constant $c_3 = \frac{1}{8\pi}$ (P1) and the carrier rank $g_{\text{car}} = 5$ (P2). They force the lattice completion $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$, and the readouts of the Standard-Model skeleton are taken off that completion. None of this is about number theory. The prime line begins with an accident of the *glue*.

The completion is not $D_5 \oplus A_3$ but $D_5 \oplus A_3$ *plus* a glue group μ_4 — a cyclic group of order four. A cyclic group of order four carries a quartic character, and the quartic character on $\mathbb{Z}$ is χ_{-4} , the non-trivial character mod 4. Once the census of glue vectors is counted *with that sign*, three things happen at once, and all three are elementary consequences rather than choices:

1. The signed glue count is a **theta series**. Signed counting of a quadratic form is a theta series by definition; here the compiler’s own count θ_3^2 is exactly the Gaussian-integer theta, $r_2(n)$, so the census counts $\mathbb{Z}[i]$ -ideals (T85).
2. The character makes the count a **Dirichlet object**. χ_{-4} is the character of $\mathbb{Q}(i)$; the associated L -function is the first Dirichlet L -function, and the split law “ $p = a^2 + b^2$ ” becomes a compiler-visible fibre property of each prime (T21, channel B: χ_4 fibre $\iff p = a^2 + b^2$, verified 81/0/87/0 on $p < 1000$; χ_3 fibre $\iff p = a^2 + 3b^2$, 80/0/86/0).
3. The shell counts of the census carry **Hecke eigenvalues**. That is the surprise, and it is the content of the first load-bearing module.




Figure 1: The chain by which the compiler acquires arithmetic. Nothing here is chosen for number theory: the glue group is forced by the completion, the quartic character is the character of a cyclic group of order four, and the signed count of a quadratic form is a theta series. The green nodes are where the line becomes load-bearing (Section 3).

The honest first question of the series was therefore not “does this prove anything about primes” but “which of these links are *mechanism* and which are pretty coincidences”. Parts T11–T13 established the signed glue theta, its L -series and the glue-character atlas (19/19, 16/16, 11/11); T14–T19 tested six candidate contracts in parallel and produced two self-corrections against the project’s own earlier readings (T14 DEFLATED: the measured 2π is the universal Bisognano–Wichmann/Unruh conversion, not “the self-dual width”). The cross-finding of that batch is worth stating because it survived the entire series: the structure is exact at the census/lattice level (weight 4), and the gap is always the *same* gap — the canonical transport to the $GL(1) / \xi(s)$ world (multiplicity one, weight 1/2, the archimedean place, the derivation of the Hecke action). One meta-contract, ZETA.CENSUS.T0.GL1 [O].

Parts T20–T25 then killed, in four stages, the slogan “the archimedean term comes from the seam”: the raw seam density of states is flat/slightly decreasing (loglog slope ≈ -0.40) while the classical digamma kernel grows logarithmically (measured 0.15917 against $1/(2\pi) = 0.159155$, rms 3×10^{-6} for $t \geq 20$) — KILLED-AS-NAIVE (T20, 25/25); the non-DOS successor located the missing logarithm in the interval cut but failed the one-constant dictionary at $2/\pi$ (T22, PARTIAL); and both dictionary closures plus the scattering-phase observable died (T23–T25, 23/23 + 25/25 + 29/29). This matters for the rest of the document: the archimedean place was never going to come for free, and when it finally arrived (T82) it arrived from *inside* the same family, not from the seam.

3 The verified layer: seven modules

This section is entirely [ledger]. Each subsection states the claim as the ledger states it, cites the module, and quotes the ledger’s own status markers. Nothing here depends on Sections 4–6.

The seven modules and their ledger markers			
Module	Claim id	Content	Markers
v535	HECKE.GEOM.01	Hecke from geometry on the E_8 census channels	[E] + [C] + [O]
v536	HECKE.GEOM.EICHLER.01	Eichler trace layer at the E_8 lattice	[E] + [C] + [O]
v537	HECKE.GEOM.HALFINT.01	half-integral bridge	[E] + [C]
v538	HECKE.GEOM.RTF.01	compiler relative trace identity	[E] + [C]
v539	RTF.GNS.WEIL.01	Weil structure of the compiler family	[E] + [C]
v540	RTF.GNS.AMP.01	amplitude route and positive linear carrier	[E] + [C]
v541	RTF.GNS.LEDGER.01	matching lemma and transport ledger package	[E] + [C]
Marker key: [E] exact / machine-proven · [C] conditional (holds under named hypotheses) · [O] open / axiom · [X] falsifiable kill test. Per-claim fine type in verification/status_ledger.csv. Every row carries NO marker moves in the ledger. The [O] on v535 and v536 is the same open gate throughout: $\text{weight-4} \rightarrow \text{GL}(1) \text{ transport}$.			

3.1 Hecke from geometry (v535)

The Hecke structure of the μ_4 -glue census is lattice-native: it comes out of Kneser p -neighbour geometry, not out of modular input [verification/v535_hecke_from_geometry.py] (25 checks, ~ 11 s; consolidated promotion of T27/T31/T32).

- **Kneser carries Hecke** [E]: $\#\text{iso_pts} = p^7 + p^4 - p^3$ and $\#\text{iso_lines} = \sigma_3(p) \cdot \#\mathbb{P}^3(\mathbb{F}_p)$ identically and by full enumeration at $p = 2, 3, 5, 7$ (135/1120/19656/137600 lines). The Eisenstein eigenvalue $\sigma_3(p) = 1 + p^3$ is an exact factor of a lattice-native correspondence count.
- **The marked neighbour sum is an affine Hecke element** [E]: $\nu_p = a \text{Id} + b T_p$ exactly (overdetermined, residual 0), with $b = \sigma_3 + a_p$; the cusp rule $a_p = b - \sigma_3$ recovers $a_3 = -4$ and $a_5 = -2$.
- **Census redundancy is 2-adic oldform structure** [E]: $\dim V = 7 = 5+2$, $\pi_{\text{cusp}} = (28 - T_3)/32$, only 2-adic levels occur.
- Fences carried *inside* the claim: the $p = 5$ geometric $S[1]$ block is a T31-validated *freeze* [C]; the $\text{weight-4} \rightarrow \text{GL}(1) \text{ transport}$ is [O].

3.2 The Eichler trace layer (v536)

The smooth (Eisenstein) background and the coherent (cusp) interference separate exactly [verification/v536_eichler_trace_layer.py] (23 checks, ~ 30 s; promotion of T33/T36/T37/T42). The Witt layer gives $\lambda_{\text{Eis}} = \sigma_3(\#\mathbb{P}^3 - \sigma_3) = L - \sigma_3^2$ identically (anchors 336, 3780, 19264 at $p = 3, 5, 7$) [E], and the Eichler identity $\lambda_{\text{geom}} = \lambda_{\text{Eis}} + a_p^2$ holds two-sidedly (352/3784/19840). The signed cusp reading is $a_p = -c(p)/8 \Rightarrow (-4, -2, +24)$, and the assembler $R(p) = a_p^2$ is exact for $p \leq 31$. The $p = 5$ type-(ii) block is again a validated freeze [C]; the transport gate stays [O].

3.3 The half-integral bridge (v537)

In the 70-element compiler weight-5/2 theta monoid there is *exactly one* object g with $\text{Sh}_{t=2}(g) = -8f_8$ coefficientwise [E], it is an exact $T(p^2)$ -eigenform with the weight-4 eigenvalues $(-4, -2, 24, -44, 22)$ [E], and $U_4(g) = 0$ places it outside the Kohnen plus space — a scope fence, not a defect [E] [verification/v537_halfintegral_bridge.py] (20 checks, ~ 90 s; promotion of

T38/T41/T44/T45). The Waldspurger/Baruch–Mao constant $R = 23.1873585645\dots$ is constant on ten discriminants $d \equiv 1 \pmod 8$ (spread $\leq 10^{-12}$), typed [C] because it is an AFE-numeric measurement.

3.4 One finite relative trace identity (v538)

The strongest structural statement of the verified layer: v535, v536 and v537 are three *projections* of one finite relative-trace identity of Jacquet type [verification/v538_relative_trace_identity.py] (18 checks, ~ 14 s; promotion of T53). $\text{Tr}_V(\nu_p \circ \pi)$ is computed twice, independently — geometrically from the glue census with no modular input, and spectrally from $\eta(2\tau)^4 \eta(4\tau)^4$ with no geometry — and the two agree exactly for all $(p, \pi) \in \{3, 5, 7\} \times \{\text{Id}, \pi_{\text{Eis}}, \pi_{\text{cusp}}, \pi_{E_4}, \pi_{f_8}\}$ (anchors, e.g. $p = 7$: full trace 727680, f_8 channel 19840) [E]. The Eichler split *is* the orbit decomposition: principal orbit $a \text{Id} \rightarrow \lambda_{\text{Eis}}$, elliptic orbit $b T_p \rightarrow a_p^2$, cross terms cancel exactly. And the period side is literally compiler geometry: g is the quaternary lattice form $n = (x^2 + y^2)/2 + 2z^2 + u^2 + 2w^2$ with sign $(-1)^{|u|+|w|}$, giving $[\text{signed lattice count}]^2 = R |d|^{3/2} L(f_8 \times \chi_d, 2)$ at relative 10^{-12} [C]. The identity is *finite*; the infinite RTF is explicitly open.

3.5 The Weil structure of the family (v539)

The family carries a fully identified Weil structure *up to two explicitly isolated obstructions*, and the obstructions are promoted content, not a footnote [verification/v539_weil_structure_family.py] (25 checks, ~ 6 s; promotion of T55/T61/T62/T63). The canonical GNS pairing is a direct integral over the Gelfand spectrum with fibre mixing $\leq 5.04 \times 10^{-16}$; the trivial Sato–Tate isotype is algebraically a $\text{GL}(1)$ core, $G_0 = (1 + Y)/(1 - Y)$; the prime side is an exact linear relation to shifted ζ -Weil forms at relative $\leq 9 \times 10^{-16}$. The two obstructions are (i) a sign/doubling residue (metaplectic, sym^2) that survives every preregistered canonical projection, and (ii) a Plancherel correction which is irreducibly non-automorphic. Finite-class positivity of Q_{fam} is measured [C] ([4.369, 11.486]); dense-class positivity is open. The ledger row says it in one line: *not almost-RH*.

3.6 The amplitude route and the positive linear carrier (v540)

The route out of the square plane [verification/v540_amplitude_linear_carrier.py] (34 checks, ~ 3 s; promotion of T67–T72). An amplitude Dirac operator $D = \begin{pmatrix} 0 & V \\ V^\dagger & 0 \end{pmatrix}$ with $V[d, m] = \sqrt{w_d} b(d) \chi_d(m)$ exists and is Hecke-equivariant [E]; the geometric polarisation $b = N_+ - N_-$, $\Theta = N_+ + N_-$ is lattice-exact with $N_\pm \geq 0$ for all $n \leq 50\,000$ [E]; the Cohen seed $\Theta(d) = -48L(-1, \chi_d)$ is exact-rational on 159 live d [E]; the prime side is a *pure plus* combination $Q = Q_\zeta(g_-) + Q_\zeta(g_+)$ at relative 2×10^{-15} [E]; the functional equation of the linear family holds at relative 1.4×10^{-40} [E]. The open boundary is *inside* the claim: the gap functional λ^* on $n \equiv 6 \pmod 8$, with an exact Farkas witness (0.0511), is a named limit, and positivity is asserted only in the Euler region.

3.7 The matching lemma and the transport ledger (v541)

The largest module of the line [verification/v541_matching_lemma_ledger.py] (33 checks, ~ 10 s; promotion of T78–T82 and T85; every core check recomputed, not cited).

- **Window proof** [E]: the matching lemma is proved exact-integer on $[4, 10^6]$ — $7S(j) < 40A(j)$ at every atom, 0 violations over 939 870 clash atoms, no-overflow proven in exact integers, 268 big-integer rechecks; exact margin $X = 22242575693/270725400000 = 0.082159$ at $j^* = 554400$; $\rho_{\text{crit}} = 1.143989 > 21/20$. A maximal-greedy identity makes the certificate uniform in the target set and the greedy weights.
- **Transport ledger** [E]: $Q_{\text{Weil}} = Q_{\text{cert}} + \Delta_{\text{arch}} + \Delta_2$ closes with residual 4.4×10^{-16} on a 24-row battery, and $\Delta_{\text{pole}} \equiv \Delta_{\text{conv}} \equiv 0$ are *proven*, not measured (kernel collapse in **sympy**, exact scale invariance).

- **Character-exact envelope [E]:** $-\psi = (\chi_{-4} + \frac{1}{4}\chi_8 + \frac{1}{4}\chi_{-8})\Theta$ coefficient-exact for all odd $n \leq 50000$; confinement holds as a *set equality* on 10^6 .
- **Archimedean term internal [E]:** the Legendre duplication bridge $(2\pi)^{-s}\Gamma(s) = \frac{1}{2}\Gamma_{\mathbb{R}}(s)\Gamma_{\mathbb{R}}(s+1)$ and the kernel identity $k_{\zeta} = K_{\text{fam}} - K_{\text{shift}}$ at 7.9×10^{-31} pointwise.
- **Frontier facts [C]:** unlifted coherent chain crossing at $k^* = 14$ ($\log_{10} N^* = 23.1$); Robin tail factor 6.16 at $J = 10^6$ and divergent — the constant route stays open.
- **Named limits as content.** Two limits are part of the claim rather than hidden by it: one classically-shaped open lemma (correlated cancellation on credit-rich non-coherent tail atoms), and the I5 equivalence typing discussed next. The row states explicitly: *NOT “almost RH”*; ZETA.HP.CARRIER untouched.

4 The localization program (sandbox)

Provenance

Everything from here to Section 8 is [sandbox]: exploratory probes in `experiments/tfpt-discovery/`, no ledger row, no marker, no authority. The findings are reported because the *shape* of the residual is the useful output, not because any of them is established.

4.1 I5: the ledger closes and one inequality is left (T79)

The transport wall — the thing that had blocked the line since T20 — was fully decomposed in T79 (`transport_ledger_probe.py`, 22/22). The key observation is that the odd-prime side of the Weil functional agrees *exactly* with the certified plus combination (relative 2.6×10^{-16}): the prime side was never the wall. What remains after the ledger closes is a minimal transport set of two exact identities (proved), two classically-provable bounds, and exactly one coupled inequality:

$$(I5) \quad Q_{\text{cert}}(h) + \Delta_2(h) + A_{\text{fam}}(h) - A_{\text{shift}}(h) \geq 0 \quad \text{for all autocorrelations } h.$$

I5 is the prime $\leftrightarrow$ archimedean coupling. By the closed ledger it is, as an identity, equivalent to Weil positivity and hence to RH. The diary is explicit about what that means: *the built-in safeguard fired*. RH does not follow from finite bookkeeping; the hidden infinite step is exactly I5 — and the value of T79 is that the step was *found and named* rather than concealed. This is an equivalence typing, not progress.

T82 then changed I5’s *type* (`heat_arch_probe.py`, 22/22, ARCH-INTERNAL): Δ_{arch} is exactly the internal Γ difference $A_{\text{fam}} - A_{\text{shift}}$ via Legendre duplication, verified on 18/18 battery rows at max relative 6.6×10^{-15} . I5 stopped being a coupling between two worlds and became a *self-consistency statement about one heat family*. Again: a type change is not a proof.

4.2 The Connes dictionary (T87) and where positivity is anchored

T87 (`i5_connes_dictionary_probe.py`, 22/22, DICTIONARY-EXACT-CORE) identified the I5 one-family form as a compiler instance of the semi-local Connes structure. Four exact matches: the two-shell orbit route equals Connes’ expression at relative 6.3×10^{-16} ; the orbit kernel of Connes–Consani (2021) is literally the compiler’s Poisson ladder at $\rho = e^u$ (`sympy` plus 1.7×10^{-50}); h_+ is by definition the compiler kernel $k_{\zeta} = K_{\text{fam}} - K_{\text{shift}}$; and $k_{\zeta} = 2\theta'_{\text{RS}}$ — the derivative of the *Riemann–Siegel phase*, at relative 0.0. The dihedral shift is the scaling group $\mathbb{R}_+^*$.

The most useful part of T87 is not a match but a *boundary*. The two positivity programs are complementary: Connes–Consani anchor positivity *before* the primes (the prime-free Sonin window, support $(\frac{1}{2}, 2)$, i.e. $u \in (-\log 2, \log 2)$), the compiler anchors it *after* the primes (atom certificates, all supports). The window boundary is exactly the first prime atom $u = \log 2$ (`sympy-exact`; on narrow rows the prime side, Q_{cert} and Δ_2 all vanish identically). The sectors *touch* and do not overlap; the I5 coupling lives in the crossing region between them, outside both proved sectors.

4.3 The tightness map (T88) and the crossing (T89–T91)

T88 (`i5_tightness_probe.py`, 27/27) mapped where I5 is actually tight: a band-limited zero plateau (classical), safe zones, and *eight* near-vertical tight curves whose spacing follows the Γ -side density law (ratio 1.01 ± 0.08 , computed zero-free — no spectral identification is claimed or made). There are no negatives on genuine autocorrelations; the earlier T76 negatives were the shadow of missing archimedean/ $p = 2$ bookkeeping, exactly as the ledger predicted.

T89 (`sonin_crossing_probe.py`, 19/19) established that the window compression of the Weil form *is* Bombieri’s classical object, and answered the sharpness question honestly: the boundary at $\log 2$ is *not* sharp. The margin falls smoothly from $a \approx 0.5$ towards the floor with no structure at $\log 2$; the prime-free sector still has margin 0.557 at its own boundary and keeps ≥ 0.25 of the reference comfort to $a \approx 0.75$ (+0.057 beyond the proved boundary). The residual subspace grows softly $0 \rightarrow 1 \rightarrow 2 \rightarrow 3 \rightarrow 4$ (out of 32) and carries the global minimum. T90 (`residual_dissection_probe.py`, 17/17) made those residual vectors explicit — n -stable to $\leq 2.1^\circ$, closed Gaussian $\times$ cos/Gaussian $\times$ sin fits with 99+% capture — and decided the core question: the residual is *not* just pole coupling (3/10 vectors are controlled by no structure at all; beyond $a \approx 0.85$ genuinely pole-free atom $\leftrightarrow$ arch content remains).

T91 (`thin_band_analytic_probe.py`, 19/19) identified the band as the *one-atom zone* and found the mechanism that dominates the rest of the document: beyond an inner edge the prime atom $u = \log 2$ becomes *load-bearing* — the prime *rescues* the form where the archimedean margin is exhausted (rescue term $+0.22 \dots + 0.41$, purely odd sector). Two uncertainty constants were decided in closed form: $a \cdot t_{\text{rms}} \rightarrow \pi$ (Wirtinger, 0.3%) and $a \cdot t_{\text{cent}} \rightarrow 2 \text{Si}(\pi) - 4/\pi = 2.4306$ (two independent routes, 10^{-25}).

Data integrity: the T92/T93 correction — reported because it happened

T92 (`zone_extension_proof_probe.py`, 36/36) failed to prove the lattice zone extension and said so (T-SKELETON): what it certified is small and real ($Q \geq 6.7 \times 10^{-12} > 0$ on an 8-dimensional sine-window subspace across the whole band, with an error certificate and a 1057-point covering), and it named two independent walls — $\lambda_{\min}$ collapses geometrically (factor 11.4 per mode), and the high-mode complement would need $N^* \gtrsim 5069$ modes. Verdict: *the finite-block route is structurally the wrong instrument, not under-resourced*. In the same run T92 raised a calibration discrepancy against T89/T91, and the entire band map was put under reservation. T93 (`band_reconciliation_probe.py`, 41/41) resolved it with a third independent implementation: one diary prose value (6.98) had been stale, one discrepancy was a factor-2 convention plus a 1% grid interpolation artefact, and exactly one real bug sat in T92 itself. Net effect: the T89/T91 map survives, with a_{neg} recalibrated $0.7410 \rightarrow 0.7486$ and T92’s own band scoping withdrawn. The corrected band map: classical edge $0.34657 \cdot$ prime-free crossing $0.3723\text{--}0.3763 \cdot$ band top $\alpha^* = 0.46265 \cdot$ second atom $\log 3/2 = 0.54931$.

4.4 The blind prime demonstration (T94)

A separate, deliberately modest exercise. T94 (`prime_blind_demo_probe.py`, 16/16, BLIND-100) used the exact geometric criterion “odd n is prime $\iff r_4(n) = 8(n+1)$ ” (Jacobi 1834, on the $\mathbb{Z}^4 = \text{rank-}|\mu_4|$ theta tower) to predict all 753 primes of the never-seen window $[1\,000\,001, 1\,010\,000]$ — count and MD5 declared *before* any ground truth, then evaluated: exactly 753, zero false positives, zero false negatives, 100.00% on all 10^4 integers, with an AST-enforced prediction path containing no divisibility test (no `isprime`, no sieve, no `%`, `//` or `/`). The cost honesty is part of the result: a sieve wins by $\sim 820\times$ (1 ms against 0.83 s). This is structure completeness, not an algorithmic advance — and the *spectral* prediction (zeros $\rightarrow$ primes) remains bound to I5/RH and is not claimed.

5 The handover induction (T95–T101)

The band work left a picture: each prime atom rescues its own zone and hands over to the next. T95–T101 turned that picture into an induction, and then measured whether the induction can

run.

5.1 The relay is real (T95, T96)

T95 (`continuum_extension_probe.py`, 28/28) proved the C1 chain unconditionally ($|h_f(\log 2)| \leq 1/2$ by disjoint supports with margin 0.2305; the windowed symmetrised shift has $\|S\| = 1/2$ exactly, 1.1×10^{-16}) and found that the binding mechanism is *not* the two-bump extremal (share 0.046) but the atom rescue: above $\alpha_{\text{free}} = 0.371654$ the prime-free minimiser has $h(\log 2) < 0$ down to -0.328 , so the atom adds positivity exactly in the losing direction.

T96 (`relay_test_probe.py`, 21/21) is the model of a self-correcting run. The apparent “edge” of T95 at α^* was a map artefact: the value $+7.8 \times 10^{-6}$ reproduced exactly, but it sits on a smooth, strictly positive curve with no feature at all ($\lambda_{\min} > 0$ on the whole of $[0.38, 0.86]$, decay law $\lambda_{\infty} \sim \exp(-49.2\alpha)$). The *relay*, however, is real — and it lives entirely in the counterfactual: remove the $\log 3$ atom and $\lambda_{\min}$ crashes to -0.445 , with the loser being exactly the anti-double-bump at distance $\log 3$ (alignment -0.99) and a rescue identity at 5×10^{-15} . All four handover windows are positive: each atom arrives strictly *before* it is needed ($+0.025/ +0.009/ +0.011/ +0.007$ for $k = 1.4$). Three consequences were recorded: it is a margin problem, not an edge problem; numerics as a witness is exhausted beyond $\alpha \approx 0.55$; and the counterfactual is the proof target, because it works on $O(0.1)$ -sized quantities instead of 10^{-6} .

5.2 The induction step and its one missing lemma (T97, T98)

T97 (`relay_induction_probe.py`, 105/105) gave the induction step proof shape. Alignment is true and sharp with a constant law; the $t = 0$ killer loss is proved by three structure identities (old atoms vanish identically on E_- ; the pole rescue flips with factor $1 - \cosh(u/2)$; $|\hat{f}|^2 = 2(1 - \cos(tu))|\hat{g}|^2$), giving $k_{\text{eff}} = (1 - \cos(tu))k(t)$ with infimum $-1.11 \dots -2.59$ instead of $k(0) = -5.37$. The structural pearl: E_0 is *literally* the same form at a smaller window (7×10^{-14}) — the induction hypothesis, by self-similarity. What remained was a single cross-block lemma about a single vector.

T98 (`cross_lemma_probe.py`, 44/44) attacked it and produced the most instructive negative result of the series: the measured $\sqrt{\lambda_0}$ law is Douglas (1966) range inclusion, i.e. a *tautology* of the positivity one is trying to prove — the identification *dissolves* the lemma rather than proving it. Three T97 premises were refuted in the same run. The replacement target is an exact scalar inequality with no constants and no near-null vector:

$$D_k(\alpha) \quad := \quad -\lambda_{\min}(Q|_{E_-} - Q_{-0}(Q|_{E_0})^{-1}Q_{0-}) \quad \leq \quad \frac{\mu_k}{2}.$$

T99 (`dk_recursion_probe.py`, 23/23) then found the a-priori structure: an exact mirror-parity selection rule ($J_-Q_{-0}J_0 = -Q_{-0}$ at 2×10^{-15}) giving $D_k = \max(D_{\text{even}}, D_{\text{odd}})$, with the binding even channel coupling only to the *odd* spectrum of the smaller window — so the fragile near-null mode is identically absent from the binding channel. T100 (`bulk_remainder_probe.py`, 27/27) closed the bulk remainder from 11/24 to 24/24 samples and showed the 10.8% drift was a grid artefact.

5.3 The race (T101)

T101 (`k_asymptotics_probe.py`, 31/31, CROWDING-TRENDS) resolved 16 zones ($n = 2 \dots 29$) and separated two questions that had been fused. The *mathematics* trends self-supportingly: the primitive quantities are flat over 16 zones (w/g slope $+0.002 \pm 0.178$; the relative D_k margin does not trend; $D_k \leq \mu_k/2$ holds 64/64 and never fails). Atom *strengths* do not fall ($\mu \sim n^{-0.087}$), only the *spacings* do ($n^{-0.615}$, exact) — pure arithmetic. What loses the race is the *closure instrument*: the demand ratio r_k decays like $\exp(-0.1622k)$ (fit, ± 0.0562). The collapsed one-parameter law $w_k = 0.0838 \cdot (\text{atom spacing})/\mu_k$ reduced the scatter from $8.85\times$ to $2.66\times$ with zero residual trend. T101 also stated what an asymptotic theorem would need, and that list has not changed since: (A) the collapsed lower bound $w_k \geq c g_k/\mu_k$ with a fixed $c > 0$ — a sharp, non-perturbative, *arithmetic*




Figure 2: *Schematic*. The instrument race over the 16 resolved zones. *Literal*: the two decay slopes (-0.1622 ± 0.0562 measured in T101; -0.0748 ± 0.0116 measured in T103, a factor 2.2 flatter), and the T103 endpoints $r_1 = 9.33 \rightarrow r_{16} = 2.70$ with the statement that the improved instrument never leaves the spectrum. *Schematic*: the straight-line interpolation in $\log_{10}$, the common anchor of the two lines, and the position of the “spectrum floor”. Both slopes are declared fits in the diary. What loses the race is the instrument, not the inequality: the true inequality $D_k \leq \mu_k/2$ holds 64/64 and its relative margin does not trend.

lower bound at a Weil-positivity edge, with Taylor arguments forbidden by the $\exp(-c/\delta')$ onset; (B) a uniform relative margin; (C) a replacement for the bulk remainder; (D) a finite check of the low zones. If crowding ever wins, the hardness sits entirely in (A).

6 The certification arc (T102–T108)

The last seven parts stopped improving measurements and started converting them into certificates — and, where that failed, into explicit refutations.

6.1 The mechanism: a crossing, not a singularity (T102)

T102 (`arithmetic_bound_probe.py`, 42/42, MECHANISM-IDENTIFIED) computed the mechanism exactly. The k -th atom acts on $E_-/E_0/E_+$ as $\text{diag}(-\frac{1}{2}, 0, +\frac{1}{2})$, so $Q_{k-1} = Q_{\text{full}} - \frac{\mu_k}{2}P_- + \frac{\mu_k}{2}P_+$: the atom strength μ_k enters exactly once, linearly. That gives a two-sided sandwich through the Schur profile $\sigma_k(\delta)$, with $\mu_k/2 \leq \sigma_k$ implying that the handover holds, and $2w_k$ lying between the crossings in 16/16 zones.

Two corrections came out of the same run. First, the onset is *anchored* at $\delta_c = 2w_k$ — a crossing of two finite quantities ($R^2 = 0.968$) — and exponential-singular versus power behaviour is not separable; T96’s essential-singularity reading is compatible but no longer distinguished. Second, and more consequentially, the binding constraint *flips*: no concentration condition binds. The bare E_- form is strongly positive ($2.65 \dots 3.52$, i.e. $4\text{--}14 \times \mu_k/2$), the pole flip and band mass saturate their classical ceilings (96.9–97.1% of Cauchy–Schwarz and Landau–Pollak/prolate respectively), and the onset is manufactured *entirely* by the Schur dressing against $E_0 \oplus E_+$, which eats 35.7%–97.3% of the bare eigenvalue. The loser does not live in E_- .

The same run performed the most useful kill of the arc: a triple decomposition test showed that the T101 law in C/g was a *proxy*. g_k is causally impossible (the form Q_{k-1} is blind to the next atom), statistically dispensable (a causal law $C \cdot u$ has CV 23.3% against 27.6%), and arithmetically only a ceiling ($C_k \leq g_k \mu_k$ exactly; the 16-zone extrapolation violates the ceiling from $k = 69$; checked over 18 120 prime-power atoms to $n = 200\,000$).

6.2 The instrument, rebuilt (T103), and the two doors

T103 (`instrument_probe.py`, 29/29, INSTRUMENT-IMPROVED) rebuilt the closure instrument with certified weights $w_j \geq 1/\lambda_j$ and a θ -weighted band sum, reproducing both anchors exactly. With




Figure 3: *Schematic*. The crossing mechanism of T102. *Literal*: μ_k enters exactly once and linearly, so the condition is a two-sided sandwich through the Schur profile $\sigma_k(\delta)$ against the flat level $\mu_k/2$; the window width $2w_k$ lies between the crossings in 16/16 zones; the onset is anchored at $\delta_c = 2w_k$ with $R^2 = 0.968$; the candidate bound $w_k \geq A_k \mu_k^{1/q_k}$ holds 16/16 with ratio 0.749–0.940. *Schematic*: the curve shapes and all absolute scales.

one fixed, k -uniform instrument ($\Lambda_0 = 3$, $r = 2$, no per-zone tuning) the closure map jumps from 7/64 to 44/64 and the race slope halves (Figure 2). The honest price is explicit: the number of explicit modes grows $2 \rightarrow 232$. And the verdict relocated the remainder: it is no longer the bulk (the bulk is measurably not low-rank — effective rank up to $0.579 \dim E_-$) but the *wing slack* $S = 1 - \rho$, which falls $0.2091 \rightarrow 0.0392$.




Figure 4: *Schematic*. The closure map. *Literal*: the grid is 16 zones $\times$ 4 wing fractions = 64 cells; the old instrument closes 7 of them, the rebuilt one closes 44 with a single fixed k -uniform setting ($\Lambda_0 = 3$, $r = 2$) and no per-zone tuning. *Schematic*: which cells are shaded — the diary records the totals, not the pattern; the shading is drawn as a monotone wedge purely to visualise the counts.

The convergence recorded at the end of T102 is the pivot of the whole arc: T103's wing slack $S = 1 - \rho$ and T102's Schur dressing are *the same object seen from two sides*. Both doors lead to the wing near-null direction.

6.3 Inductive chains, and the death of the margin route (T104)

T104 ran as *two independent arms* after a file collision between two parallel workers was resolved by keeping both implementations of the same contract — `schur_profile_bound_probe.py` (21/21) and `schur_profile_chain_probe.py` (47/47). All core findings converged, which makes it a de-facto independent replication.

- **The naive margin route is dead:** $M \geq m \cdot \text{Id}$ holds in 0/16 zones; the coupling mass to margin ratio is 10^3 – 10^6 ; m vanishes in the continuum like $M^{-1.7}$ (fit).
- **The mechanism is avoidance:** the coupling B_- decouples from the soft bulk modes — the softest mode carries 0.00% of the coupling mass for $n \geq 3$, and the mass sits in mid/high bands (lowest band only 2–12%). Positivity is not saved by a margin but by structural avoidance.




Figure 5: Two doors, one object. Both entries are literal diary findings; the convergence statement (“the same object from two sides”) is quoted from the T102 closing note.

- **Exact spectral-split chains close 16/16** with finite data: arm A via the exact block-inverse identity with matrix caps in PSD order (headroom 1.15–1.44, reach $\gamma_{\max} = 0.50$ –0.90); arm B via a threshold split with an $O(1)$, resolution-stable threshold $w = 2.47$ –5.90 — but requiring $r_{\min} = 64 \dots 1024$ soft modes $\propto M$, which makes the continuum statement a spectral-density condition.
- A recommendation from T96/T103 was tested and *rejected*: the prolate wing basis loses against the raw basis in both arms.

Consequence: the hard core migrated from σ_k as a whole to four named ingredients — a lower bound on $\text{bare}_k = \lambda_{\min}(Q_{\text{full}}|_{E_-})$, the soft dressing scalar L , finite induction data, and a one-sided edge estimate instead of a fit. The chain around them is classical (Schur complement, exact block inverse, Parseval/Bessel, Weyl) and fit-free.

6.4 The certified bare bound and the avoidance theorem (T105)

T105 (`bare_wing_bound_probe.py`, 28/28, ONE-OF-TWO) delivered one of the two missing ingredients in closed form. First it resolved the T104 arm discrepancy: *bare* depends only on (u_k, δ) — the window centre cancels ($\leq 0.74\%$ drift under $8\times$ refinement) — with the exact split $\text{bare} = \mu_k/2 + b_0$. Then three classical steps (Bessel; a Legendre/level-set optimisation at $k(\pi/\delta)$; Cauchy–Schwarz at the pole pair) collapse to a closed lower bound:

$$b_0 \geq \frac{\delta}{\pi} \int_0^{\pi/\delta} k(t) dt - \delta K_{\max} - 4 \left(\cosh \frac{u}{2} - 1 \right) \sinh \frac{\delta}{2}.$$

This is the band average of the archimedean kernel over the band conjugate to the wing, of size $\sim \log(1/2\delta) - 1$. It is positive in 16/16 zones (1.27–3.54) at 81.1–92.7% of the measured value (median 92%), it sharpens as the wing flattens, and it needs no eigenvalue or induction data.

Second, the *avoidance law* became a theorem rather than an observation: $Q_{\text{full}} \geq 0 \Rightarrow \|B^T v_i\|^2 \leq w_i \Lambda_-$ (a semi-inner-product argument; coupling proportional to the mode eigenvalue), verified on every mode of every zone with worst ratio 0.377. That explains the measured 0.00% avoidance exactly. Alongside it, an exact parity superselection: $JQJ = Q$ (5.7×10^{-15}) and $JB = -BR$ (3.4×10^{-15}) — the two channels do not mix, and the softest mode is *J*-even in 16/16 zones. Notably the *Euclidean* angles are not small (cosine up to 1.0): the avoidance lives in the form metric, as a Friedrichs angle.

What T105 left is one Loewner statement: $Q_{\text{full}}(\alpha) \succeq (\mu_k/2)P_-^{(k)} \iff \rho \leq 1 - (\mu_k/2)/\text{bare}_k$ — a Friedrichs-angle / spectral-density condition, explicitly *not* a margin condition. The free cap

$L \leq \Lambda_- = 5.63\text{--}7.52$ diverges like $\log(1/D)$, so the certificate has to come from the spectral density near the soft end.

6.5 Parity superselection: the pole splits (T106)

T106 (`friedrichs_angle_probe.py`, 32/32, DENSITY-MAPPED) killed two routes and found the structural result of the arc. The kills are in Section 7. The finding:

$$ab^\top + ba^\top = ss^\top - tt^\top \quad (10^{-18}; J a = b)$$

The rank-2 Weil pole splits *exactly* into a positive rank-one lift in the J -even channel and a negative rank-one pressure in the J -odd channel. The Toeplitz part has exactly one negative eigendirection; it is J -even, and $ss^\top$ removes it precisely there. The odd channel never needed the pole at all.




Figure 6: Parity superselection (T106). All numbers are literal; the layout is illustrative. The practical consequence: it is not the density that decides the angle but the *avoidance* — the dangerous channel is the one with the better density. T105’s certified bare bound sits 16/16 exactly in the hard channel, and ρ_- is resolution-stable ($\leq 17.6\%$).

The channel-wise angle chain closes 16/16 in the even channel and 9/16 in the odd channel, against only 3/16 unsplit; all 16 handovers close at operating depth in both channels under the exact Schur criterion, with certified b_0 at 89–93%. Certified after T106: bare, the parity split, the pole splitting, the channel inertia, the avoidance law, the growth inequality, the accumulation bounds. Measured: ρ_- .

6.6 One scalar ratio (T107)

T107 (`odd_channel_closure_probe.py`, 30/30, SCALAR-TRACTABLE) reduced the matrix statement (R) to a scalar one. Sherman–Morrison gives exactly (R) $\iff s^* \leq 1$ (identity at 3.4×10^{-11} ; equivalence confirmed 16/16 against Rayleigh–Ritz; precondition $G \succ 0$ via Cholesky 16/16) — but s^* is *saturated* (0.99885...0.99999, with only $n = 2$ at 0.2569), so no uniform s^* chain can work. The Woodbury decomposition rescues it: $s^* = \tau + \kappa$ and

$$(R) \iff r := \frac{\kappa}{\varepsilon} \leq 1, \quad \varepsilon := 1 - \tau, \quad \text{measured } r = 0.0053 \dots 0.1814.$$

Two orders of magnitude of headroom instead of five decimal places, and s^* is resolution-stable (spread $\leq 2.60\%$ over $M = 600/900/1200$). The synthesis is stated with the usual asymmetry: certified-closed 0/16, measured 16/16 (with $q = 1.31\text{--}4.16$ at $M = 1200$, replacing T106’s 0.747–1.156 — a factor 1.7–2.5 gain). Exactly one object is missing: a certified positive lower bound for $\varepsilon = 1 - \tilde{t}^\top T_{\text{odd}}^{-1} \tilde{t}$ — how far the pole vector is from exhausting the odd channel — or, since ε and κ

both fall with M ($\varepsilon \sim M^{-1.99\dots-1.69}$, fit), a certificate for the *ratio* r itself. The named route is via a wing statement of the T105 type plus an avoidance norm, explicitly not via Szegő.

6.7 The ε identity, and two scalars (T108)

T108 (`ratio_certificate_probe.py`, 44/44, EPSILON-IDENTITY) turned the missing positivity of ε from a bound to be won into an *identity*. Everything hangs on the pole response $x := T_{\text{odd}}^{-1}\tilde{t}$, and three exact identities (verified at 10^{-11} – 10^{-13}) pin it down: $\varepsilon = x^T Q|_{\text{odd}} x / \tau$ — so ε is a $Q|_{\text{odd}}$ -energy, not an accident of subtraction — $1/\varepsilon = 1 + \tilde{t}^T Q|_{\text{odd}}^{-1} \tilde{t}$, and the overlap of ε and κ is exactly one rank-one term hh^T/ε , so that (R) $\iff \hat{r} = (\mu/2)\lambda_{\max}(\Gamma + hh^T/\varepsilon) \leq 1$. The third route supplies the classical reading: $\tilde{t}$ is exactly two-term geometry, τ a 2×2 Christoffel–Darboux form, and ε exactly the square of the last Cholesky pivot — the Szegő/Levinson prediction error. *The positivity of ε coincides with the induction positivity in the odd channel*; it is not a second, independent thing to certify.

Two consequences, and one correction of the previous picture. The consequence: (R) reduces from an $M/2$ -dimensional Loewner statement to **two scalars** — $\omega < 1$ (measured 0.2366–0.7531, 16/16; the T105 structure survives the parity split verbatim, $V^T S|_{\text{odd}} V = -\frac{1}{2}I$ at 4.9×10^{-14}) and one statement at the vector x . The chain (R) $\Leftarrow [\omega < 1] \wedge [\lambda_{\min}(Q|_{\text{odd}}) \geq (\mu/2)\tau\theta/(1-\omega)]$ has C1 closing 16/16 (margin 5.5–178; ≥ 1 at all 48 ladder points, 2.7–343), C2 15/16, C3 0/16 — the first time a margin chain in this arc is *not vacuous*: the induction margin it needs is explicitly 9.3×10^{-8} – 3.7×10^{-3} , i.e. two to six orders of magnitude *below* $\mu/2$. The correction: the soft-edge picture of T107 was wrong in its mechanism — τ sits to 99.66% on modes $\lambda \geq 1$, and $\|h\|^2$ is small by sign cancellation ($\sum |k| = 401$ versus $\sum k = 1$), not by softness. T107’s Cauchy–Schwarz chain turns out to be exactly the Weyl bound (slack 1.0000).

What is exactly open after T108 is one boundary value. The remaining object is a certified upper bound for the avoidance norm $\|V^T T_{\text{odd}}^{-1} \tilde{t}\|^2$, and on 8 zones that is literally the single edge entry x_0 of an explicit vector (suppressed at $|x_0|/\max |x| = 2.9 \times 10^{-3}$ – 5.7×10^{-3} , with a linear edge profile $r^{1.01}$ quoted as a fit) — a boundary-decay statement about an explicit vector, the same object class as the T105 carrier separation. $\hat{r}$ is flat in M ($6.1\times$ more stable than ε) and $r, \hat{r} < 1$ at all 92 (zone, M) points up to $M = 3000$, tightest at 0.9648: measured, on resolved zones, as always. T109 (`boundary_decay_probe.py`, contract BOUNDARY.DECAY — Combes–Thomas/Green decay, a Szegő recursion bound, ω via PSD matrix caps) was running at the time of writing and is *not* reported here.

7 The kill list

A program that only reports what survived is not reporting. These are the readings and routes that were refuted *by the program’s own probes*, each with the measurement that killed it. They are listed as a section rather than as footnotes because they are the more transferable output: each one tells a later attempt where not to go.

- K1. The essential-singularity reading of the onset** (T96, refuted T102). T96 read the counterfactual onset as super-polynomial, compatible with $\exp(-c/\delta')$, i.e. an essential singularity. T102 showed the onset is *anchored* at the crossing $\delta_c = 2w_k$ of two finite quantities ($R^2 = 0.968$), that exponential versus power behaviour is not separable at this resolution, and that the scatter of c (CV 103%) collapses to $c' = 1.52$ in the scale-free variable. The essential-singularity reading is *compatible but not distinguished* — and it is no longer used.
- K2. The C/g law as a causal law** (T101, refuted T102). Triple negative: g_k is causally impossible (Q_{k-1} is blind to atom $k+1$), statistically dispensable (causal law $C \cdot u$: CV 23.3% versus 27.6%), and arithmetically only a ceiling ($C_k \leq g_k \mu_k$ exactly; the 16-zone extrapolation violates its own ceiling from $k = 69$; checked on 18 120 prime-power atoms to $n = 200\,000$). *It was a proxy.*




Figure 7: The reduction cascade, with its arrow types made explicit. The top line is an *equivalence typing* produced by an exact identity (Section 4); the arrows below are the architecture of a would-be sufficient route, verified on finitely many windows only. The dashed box is the reason this figure is not a proof sketch.

- K3. The naive margin route** (T104). $M \geq m \cdot \text{Id}$ holds in 0/16 zones; the coupling-mass to margin ratio is 10^3 – 10^6 ; $m \sim M^{-1.7}$ vanishes in the continuum (fit). The genuine form is nearly singular on the induction window and no margin rescues it. (The prolate wing basis recommended by T96/T103 was tested in both arms and also rejected.)
- K4. The density chain** (T106). It would need a hard gap $w_0 \geq 0.55$ – 2.44 ; measured 5.3×10^{-6} – 1.1×10^{-3} — short by a factor 5×10^2 to 3.6×10^5 . Dead for good.
- K5. The Landau–Widom anchor** (T106). The wrong classical anchor for this problem (rms 0.50–1.55). The right one is Grenander–Szegő on the Planck-coarsened symbol — averaging over the mode-spacing cell π/M — which hits 0.92–2.43 and is better on 67/72 pairs. This also explains why a negative symbol (dip $-21.3 \dots -3.4$) still gives $Q \succeq 0$: the atoms oscillate exactly at the window’s Planck scale.
- K6. The accumulated invariant with self-propagation** (T106). A no-go *with a measured mechanism*. The wing demand gives $\beta_{\text{wing}} = 2.21$ – $14.43 > 1$ on 16/16, but inherited demands give $\beta_{\text{int}} \sim 10^{-2}$ on 15/15: an accumulated invariant with absolute anchoring is false. The reason is transport — in a larger window an old wing pair becomes an interior pair and draws 99.7–99.95% of its demand from the 64 softest modes (coupling 751–2653× stronger). No self-propagation either ($\beta_{\text{on}} < 1$ on 15/15, loss 13–320×, $\theta_k = 0.9866$ – 0.9998). *The wing margin is an edge property of the window, not a property of the pair*. What survives is the window-wise family $\sigma_k(\delta_0) \geq 1.31(\mu_k/2)$, uniform in M with drift $\leq 2.01 \times$.

- K7. The symbol route for the odd channel (T107).** Structurally dead, not merely weak: the symbol lower bound for T_{odd} is certified but *negative* ($\lambda_{\text{cert}} = -2.54 \dots -0.81$) against a measured $\lambda_{\text{min}}(T_{\text{odd}}) = 4.7 \times 10^{-5} \dots 6.4 \times 10^{-2}$ and a symbol infimum of $-22 \dots -6$: five orders of magnitude and a sign change. The soft edge of T_{odd} is a pure finite-section effect that no symbol can see. Coarse certified chains (Cauchy–Schwarz against λ_{min}) close 0/16 — and fail even with the *measured* λ_{min} , so it is a category error, not a precision problem.
- K8. Three certificate candidates for the avoidance norm (T108).** Killed *quantitatively*, which is the useful form: the cancellation route ($\sum |k| = 401$ against $\sum k = 1$), the Bessel route (off by 10^7) and the pole Cauchy–Schwarz route (off by 10^6). ω_{cert} is blocked by the compression Schur distance alone (0.18–0.52). What replaced them is not a fourth bound of the same kind but a different object class: a boundary-decay statement about one explicit vector.
- K9. Two earlier structural kills**, recorded here because they still bound the design space: the cross-block $\sqrt{\lambda_0}$ law is Douglas range inclusion and therefore circular (T98), and the cone-library route saturates at 5/24 coverage and is dead as a covering strategy (T72, promoted into v540 as a named limit). Likewise the seam route to the archimedean place (T20–T25) and the Kohnen-plus route to the central values (T44, promoted into v537 as a scope fence).

8 Honest status and outlook

8.1 What is certified, what is measured, what is hypothesis input

The three-way split, stated once

Certified (exact identities or closed-form bounds, machine-checked): the seven ledger modules of Section 3; and, in the sandbox arc, the bare bound b_0 in closed form, the parity superselection $JQJ = Q$, $JB = -BR$, the pole splitting $ab^T + ba^T = ss^T - tt^T$, the channel inertia, the avoidance law $Q_{\text{full}} \succeq 0 \Rightarrow \|B^T v_i\|^2 \leq w_i \Lambda_-$, the growth inequality, the accumulation bounds, the Sherman–Morrison/Woodbury reduction of (R) to $r \leq 1$, and the three T108 identities (ε as a $Q|_{\text{odd}}$ energy, the rank-one overlap hh^T/ε , and ε as the square of the last Cholesky pivot).

Measured (holds on the resolved windows; no uniform constant): ρ_- ; the ratio $r = 0.0053\text{--}0.1814$ and $\hat{r} < 1$ at all 92 resolution points; $\omega = 0.2366\text{--}0.7531$; the closure map 44/64; the handover margins; the two race slopes; every scaling exponent quoted in this document.

Hypothesis input / classical (used, named, not proved here): Kneser, Witt/Eichler, Shimura, Waldspurger/Baruch–Mao, Cohen, Weil–Guinand, Bombieri, Yoshida, Connes–Consani, Douglas, Slepian–Landau–Pollak, Gronwall/Robin (unconditional form only), Grenander–Szegő, Sherman–Morrison/Woodbury, Christoffel–Darboux, Szegő–Levinson, the Weyl bound.

8.2 The asymptotic mountain

Three separate reasons why the cascade of Figure 7 is not a route to a theorem as it stands, and why the document does not present it as one:

- Sixteen zones are sixteen zones.** Everything below the top line of the cascade is established on the resolved zones $n = 2 \dots 29$. Not one constant in the chain is certified uniformly in k . The T101 requirement list (A)–(D) is unchanged: the hardness would sit in (A), a sharp, non-perturbative arithmetic lower bound at a Weil-positivity edge.
- Finite resolution is a wall, and a measured one.** Numerics as a witness is exhausted beyond $\alpha \approx 0.55$ ($O(M^{-1.8})$, T96); the finite-block route was judged the structurally wrong instrument, not an under-resourced one (T92); λ_{min} collapses geometrically by a factor 11.4 per mode, so double precision ends at $N \approx 10$ while $N = 24$ would need 10^{-27} entries.
- The induction base is not free either.** The recursion terminates arithmetically (240/240 in ≤ 4 steps to the classical zone, T99; 960/960 with longest chain 13 steps, T101), but the

zone peaks need percent-level bounds, not 0.1% (T99: extrapolated peak margins $+2.7 \times 10^{-2}$, $+1.4 \times 10^{-2}$, $+1.7 \times 10^{-2}$, i.e. 2.2–5.5% of $\mu_k/2$, with zone 5 saturating exactly).

8.3 Extraction candidates: the community-checkable path

The useful question at this point is not “how close is the program to RH” — it is not close, and no such claim is made — but “which pieces of it are unconditional theorems that someone else can check”. Two candidates are named in the diary and are repeated here as the honest outlook:

- **Zone extension beyond $\log 2$.** Unconditional Weil positivity is classically known exactly up to support width $\log 2$ (Yoshida 1992; Bombieri 2000). T91’s target inequality — for $a \in (\log 2, a^*]$ and $\|f\| = 1$, $(P_{\text{pole}} + A_{\text{arch}})(f) \geq \sqrt{2} \log 2 \cdot h_f(\log 2)$ — is typed as *provable-shaped* and would, if proved, be a standalone classical result: a zone extension past Bombieri’s boundary. Its typing is explicit and unflattering: $\text{RH} \Rightarrow (\text{T})$, but $(\text{T}) \not\Rightarrow \text{RH}$. T92 showed the finite-block route to it is the wrong instrument; that narrows, rather than closes, the target.
- **The matching lemma.** Already proved exact-integer on $[4, 10^6]$ with an explicit margin inside v541, and reduced by T77 to a restricted σ_{-1} divisor bound in the Gronwall regime — with the honest note that Robin’s *unconditional* 1983 bound suffices at the measured headroom, so the RH-equivalent sharpening is not needed. Closing the tail (the correlated-cancellation lemma on credit-rich non-coherent atoms) would be a self-contained classical statement.

Final fence

The prime line of TFPT has produced seven machine-checked modules about lattice-native Hecke structure, a finite relative-trace identity, and an exactly closing transport ledger. It has *not* produced evidence for the Riemann Hypothesis, and the one place where RH appears — I5 — appears as an equivalence typing that names the residual difficulty rather than reducing it. The corresponding open gates in the ledger (ZETA.CENSUS.TO.GL1, ZETA.HP.CARRIER) are untouched by everything in Sections 4–6, and no marker moves anywhere on account of this document.

A Probe index, diary parts T11–T108

Every row is one exploratory probe in `experiments/tfpt-discovery/`; the file column omits the common suffix `_probe.py`. “Checks” is the probe’s own assertion count, all green. Verdict strings are the diary’s preregistered verdict enums. Rows marked “→ vN” were later consolidated into the corresponding load-bearing module; the promotion, not the probe, is the load-bearing artefact. The index covers the 98 parts T11–T108 that the current diary file records (99 runs — T104 was a double run with two independent arms — and 2509 green checks); the ten earlier probes T1–T10 predate that file and are not indexed here. Series totals as recorded by the diary at T108: 108 **probes**, 2701/2701 **sandbox checks**, plus v535–v541 promoted. T109 (`boundary_decay_probe.py`, contract `BOUNDARY.DECAY`) was still running when this document was written and is deliberately absent.

Part	Probe	Contract / finding	Verdict	Checks
T11	e8_glue_chi4_signed_theta	the χ_4 -signed glue theta welds parts 6 and 7	NO PROMOTION	19/19
T12	e8_glue_lseries_selfdual	L-series of the signed glue count; Apéry/ $\zeta(3)$ bridge; Poisson self-dual width 2π	NO PROMOTION	16/16
T13	e8_glue_character_atlas	glue-character atlas of E_8	NO PROMOTION	11/11
T14	seam_selfdual_width	SEAM.SELFDUAL.WIDTH; unimodularity gate	DEFLATED	13/13
T15	zeta_arith_completion	ZETA.ARITH.COMPLETION; shell Hamiltonian	PARTIAL	12/12
T16	zeta_local_euler	ZETA.LOCAL.EULER; one weight-4 Hecke polynomial	PARTIAL	16/16
T17	zeta_funceq_duality	ZETA.FUNCEQ.DUALITY; Poisson involution	PARTIAL	9/9

Part	Probe	Contract / finding	Verdict	Checks
T18	zeta_weil_recovery_trace	ZETA.TRACE.COMPLETENESS; Guinand–Weil balance	BENCHMARK-STANDING	17/17
T19	prime_thinning_seam_truncation	PRIMON.SEAM.TRUNCATION; thinning vs. the finite tower	MIXED	19/19
T20	seam_archimedean_kernel	archimedean kernel from the seam density of states	KILLED-AS-NAIVE	25/25
T21	prime_prediction_mechanics	four prediction channels quantified	MAPPED	19/19
T22	seam_halfcut_dilation_arch	non-DOS successor of the T20 kill	PARTIAL (K1, K2 pass; K3 fails)	25/25
T23	seam_boost_arch_dictionary	arch dictionary via the seam boost	DEAD	23/23
T24	seam_truncated_trace_rvm	truncated trace / Riemann–von Mangoldt route	DEAD	25/25
T25	seam_tate_phase	scattering phase as the last observable	DEAD (arch route closed)	29/29
T26	shell_multiplicity_one_quotient	shell quotient towards multiplicity one	PARTIAL	18/18
T27	e8_pneighbor_hecke	Kneser p -neighbours carry the Eisenstein half	PARTIAL $\rightarrow$ v535	27/27
T28	e8_kneser_transport_holonomy	cusp coefficient from mod-3 lattice geometry	TRANSPORT-VISIBLE	22/22
T29	primon_transport_category	primon monoid relations	RELATIONS-VERIFIED	16/16
T30	e8_transport_cusp_uniformity	uniformity of the T28 reading at $p = 5, 7$	BROKEN (reading); transport holds	32/32
T31	e8_neighbor_operator_decomposition	cusp extraction is a rule $\nu_p = a \text{ Id} + b T_p$	REPAIRED-BY-DICTIONARY $\rightarrow$ v535	49/49
T32	census_newform_recovery	census redundancy is 2-adic oldform structure	REDUNDANCY-IS-OLDFORM $\rightarrow$ v535	24/24
T33	e8_nu_rule_many_primes	hardening of the ν -rule to $p \leq 100$; Eichler finding	HARDENED $\rightarrow$ v536	26/26
T34	compiler_functor_theta	project ZETA.COMPILER.FUNCTOR founded	FOUNDED-WITH-ANCHOR	13/13
T35	rankin_selberg_functor	Rankin–Selberg translates the abelian shadow	RS-TRANSLATES-EISENSTEIN	13/13
T36	e8_lambda_charsum_evaluator	cusp remainder isolated geometrically	TWO-SIDED $\rightarrow$ v536	29/29
T37	e8_salie_signed_cusp	signed cusp channel, scalable	SCALABLE-AND-SIGNED	37/37
T38	cuspidal_bridge	Shimura preimage of f_8 inside the compiler monoid	BRIDGE-FOUND $\rightarrow$ v537	13/13
T39	zeta_center_atlas	the abelian drop	ABELIAN-DROP-CLOSED	13/13
T40	weil_recovery_compiler_sector	Weil-recovery sector terrain	TERRAIN-MAPPED	24/24
T41	functor_wiring_halfintegral	the half-integral bridge is Hecke-equivariant	FUNCTOR-WIRED $\rightarrow$ v537	13/13
T42	e8_local_densities	local densities closed	CLOSED $\rightarrow$ v536	25/25
T43	kohnen_plus_waldspurger	Kohnen plus space	PARTIAL	12/12
T44	kohnen_operator_projection	central-value route via Kohnen plus	OUTSIDE-KOHNEN-SCOPE $\rightarrow$ v537	17/17
T45	baruch_mao_metaplectic	central-value family as $b(d)^2$; R constant	CENTRAL-VALUES-WIRED $\rightarrow$ v537	13/13
T46	stage4_prolate_prototype	stage-4 prolate prototype	IMPL-OPEN	16/16
T47	stage4_prolate_uv_dirac	pointwise UV/Dirac prototype	UV-NO-SIGNAL	17/17
T48	lambda_scale_uniqueness	does the compiler fix the Λ scale?	SCALE-TORSOR (claim killed)	17/17
T49	rankin_positivity_miniature	the Deligne/Rankin mechanism runs in-suite	MINIATURE-RUNS	28/28

Part	Probe	Contract / finding	Verdict	Checks
T50	half_integral_ selfconvolution	the central-value family aggregates exactly	FAMILY- AGGREGATED	17/17
T51	waldspurger_family_ kernel	first non-collapsing infinite carrier candidate	INFINITE-CARRIER- CANDIDATE	22/22
T52	hecke_orbit_transfer	path space of Hecke words; fibre determinants	TRANSFER-REALIZED	16/16
T53	relative_trace_ compiler	v535/v536/v537 are three projections of one RTF identity	ONE-FORMULA → v538	16/16
T54	waldspurger_ canonical_measure	the “canonical discriminant measure” claim	MARGINAL-WEIGHT (killed)	26/26
T55	rtf_period_pairing	canonical distributional limit of the finite RTF	DISTRIBUTIONAL- RTF-TYPED → v539	36/36
T56	rtf_pairing_ sharpening	all T55 identifications to sub-percent	SHARPENED	27/27
T57	infinite_rtf_packing	both infinity directions in one double series $Z(s, w)$	PACKED	24/24
T58	zeta_rtf_xi_residue	GL(1) shadows of $Z(s, w)$	GL1-SHADOW- CLASSICAL (route A closed)	24/24
T59	weil_gns_ identification	residual factor in closed form	TRANSFORM- REQUIRED	19/19
T60	plancherel_descent	Sato–Tate moment contraction	PARTIAL	21/21
T61	st_isotype_gl1_core	the trivial Sato–Tate isotype is a GL(1) core	GL1-CORE-EXACT → v539	24/24
T62	core_isolation	d -twist mixing vanishes fibrewise	CORE-ISOLATED → v539	24/24
T63	weil_core_completion	Plancherel correction irreducibly non-automorphic	EXTRA-TERMS → v539	33/33
T64	rtf_stabilization	$\det_2$ absorbs obstruction 2; the minus stays	HALF-STABLE	34/34
T65	pi_digits_prime_ distribution	prime distribution in the digits of π	PI-NULL	16/16
T66	pi_prime_four_level	the π /prime question on four levels	FOUR-LEVEL-NULL	23/23
T67	amplitude_dirac_sqrt	Dirac square root of the amplitude plane	DIRAC-SQRT-EXACT → v540	27/27
T68	geometric_ polarization	geometric polarisation $b = N_+ - N_-$	RESHUFFLE-ONLY → v540	34/34
T69	mixed_channel_full_ weight	even- k deletion is a determinant invariant	DELETION- UNIVERSAL → v540	33/33
T70	linear_measure_weil	positive linear Θ carrier; pure plus combination	LINEAR-PLUS- COMBINATION → v540	29/29
T71	transport_terrain	FE of the linear family exactly anchored	TERRAIN-MAPPED → v540	40/40
T72	cone_enlargement	the cone library saturates at 5/24	COMPLEMENTARY- PAIR → v540	34/34
T73	continuous_cone	per-direction hybrid cones in the continuum	HYBRID-GAINS	28/28
T74	door_a_spectral_ polarization	the spectral side is provably sign-blind	VACUUM-STRUCTURE	32/32
T75	door_b_lambda_ transport	the gap functional λ^* has its own structure	LAMBDA- STRUCTURED	28/28
T76	hybrid_universality	the recipe certifies 91/91 adversarial Weil directions	RECIPE-UNIVERSAL- ON-BATTERY	22/22
T77	matching_lemma_ structure	the matching lemma reduces to a restricted divisor bound	LEMMA-CLASSICAL- SHAPED	21/21
T78	lemma_window_proof	matching lemma exact-integer on $[4, 10^6]$	WINDOW-PROVED → v541	25/25
T79	transport_ledger	the transport ledger closes; I5 named and typed	LEDGER-CLOSES → v541	22/22
T80	tail_correlation_ lemma	character-exact signed envelope; tail to $N^* \approx 10^{23}$	RESERVE-PARTIAL → v541	29/29
T81	recipe_coherent_ avoidance	reachability theorem: coherent targets need coherent m	AVOIDANCE-FAILS → v541	31/31
T82	heat_arch	Δ_{arch} is the internal Γ difference	ARCH-INTERNAL → v541	22/22
T83	fe_symmetrization	symmetrisation absorbed; the wall is FE-transversal	INVARIANT-NULL	27/27

Part	Probe	Contract / finding	Verdict	Checks
T84	gaussian_lift	Größencharacter phases give $L(1, \lambda_k)$ convergence	LIFT-WORKS-UNANCHORED	29/29
T85	lambda_equivariant_design	the λ design closes the coherent class	LEMMA-CLOSES-LAMBDA $\rightarrow$ v541	24/24
T86	correlated_cancellation	the Q -pairing involution closes the non-coherent tail	LEMMA-FULLY-CLOSED	29/29
T87	i5_connes_dictionary	I5 one-family form vs. the semi-local Connes structure	DICTIONARY-EXACT-CORE	22/22
T88	i5_tightness	eight near-vertical tight curves; Γ -side spacing	TIGHT-SET-PARAMETRIZED	27/27
T89	sonin_crossing	window compression is Bombieri's classical object	CROSSING-MAPPED	19/19
T90	residual_dissection	the residual vectors are explicit Gaussian modes	CORE-DISSECTED	17/17
T91	thin_band_analytic	the band is the one-atom zone; atom rescue	BAND-PARTIAL (3/4 closed)	19/19
T92	zone_extension_proof	the finite-block route is the wrong instrument	T-SKELETON	36/36
T93	band_reconciliation	data-integrity reconciliation of the T89/T91 map	MIXED (map recalibrated)	41/41
T94	prime_blind_demo	753/753 primes of an unseen window from lattice counting	BLIND-100	16/16
T95	continuum_extension	C1 chain proved; band top behaves as a positivity edge	T-CONTINUUM-NUMERIC	28/28
T96	relay_test	the T95 edge was a map artefact; the relay is real	EDGE-ARTIFACT + RELAY-CONFIRMED	21/21
T97	relay_induction	the induction step has proof shape; one cross-block lemma	ALIGNMENT-ONLY	105/105
T98	cross_lemma	the $\sqrt{\lambda_0}$ law is Douglas range inclusion (circular)	LAW-CONFIRMED-MECHANISM-OPEN	44/44
T99	dk_recursion	exact mirror-parity selection rule; recursive inequality	DECAY-LAW-FOUND	23/23
T100	bulk_remainder	closure 11/24 $\rightarrow$ 24/24 samples; the drift was a grid artefact	REMAINDER-CLOSES-ZONES	27/27
T101	k_asymptotics	16 zones resolved; primitives flat, instrument loses the race	CROWDING-TRENDS	31/31
T102	arithmetic_bound	the onset is manufactured by the Schur dressing; C/g was a proxy	MECHANISM-IDENTIFIED	42/42
T103	instrument	slope halves; closure map 7/64 $\rightarrow$ 44/64	INSTRUMENT-IMPROVED	29/29
T104	schur_profile_bound	arm B: naive margin route dead, threshold split closes 16/16	CHAIN-PARTIAL	21/21
T104	schur_profile_chain	arm A: independent twin of the same contract	CHAIN-PARTIAL	47/47
T105	bare_wing_bound	bare bound certified in closed form; avoidance is a theorem	ONE-OF-TWO	28/28
T106	friedrichs_angle	parity split of the Weil pole; hardness in the odd channel	DENSITY-MAPPED	32/32
T107	odd_channel_closure	Sherman–Morrison/Woodbury reduction to $r = \kappa/\varepsilon$	SCALAR-TRACTABLE	30/30
T108	ratio_certificate	ε is an exact $Q _{\text{odd}}$ energy; (R) on two scalars; first non-vacuous margin chain	EPSILON-IDENTITY	44/44

B Symbols used in Sections 4–6

All of these are sandbox objects; they are listed so the figures and formulas above can be read without the diary.

Symbol	Meaning
$Q_{\text{Weil}}, Q_{\text{cert}}, \Delta_2, \Delta_{\text{arch}}$	the four terms of the transport ledger; $Q_{\text{Weil}} = Q_{\text{cert}} + \Delta_{\text{arch}} + \Delta_2$ closes at 4.4×10^{-16} (v541)
$A_{\text{fam}}, A_{\text{shift}}$	the two archimedean halves of the heat family; $\Delta_{\text{arch}} = A_{\text{fam}} - A_{\text{shift}}$ (T82)
I5	the one remaining coupled inequality, $Q_{\text{cert}} + \Delta_2 + A_{\text{fam}} - A_{\text{shift}} \geq 0$ (T79)
μ_k, w_k, g_k	atom strength, handover window width, atom spacing at zone k
$D_k(\alpha)$	the Schur-complement defect of the induction step; target $D_k \leq \mu_k/2$ (T98)
$\sigma_k(\delta)$	the E_- Schur profile of the genuine Weil form just above atom entry (T102)
E_-, E_0, E_+	the three atom eigenspaces; the k -th atom acts as $\text{diag}(-\frac{1}{2}, 0, +\frac{1}{2})$
bare_k, b_0	$\lambda_{\min}(Q_{\text{full}} _{E_-})$ and its certified excess over $\mu_k/2$ (T105)
L, Λ_-	the soft dressing scalar and its free cap (5.63–7.52, divergent) (T105)
J	the mirror-parity involution; $JQJ = Q, JB = -BR$ (T105)
$\rho_{\pm}$	the Friedrichs-angle cosines of the even / odd channel (T106)
(R)	$Q_{\text{full}}(\alpha_k) _{J=-1} \succeq (\mu_k/2)P_- _{J=-1}$, the remaining Loewner statement (T106)
$s^*, \tau, \kappa, \varepsilon$	Sherman–Morrison scalar, its Woodbury split $s^* = \tau + \kappa$, and $\varepsilon = 1 - \tau$ (T107)
r	the residual ratio κ/ε ; (R) $\iff r \leq 1$; measured 0.0053–0.1814 (T107)