

TFPT — Architecture and the E_8 Compiler

The two axioms $\{c_3, g_{\text{car}}\}$, the derivation map, and the Coxeter–cyclotomic $D_5 \times A_3 \rightarrow E_8$ construction

Stefan Hamann

Alessandro Rizzo

July 13, 2026 · v5.4 (rev 363)

What this document is about

The architecture layer: how the two axioms $c_3 = \frac{1}{8\pi}$ and $g_{\text{car}} = 5$ build the Coxeter–cyclotomic compiler — the carrier $C^+ = D_5$, the family geometry $\mathbb{P}^1 \setminus \mu_4 = A_3$, the μ_4 glue $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$, the EM fixed point α^{-1} (with its ablation), and the whole number alphabet 16, 40, 41, 48, 240, 248 as carrier traces.

The TFPT document set — what is treated where

Plain language: TFPT (Topological Fixed-Point Theory) is a small discrete compiler. Two inputs — the seam constant $c_3 = \frac{1}{8\pi}$ and the carrier rank $g_{\text{car}} = 5$ — build $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ and read off the Standard-Model skeleton, the constants and the scale grammar. The series is **the introduction plus five numbered papers and three companions**, best read in order (with **four** companions):

0. **introduction** — reading guide, compiler closure, predictions, the dependency DAG and the single proof ledger (`verification/status_ledger.csv`). (*entry point*)
1. **tfpt_1_architecture_e8** — the two axioms, the derivation map, the EM fixed point α^{-1} , the $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ construction, the QBL theorem chain.
2. **tfpt_2_standard_model** — the SM in one φ_0 -ladder formula, flavor from parabolic transport (sheet diamond, dual anchor, Q_+ cohomology), the worked closures.
3. **tfpt_3_e8_audit_bootstrap** — the seven E_8 slices as an audit raster, the cascade bridge, the Möbius bootstrap.
4. **tfpt_4_frontier** — honest status of η_B , m_p/m_e , Koide, dark matter, full QG.
5. **tfpt_5_redteam** — the adversarial audit: declared attacks, what survives, what each reduces to, and the kill tests.

Companions: **R.** **tfpt_research_contracts** — the open interfaces $(v_{\text{geo}}, G_{\text{net}}, F_{\text{transfer}})$ as numbered contracts; **H.** **tfpt_horizon_readouts** — Appendix H (unit-system reframe): one seam constant as the universal horizon thermal code, the Nariai/anchor surface, the resummed seam clock; **O.** **origin_theory** — why no free fundamental number remains (exact core + one honestly-typed cyclic interpretation); **S.** **tfpt_safeguards** — the verification discipline: every mechanism that defends a claim against coincidence and numerology (status calculus, no-free-pattern, the over-determination map, the firewall, frozen predictions + null model, the independent Wolfram and Lean paths, the red team).

You are reading document 1 (Architecture).

TFPT status at a glance — read this card the same way in every document

Two inputs, one compiler. From the boundary constant $c_3 = \frac{1}{8\pi}$ (axiom P1) and the five-slot carrier $g_{\text{car}} = 5$ (axiom P2) — jointly the elementary symmetric data of one anchor $a = (1, 1, 2)$ — the discrete compiler $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ is *built*, and the Standard-Model skeleton (gauge group, three families, hypercharges, the flavor matrix), the dimensionless constants ($\alpha^{-1} = 137.0359992$) and the scale grammar are read off as fixed points. The algebraic core is machine-checked; physical readouts run through *named bridges* whose status is typed below.

What is closed, and what genuinely remains. The discrete/algebraic layer is closed ([E]); the everything-open residual reduces to *three named interface problems*:

Interface	Question	Status
v_{geo}	the one metrology unit ($= 1/\sqrt{G} = m/\mu$); No-Unit Thm: no compiler scale	primitive [O]
G_{net}	simple-current extension $(D_5)_1 \otimes (A_3)_1 \rtimes \langle (1, 1) \rangle \cong (E_8)_1$	[C]
F_{transfer}	one functor, four typed interfaces (Koide, η_B , axion, m_p/m_e)	[C]

Gravity is parameter-free. The classical metric-sector field equation is no longer only an $R+R^2$ readout: the fixed-volume entanglement equilibrium (Jacobson; Faulkner et al.), run with TFPT's atoms, gives the *full* covariant Einstein equation $G_{ab} + \Lambda g_{ab} = c_3^{-1} T_{ab}$ with *both* coefficients fixed — $c_3^{-1} = 8\pi$ (no free dimensionless Newton dial; the thermodynamic origin $2\pi/\eta$ *coincides* with the geometric one $|\mathbb{Z}_2|2\pi\chi$ via $|\mu_4| = |\mathbb{Z}_2|\chi(S^2) = 4$, so c_3 is triply over-determined — anchor, geometry, thermodynamics, v358) and Λ from α ($\rho_\Lambda = (3/4\pi^2)e^{-2\alpha^{-1}}$, v60); the Einstein tensor (not Ricci) is forced by Lovelock, so matter conservation is an output (v359). The residual here is only the equation-of-state status (an external candidate action — Bianconi's entropic action, arXiv:2408.14391 — is quantified in v473 without changing the typing) and the one unit v_{geo} ; the global measure (C7 = QG.AMB.01) is now a [C] redundancy (v369), a certification object rather than missing dynamics, conditional on SEAM.EQUIV.01 and Bisognano–Wichmann.

The seam-net structural theorem G_{net} (= SEAM.EQUIV.01: the raw seam *is* the holomorphic $(E_8)_1$ net) is now *closed modulo a cited theorem*. The explicit lattice model (v367/v368) and the S_3 closure stack pin the target at every computable level — central charge $c = 8$ (v376), the $(E_8)_1$ character with 248 currents and one primary (v377), genus-one torus $\text{GSD} = 1$ (v378) and reflection positivity (v379) — and it is Lean-formalised as FORM.SEAM.MMST.01: the collar's MMST hypotheses are kernel-proved, the MMST scaling-limit and Adamo–Moriwaki–Tanimoto OS-reconstruction theorems enter as named cited axioms, and the #print axioms check is clean. The *only* residual that stays [O] is the abstract continuum existence of the scaling limit (v336) — its 128-spinor extension leg since certified at net level by the peer-reviewed crossed-product package ($h_s=16/16=1 \in \mathbb{Z}$, the Longo–Rehren locality criterion; Böckenhauer; KLM $\mu=4/2^2=1$; AGT/AMT the independent second witness) with the realisation input reduced to invariant level and the parallel Lean route `seamResidualClosed'` (v469); so this is *closed modulo a cited theorem*, not solved. QGEO.SYM.01 (the μ_4 deck acting geometrically) is its *corollary*: a conformal net's vacuum is rotation-invariant by axiom, so the deck follows with no extra premise (v335, Lean FORM.QGEO.BW.01). The nonperturbative quantum-gravity measure QG.AMB.01 is no longer carried as an open item: it is a [C] *redundancy* (v369), a certification object rather than missing dynamics, conditional on SEAM.EQUIV.01 and Bisognano–Wichmann (and TFPT is moreover rigorously gap-decoupled from the general Euclidean-QG conformal-factor problem, margin $1.648 > 0$, v76/v330). So the honest residual is the one unit v_{geo} plus the typed F_{transfer} interfaces, above the cited-theorem ceiling on the seam and with QG.AMB.01 a [C] redundancy. Everything carries a claim ID resolving to a row of `verification/status_ledger.csv` (the single source of truth); **if the text and the ledger ever disagree, the ledger wins**. The historical labels ($U_{\text{wall}}, G_{\text{metric}}, F_{\text{frontier}}$) are retained only for ledger continuity — the live residual is $v_{\text{geo}} \oplus F_{\text{transfer}}$, with the seam G_{net} closed modulo a cited theorem and QG.AMB.01 a [C] redundancy.

Marker key: [E] exact / machine-proven · [C] conditional (holds under named hypotheses) · [O] open / axiom · [X] falsifiable kill test. Per-claim fine type in `verification/status_ledger.csv`.

[E] exact / machine-proven

[C] conditional

[O] open / axiom

[X] kill test



Contents

I	Architecture and the derivation map	5
1	How to read this document	6
2	Headline: the Pascal compiler on five carrier slots	9
2.1	$g_{\text{car}} = 5$ as the Pascal closure	10
2.2	One master moment generates 40, 41, 240, δ_2	10
2.3	Compact E_8 compression and the role of ν^c	11
3	The $D_5 \times A_3$ compiler: E_8 as a closure, not an input	11
3.1	Carrier = $D_5 = \mathfrak{so}_{10}$ (proved)	12
3.2	Family = $A_3 = \mathfrak{su}_4$ from $\mathbb{P}^1 \setminus \mu_4$ (proved, with the metric settled)	12
3.3	The μ_4 glue and the $5/4 + 3/4 = 2$ discriminant theorem (proved)	13
3.4	Glue-norm arithmetic, and three readings of 41, 60, 120	14
3.5	What the compiler outputs for the Standard Model and our reality	15
4	Grammar I — the carrier grammar (3 + 2)	17
4.1	Dual-root dictionary (everything from two charges)	17
4.2	Carrier arithmetic (proved identities)	18
5	Grammar II — the seed grammar (one decoder u)	18
6	Grammar III — the action grammar (exponentials of $\alpha_\star^{-1}$)	19
6.1	The EM closure: $\alpha_\star^{-1}$ as an explicit fixed point (made explicit here)	19
6.2	The decuple bridge (new, exact headline)	23
6.3	The full power tower: x^1, x^5, x^{10} on the carrier graph K_5	23
6.4	Hubble is the square root of Λ (corollary, exact)	23
6.5	The ~ 123 orders of the cosmological constant	24
6.6	The ladder is binomial; $2g_{\text{car}} = \binom{g_{\text{car}}}{2}$ singles out $g_{\text{car}} = 5$	24
6.7	Discriminant action theorem: the ladder reads the hypercharge discriminant	24
6.8	A two-dimensional action lattice	25
7	The Einstein normaliser as a corrected tree value	25
8	Λ-anchored metrology closure, and two anchors	25

9 The landscape: $g_{\text{car}} = 5$ is overdetermined	26
10 The E_8 carrier atlas — stated unambiguously	27
10.1 Level 1 — original E_8 data of the series ([E])	27
10.2 Level 2 — new exact counting identities ([E])	27
10.3 Level 2.5 — the even-flip transition atlas (a falsifiable program)	27
10.4 Level 2.7 — the carrier Coxeter theorem (the strongest new bridge)	28
10.5 Level 2.8 — positive-root moment: $ R^+(E_8) = \text{Tr}_{S^+} X^2 = 5!$	28
10.6 Level 2.9 — the 2·3·5 Coxeter cyclotomic compiler (meta-theorem)	29
10.7 Level 3 — conjectural geometric readings ([C])	29
11 $\mathbb{Z}_6$ monodromy, flavor and CP readouts	29
11.1 CP as triple seed variance; a closed observable quadrilateral	30
12 Mass hierarchy as a hexagon with one extra turn	30
12.1 Lepton flavor decouple: the charged-lepton determinant carries u^{10}	31
12.2 Why the top Yukawa is ≈ 1 : the $L = 0$ anchor (Froggatt–Nielsen ladder)	31
13 Inflation interface	32
14 Further simple bridges	32
15 Closures that the architecture rests on	32
16 Status summary and open audit points	33
17 Consolidated picture	35
 II The Coxeter–cyclotomic compiler and the explicit E_8 construction	 36
Theorem 1	36
Theorem 2	37
Theorem 3	38
Theorem 4	38
Theorem 5	38
Theorem 6	39
Theorem 7	40
Theorem 8 — anchor microcode	43
Master statement	44
Appendix: explicit E_8 construction	44
 III Appendix: ablation of the EM fixed point	 46
Appendix: computational verification	47

Part I

Architecture and the derivation map

Abstract

This document develops the *architecture layer* of TFPT: the two axioms, the derivation map, and the explicit E_8 construction. The thesis is that the bridge layer of TFPT is governed by *three nested grammars*: a **carrier grammar** (from $3 + 2$ come charges, families, hypercharge, $\Omega_{\text{adm}} = 48$, $b_1 = 41/10$), a **seed grammar** (one seed $u = \varphi_0^{\text{ret}}$ generates the Cabibbo angle, birefringence, the baryon fraction and the reactor angle), and an **action grammar** (the large scale ratios are exponential actions of $\alpha_\star^{-1}$ with rungs $1/g_{\text{car}}$, 1 , 2). Two new exact results sharpen the action grammar: the *decuple bridge* $\rho_\Lambda/\bar{M}_{\text{Pl}}^4 = \frac{3}{4\pi^2} e^{2\delta_{\text{ph}}} [v_{\text{geo}}/(5\beta_{\text{rad}}^2 \bar{M}_{\text{Pl}})]^{10}$ (the cosmological constant is, in essence, the tenth power of the stripped electroweak scale, with $10 = 2g_{\text{car}}$), and the Hubble corollary $H_0/\bar{M}_{\text{Pl}} = e^{-\alpha_\star^{-1}}/(2\pi\sqrt{\Omega_\Lambda})$. We also give the exact corrected tree form of the Einstein normaliser $\xi = \frac{3}{4}/(1 + 9/128\pi^3)$, the dual-root dictionary that generates all hypercharges and transport cusps from $y_\pm = \{1/2, -1/3\}$, the Λ -anchored metrology closure, the overdetermination of $g_{\text{car}} = 5$, and the E_8 carrier atlas in three clearly separated levels. Section 3 adds the strongest structural result: the carrier code is the $D_5 = \mathfrak{so}_{10}$ half-spinor ($\text{Aut} = W(D_5)$, order 1920; $R_4 = \text{Clebsch graph}$), the family geometry $\mathbb{P}^1 \setminus \mu_4$ is an A_3 , and E_8 is the explicit unimodular glue closure of $D_5 \oplus A_3$ — with the discriminant-form norms $q(D_5) = 5/4$ and $q(A_3) = 3/4$ summing to the E_8 root norm 2 and coinciding with the TFPT constants $\delta_2/\delta_{\text{top}}^2$ and ξ_{tree} . Status markers used throughout: **[E]** exact identity, **[E]** Lie/lattice theorem, **[E]** formalised, **[E]** numerical fixed point, **[C]** physical/conditional, **[O]** axiom/open. (All markers are explicit; the legacy **[M]**/**[C]** compatibility aliases have been removed.)

1 How to read this document

Notation convention. We write $D_5 \oplus A_3$ for the *root-lattice* direct sum (used in the glue/lattice statements: discriminant forms, μ_4 glue, E_8 as an even unimodular lattice), and $D_5 \times A_3$ for the *group/representation* branching (used for subalgebra embeddings and E_8 -rep decompositions). The objects are the same atoms; the symbol marks whether the statement is lattice-theoretic ($\oplus$) or branching/representation-theoretic ($\times$).

Nothing here introduces a new primitive assumption; it reorganises existing outputs and proves the cross-links. The whole bridge layer rests on two inputs,

$$c_3 = \frac{1}{8\pi} \quad (\text{axiom P1}), \quad g_{\text{car}} = 5 \quad (\text{axiom P2}). \quad (1)$$

Foundational postulates — the two genuine axioms (hardenable, not derivable)

Everything downstream is a theorem *given* two structural inputs that are not reduced further and are honestly flagged as the theory’s axioms:

(P1) Boundary kernel.

A single oriented seam Σ carries a primitive, reflection-positive boundary kernel with unit winding $[u_\Sigma] = 1$; its self-linking normalisation fixes $c_3 = 1/(8\pi)$. *Status:* primitive postulate. *Hardenable by:* the Calderón projector / determinant-character chain (the P1 hardening) and the Lean carrier-rigidity development (`TfptCarrier.*`, boundary-polarization and Calderón interfaces).

(P2) Carrier interface.

This splits into an *axiom* and a *theorem*, which must not be conflated:

- **P2_{interface} [O] (axiom):** the seam is read out by a five-slot carrier $g_{\text{car}} = 5$ (3 colour + 2 weak) — a postulate (the readout interface).
- **P2_{algebra} [E] (theorem):** *given* the carrier, hypercharge ($y_- = -\frac{1}{3}$, $y_+ = \frac{1}{2}$), rigidity and the family count follow from the idempotent/lattice axioms — machine-verified in Lean (`Hypercharge.lean`, `LatticeRigidityGeneral.lean`, `Rigidity.lean`).

So P2 is an *interface axiom* whose *algebraic consequences* are a *formalised theorem* — “axiom or theorem?” has the precise answer: interface axiom, algebra theorem.

These two are *inputs*: they can be formalised and stress-tested (more Lean, more interface theorems) but not computed away. Every other number in this note — $N_{\text{fam}} = 3$, $\Omega_{\text{adm}} = 48$, $b_1 = 41/10$, $\alpha_\star^{-1}$, the flavor matrix, the E_8 glue, A_s — is a consequence of (P1),(P2).

Anchor-first refinement. The two inputs are themselves the elementary symmetric polynomials of the single parabolic anchor $a = (1, 1, 2)$ (the exponents-at-infinity of the splitting type $\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$): $e_1(a) = 4 = |\mu_4|$, $e_2(a) = 5 = g_{\text{car}}$, $e_3(a) = 2 = |\mathbb{Z}_2|$, with $c_3 = 1/(2e_1(a)\pi) = 1/(8\pi)$. The power sums $p_n(a) = 2 + 2^n$ then give $|R(E_8)| = p_1 p_2 p_3 = 240$, $\dim E_8 = 240 + (p_4 - p_3) = 248$ and the binary ladder $p_{n+1} - p_n = 2^n$. So the input set reduces to $\{a = (1, 1, 2), \pi\}$. [`verification/v23_anchor_generator.py`]

Seed normal form, the factorial spine, and the degree-2 pattern [E]/[C] ([`verification/v106_review_validation.py`])

Three review-validated sharpenings of the anchor picture. **(1) Seed normal form [E].** The seed formula is exactly

$$\varphi_0^{\text{ret}} = \frac{|\mu_4|}{N_{\text{fam}}} c_3 + \Omega_{\text{adm}} c_3^{|\mu_4|} \quad \left(\frac{4}{3} = \frac{|\mu_4|}{N_{\text{fam}}}, \quad 48 = \Omega_{\text{adm}} = N_{\text{fam}} \dim S^+ \right),$$

a *linear seam term* plus a *topological fourth-order correction* over all admissible fermion states — a structured anchor readout, not a fitted expression. **(2) Factorial spine [E].** With $X = 6Y$ on one 16-state generation, $\text{Tr } X = 0$ and $\text{Tr } X^2 = 120 = 5!$ (computed from the explicit charge table); the *same* factorial moment is $|R^+(E_8)| = \sum (E_8 \text{ exponents}) = 120$, doubles to $|R(E_8)| = 240$, and scales to $|W(D_5)| = 2^4 \cdot 5! = 1920$ (the Hawking-power fingerprint) — one moment, four projections (hypercharge, roots, Weyl group, horizon). **(3) The degree-2 inventory.** The compiler repeatedly selects *degree two*: the Pascal closure $2^{g-1} = \sum_{k \leq 2} \binom{g}{k}$ truncates at $K=2$ with the *unique* solution $g = 5$ (scan 1..40);

the glue norms add to the root norm, $\frac{5}{4} + \frac{3}{4} = 2$; $c_3 = \frac{1}{2} \times \frac{1}{4\pi}$ (variational half $\times$ two-dimensional-slice Gauss–Bonnet); the pair sector $\binom{5}{2} = 10 = \mathcal{A}_\Lambda$; the hypercharge moment is a *second* moment; the seam is codimension *two*. *Named hypothesis [C] (recorded, not claimed): Quadratic Boundary Locality* — the seam supports only bilinear data, so the truncation $K = 2$ is *forced* rather than chosen; if proven, the Pascal *selection* (red-team Target B residual, typed [O]/[C]) upgrades to [E] and the chain locality $\Rightarrow K=2 \Rightarrow g=5 \Rightarrow 16 \Rightarrow D_5 \Rightarrow E_8$ closes the carrier choice. *Audit*: the even-code transition reading $240 = 16 \times 15$ is exact arithmetic but non-unique ($240 = 16 \times 5 \times 3$ is the established glue factorisation) and stays recorded, not promoted.

The Sheet-Pairing Lemma ([verification/v109_sheet_pairing.py]) — **the first exact anchor of the hypothesis**. Character-exact weight-multiset combinatorics of the D_5 spinor weights $(\pm \frac{1}{2})^5$ gives: (i) **no scalar within a sheet** — the zero-weight multiplicity of $S^+ \otimes S^+$ is 0, because the odd slot count $g = 5$ flips the sign parity of $-w$ (a parity theorem of the five-slot code); (ii) **the scalar lives across the sheets**:

$$S^+ \otimes S^- = \Lambda^0 \oplus \Lambda^2 \oplus \Lambda^4 \quad (256 = 1 + 45 + 210),$$

exact as weight multisets, with zero-mode grading $(1, 5, 10)$ = the carrier code; (iii) **sheet-diagonal = odd forms**: $(S^+ \otimes S^+) \cup (S^- \otimes S^-) = 2\Lambda^1 + 2\Lambda^3 + \Lambda^5$ exactly — within-sheet pairings are odd-form-valued, never scalar; (iv) the scalar-bearing tower **tops at** Λ^{2K} with $K = 2$ — bilinear seam data reach exactly the Pascal-ladder degree (the source-layer transport theorem; document 3). *Consequence [C]*: the seam’s single scalar two-point kernel is necessarily a **sheet pairing** (chirality-off-diagonal — the same $\mathbb{Z}_2$ as the glue chiralities and the branch-kernel selection); the remaining QBL step is to show the Calderón kernel supplies exactly this pairing and nothing beyond.

The Calderón-sheet selection theorem ([verification/v110_calderon_sheet.py]) — **which involutions certify a scalar kernel**. On $H = S^+ \oplus S^-$ the certified bilinear data of a Calderón involution ε ($\varepsilon^2 = 1$) are the matrix elements $\langle f, \varepsilon g \rangle$. The lemma’s singlet counts per sheet block, $(0, 0, 1, 1)$, give the exact selection statement [E]:

$$\text{a scalar two-point datum exists} \iff \varepsilon \text{ is sheet-odd}$$

— a sheet-odd involution (swapping $S^+ \leftrightarrow S^-$) certifies exactly $1 + 1 = 2 = |\mathbb{Z}_2|$ invariant scalar kernels (one per orientation), a sheet-even involution certifies *none*. The kernel count $2 = |\mathbb{Z}_2|$ matches exactly the *two Lagrangian glues* (swapped by single spinor conjugation): the Calderón scalar channel carries precisely the glue ambiguity, resolved by the same sheet choice. *Anti-overclaim control [E]*: the parity theorem is generic over the odd ladder ($g = 3, 5, 7$: within-sheet count 0, cross count 2^{g-1}) — sheet-oddness forces the *half-spinor relation* $K = (g - 1)/2$, **not** the value $g = 5$; the g -selection stays with the independent rank-8/integrality closure. *Consequence*: the QBL residue splits into two named parts [C] (recorded, not claimed): (a) **interface** — “the seam Calderón involution is sheet-odd” (structurally natural: the one-sided collar/double-cover deck is the *same* $|\mathbb{Z}_2|$ that halves $c_3 = \frac{1}{2}(4\pi)^{-1}$, and exchanging cover sides is sheet-odd by construction); (b) **analytic core** — “the two-point kernel certifies only slot-degrees ≤ 2 ”. Proving both closes $\text{QBL} \Rightarrow K=2 \Leftrightarrow g=5$.

The Quadratic-Transport Theorem ([verification/v111_quadratic_transport.py]) — **the transport half of the analytic core is a theorem**. Grade seam transport by Clifford degree in the ten field operators on the five-slot Fock space ($S = \Lambda^\bullet(\mathbb{C}^5)$, S^+ the 16-state code; integer Jordan–Wigner model, spanning claims by full-rank certificates mod p , valid over $\mathbb{Q}$). *Parity [E]*: all linear words are sheet-odd, all 45 quadratic words sheet-even — code-preserving transport has *even* Clifford degree (the same $\mathbb{Z}_2$ again). *Minimality [E]*: sheet-even words of degree ≤ 1 are the scalars alone. *Generation and completeness [E]*: the 45 quadratics (= $\mathfrak{so}(10)$, exactly the Λ^2 term of the certified tower $1 + 45 + 210$) generate the whole code from the vacuum *and* span the full operator algebra $\text{End}(S^+)$ ($\dim = 256$) at word length two — every code operation whatsoever is a word in pair transport, and all higher even-degree elements (e.g. the 210 quartics) are redundant. Hence

$$\text{the transport degree is exactly 2}$$

— degree ≤ 1 generates nothing, degree 2 generates everything: a theorem, not a hypothesis. *Anti-overclaim control [E]*: the $g = 3$ world behaves identically (15 quadratics generate $\text{End}(S^+) = 16$) — the theorem selects the *degree*, not the rank; the g -selection stays with Pascal closure + rank-8. *Residue after v109–v111 [C]*: QBL’s remaining content is two *interface* statements — (a) the seam Calderón involution is sheet-odd; (b’) the certified *state* inventory is the ≤ 2 -slot tower. Nothing about transport

remains hypothetical.

The Self-Counting Channel (`[verification/v112_selfcounting_channel.py]`) — **the Pascal closure is an identity of the certified channel.** For odd g , negation $w \mapsto -w$ maps the S^+ weights bijectively into the opposite sheet, so the Cartan-neutral kernels of the certified channel are exactly the pairs $(w, -w)$ — *one per code state*: $\#\{\text{neutral kernels}\} = \dim S^+ = 2^{g-1}$ exactly [E]. Grading the *same* neutral set by pair-degree m gives sizes $\binom{g}{m}$, $m = 0..K$ — for $g = 5$ the triple $(1, 5, 10)$ [E]. Two countings of one set:

$$2^{g-1} = \sum_{m \leq K} \binom{g}{m} \text{ is the channel counting itself}$$

(symbolically verified for all odd $g = 3..13$) — the Pascal *closure* (v2/v108) is not an extra requirement but an identity of the certified channel. *Hodge fold* [E]: the code’s minus-sign grading $\binom{g}{2j}$ equals the folded binomial $\binom{g}{\min(2j, g-2j)}$ with $\min(2j, g-2j) \leq K$ — every code sector appears at pair-degree $\leq K$. *Anti-overclaim*: the identity holds for every odd g and carries no g -selection; the selection stays with the rank-8 closure $g + N_{\text{fam}} = 8$ and ladder integrality. *Residue re-type* [C] (*recorded, not claimed*): clause (b’) loses its closure burden — given the sheet-odd kernel, the channel *automatically* contains one neutral kernel per code state at pair-degree $\leq K$; what remains physical is the single input “the seam certifies through one scalar two-point kernel” (for the established free $c = 8$ seam net, Wick extends the one kernel to the entire even tower by determinants). QBL is thereby the structural consistency frame of the carrier choice; the *selection* is carried by rank-8 + integrality.

One kernel is the whole net (`[verification/v113_quasifree_kernel.py]`) — **the QBL input merges with the holomorphy premise.** The remaining input is now anchored in the established Gate-A objects. *Bookkeeping* [E]: $c(D_5)_1 = \frac{45}{9} = 5 = g_{\text{car}}$, $c(A_3)_1 = \frac{15}{5} = 3 = N_{\text{fam}}$, and the carrier net $(D_5)_1 \times (A_3)_1 = SO(10)_1 \times SO(6)_1$ is $10 + 6 = 16$ *free Majorana fermions* ($c = \frac{16}{2} = 8$); the whole extension tower carrier $(\mu=16) \rightarrow SO(16)_1(\mu=4) \rightarrow (E_8)_1(\mu=1)$ carries the *same* sixteen fermions — only the certified glue grows (index $2 \times 2 = 4 = |\mu_4|$). *Wick/Pfaffian* [E]: in the exact ten-Majorana model *all* 210 vacuum four-point and all 210 six-point functions equal the Pfaffian of the single two-point kernel — one kernel determines all correlations. *The kernel is a Calderón involution of rank g* [E]: $M = \mathbf{1} + iA$ with $A^2 = -\mathbf{1}$, so $P = M/2$ is a projection with rank $P = 5 = g_{\text{car}}$ and $\varepsilon = iA$ an involution; at seam level (sixteen Majoranas) the rank is $8 = \text{rank } E_8$:

the central charge is the rank of the one kernel

($c = 5$ carrier block, $c = 8$ seam hull); and the polarization fixes the vacuum *uniquely* (joint annihilator kernel one-dimensional). *Premise merge* [C] (*recorded, not claimed*): “single scalar two-point kernel” = “the seam state is quasi-free” = the defining property of the free-fermion net the holomorphy gate already posits — QBL adds *no* assumption beyond Gate A; when the gate closes, the carrier choice closes [E] along free net $\Rightarrow$ one kernel $\Rightarrow$ sheet-odd $\Rightarrow$ self-counting tower $\Rightarrow$ complete transport $\Rightarrow$ carrier. Gate typing [C]/[O] unchanged.

Hardening (P1): the Gauss–Bonnet origin of $c_3 = \frac{1}{8\pi}$

$c_3 = 1/(8\pi)$ is not an arbitrary number: it is the *one-sided Gauss–Bonnet normalisation of the seam*. The seam construction compactifies the oriented two-dimensional normal slice N_Σ to a sphere S^2 , so Gauss–Bonnet gives the seam curvature integral

$$\oint_{S^2} K dA = 2\pi \chi(S^2) = 4\pi \quad (\chi(S^2) = 2),$$

and the boundary datum is *one-sided* (the Calderón/double-cover construction uses a single collar), contributing a factor $1/|\mathbb{Z}_2| = \frac{1}{2}$. Hence

$$c_3 = \frac{1}{|\mathbb{Z}_2| \oint_{S^2} K dA} = \frac{1}{2 \cdot 4\pi} = \frac{1}{8\pi}$$

equivalently $c_3 = \frac{1}{2}(4\pi)^{-1}$, the one-sided 2D heat-kernel coefficient. This locates both factors precisely:

- the *discrete* $8 = |\mathbb{Z}_2| \cdot 2 \cdot \chi(S^2) = 2 \cdot 2 \cdot 2$ *coincides with* $h(D_5) = \text{rank } E_8 = \varphi(30)$ — the geometric and the bootstrap readings of the “8” agree;

- the *continuous* π is the irreducible Gauss–Bonnet / 2D-Gaussian primitive ($\int e^{-x^2} = \sqrt{\pi}$), the single primitive the self-consistency loop also isolates.

Status: this *hardens* (P1) — it gives c_3 a named geometric origin and reconciles the “8” with E_8 — but does *not* eliminate π ; π remains the one continuous primitive. [C]

The one-paragraph version. There is a boundary (the seam). It yields $c_3 = 1/(8\pi)$ and a five-slot carrier $g_{\text{car}} = 5$, split 3 colour + 2 weak. Two base charges $y_- = -1/3$ and $y_+ = 1/2$ generate the whole fermion family; the 16 even-occupation states are one generation, three families give $3 \times 16 = 48$. One seed $u = \varphi_0^{\text{ret}}$ drives four low-energy observables. Finally $\alpha_\star^{-1} \approx 137$ is the exponential engine: the electroweak scale takes one fifth of it ($e^{-\alpha_\star^{-1}/5}$), the cosmological constant takes it twice ($e^{-2\alpha_\star^{-1}}$), and the Hubble scale is the square root of Λ . A small grammar generates many sectors.

2 Headline: the Pascal compiler on five carrier slots

The strongest new statement is that three of the theory’s separate structures are the *same* truncated row of Pascal’s triangle on the five-slot carrier.

Family = action ladder = transition count (one Pascal row)

The even exterior algebra of E ($\dim E = g_{\text{car}} = 5$), the action ladder, and the E_8 root count are one Pascal row:

$$S^+ = \Lambda^0 E \oplus \Lambda^2 E \oplus \Lambda^4 E, \quad (\dim \Lambda^0, \dim \Lambda^2, \dim \Lambda^4) = (1, 10, 5), \quad \dim S^+ = 1 + 10 + 5 = 16,$$

$$A_{\text{EW}} : A_H : A_\Lambda = \binom{5}{0} : \binom{5}{1} : \binom{5}{2} = 1 : 5 : 10 = (\dim \Lambda^0 E, \dim \Lambda^4 E, \dim \Lambda^2 E),$$

$$|R(E_8)| = \dim S^+ (\dim S^+ - 1) = 16 \cdot 15 = 240.$$

Family (exterior degree), cosmology (action charge) and E_8 (root transitions) read the same $\{1, 5, 10\}$. [E]




Figure 1: Exterior–Action duality. The Standard-Model family is the valency side of the even Hamming scheme on the five-slot code; the cosmological action ladder is its spectral (multiplicity) side. The Λ (pair) sector $\Lambda^2 E$ has dimension $\binom{5}{2} = 10$, the origin of the decuple exponent.

Natural sector assignment:

$$EW \leftrightarrow \Lambda^0 E \text{ (singlet)}, \quad H \leftrightarrow \Lambda^4 E \cong E^\vee \text{ (dim 5)}, \quad \Lambda \leftrightarrow \Lambda^2 E \text{ (dim 10)}.$$

The cosmological constant lives on the *pair* sector $\Lambda^2 E$, $\dim = \binom{5}{2} = 10$; this is why the decuple bridge has exponent 10 — Λ is a product over all $\binom{5}{2}$ carrier pairs (K_5 edges), not over the full 16-state family.

2.1 $g_{\text{car}} = 5$ as the Pascal closure

A family is exactly the complete Pascal truncation to degree two:

$$\dim S^+ = 2^{g_{\text{car}}-1} = \binom{g_{\text{car}}}{0} + \binom{g_{\text{car}}}{1} + \binom{g_{\text{car}}}{2} \iff g_{\text{car}} = 5 \quad (16 = 1 + 5 + 10), \quad [\text{E}], \quad (2)$$

the unique positive-integer solution of $f(g) := 2^{g-1} - 1 - \frac{g(g+1)}{2} = 0$. *Proof of uniqueness.* Direct evaluation gives $f(1)=-1$, $f(2)=-2$, $f(3)=-3$, $f(4)=-3$, $f(5)=0$, $f(6)=+10$; so $g=5$ is the only root in $\{1, \dots, 6\}$. For $g \geq 6$ the forward difference is $f(g+1) - f(g) = 2^{g-1} - (g+1) \geq 2^5 - 7 = 25 > 0$ and increasing, so f is strictly increasing on $[6, \infty)$ from $f(6)=10 > 0$; hence $f(g) > 0$ for all $g \geq 6$ and there is *no* further root. Thus $g_{\text{car}} = 5$ is the unique solution (no finite-range caveat). [\[verification/v2_carrier_pascal.py\]](#) This is sharper than $2g_{\text{car}} = \binom{g_{\text{car}}}{2}$: the family code (even states, 2^{g-1}) equals the action code (sum of the first three Pascal entries) only at $g_{\text{car}} = 5$.

Carrier uniqueness theorem: $g_{\text{car}} = 5$ and the 3+2 split are forced [E]

The carrier rank and its colour/weak split are not chosen — they are the unique solutions of two integer conditions tied to the E_8 closure and the SM gauge algebra:

$$g_{\text{car}} = 5 \text{ is the unique } g \text{ with } N_{\text{fam}} = \frac{2^{g-1}-1}{g} \in \mathbb{Z}^+ \text{ and } \text{rank } E_8 = g + N_{\text{fam}} = 8 \quad (3)$$

$$(b, s) = (3, 2) \text{ is the unique split of } b + s = g_{\text{car}} = 5, \quad b^2 + s^2 = 13 = |R(A_3)| + 1 \quad (4)$$

The integer-family carriers are $g \in \{3, 5, 7, \dots\}$ (rank 4, 8, 16), so $\text{rank } E_8 = 8$ picks $g_{\text{car}} = 5$ uniquely; and $b^2 + s^2 = 13$ with $b+s = 5$ forces (3, 2), reproducing $\dim \mathfrak{g}_{\text{SM}} = (b^2-1) + (s^2-1) + 1 = 12 = |R(A_3)|$. The induced hypercharge is anomaly-free ($\text{Tr } Y = \text{Tr } Y^3 = 0$ over the 16 states). This hardens P2: the *rank* and the *split* are forced; only the seam→carrier interface itself remains a declared input. [\[verification/v14_carrier_uniqueness.py\]](#)

The Pascal grammar: one $\Lambda^\bullet$ mechanism in four places (anti-numerology) [E]. The recurring small integers are *not* coincidences — they are the *same* exterior-algebra (Pascal) mechanism read in four spaces, which is why the binomials of rows 4 and 5 keep reappearing:

space	Pascal reading
carrier (one generation)	$\dim S^+ = \Lambda^{\text{even}}(5) = \binom{5}{0} + \binom{5}{2} + \binom{5}{4} = 1 + 10 + 5 = 16$
scale ladder (cosmology)	$A_{\text{EW}} : \mathcal{A}_H : \mathcal{A}_\Lambda = \binom{5}{0} : \binom{5}{1} : \binom{5}{2} = 1 : 5 : 10$
flavor u/d readout	$\ \text{Pl}(K) \ _1 = \binom{4}{0} + \binom{4}{1} + \binom{4}{2} = 1 + 4 + 6 = 11 = 16 - g_{\text{car}}$
mass volume	$\det M_{\text{SM}} \sim (\varphi_0^{\text{ret}})^{6+9+10} = (\varphi_0^{\text{ret}})^{25} = (\varphi_0^{\text{ret}})^{g_{\text{car}}^2}$

So the carrier is the even row 5, the scale ladder is the lower row 5, the u/d leg is the cumulative row 4 up to $\deg u^c = 2$, and the mass volume closes on g_{car}^2 . *One* grammar (Λ^k of the 5-slot carrier and the $4 = |\mu_4|$ family fundamental), not four lucky hits. [\[verification/v44_carrier_exterior.py\]](#) [\[verification/v45_family_exterior.py\]](#) [\[verification/v46_grand_mass_volume.py\]](#)

2.2 One master moment generates 40, 41, 240, δ_2

With $X = 6Y$, the quadratic hypercharge trace on one family is $\text{Tr}_{S^+} X^2 = 120 = 5!$, and it generates the whole integer chain:

$$\sum_{f,j} L_{f,j} = \frac{\text{Tr}_{S^+} X^2}{3} = 40, \quad 10b_1 = \frac{\text{Tr}_{S^+} X^2}{3} + 1 = 41, \quad (5)$$

$$|R(E_8)| = 2 \text{Tr}_{S^+} X^2 = 240, \quad \delta_2 = 4! \text{Tr}_{S^+} X^2 c_3^8.$$

The flavor budget, the $U(1)$ budget, the E_8 count and the second seam defect all hang on the second hypercharge moment $5!$. [E]

2.3 Compact E_8 compression and the role of ν^c

$$\begin{aligned} |R(E_8)| &= \dim S^+ \cdot g_{\text{car}} \cdot N_{\text{fam}} = 16 \cdot 5 \cdot 3 = 240, \\ \dim E_8 &= \dim S^+(\mathcal{A}_H + \mathcal{A}_\Lambda) + (g_{\text{car}} + N_{\text{fam}}) = 240 + 8 = 248 \end{aligned} \quad (6)$$

with $N_{\text{fam}} = (2^{g_{\text{car}}-1} - 1)/g_{\text{car}} = 15/5 = 3$ and $\text{rank } E_8 = g_{\text{car}} + N_{\text{fam}} = 8$. [E]. The right-handed neutrino ν^c is the Pascal singleton $\Lambda^0 E = (1, 1)_0$ that closes $16 = 1 + 5 + 10$; removing it breaks the truncation and shifts α^{-1} by 2.16×10^{-3} .

3 The $D_5 \times A_3$ compiler: E_8 as a closure, not an input

This section upgrades the E_8 atlas (Section 10) from a counting coincidence to an explicit lattice construction. The whole content is consolidated and verified in the standalone companion note “*The Even Carrier Code and the $\mathbb{Z}_{30}$ Coxeter Compiler*.” The thesis: the carrier code is the D_5 half-spinor, the family geometry is an A_3 , and E_8 is the standard unimodular glue closure of $D_5 \oplus A_3$ — with the four punctures μ_4 as the glue code.



$D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ — the unique even unimodular rank-8 glue [E] exact (lattice).

The compiler in one line:


five carrier slots $\Rightarrow D_5$, four family corners $\mu_4 \Rightarrow A_3$, $(D_5 \oplus A_3) + \mu_4\text{-glue} \Rightarrow E_8$.



Figure 2: The explanation hierarchy. The boundary fixes the seam seed c_3 ; the five-slot even carrier code is the D_5 half-spinor; the four punctures μ_4 give A_3 ; their common $\mathbb{Z}_4$ discriminant (μ_4) glues $D_5 \oplus A_3$ into the unimodular E_8 . E_8 is the *closure*, not the input; c_3 dresses the continuous observables.

The glue is one operation in two categories. The same order- $|\mu_4|=4$ discriminant glue closes $D_5 \oplus A_3$ into E_8 *twice over*: once on *root lattices* (the unimodular pushout $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$, v1) and once on *chiral conformal nets* (the simple-current / anyon-condensation extension $(D_5)_1 \otimes (A_3)_1 \rtimes \mu_4 \cong (E_8)_1$, v92/v281). These are not two coincidences but the two functorial faces of a single commuting square (v234/v277/v260); the boundary realisation on which the strict-TOE residual turns (SEAM.EQUIV.01) is exactly the statement that the raw seam realises

this square — now *closed modulo a cited theorem* (the lattice/ S_3 stack pins it at every computable level, Lean `FORM.SEAM.MMST.01`; only the cited MMST/OS continuum-existence theorems stay [O], v367/v368/v376–v379/v336). The extension leg of that citation package is itself re-founded at net level ([verification/v469_seam_crossedproduct_route.py]): the 128-spinor step is the *local* $\mathbb{Z}_2$ simple-current crossed product ($h_s(SO(16)_1) = 16/16 = 1 \in \mathbb{Z}$, the Longo–Rehren locality criterion; KLM $\mu = 4/2^2 = 1$), on 1995–2001 peer-reviewed subfactor theory, with the lattice-VOA preprint route demoted to an independent second witness, and the realisation input reduced to invariant level (R1'). `SEAM.EQUIV.01` stays [O].



One operation, two categories: the order- $|\mu_4|=4$ glue is the lattice pushout $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ and the simple-current / anyon-condensation extension $(D_5)_1 \otimes (A_3)_1 \rtimes \mu_4 \cong (E_8)_1$; the square commutes. [E] exact.

3.1 Carrier = $D_5 = \mathfrak{so}_{10}$ (proved)

The 16 even-occupation states $C^+ = \{x \in \mathbb{F}_2^{g_{\text{car}}} : |x| \text{ even}\}$ are exactly the weights of the D_5 half-spinor $\frac{1}{2}(\pm 1, \dots, \pm 1)$ (even number of minus signs). The flip-weight-4 relation is the **Clebsch graph** $\text{SRG}(16, 5, 0, 2)$ and the flip-weight-2 relation is its complement $\text{SRG}(16, 10, 6, 6)$; the *affine* automorphism group of the scheme (translations $\rtimes$ coordinate permutations) is the D_5 Weyl group,

$$\boxed{\text{Aut}_{\text{aff}}(C^+) = C^+ \rtimes S_5, \quad |\text{Aut}_{\text{aff}}(C^+)| = 2^{g_{\text{car}}-1} g_{\text{car}}! = 1920 = |W(D_5)| = r |R(E_8)|} \quad [\text{E}], \quad (7)$$

(the linear code automorphism group alone is only S_5 ; the order-1920 group is the affine/scheme one) and the Clebsch edge count is the flavor budget $|E(R_4)| = \frac{16 \cdot 5}{2} = 40 = |R(D_5)| = \sum_{f,j} L_{f,j}$.

3.2 Family = $A_3 = \mathfrak{su}_4$ from $\mathbb{P}^1 \setminus \mu_4$ (proved, with the metric settled)

The carrier geometry fixes $X_f^\circ \cong \mathbb{P}^1 \setminus \mu_4$, $H_1 = \mathbb{Z}^3$, rigid D_4 generated by $z \mapsto iz$, $z \mapsto 1/z$. The twelve puncture-difference cycles $\gamma_i - \gamma_j$ carry the residue pairing $\text{Res}(\omega_i, \gamma_j) = \delta_{ij}$ and form the A_3 root system:

$$\boxed{\#\{\gamma_i - \gamma_j\} = 12 = |R(A_3)| = \dim \mathfrak{g}_{\text{SM}}, \quad \text{rank} = 3 = N_{\text{fam}}, \quad \text{Gram} = \text{Cartan}(A_3), \quad \det = 4} \quad [\text{E}]. \quad (8)$$

The corner rotation $z \mapsto iz$ acts on $H^1 = \mathbb{C}^3$ as the A_3 Coxeter element (order $4 = h(A_3) = |\mu_4|$, spectrum $\{i, -1, -i\}$, $\chi = \Phi_4 \Phi_2$), an isometry of the Cartan form — the exact analogue of the E_8 Coxeter element reading Φ_{30} .

The μ_4 clock is the order-4 character of the E_8 cycle, and $(2, 3, 5)$ is the icosahedron ([verification/v223_coxeter_totative_clock.py], [verification/v219_icosahedral_mckay.py]). The two “Coxeter” clocks are one object read at two scales. The E_8 exponents are the totatives $(\mathbb{Z}/30)^\times = \{1, 7, 11, 13, 17, 19, 23, 29\}$, and the multiplicative order-4 element 7 ($7^2=19$, $7^4=1 \pmod{30}$) generates a literal μ_4 clock $\langle 7 \rangle = \{1, 7, 13, 19\}$ that permutes the four conjugate Coxeter 2-planes — the same μ_4 as the seam corner rotation. And the period $30 = 2 \cdot 3 \cdot 5$ itself is not a free factorisation: by the McKay correspondence E_8 is the exceptional *top* of the tower of finite $SU(2)$ subgroups ($2T \rightarrow \hat{E}_6$, $2O \rightarrow \hat{E}_7$, $2I \rightarrow \hat{E}_8$), so its branch data is the *icosahedron* and the three atoms $(|\mathbb{Z}_2|, N_{\text{fam}}, g_{\text{car}}) = (2, 3, 5)$ are its rotation-axis orders. The binary icosahedral group $2I$ (order $120 = |R^+(E_8)|$) has nine

irreducible-representation degrees $\{1, 2, 2, 3, 3, 4, 4, 5, 6\}$ equal to the affine- E_8 Kac marks, with $\sum d_i = 30 = h(E_8)$ and $\sum d_i^2 = 120$. This is a *backward certificate* of the closed E_8 [E] (it explains the $(2, 3, 5)$ choice [C]; it does *not* replace the P2 input).

The metric, settled (former [C]). The transcendental geometric (Green) metric $G_{ij} = -\log |p_i - p_j|$ on μ_4 is only D_4 -symmetric, with a regularisation-independent anisotropy $-\log 2$ on the root space, so it is *not* proportional to the A_3 Cartan form. But it does not need to be: the E_8 -relevant A_3 is the canonical *residue* pairing (integral, S_4 -symmetric, $= \text{Cartan}(A_3)$), and the geometric metric is a D_4 -equivariant deformation preserving the root system, Weyl group and Coxeter element. So the lattice structure is exact [E]; the analytic metric is a characterised deformation, not an open gap.

3.3 The μ_4 glue and the $5/4 + 3/4 = 2$ discriminant theorem (proved)

The two factors have equal discriminant groups and E_8 is unimodular, so the glue index is exactly $|\mu_4|$:

$$\text{disc}(D_5) = \text{disc}(A_3) = \mathbb{Z}_4, \quad [E_8 : D_5 \oplus A_3] = \sqrt{\frac{\det D_5 \det A_3}{\det E_8}} = \sqrt{\frac{4 \cdot 4}{1}} = 4 = |\mu_4|. \quad (9)$$

The discriminant-form norms sum to the E_8 root norm, and — strikingly — they are pre-existing TFPT constants:

$$\boxed{\begin{aligned} q(D_5) &= \frac{5}{4} = \frac{\delta_2}{\delta_{\text{top}}^2}, & q(A_3) &= \frac{3}{4} = \xi_{\text{tree}}, \\ q(D_5) + q(A_3) &= 2 = |E_8 \text{ root}|^2 \end{aligned}} \quad [\text{E}] \quad [\text{verification/v1_e8_glue.py}] \quad (10)$$

This is literally the E_8 spinor-root split $\frac{1}{2}(\pm)^8 = \frac{1}{2}(\pm)^5 \oplus \frac{1}{2}(\pm)^3$, $5 \cdot \frac{1}{4} + 3 \cdot \frac{1}{4} = 2$.




Figure 3: The discriminant-norm glue. The E_8 spinor root $\frac{1}{2}(\pm)^8$ (norm 2) splits across $D_5 \times A_3$ as $5 \cdot \frac{1}{4} + 3 \cdot \frac{1}{4}$; the two halves are the pre-existing TFPT constants $\delta_2/\delta_{\text{top}}^2 = \frac{5}{4}$ and $\xi_{\text{tree}} = \frac{3}{4}$.

The standard branching is realised explicitly:

$$\mathfrak{e}_8 = (45, 1) \oplus (1, 15) \oplus (10, 6) \oplus (16, 4) \oplus (\overline{16}, \overline{4}), \quad 248 = 45 + 15 + 60 + 64 + 64, \quad 240 = 40 + 12 + 60 + 64 + 64, \quad (11)$$




Figure 4: The $E_8 \supset D_5 \times A_3$ branching as norm-2 blocks. The adjoint half is the hypercharge moment $\text{Tr}_{S^+} X^2 = 120 = \dim \mathfrak{so}_{16}$; the spinor half is two sheets over the four corners μ_4 .

and assembling $D_5 \oplus D_3 \subset D_8$ with the even spinor coset produces all 240 roots (norm 2, Cartan $\det = 1$, every root an integer combination of one simple-root basis): *the 240-metrisation is no longer blind — E_8 is built.* [E]

3.4 Glue-norm arithmetic, and three readings of 41, 60, 120

The two glue norms generate the carrier integers through sum, difference and product. With $\dim S^+ = 16$:

$$\boxed{16(q_D + q_A) = 32 = 2^{g_{\text{car}}} = h(D_5)h(A_3), \quad 16(q_D - q_A) = 8 = r(E_8) = h(D_5), \quad 16 q_D q_A = 15 = \dim S^+ - 1} \quad [\text{E}], \quad (12)$$

so the glue already contains the full carrier exhaustion $2^{g_{\text{car}}}$, the E_8 rank, and the non-trivial transition count $\mathcal{A}_H + \mathcal{A}_\Lambda = 15$; and $|\mu_4| = (q_D - 1)^{-1} = (1 - q_A)^{-1} = 4$ is the inverse displacement of the two norms from the unit norm.

Spine Quotient Lemma: the recurring factors are adjacent quotients of (2,3,4,5) [E]
(`[verification/v74_compiler_micro_lemmas.py]`)

The integers of the spine are $(|\mathbb{Z}_2|, N_{\text{fam}}, |\mu_4|, g_{\text{car}}) = (2, 3, 4, 5)$. Every “mysterious” $O(1)$ factor in TFPT is an *adjacent quotient* of this chain, not a separate special number:

$$\frac{|\mathbb{Z}_2|}{N_{\text{fam}}} = \frac{2}{3}, \quad \frac{N_{\text{fam}}}{|\mu_4|} = \frac{3}{4} = q(A_3), \quad \frac{|\mu_4|}{g_{\text{car}}} = \frac{4}{5}, \quad \frac{g_{\text{car}}}{|\mu_4|} = \frac{5}{4} = q(D_5), \quad \frac{|\mu_4|}{N_{\text{fam}}} = \frac{4}{3}.$$

These are exactly the factors that recur across the theory: $\lambda_2 = (\frac{2}{3})^6$ (the transport gap = Page recovery rate, and the Koide target $Q_\star = \frac{2}{3}$); $\varepsilon = \frac{3}{4}\varphi_0 = q(A_3)\varphi_0$ (the solar misalignment); $\varphi_0 = \frac{4}{3}c_3 + 48c_3^4$ with leading $\frac{4}{3}c_3 = \frac{1}{6\pi}$; and $q(D_5) + q(A_3) = \frac{5}{4} + \frac{3}{4} = 2$ (the E_8 root norm). So $(\frac{2}{3}, \frac{3}{4}, \frac{4}{5}, \frac{5}{4}, \frac{4}{3})$ are five projections of one discrete chain — not scattered constants. The spine is moreover *one* finite object: as a tetrahedron on $T = \{e_3(a), p_0(a), e_1(a), e_2(a)\}$ its edge and face products and its volume $120 = |R^+(E_8)| = 5!$ reproduce the central integer grammar, with the honest negative control that the graph reading $240 = |\mu_4| \cdot |E(K_4)| \cdot |E(K_5)|$ breaks at K_6 (`[verification/v91_spine_tetrahedron.py]`).

The carrier adjoint $\dim D_5 = |R(D_5)| + g_{\text{car}} = 45$ then gives a second reading of the $U(1)$ integer,

$$\boxed{10b_1 = \dim D_5 - |\mu_4| = 45 - 4 = 41 = \underbrace{40}_{\sum L} + \underbrace{(g_{\text{car}} - |\mu_4|)}_{=N_\Phi=1}} \quad [\text{E}], \quad (13)$$

i.e. the single Higgs doublet is the carrier rank minus the glue corners, $5 - 4 = 1$.

A fourth, metric reading of 41 (`[verification/v222_cm_norm_duality.py]`). On the square seam ($j=1728$, complex multiplication by the Gaussian integers $\mathbb{Z}[i]$) the $U(1)$ index is the Gaussian norm of the carrier-glue vector,

$$10b_1 = N_{\mathbb{Z}[i]}(g_{\text{car}} + i|\mu_4|) = g_{\text{car}}^2 + |\mu_4|^2 = 5^2 + 4^2 = 41,$$

and the hexagonal partner ($j=0$, Eisenstein $\mathbb{Z}[\omega]$) reads the same (3,2) carrier split as the scalaron exponent $N_{\mathbb{Z}[\omega]}(3+2\omega) = 3^2 - 3 \cdot 2 + 2^2 = 7$. So 41 (EM) and 7 (scalaron) are the two exceptional CM norms of the one carrier, square vs. hexagonal — not two unrelated $O(1)$ numbers [E]. The start defect 60 and the hypercharge moment 120 are “bilingual”:

$$\begin{aligned} D_{\text{start}} &= \dim D_5 + \dim A_3 = 45 + 15 = 60 = \mathcal{A}_\Lambda \cdot |R^+(A_3)| = 10 \cdot 6, \\ \text{Tr}_{S^+} X^2 &= 120 = \underbrace{60}_{\text{adjoint}} + \underbrace{60}_{\text{mixed (10,6)}}, \end{aligned} \quad (14)$$

and $\dim E_8 = 248 = \text{Tr}_{S^+} X^2 + 2 \dim S^+ |\mu_4| = 120 + 2 \cdot 16 \cdot 4$. Finally the hypercharge norm and the EW exponent are A_3 -rational,

$$\boxed{\gamma_{\text{car}} = \frac{\mathcal{A}_\Lambda}{|R(A_3)|} = \frac{10}{12} = \frac{5}{6}, \quad I_1^{\text{EW}} = \frac{|E(K_5)|}{|R(A_3)|} + \frac{1}{|\mu_4||R(A_3)|} = \frac{10}{12} + \frac{1}{48} = \frac{41}{48}} \quad [\text{E}]. \quad (15)$$

Transport reduction. The flavor matrix L has its mod-6 content fixed by a single combinatorial datum: $\{0, 1, 3\}$ is the *unique* (up to D_6) 3-subset of $C_6 = |R^+(A_3)|$ with all distinct cyclic distances

$\{1, 2, 3\}$, and the three sectors are its D_6 orbit ($u \equiv \{0, 1, 3\}$, $d \equiv 2 - \{0, 1, 3\}$, $e \equiv 2 + \{0, 1, 3\}$). The total is fixed by the carrier, $\sum L = 40 = |R(D_5)|$, with winding unit $6 = |R^+(A_3)|$. Given the residue *set* of a sector, the full ordered packet is reproduced by two universal rules — the lightest generation carries the single $+6$ winding, and the entries are sorted in decreasing length (monotone hierarchy). The *only* remaining datum is then, per sector, which residue is the first-generation anchor that receives the winding:

$$\boxed{\text{anchors } (a_u, a_d, a_e) = (1, 1, 2) \Rightarrow L_u = (7, 3, 0), L_d = (7, 5, 2), L_e = (8, 5, 3)} \quad [\mathbf{E}] \text{ (given anchors).} \quad (16)$$

So the open step of a full “Clebsch walk” theorem is sharply isolated: it is exactly these three per-sector anchors. They are *not* free: the three transport cusps are exactly the D_5 dual-root combinations, and each sector is tied to its cusp by the right-handed hypercharge,

$$\boxed{\{1, \frac{2}{3}, \frac{1}{3}\} = \{2y_+, -2y_-, 2(y_+ + y_-)\}, \quad \text{lepton} \leftrightarrow 2y_+, \text{ up} \leftrightarrow -2y_-, \text{ down} \leftrightarrow 2(y_+ + y_-)} \quad [\mathbf{E}]. \quad (17)$$

The canonical-transport-kernel theorem (document 3, Part II) derives the residue packets — and hence the anchors — as the *word-lengths of the family-connection holonomy* $\text{Hol}_{\mathcal{G}_{D_4}}(\nabla_F^*)$ along the D_4 -invariant geodesic spine of $\mathbb{P}^1 \setminus \mu_4$ with these cusps as boundary data. The anchors are therefore a genuine $D_5 \times A_3$ readout (cusps from the D_5 carrier, spine holonomy from the A_3 four-punctured sphere), not an extra parameter. A compact candidate closed form is the nearest-integer cusp map

$$\boxed{a_f = \text{round}(2 \text{ cusp}_f) : (a_u, a_d, a_e) = \text{round}(\frac{4}{3}, \frac{2}{3}, 2) = (1, 1, 2)} \quad [\mathbf{C}], \quad (18)$$

which reproduces the matrix exactly. The remaining open step is now precise and purely geometric: an *independent* closed-form evaluation of the D_4 -spine holonomy word-lengths from the lattice data alone (the same $\mathbb{P}^1 \setminus \mu_4$ whose residue pairing is $\text{Cartan}(A_3)$ and whose corner rotation is the A_3 Coxeter element), rather than from the family-connection theorem (document 3).

Riemann–Hilbert check, and what it does (and does not) fix. Setting up the rigid $\mathbb{Z}_4$ -symmetric rank-3 $SU(3)$ local system explicitly — four puncture monodromies $M_k = R^k C R^{-k}$ with the order-3 cusp class $C \sim \text{diag}(1, \omega, \omega^2)$ ($\omega = e^{2\pi i/3}$, the three cusps as eigenvalue-exponents) and the rotation R ($R^4 = \mathbb{K}$) — the closure $M_0 M_1 M_2 M_3 = \mathbb{K}$ has a solution to machine precision, and the unique R has eigenvalues $\{i, -1, -i\}$: the A_3 Coxeter element. This *independently* confirms the rigidity of the family connection and the corner-rotation = Coxeter identity. However, the flat monodromy *alone* does not fix the integer word-lengths $\{0, 1, 3\}$: those are *parabolic degrees* (the cusp exponents as parabolic weights, together with the $\mathbb{Z}_2$ sheet parity), i.e. Hodge/metric data on the spine, not topological monodromy. *Status update (the computation has since been carried out)*: the exact residue normal form A_0^* with $\text{spec} = \{0, \frac{1}{3}, \frac{2}{3}\}$ is pinned algebraically ([verification/v115_anchor_residue.py], [verification/v116_resonance_uniqueness.py]), the unitarised holonomy is the exact 24-element $W(A_3) = S_4$ ([verification/v117_monodromy_weyl_a3.py]), and the parabolic weights are the μ_4 -character degrees of $H^1(\mathbb{P}^1 \setminus \mu_4)$ — one spectrum in four readouts ([verification/v137_qplus_cohomology.py]); the deck itself is *derived* — the integer model selects the sheet-twisted class, no discrete freedom left ([verification/v141_deck_selection.py]), and the full Möbius D_4 of the curve matches the integer model parity by parity ([verification/v146_moebius_d4.py]). What remains of the old residual is the much smaller selector establishment $R4'$ (research contracts): derive the *one* lift map of the torsion normal — the pairing values are atom identities ([verification/v142_frame_integrality.py]/[verification/v145_pairing_atoms.py]). $[\mathbf{E}]/[\mathbf{C}]$

3.5 What the compiler outputs for the Standard Model and our reality

Every entry below is an *output* of (c_3, g_{car}) through the $D_5 \times A_3$ compiler; none is an independent input. “Reads” names the lattice object the observable comes from.




Figure 5: Flavor transport on the hexagon $C_6 = Z_6 = |R^+(A_3)|$. The unique distinct-distance triple $\{0, 1, 3\}$ (cyclic distances $\{1, 2, 3\}$) and its D_6 orbit give the three sectors' residues mod 6; the total $\sum L = 40 = |R(D_5)|$ and the first-generation +6 winding come from the carrier.

SM / cosmology observable	Compiler output	Reads	St.
Number of generations	$N_{\text{fam}} = 3 = \text{rank } A_3 = \dim H^1(\mathbb{P}^1 \setminus \mu_4)$	A_3	[E]
Gauge algebra dimension	$\dim \mathfrak{g}_{\text{SM}} = 12 = R(A_3) = \gcd(6r, 2h)$	A_3	[E]
One generation (16, incl. ν^c)	$S^+ = D_5$ half-spinor $= \Lambda^{0,2,4} E$	D_5	[E]
Hypercharge spectrum	all Y from $y_{\pm} = \{\frac{1}{2}, -\frac{1}{3}\}$; $\text{Tr } X^2 = 120$	D_5 root data	[E]
Abelian index b_1	$10b_1 = 1 + \mu_4 E(K_5) = 1 + 4 \cdot 10 = 41$	$\mu_4 \times K_5$	[E]
Fermion family occupancy	$\Omega_{\text{adm}} = N_{\text{fam}} \dim S^+ = (N_{\text{fam}} + N_{\Phi}) R(A_3) = 48$	A_3	[E]
Fine structure $\alpha_{\star}^{-1} = 137.0359992168$	root of $F_{U(1)}(\alpha) = 0$ (explicit, unique)	EM closure	[E]
Cabibbo angle	$\lambda_C = \sqrt{u(1-u)}$, $u = \frac{4}{3}c_3 + 48c_3^4$	seed	[E]
CKM CP phase	$\delta_{\text{CKM}} = \frac{2\pi}{ R^+(A_3) } + N_{\text{fam}} u(1-u) = \frac{\pi}{3} + 3\lambda_C^2$	A_3^+ , seed	[E]/[C]
PMNS angles/phase	$\sin^2 \theta_{12} = \frac{1}{3} - \frac{u}{2}$, $\sin^2 \theta_{13} = e^{-5/6} u$, $\delta^\nu = \frac{4\pi}{3}$	A_3^+ , seed	[E]/[C]
Charged-lepton hierarchy	$\det M_\ell \propto u^{h/N_{\text{fam}}} = u^{10}$; $(5, 3, 2)$	decuple	[E]/[C]
Mass-hierarchy transport	$\sum L = 40 = R(D_5) $; entries on $C_6 = R^+(A_3) $	$D_5 \times A_3$	[E]
Strong CP	$\theta_{\text{eff}} = 0$ (determinant erasure, given RP; document 3, Part III)	—	[E]/[C]
Weak CP	holonomy: $\delta_{\text{CKM}} - \frac{\pi}{3} = N_{\text{fam}} u(1-u)$	A_3^+	[C]
Gravity tree normaliser	$\xi_{\text{tree}} = \frac{\dim \mathfrak{g}_{\text{SM}}}{\dim S^+} = \frac{3}{4} = q(A_3)$	A_3 glue	[E]
Cosmological constant	$\rho_\Lambda / \bar{M}_{\text{Pl}}^4 = \frac{3}{4\pi^2} e^{2\delta_{\text{ph}}} [v_{\text{geo}} / (5\beta_{\text{rad}}^2 \bar{M}_{\text{Pl}})]^{10}$	decuple	[E]
Hubble anchor	$H_0 \sqrt{\Omega_\Lambda} = \bar{M}_{\text{Pl}} e^{-\alpha_{\star}^{-1}} / (2\pi)$	action	[E]
Baryon density	$\omega_b = \lambda_C / \mathcal{A}_\Lambda = \lambda_C / 10$	decuple	[E]/[C]
Inflation (scalaron)	$M_{\text{scal}}^2 / \bar{M}_{\text{Pl}}^2 = c_3^{\Omega_{\text{adm}} - 10b_1} = c_3^7$	deficit $r - 1$	[E]
Inflation amplitude	$A_s = N_{\star}^2 c_3^7 / 24\pi^2 \approx 2.2 \times 10^{-9}$	deficit	[C]/[O]

Marker key: [E] exact / machine-proven · [C] conditional (holds under named hypotheses) · [O] open / axiom · [X] falsifiable kill test. Per-claim fine type in verification/status_ledger.csv.

Reading. The discrete sector — generations, gauge dimension, family content, hypercharge, b_1 , occupancy — is now *group-theoretic output* of $D_5 \times A_3$. The continuous sector — α^{-1} , λ_C , the mixings, ρ_Λ , H_0 , inflation — is the seam seed c_3 and the action exponentials read on the *same* carrier graphs (K_5 pairs, the C_6 hexagon $= |R^+(A_3)|$, the decuple $\mathcal{A}_\Lambda = \binom{g_{\text{car}}}{2} = 10$). What used to be a list of separate hits is, structurally, one machine: D_5 carries the family and the flavor budget, A_3 carries the generations and the gauge dimension, μ_4 glues them (and reappears as the 41 chain, the discriminant norms, and the CP phase step $2\pi/6$), and c_3 dresses everything.

4 Grammar I — the carrier grammar $(3 + 2)$

Machine checks for this section: [verification/v2_carrier_pascal.py]

From $g_{\text{car}} = 5$ and the split $(b, s) = (3, 2)$ the carrier data follow:

$$\begin{aligned} \gamma_{\text{car}} &= \frac{g_{\text{car}}}{g_{\text{car}} + 1} = \frac{5}{6}, & N_{\text{fam}} &= \frac{g_{\text{car}} + 1}{2} = 3, & \dim S^+ &= 2^{g_{\text{car}}-1} = 16, \\ \Omega_{\text{adm}} &= N_{\text{fam}} 2^{g_{\text{car}}-1} = 48, & b_1 &= \frac{g_{\text{car}} 2^{g_{\text{car}}-2} + 1}{10} = \frac{41}{10}. \end{aligned} \tag{19}$$

4.1 Dual-root dictionary (everything from two charges)

Two carrier roots generate the family and the transport cusps [E]. With $y_+ = \frac{1}{2}$ and $y_- = -\frac{1}{3}$ (the determinant-normalised carrier eigenvalues), the full one-family hypercharge spectrum and the flavor-transport cusps are sums of the two roots:

$$\begin{aligned} Y(Q_L) &= y_+ + y_- = \frac{1}{6}, & Y(u^c) &= 2y_- = -\frac{2}{3}, & Y(d^c) &= -y_- = \frac{1}{3}, \\ Y(L_L) &= -y_+ = -\frac{1}{2}, & Y(e^c) &= 2y_+ = 1, & Y(\nu^c) &= 0, \\ \{1, \frac{2}{3}, \frac{1}{3}\} &= \{2y_+, -2y_-, 2(y_+ + y_-)\}, & \gamma_{\text{car}} &= \text{Tr}_E Y^2 = 3y_-^2 + 2y_+^2 = \frac{5}{6}, \\ \varphi_{\text{base}} &= \frac{1 - \gamma_{\text{car}}}{\pi} = \frac{1}{6\pi}. \end{aligned}$$

All residuals are exactly zero. The Standard-Model charge table is not assumed; it is built by addition from $\{1/2, -1/3\}$.

The $X = 6Y$ quadratic and the hypercharge Pascal-sign duality [E]

Clearing denominators with $X = 6Y$ turns the two base charges into the integer roots $x_+ = 3$, $x_- = -2$ of

$$\boxed{x^2 - x - 6 = 0} : \quad x_+ + x_- = 1 = N_{\Phi}, \quad x_+ - x_- = 5 = g_{\text{car}}, \\ -x_+ x_- = 6 = |R^+(A_3)|, \quad \sqrt{\Delta} = 5,$$

so the Higgs singleton, the carrier rank and the A_3 hexagon all fall out of one quadratic. Counting the signs of X over a generation $(X = 1^{\times 6}, 2^{\times 3}, 6^{\times 1}, (-4)^{\times 3}, (-3)^{\times 2}, 0^{\times 1})$ gives the *Pascal triple* again:

$$\boxed{(\#X=0, \#X<0, \#X>0) = (1, 5, 10) = \binom{5}{0}, \binom{5}{1}, \binom{5}{2}},$$

$$10-5=g_{\text{car}}, \quad 5-1=|\mu_4|, \quad 10-1=\text{tr } R,$$

with $\nu^c \leftrightarrow \binom{5}{0}$ the neutral singleton — the right-handed neutrino is the Pascal zero with a job. The moment ladder adds $\text{Tr } X^2 = 120 = 5!$ and $\text{Tr } X^4 / \text{Tr } X^2 = 2280/120 = 19$ (an E_8 exponent; audit). [E]

4.2 Carrier arithmetic (proved identities)

The discrete coefficients are not isolated numbers; they are exact carrier traces:

$$10b_1 = 41 = g_{\text{car}}^2 + (g_{\text{car}} - 1)^2 = \Omega_{\text{adm}} \gamma_{\text{car}} + 1 = 1 + h_{E_8} \frac{\dim S^+}{\dim \mathfrak{g}_{\text{SM}}} = 1 + 30 \cdot \frac{16}{12} = 40 + 1, \quad [\text{E}] \quad (20)$$

$$I_1^{\text{EW}} = \frac{41}{48} = \text{Var}_{S^+}(Y) \cdot b_1 = \frac{5}{24} \cdot \frac{41}{10} = \gamma_{\text{car}} + \frac{1}{\Omega_{\text{adm}}}, \quad [\text{E}] \quad (21)$$

$$\text{Tr}_{S^+}(X^2) = 120 = 5! \quad (X = 6Y), \quad \delta_2 = \frac{5}{4} \delta_{\text{top}}^2 = 2880 c_3^8 = 4! \cdot 5! c_3^8 = 4! \text{Tr}_{S^+}(X^2) c_3^8, \quad [\text{E}] \quad (22)$$

$$\frac{5}{4} = \underbrace{\frac{3}{2} \cdot \frac{5}{6}}_{B\gamma} = \underbrace{\frac{D_{\text{start}}}{\Omega_{\text{adm}}}}_{60/48} = \underbrace{\frac{g_{\text{car}}}{g_{\text{car}}-1}}_{5/4} = \frac{\delta_2}{\delta_{\text{top}}^2}. \quad [\text{E}] \quad (23)$$

So: b_1 is a Pythagorean carrier identity ($5^2 + 4^2$); equivalently the anchor-product form $10b_1 = p_1 p_3 + (g_{\text{car}} - p_1) = 4 \cdot 10 + (5 - 4) = 41$ reads “all four glue corners times all ten carrier pairs, plus the one leftover Higgs slot”; the electroweak exponent $41/48$ is the hypercharge variance times the $U(1)$ budget; the second seam defect $2880 c_3^8$ is a product of factorial traces $4! \cdot 5!$; and the universal compression $5/4 = g_{\text{car}}/(g_{\text{car}} - 1)$ recurs whenever the five-slot carrier projects onto four effective directions. As a minor curiosity, the $SU(5)$ -tree weak mixing angle rewrites as $\sin^2 \theta_W^{\text{tree}} = \frac{3}{8} = N_{\text{fam}}/(g_{\text{car}} + N_{\text{fam}}) = N_{\text{fam}}/\text{rank}(E_8)$ (the value is the standard one; the rewrite is a coincidence to note, not a claim). [C]

5 Grammar II — the seed grammar (one decoder u)

Machine checks for this section: [\[verification/v106_review_validation.py\]](#)

The seed is, exactly, a pure function of c_3 :

$$u = \varphi_0^{\text{ret}} = \frac{4}{3} c_3 + \Omega_{\text{adm}} c_3^4 = \underbrace{\frac{1}{6\pi}}_{\varphi_{\text{base}}} + \underbrace{\frac{3}{256\pi^4}}_{\delta_{\text{top}}} = 0.0531719521768 \quad [\text{E}]. \quad (24)$$

Its two coefficients are themselves carrier data: $\frac{4}{3} = \dim S^+ / \dim \mathfrak{g}_{\text{SM}} = 16/12 = 8/(g_{\text{car}} + 1)$ (so $\varphi_{\text{base}} = 1/((g_{\text{car}} + 1)\pi)$) and $48 = \Omega_{\text{adm}}$. The seed is thus a pure function of c_3 and the carrier counts.


One seed, four readouts (and their inverses) [E].

$$\begin{aligned} \lambda_C^2 &= u(1-u), \quad \beta_{\text{rad}} = \frac{u}{4\pi}, \quad \Omega_b = \left(1 - \frac{1}{4\pi}\right)u, \quad \sin^2 \theta_{13} = e^{-5/6} u, \\ u &= \frac{1 - \sqrt{1 - 4\lambda_C^2}}{2} = 4\pi \beta_{\text{rad}} = \frac{\Omega_b}{1 - 1/(4\pi)} = e^{5/6} \sin^2 \theta_{13} \\ &= \sum_{n \geq 0} C_n \lambda_C^{2n+2} = \lambda_C^2 + \lambda_C^4 + 2\lambda_C^6 + 5\lambda_C^8 + \dots \end{aligned}$$

One knob u turns four dials simultaneously: Cabibbo mixing (the Bernoulli width $\sqrt{u(1-u)}$), cosmic birefringence, the baryon fraction, and the reactor angle. Turning one forces the others; there is no free per-observable tuning. The exponent $5/6$ is the carrier trace $\gamma_{\text{car}} = \text{Tr}_E Y^2$, *not* the Euler–Mascheroni constant $0.5772\dots$ (audit [O]).

6 Grammar III — the action grammar (exponentials of $\alpha_\star^{-1}$)

The relation in plain words. The fine-structure constant is not just “a number near 137”: it is the *one exponential engine* behind three wildly separated scales. Take the same $\alpha_\star^{-1} = 137.036$ and divide the carrier into it: $\frac{1}{5}$ of it sets the *electroweak* scale, $1\times$ it sets the *Hubble* scale, $2\times$ it sets the *cosmological constant*. The three action charges are the Pascal row $1 : 5 : 10$ of the five-slot carrier, and the Hubble rung is just the square root of the vacuum energy. So one constant fixes the electroweak hierarchy, dark energy and the expansion rate at once.



action charges $1 : g_{\text{car}} : 2g_{\text{car}} = 1 : 5 : 10 = \binom{5}{0} : \binom{5}{1} : \binom{5}{2}$ (Pascal row of K_5)

Read every dimensionless ratio through its *action* and ask for the action charge:

$$x = R_x e^{-A_x}, \quad A_x = q_x \alpha_\star^{-1} + \Delta_x, \quad q_x \in \left\{ \frac{1}{g_{\text{car}}}, 1, 2 \right\}. \quad (25)$$

The robust ladder and its residues are

sector	action A_x	charge q_x	residue R_x	value of $-\ln x$
electroweak $v_{\text{geo}}/\bar{M}_{\text{Pl}}$	$(\alpha_\star^{-1} + \delta_{\text{ph}})/g_{\text{car}}$	$1/g_{\text{car}}$	$g_{\text{car}}\beta_{\text{rad}}^2$	36.85
Hubble $H_0/\bar{M}_{\text{Pl}}$	$\alpha_\star^{-1}$	1	$1/(2\pi\sqrt{\Omega_\Lambda})$	138.68
cosm. const. $\rho_\Lambda/\bar{M}_{\text{Pl}}^4$	$2\alpha_\star^{-1}$	2	$3/(4\pi^2)$	276.65

$$A_{\text{EW}} = \frac{\alpha_\star^{-1} + \delta_{\text{ph}}}{g_{\text{car}}} = 27.5259, \quad \mathcal{A}_\Lambda = 2\alpha_\star^{-1} = 274.072, \quad \boxed{A_{\text{EW}}:\mathcal{A}_H:\mathcal{A}_\Lambda = 1:5:10}. \quad (26)$$

The leading ratio is exact: $\mathcal{A}_\Lambda/(\alpha_\star^{-1}/g_{\text{car}}) = 2g_{\text{car}} = 10$ (with the transport correction $\mathcal{A}_\Lambda/A_{\text{EW}} = 9.957$). The claim is that the *action charges*, not the raw logarithms, form the ladder.

6.1 The EM closure: $\alpha_\star^{-1}$ as an explicit fixed point (made explicit here)

The engine $\alpha_\star^{-1}$ is not an input — it is the unique root of an explicit closure equation (the exact electromagnetic closure theorem). With $c_3 = 1/(8\pi)$, $\varphi_{\text{base}} = 1/(6\pi)$, $q(\alpha) = \delta_{\text{top}}e^{-2\alpha}$ and the seam response $\varphi_{\text{seam}}(\alpha) = \varphi_{\text{base}} + q(\alpha)(1 - q(\alpha))^{-5/4}$,

$$\boxed{F_{U(1)}(\alpha) = \alpha^3 - 2c_3^3\alpha^2 - \frac{4}{5}c_3^6\left(\sum_{f,j} L_{f,j} + N_\Phi\right) \log \frac{1}{\varphi_{\text{seam}}(\alpha)} = 0} \quad [\mathbf{E}]/[\mathbf{C}], \quad (27)$$

where the coefficient is $\frac{4}{5}(\sum L + N_\Phi)c_3^6 = \frac{4}{5} \cdot 41 c_3^6 = 8b_1 c_3^6$, so the fine-structure constant reads the *flavor transport budget* $\sum L = 40$ and the $U(1)$ index $b_1 = 41/10$.

Lemma 1 (Existence and uniqueness of the EM fixed point). $F_{U(1)}$ has *exactly one* positive root, and it is simple.




Figure 6: Action tower (data plot, `verification/make_figures.py`). The three action charges $A_x = -\ln x$ for the electroweak, Hubble and cosmological-constant ratios fall on the line $A = (\alpha_\star^{-1}/5)\binom{5}{k}$: one engine $\alpha_\star^{-1}$ generates all three scales with the Pascal exponents 1 : 5 : 10.

Proof. Write $F_{U(1)}(\alpha) = h(\alpha) - C g(\alpha)$ with $h(\alpha) = \alpha^3 - 2c_3^3\alpha^2$, $C = \frac{4}{5}c_3^6 M > 0$ ($M = 41$), and $g(\alpha) = -\log \varphi_{\text{seam}}(\alpha)$. Since $q(\alpha) = \delta_{\text{top}} e^{-2\alpha} \in (0, 1)$ is strictly decreasing, $\varphi_{\text{seam}} = \varphi_{\text{base}} + q(1 - q)^{-5/4}$ is C^1 and decreasing, so g is C^1 , bounded, and slowly increasing with $g'(\alpha) = O(q)$ tiny; numerically $g \in [2.9342, 2.9343]$ on the admissible window. The cubic h has its interior critical point at $\alpha_c = \frac{4}{3}c_3^3 = 8.399 \times 10^{-5}$, with $h(0) = 0$ and $h < 0$ on $(0, \alpha_c)$. *Monotonicity (corrected endpoint).* Since $F'_{U(1)}(\alpha) = 3\alpha^2 - 4c_3^3\alpha - C g'(\alpha)$ and at α_c the first two terms *cancel* ($3\alpha_c = 4c_3^3$), one has $F'_{U(1)}(\alpha_c) = -C g'(\alpha_c) \approx -5.89 \times 10^{-10} < 0$: the derivative is *not* yet positive at α_c . Its first zero lies slightly above, at $\alpha_0 = 8.6264 \times 10^{-5}$ (where $3\alpha_0^2 - 4c_3^3\alpha_0 = C g'(\alpha_0)$), with $F'_{U(1)} < 0$ on (α_c, α_0) and $F'_{U(1)} > 0$ on (α_0, ∞) . Now $F_{U(1)}(0) = -C g(0) < 0$, and $F_{U(1)}$ *decreases* on $(0, \alpha_0]$ (both h and $-Cg$ decrease on $(0, \alpha_c]$, then $F'_{U(1)} < 0$ on $[\alpha_c, \alpha_0]$), so $F_{U(1)}(\alpha_0) \approx -3.82 \times 10^{-7} < 0$; thereafter $F_{U(1)}$ is strictly increasing to $+\infty$. Hence $F_{U(1)} < 0$ on $(0, \alpha_0]$ and strictly increasing on $[\alpha_0, \infty)$, so it crosses zero *exactly once*. The crossing has $F'_{U(1)}(\alpha_\star) = 1.58 \times 10^{-4} > 0$, so the root is simple. A sign scan on $(0, 0.05)$ confirms a single change. *Rigorous version:* an **interval-arithmetic** enclosure (`mpmath.iv`) brackets the root in $[0.00729735256220970, 0.00729735256221000]$ with $F_{U(1)} < 0$ on the lower and > 0 on the upper endpoint (definite signs), and a 200-cell interval partition of $(\alpha_c, 0.05)$ contains exactly *one* indeterminate (root-bearing) band — a verified uniqueness, not merely a sampled one. [`verification/v3_em_alpha.py`] $\square$

Solving:

$$\alpha_\star = 0.00729735256220985\dots, \quad \alpha_\star^{-1} = 137.0359992168407\dots \quad [\text{E}] \quad (28)$$

[`verification/v3_em_alpha.py`] versus CODATA-2022 $\alpha_\star^{-1} = 137.035999177(21)$ — a relative deviation 2.9×10^{-10} ($\approx 1.9\sigma$ on the tiny experimental error; the older 2018 value 137.035999084 gave 9.7×10^{-10} , so the agreement *improved* in relative terms). The exponent $-5/4$ in φ_{seam} is the carrier discriminant norm $q(D_5)$, and $\delta_{\text{top}} = 48c_3^4$, so the closure is built only from c_3 , the carrier, and the flavor budget. This discharges the former open audit point: the EM closure is a stated, existence-/uniqueness-proven, numerically reproduced equation, not a fit.

Why *this* function: the boundary $U(1)$ Ward reading (Theorem C) [E]/[C]. $F_{U(1)}$ is not a chosen ansatz but the reduced Ward identity of the boundary $U(1)$ partition function $Z_\partial = Z_{\text{Maxwell}} Z_{\text{Calderón}} Z_{\text{transport}}$, with three terms whose coefficients are compiler atoms, not fits:

$$\underbrace{\alpha^3}_{\text{Maxwell moment}} - \underbrace{2c_3^3\alpha^2}_{\text{Calderón sheet } (2=|\mathbb{Z}_2|)} - \underbrace{8b_1c_3^6 \log \frac{1}{\varphi_{\text{seam}}}}_{C_6 \text{ transport det.}} = 0, \quad 8b_1 = \frac{4}{5}M = \frac{4}{5}(\sum L + N_\Phi) = \frac{164}{5}.$$

The Calderón factor $2 = |\mathbb{Z}_2|$ (sheet), the transport power c_3^6 is the C_6 hexagon, and the seam exponent $-\frac{5}{4} = -q(D_5)$ is the carrier discriminant norm. **Typing:** the three-term decomposition and the coefficient identities ($8b_1 = \frac{4}{5}M$, $-\frac{5}{4} = q(D_5)$) are exact [E]; the *physical origin* as a boundary Ward closure (the lemmas $\partial_\tau \log Z_{\text{Maxwell}} = \alpha^3$, $\partial_\tau \log Z_{\text{Calderón}} = -2c_3^3\alpha^2$, $\partial_\tau \log \det_\zeta(1 - \mathcal{T}_{C_6, U(1)}) = -8b_1c_3^6 \log \frac{1}{\varphi_{\text{seam}}}$) is a conjectured [C] interpretation, not yet machine-proven. [verification/v48_em_ward.py]

The determinant-line (Quillen) form. ([verification/v341_alpha_quillen.py]) The three terms assemble into a single *stationarity* statement for the boundary $U(1)$ determinant line,

$$\underbrace{[\alpha^3 - 2c_3^3\alpha^2]}_{\delta \log \det \Delta_{U(1)}} = \underbrace{[8b_1c_3^6]}_{\text{ctr.}} \underbrace{[\log \frac{1}{\varphi_{\text{seam}}}]_{\text{seam resp.}}}$$

so $F_{U(1)} = 0$ is $\delta \log \det \Delta_{U(1)} + \text{counterterm} + \text{seam response} = 0$. Every factor is a *named* object, not a knob: the counterterm coefficient is $8b_1c_3^6 = (\text{rank } E_8) b_1 c_3^6$ (index $\times$ heat-kernel), $c_3 = 1/(8\pi)$ is the Gauss–Bonnet boundary coefficient, the exponent $-\frac{5}{4} = -q(D_5)$ is one carrier discriminant norm and the seam base $\varphi_{\text{base}} = 1/(6\pi) = \frac{4}{3}c_3$ carries the *other*, $c_3/\varphi_{\text{base}} = \frac{3}{4} = q(A_3)$ — both glue forms appear, and the split is verified at the root. **Typing (and: no new gate).** The determinant-line assembly and all coefficient identities are exact [E]; $\alpha_\star^{-1}$ itself stays closed [E] (unique root + ablation). This adds *no* open item — it *sharpens* the pre-existing [O]/physical-origin obligation EM.WARD.01 (the “why *this* function” residual, already conjectural) by naming every coefficient. The one still-open part *is* that residual, now *named* as its own tracked target ALPHA.QUILLEN.EXACT.01 ([verification/v382_alpha_quillen_exact.py]): the from-first-principles AQFT/ ζ proof that $\delta_\tau(\log \det_\zeta \Delta_{U(1)} + 8b_1c_3^6 \log \varphi_{\text{seam}}) = 0$ — i.e. that $F_{U(1)}$ is the *exact* Quillen functional of the seam $U(1)$, the variational principle *derived*, not assembled. It is a *face* of SEAM.EQUIV.01 (v336) — the seam $U(1)$ determinant line lives on the same holomorphic net, so once the net is $(E_8)_1$ its $U(1)$ sub-determinant is fixed — hence no second independent gate, just the sharpest remaining EM-Ward obligation, now citable on its own. *Honest progress towards it* ([verification/v391_alpha_quillen_progress.py]), *without closing it*: a solvable model $\det_\zeta(-\partial_\tau^2 + m^2)$ on a circle $= (2 \sinh(\beta m/2))^2$ has the closed m -variation $\beta \coth(\beta m/2) = 2/m + (\beta^2/6)m - (\beta^4/360)m^3 + \dots$, a heat-kernel/Laurent series — so a ζ -determinant’s variation is *structured* heat-kernel data, and the counterterm $8b_1c_3^6$ (a Gauss–Bonnet heat-kernel power) is the right *kind* of object. This *reduces* ALPHA.QUILLEN.EXACT.01 to the named sub-target “the seam $U(1)$ Seeley–DeWitt coefficient $= 8b_1c_3^6$ ”, but the target itself stays [O] (an external math proof, v384); the value $\alpha_\star^{-1}$ stays [E]. *A second honest step* ([verification/v433_alpha_quillen_heatkernel.py]) *bridges this reduction to the heat-kernel origin below, without closing it*: v391’s circle reached only the low orders a_0, a_1 , whereas a solvable $4D$ model — the flat $\Delta = -\partial^2 + m^2$ on T^4 , $\text{Tr } e^{-t\Delta} = V/(4\pi)^2 t^{-2} e^{-tm^2}$ — has $a_4 = Vm^4/(2(4\pi)^2) \neq 0$, *reaching* the t^0 (conformal-anomaly) order the circle could not, the very order in which the quadratic term is grounded below (v342, Gilkey a_4). So a ζ -determinant’s variation is heat-kernel data *at the a_4 order*, and the measure itself is c_3 -built, $(4\pi)^{-2} = (2c_3)^2 = 4c_3^2$; the *actual* seam $a_4 = 8b_1c_3^6$ and the cubic α^3 Chern moment stay [O] (v384), the seam F -normalisation [C] (retyped by the fifth step below, v470, as the embedding index $k_Y=5/3$).

The heat-kernel origin of the terms ([verification/v342_em_ward_heatkernel.py]). The determinant-line form is more than bookkeeping: its terms are textbook heat-kernel coefficients. [E] $c_3 = 1/(8\pi)$ *is* the boundary Gauss–Bonnet coefficient ($\oint_{S^2} K = 2\pi\chi = 4\pi$, $c_3 = 1/(|\mathbb{Z}_2| \oint K)$, the same $\chi=2$ as the marks, v216). [E] the α^2 (Calderón) term *is* the Gilkey gauge-curvature coefficient: the $U(1)$ Laplacian’s Seeley–DeWitt a_4 contains $30 \Omega_{ij} \Omega^{ij}/360 = \frac{1}{12} \Omega^2$ (Gilkey, *Invariance Theory*, Thm 4.8.16), and with $\Omega = i\alpha F$ this forces an $-(\alpha^2/12)F^2$ contribution to $\log \det \Delta_{U(1)}$ — so the Calderón $-2c_3^3\alpha^2$ is the gauge-curvature (Quillen-curvature) piece, not an ansatz. [E] the c_3 -orders $\{0, 3, 6\}$ are an arithmetic ladder (step 3 = the three channels; $6 = |R^+(A_3)|$, the C_6 hexagon), and the π -power test confirms it: $2c_3^3 = 1/(256\pi^3)$ carries π^3 , i.e. *three* boundary c_3 -insertions, not a single bulk coefficient. **Typing:** the boundary structure and the gauge-curvature origin of the quadratic term are heat-kernel-forced [E]; the exact coefficient value

is [C] (the seam F -normalisation — retyped as the embedding index $k_Y=5/3$ by the fifth step below, v470); the cubic α^3 Maxwell moment (a topological/Chern level) and the full “this is the seam $U(1)$ ζ -determinant” claim stay [O] — exactly the EM.WARD.01 residual, unchanged. So the quadratic/curvature half of the EM-Ward origin is now derived; the cubic-moment half remains the honest open part. *A third honest step ([verification/v434_alpha_quillen_betafunction.py]) settles those residuals and shows they are not three independent problems.* The counterterm’s matter factor is heat-kernel-forced too: by the $\beta = a_4$ theorem the one-loop $U(1)_Y$ β is the a_4 coefficient, and the carrier hypercharge content gives $\sum \frac{2}{3}Y_f^2 + \frac{1}{3}Y_s^2 = 41/6$ (SM) $\Rightarrow b_1 = 41/10$ (GUT, v159/v246), so $8b_1c_3^6 = b_1$ (rank E_8) c_3^6 is an a_4 content coefficient times a boundary geometry — the same kind of object as the Calderón a_4 . The exact value then turns only on the seam F -normalisation, so the “actual seam a_4 ” residual is the F -normalisation [C] (one obligation, not two); and the cubic factors $a^3 - 2c_3^3a^2 = a^2(a - 2c_3^3)$, exhibiting a^3 as the first moment of the Chern density a^2 — a level whose from-first-principles origin stays [O]. The EM-Ward residual is thus one [C] + one [O], not three. *A fourth honest step ([verification/v435_alpha_quillen_chernlevel.py]) sharpens that single remaining [O] — residual (2), the cubic α^3 Chern level — without closing it.* A π -power / metric-independence test partitions the closure: the three coefficients carry π -powers exactly $\{0, 3, 6\}$ ($k_0 = 1$; $-2c_3^3 = -1/(256\pi^3)$; $-8b_1c_3^6 = -41/(327680\pi^6)$), so the Maxwell a^3 is the unique π^0/c_3^0 (metric-independent) term. A heat-kernel boundary coefficient always carries the $(4\pi)^{-d/2}$ measure; a metric-independent term carries none — the signature of a *topological* term. So the test confirms, now from a test rather than a claim, that a^3 is a level and not a Seeley–DeWitt curvature integral (the other two terms are heat-kernel boundary data). [E] *Conditional* on the Chern–Simons/ η reading of the Maxwell moment (the EM1 interpretation, v48), large-gauge invariance quantises the seam $U(1)$ level to $\mathbb{Z}$, so $k_0 = 1$ is the *unit* (minimal) level — an integer, not a tunable real. [C] What stays [O] is the EM1 lemma itself — the from-first-principles Chern–Simons-boundary proof that $\partial_\tau \log Z_{\text{Maxwell}} = a^3$ — external math (type A, v384). *A fifth honest step ([verification/v470_alpha_inflow_level.py]) upgrades both leftovers, still without closing the target.* [C] *The level is computed, not minimal:* under the same EM1 reading, the a^3 coefficient $k_0 = 1$ equals the FHS bulk Chern invariant $|C| = 1$ of the very p+ip collar that realises S3 of SEAM.EQUIV.01 (v367/v461; control $C(M=3) = 0$) — “why exactly 1” is answered by a computation plus the cited inflow identification (TKNN / Avron–Seiler–Simon quantisation; Callan–Harvey inflow; the APS/Witten η =CS reading of $\delta \log \det$; Quillen/Bismut–Freed determinant-line integrality), replacing the minimality assumption of the fourth step. [E] *The F -normalisation is the embedding index:* $k_Y = \text{tr}_5(Y^2)/\text{tr}(T_3^2) = (5/6)/(1/2) = 5/3$ (Ginsparg 1987), so $(3/5) \times \frac{41}{6} = \frac{41}{10} = b_1$ exactly — the “GUT 3/5” is the inverse embedding index, and once the seam net is $(E_8)_1$ at level 1 the $U(1)_Y$ normalisation $\langle J(z)J(w) \rangle = k/(z-w)^2$ is fixed by it (level-1 current-algebra rigidity, no continuous dial): the seam F -normalisation [C] has zero independent content — a face of SEAM.EQUIV.01, per the v382 typing. [C] *One phase, two responses:* the collar phase carries $c_- = 8$ (gravitational, feeding SEAM.EQUIV.01/S3, v456) and $C = 1$ ($U(1)$, feeding EM1) — so discharging the one realisation input feeds both named targets at once. *A sixth honest step ([verification/v472_quillen_detline_moduli.py]) exhibits that bridge lemma at the finite level.* ALPHA.QUILLEN.EXACT.01 names the level as the curvature of the determinant line over the $U(1)$ moduli (Quillen/Bismut–Freed curvature, Dai–Freed section) — and the fifth step computed the Chern invariant only over the Bloch Brillouin zone, the translation-invariant shortcut. The sixth step computes the Quillen-shaped object itself: the same collar Hamiltonian in real space with twisted boundary conditions — the twist torus (θ_x, θ_y) is the moduli space of flat $U(1)$ connections — and the Fukui–Hatsugai–Suzuki curvature of the determinant line of the occupied frame over that torus integrates to $2\pi \cdot 1$ exactly (10^{-9}), size-independently ($L = 4, 6$), with clean controls (trivial collar $\rightarrow 0$, orientation flip $\rightarrow -1$), the twist-moduli integer equal to the Bloch-BZ integer for all three phases (Niu–Thouless–Wu), and the Fermi gap open over the whole torus (the Dai–Freed section has no zero). So “ $\delta \log \det(\text{seam model}) = \text{the inflow response}$ ” holds verbatim at the finite/model level, where $\log \det$ of the occupied frame replaces $\log \det_\zeta$. [E] (computation) / [C] (the EM1 reading). What stays [O] is the *continuum* identification “ $\delta \log \det_\zeta(\text{seam}) = \text{the$

inflow response” on the abstract seam (the `SEAM.EQUIV.01` face); `ALPHA.QUILLEN.EXACT.01` stays [O], and $\alpha_\star^{-1}$ stays [E] regardless.

The full ablation — which inputs are load-bearing and which only set the last digit — is in the [Ablation appendix](#) below.

Inversion test (this round). Running the closure backwards — inserting the *measured* α and solving for the integer budget $M = \sum L + N_\Phi$ — returns $M = 41.0000001\dots$, i.e. the external measurement reconstructs the internal discrete budget $41 = 40 + 1$ to 10^{-7} . This is stronger than “we hit α ”: the equation reads the discrete 41 out of nature. [E]

6.2 The decuple bridge (new, exact headline)

Eliminating $\alpha_\star^{-1} = g_{\text{car}} A_{\text{EW}} - \delta_{\text{ph}}$ between the electroweak and the cosmological-constant lines gives a direct relation between the two scales:

Decuple bridge: Λ is the tenth power of the stripped EW scale [E]

$$\frac{\rho_\Lambda}{\bar{M}_{\text{Pl}}^4} = \frac{3}{4\pi^2} e^{2\delta_{\text{ph}}} \left[\frac{v_{\text{geo}}}{5\beta_{\text{rad}}^2 \bar{M}_{\text{Pl}}} \right]^{10} \quad (29)$$

where the exponent $10 = 2g_{\text{car}}$ and $5 = g_{\text{car}}$. Verified to a relative residual $\sim 10^{-159}$.

Proof. From $v_{\text{geo}}/\bar{M}_{\text{Pl}} = g_{\text{car}}\beta_{\text{rad}}^2 e^{-A_{\text{EW}}}$, $e^{-A_{\text{EW}}} = (v_{\text{geo}}/\bar{M}_{\text{Pl}})/(g_{\text{car}}\beta_{\text{rad}}^2)$. From $\rho_\Lambda/\bar{M}_{\text{Pl}}^4 = \frac{3}{4\pi^2} e^{-2\alpha_\star^{-1}}$ and $\alpha_\star^{-1} = g_{\text{car}} A_{\text{EW}} - \delta_{\text{ph}}$, $e^{-2\alpha_\star^{-1}} = e^{2\delta_{\text{ph}}} e^{-2g_{\text{car}} A_{\text{EW}}} = e^{2\delta_{\text{ph}}} (e^{-A_{\text{EW}}})^{2g_{\text{car}}}$. Substituting and using $2g_{\text{car}} = 10$, $g_{\text{car}} = 5$ gives the boxed form. $\square$

In words: the electroweak scale is not directly the Λ scale, but once the electroweak ratio is stripped of its small β_{rad} prefactor, Λ is its tenth power, and the “ten” is $2g_{\text{car}}$, not a fitted exponent. This is the direct algebraic compression of the action ladder.

6.3 The full power tower: x^1, x^5, x^{10} on the carrier graph K_5

The decuple bridge is one rung of a tower. Writing $x := e^{-A_{\text{EW}}} = v_{\text{geo}}/(5\beta_{\text{rad}}^2 \bar{M}_{\text{Pl}})$ for the stripped electroweak unit, all three cosmological scales are powers of the *same* x with binomial exponents (verified exactly):

$$\frac{v_{\text{geo}}}{5\beta_{\text{rad}}^2 \bar{M}_{\text{Pl}}} = x^{\binom{5}{0}} = x, \quad \frac{H_0}{\bar{M}_{\text{Pl}}} = x^{\binom{5}{1}} R_{\text{H}} = x^5 R_{\text{H}}, \quad \frac{\rho_\Lambda}{\bar{M}_{\text{Pl}}^4} = x^{\binom{5}{2}} R_{\Lambda} = x^{10} R_{\Lambda} \quad (30)$$

with residues $R_{\text{H}} = e^{\delta_{\text{ph}}}/(2\pi\sqrt{\Omega_\Lambda})$ and $R_{\Lambda} = \frac{3}{4\pi^2} e^{2\delta_{\text{ph}}}$. [E]. The exponents are the complete-graph counts of the five-slot carrier K_5 :

$$H \leftrightarrow \text{vertices of } K_5 \text{ (}\# = \binom{5}{1} = 5\text{)}, \quad \Lambda \leftrightarrow \text{edges of } K_5 \text{ (}\# = \binom{5}{2} = 10\text{)}.$$

So the cosmological constant is the product of one stripped-electroweak unit over every carrier *pair* (K_5 edge product), and the Hubble scale the product over every carrier *slot* (K_5 vertex product). This is the cleanest single-variable statement of the action ladder.

6.4 Hubble is the square root of Λ (corollary, exact)

On the comparison layer $\rho_\Lambda/\bar{M}_{\text{Pl}}^4 = 3\Omega_\Lambda(H_0/\bar{M}_{\text{Pl}})^2$; equating to the TFPT line gives

$$\frac{H_0}{\bar{M}_{\text{Pl}}} = \frac{e^{-\alpha_\star^{-1}}}{2\pi\sqrt{\Omega_\Lambda}}, \quad \mathcal{A}_H = -\ln \frac{H_0}{\bar{M}_{\text{Pl}}} = \alpha_\star^{-1} + \ln(2\pi\sqrt{\Omega_\Lambda}) = 138.68. \quad (31)$$

So the Hubble rung is *not* an independent third hit: it is the square root of the vacuum energy plus a geometric cosmology factor. The genuine TFPT content is the single $e^{-\alpha_\star^{-1}}$. (The numerical coincidence $\mathcal{A}_H - \alpha_\star^{-1} \approx \pi^2/6$ is just $\ln(2\pi\sqrt{\Omega_\Lambda})$ and is Ω_Λ -dependent; not a claim. [O])

6.5 The ~ 123 orders of the cosmological constant

$$\log_{10} \frac{M_{\text{Pl}}^4}{\Lambda_{\text{IR}}} = \underbrace{\frac{2\alpha_\star^{-1}}{\ln 10}}_{119.028} + \underbrace{\log_{10} \frac{1}{\delta_{\text{top}}}}_{3.920} = 122.948 \approx 123. \quad (32)$$

The “119” is the action $2\alpha_\star^{-1}$; the “ ≈ 4 ” is the seam defect $\delta_{\text{top}} = \Omega_{\text{adm}} c_3^4$.

High-precision cosmological constant: a $\sim 10^{-13}$ prediction [E]

Because $\alpha_\star^{-1}$ is a *derived* number (the EM fixed point, known to 13 digits) and the residue $3/(4\pi^2)$ is exact, the dimensionless vacuum energy is predicted essentially exactly:

$$\frac{\rho_\Lambda}{\bar{M}_{\text{Pl}}^4} = \frac{3}{4\pi^2} e^{-2\alpha_\star^{-1}} = 7.12533 \times 10^{-121} \quad [\text{E}] \text{ [verification/v7_gravity_cosmo.py]}, \quad (33)$$

i.e. a dark-energy scale $\rho_\Lambda^{1/4} = \left(\frac{3}{4\pi^2}\right)^{1/4} \bar{M}_{\text{Pl}} e^{-\alpha_\star^{-1}/2} = 2.23747 \text{ meV}$ (reduced Planck mass $\bar{M}_{\text{Pl}} = 2.435323 \times 10^{18} \text{ GeV}$). The *dimensionless* ratio carries no observational input — its uncertainty is that of $\alpha_\star^{-1}$, $\sim 10^{-13}$ relative; the absolute meV value inherits only the Planck-mass uncertainty ($\sim 2 \times 10^{-5}$). Both are *far* sharper than the measured vacuum energy (Planck $\rho_\Lambda^{1/4} \simeq 2.24 \text{ meV}$, known to $\sim 1\%$ and entangled with the H_0 tension). So TFPT pins the cosmological constant several orders of magnitude more precisely than it is currently measured — a genuine, falsifiable forward prediction: any future sub-percent ρ_Λ must land on 2.2375 meV.

6.6 The ladder is binomial; $2g_{\text{car}} = \binom{g_{\text{car}}}{2}$ singles out $g_{\text{car}} = 5$

In units of the electroweak action $\alpha_\star^{-1}/g_{\text{car}}$, the three rungs are the first three binomial coefficients of the five-slot carrier:

$$\frac{A_{\text{EW}}}{\alpha_\star^{-1}/g_{\text{car}}} : \frac{\mathcal{A}_H}{\alpha_\star^{-1}/g_{\text{car}}} : \frac{\mathcal{A}_\Lambda}{\alpha_\star^{-1}/g_{\text{car}}} = 1 : g_{\text{car}} : 2g_{\text{car}} = \binom{g_{\text{car}}}{0} : \binom{g_{\text{car}}}{1} : \binom{g_{\text{car}}}{2} = 1 : 5 : 10. \quad (34)$$

The first two are automatic ($\binom{g_{\text{car}}}{0} = 1$, $\binom{g_{\text{car}}}{1} = g_{\text{car}}$). The non-trivial content is the cosmological-constant charge:

$$2g_{\text{car}} = \binom{g_{\text{car}}}{2} \iff g_{\text{car}} = 5 \quad [\text{E}] \text{ [verification/v2_carrier_pascal.py]}, \quad (35)$$

since $\binom{g_{\text{car}}}{2} = \frac{g_{\text{car}}(g_{\text{car}}-1)}{2} = 2g_{\text{car}} \iff g_{\text{car}} = 5$. So the “double overlap” charge $2g_{\text{car}}$ equals the number of carrier-slot *pairs* $\binom{g_{\text{car}}}{2}$ *only* at $g_{\text{car}} = 5$: the cosmological constant is suppressed by one action unit per carrier pair, and this combinatorial reading is exclusive to the physical rank. It is an independent overdetermination of $g_{\text{car}} = 5$ from the action grammar (cf. the landscape section).

6.7 Discriminant action theorem: the ladder reads the hypercharge discriminant

The g_{car} in the action ladder is not merely a rank: it is the square root of the discriminant of the carrier (hypercharge) polynomial. For a general split $b + s = g$ the carrier polynomial is $bsY^2 + (s-b)Y - 1 = 0$ with discriminant

$$\Delta_Y = (s-b)^2 + 4bs = (b+s)^2 = g_{\text{car}}^2, \quad \text{so for } 6Y^2 - Y - 1 = 0 : \Delta_Y = 25, \sqrt{\Delta_Y} = 5 = g_{\text{car}}. \quad (36)$$

Hence the action ladder is, exactly,

$$A_{\text{EW}} = \frac{\alpha_\star^{-1}}{\sqrt{\Delta_Y}}, \quad \mathcal{A}_H = \sqrt{\Delta_Y} A_{\text{EW}}, \quad \mathcal{A}_\Lambda = \binom{\sqrt{\Delta_Y}}{2} A_{\text{EW}} \quad [\text{E}]. \quad (37)$$

The electroweak hierarchy therefore reads the *discriminant of the hypercharge polynomial*, tying the Standard-Model hypercharge derivation directly to the EW-hierarchy and Λ problems.

6.8 A two-dimensional action lattice

All scales sit on the integer lattice spanned by two generators — the gauge unit $\alpha_\star^{-1}/g_{\text{car}}$ and the flavor unit $L_0 = \ln(1/u)$ — as $S_x = a_x(\alpha_\star^{-1}/g_{\text{car}}) + b_x L_0 + (\text{residue})$:

scale	a_x	b_x	axis
$v_{\text{geo}}/\bar{M}_{\text{Pl}}$	$1 = \binom{5}{0}$	0	gauge
$H_0/\bar{M}_{\text{Pl}}$	$5 = \binom{5}{1}$	0	gauge
$\rho_\Lambda/\bar{M}_{\text{Pl}}^4$	$10 = \binom{5}{2}$	0	gauge
$m_e/\bar{M}_{\text{Pl}}$	1	5	flavor
$m_\mu/\bar{M}_{\text{Pl}}$	1	3	flavor
$m_\tau/\bar{M}_{\text{Pl}}$	1	2	flavor

The cosmological/gauge sector runs along $b = 0$ with binomial charges; the charged-fermion sector runs along $a = 1$ with the integer flavor ladder $n_f = (5, 3, 2)$. This is the compact “periodic table” of TFPT scales: two integer labels per scale. **[E]** (charges) / **[C]** (residues).

A third (seam/gravity) generator completes the lattice: $L_c = \ln(1/c_3) = \ln(8\pi)$. The scalaron lives purely on it, $M_{\text{scal}}/\bar{M}_{\text{Pl}} = c_3^{7/2} = e^{-\frac{7}{2}L_c}$, and the seam defect is $\delta_{\text{top}} = e^{-(4L_c - \ln 48)}$. So the three log-generators are $\{\alpha_\star^{-1}$ (hierarchy), $L_0 = \ln(1/u)$ (flavor), $L_c = \ln(1/c_3)$ (seam/gravity) $\}$. **[E]**

7 The Einstein normaliser as a corrected tree value

Machine checks for this section: `[verification/v1_e8_glue.py]`

The gravitational normaliser $\xi = c_3/u$ has a transparent exact form. Writing $u = \frac{1}{6\pi} [1 + \frac{9}{128\pi^3}]$,

$$\xi = \frac{c_3}{u} = \frac{3}{4} \cdot \frac{1}{1 + \frac{9}{128\pi^3}} = 0.748303 \quad \textbf{[E]}. \quad (38)$$

So $\xi_{\text{tree}} = \frac{3}{4} = \dim \mathfrak{g}_{\text{SM}} / \dim S^+ = 12/16$ is the exact leading value (the inverse of the $\frac{4}{3} = \dim S^+ / \dim \mathfrak{g}_{\text{SM}}$ family/gauge compression that sits in the seed $u_{\text{base}} = \frac{4}{3}c_3$, so $\xi_{\text{tree}} u_{\text{base}} = c_3$), and the deviation is precisely the small topological seed correction $9/(128\pi^3)$, i.e. 0.226%. Gravity does not pick up a mysterious factor: first comes $3/4$, then the seam defect shifts it minimally.

8 Λ -anchored metrology closure, and two anchors

Machine checks for this section: `[verification/v60_lambda_metrology_branch.py]`

The absolute SI scale is a calibration, not a compiler output. The dimensionless vacuum quotient lets one anchor it to the observed geometric curvature without circularity.

Theorem 1 (Λ -anchored metrology closure). With $\lambda_\Lambda := \rho_\Lambda/\bar{M}_{\text{Pl}}^4 = \frac{3}{4\pi^2} e^{-2\alpha_\star^{-1}}$ and the observed geometric curvature $\Lambda_{\text{geom}}^{\text{obs}} > 0$,

$$\bar{M}_{\text{Pl}}^2 = \frac{\Lambda_{\text{geom}}^{\text{obs}}}{\lambda_\Lambda}, \quad G_N^{\text{TFPT}} = \frac{c^3}{8\pi\hbar} \frac{\lambda_\Lambda}{\Lambda_{\text{geom}}^{\text{obs}}} = \frac{3c^3}{32\pi^3\hbar \Lambda_{\text{geom}}^{\text{obs}}} e^{-2\alpha_\star^{-1}}.$$

Proof. $\bar{M}_{\text{Pl}}^2 = 1/(8\pi G_N)$ and $\Lambda_{\text{geom}} = \rho_\Lambda/\bar{M}_{\text{Pl}}^2$; the branch sets $\rho_\Lambda = \lambda_\Lambda \bar{M}_{\text{Pl}}^4$, so $\Lambda_{\text{geom}} = \lambda_\Lambda \bar{M}_{\text{Pl}}^2$, giving $\bar{M}_{\text{Pl}}^2$ and then $G_N = \lambda_\Lambda/(8\pi\Lambda_{\text{geom}}^{\text{obs}})$; restoring $c^3/\hbar$ closes it. $\square$

Non-circularity ([O], essential). The anchor must be the curvature $\Lambda_{\text{geom}}^{\text{obs}}$ (units m^{-2}), *not* the SI vacuum energy density $\rho_\Lambda^{\text{obs}} = \Lambda_{\text{geom}} c^4/(8\pi G_N)$, which already contains G_N . The curvature follows from $\Lambda_{\text{geom}}^{\text{obs}} = 3\Omega_\Lambda H_0^2/c^2$.

Two independent anchors (crosscheck with teeth). The metrology can be pinned in two ways that should agree:

1. *Electroweak anchor*: the dimensionless $G_N v_{\text{phys}}^2 = 4.06717 \times 10^{-34}$ fixes G_N once v_{phys} is calibrated non-gravitationally.
2. *Λ -curvature anchor* (this note): $G_N = \frac{3c^3}{32\pi^3 \hbar \Lambda_{\text{geom}}^{\text{obs}}} e^{-2\alpha_\star^{-1}}$.

With the example anchor $H_0 = 67.36 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_\Lambda = 0.6847$ ($\Lambda_{\text{geom}}^{\text{obs}} = 1.08914 \times 10^{-52} \text{ m}^{-2}$) the Λ anchor gives

$$G_N^{\text{TFPT}} = 6.65071 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}, \quad G_N^{\text{TFPT}} / G_N^{\text{CODATA}} - 1 = -3.5 \times 10^{-3}. \quad (39)$$

Closure algebra [E]; SI anchor and numerical match [C] (the cosmological anchor is model dependent; the Hubble tension applies). EW crosscheck: the Λ -pinned $\bar{M}_{\text{Pl}} = 2.43964 \times 10^{18} \text{ GeV}$ gives $v_{\text{geo}} = 242.6 \text{ GeV}$, needing $Z_{\text{EW}} = (246.22/242.61)^2 = 1.030$.

Prefactor branch ([O]). This note uses $\Lambda_{\text{IR}}/M_{\text{Pl}}^4 = \delta_{\text{top}} e^{-2\alpha_\star^{-1}}$. A competing prefactor $2c_3 e^{-2\alpha_\star^{-1}}$ would differ by $2c_3/\delta_{\text{top}} = 661.5$ and misscale G_N by the same factor; the branch must be versioned.

Mandatory Λ -layer typing ([O]). Three distinct Λ objects must never be conflated:

$$\underbrace{\frac{\rho_\Lambda}{M_{\text{Pl}}^4} = \frac{3}{4\pi^2} e^{-2\alpha_\star^{-1}}}_{\text{vacuum density quotient (dimensionless)}} \neq \underbrace{\frac{\Lambda_{\text{IR}}}{M_{\text{Pl}}^4} = \delta_{\text{top}} e^{-2\alpha_\star^{-1}}}_{\text{seam-determinant / IR scale (dimensionless)}} \neq \underbrace{\Lambda_{\text{geom}}^{\text{obs}} = \frac{3\Omega_\Lambda H_0^2}{c^2}}_{\text{observed curvature [m}^{-2}\text{]}}.$$

Only $\Lambda_{\text{geom}}^{\text{obs}}$ carries a dimension and is the non-circular anchor. Keeping the three typed protects the decuple bridge and the metrology closure from a spurious prefactor objection.

Measurement-near (physical) decuple bridge. Since $v_{\text{phys}} = v_{\text{geo}} \sqrt{Z_{\text{EW}}}$ (charged-current residue, not a fit), the decuple bridge in the observable vev reads

$$\frac{\rho_\Lambda}{M_{\text{Pl}}^4} = \frac{3}{4\pi^2} e^{2\delta_{\text{ph}}} \left[\frac{v_{\text{phys}}}{5\beta_{\text{rad}}^2 \sqrt{Z_{\text{EW}}} \bar{M}_{\text{Pl}}} \right]^{10}. \quad (40)$$

This is the same identity, written in the low-energy charged-current scale rather than the geometric source scale; both forms should appear side by side. [E]/[C]

9 The landscape: $g_{\text{car}} = 5$ is overdetermined

Machine checks for this section: [verification/v6_bootstrap.py] [verification/v14_carrier_uniqueness.py]

Varying the carrier rank and rebuilding every constant from the same machinery (c_3 fixed), the EM-closure value $\alpha^{-1}(g_{\text{car}})$ is monotone and meets the observed value only at $g_{\text{car}} = 5$:

g_{car}	N_{fam}	$\dim S^+$	Ω_{adm}	$10b_1$	$\alpha^{-1}(g_{\text{car}})$	verdict
3	2	4	8	7	258.15	wrong α , wrong family count
4	2.5	8	20	17	187.31	non-integer families
5	3	16	48	41	137.036	SM, 3 families, observed α
6	3.5	32	112	97	101.30	non-integer families
7	4	64	256	225	75.61	wrong α

Several distinct requirements converge on $g_{\text{car}} = 5$: integer families (g_{car} odd); three families $(g_{\text{car}}+1)/2 = 3$; the observed α^{-1} on the monotone curve; the SM structure $(b, s) = (3, 2)$, $\dim S^+ = 16$; and $\Omega_{\text{adm}} = 48$, $10b_1 = 41$. A clean algebraic shadow: $\frac{g(g+1)}{2} = 2^{g-1} - 1$ has $g = 5$ as its non-trivial solution ([C]).

10 The E_8 carrier atlas — stated unambiguously

Machine checks for this section: [\[verification/v5_e8_cascade.py\]](#)

In the series E_8 is a *downstream scale grammar*, not a primitive cause: the boundary kernel fixes α , the carrier rank, families and admissibility without E_8 upstream. Three levels.

10.1 Level 1 — original E_8 data of the series ([E])

$$|R(E_8)| = 248 - 8 = 240 = 5 \cdot 48 = 5 \Omega_{\text{adm}} = \text{lcm}(\Omega_{\text{adm}}, D_{\text{start}}) = \text{lcm}(48, 60), \quad (41)$$

$$D_{\text{start}} = g_{\text{car}} \cdot \dim \mathfrak{g}_{\text{SM}} = 5 \cdot 12 = 60, \quad \gcd(\Omega_{\text{adm}}, D_{\text{start}}) = 12 = \dim \mathfrak{g}_{\text{SM}}, \quad (42)$$

$$\frac{|R(E_8)|}{D_{\text{start}}} = \frac{240}{60} = 4 = \frac{\Omega_{\text{adm}}}{\dim \mathfrak{g}_{\text{SM}}}, \quad 2880 = D_{\text{start}} \Omega_{\text{adm}} = |R(E_8)| \cdot \dim \mathfrak{g}_{\text{SM}}, \quad (43)$$

$$\kappa_{E_8} = \frac{\gamma_{\text{car}}}{\ln(248/60)} = 0.58723, \quad X(n, r, I_1) = \frac{\bar{M}_{\text{Pl}}}{8} \sin^2 \theta_{13} \left(\frac{60 - 2n}{60} \right)^{\kappa_{E_8}} e^{-(8-r)/64} e^{-12\pi I_1}. \quad (44)$$

Coxeter/gauge typing of the lcm–gcd pair. Writing $D_{\text{start}} = 60 = 2h_{E_8}$ and $\Omega_{\text{adm}} = 48 = 6 \text{rank}(E_8)$, the relations become $\gcd(2h_{E_8}, 6r_{E_8}) = \dim \mathfrak{g}_{\text{SM}} = 12$ and $\text{lcm}(2h_{E_8}, 6r_{E_8}) = |R(E_8)| = 240$: the local SM algebra is the gcd of the Coxeter double-period and the $\mathbb{Z}_6$ rank-occupancy, and the E_8 root count is their lcm. Likewise the EW exponent reads $I_1^{\text{EW}} = \frac{41}{48} = \frac{h_{E_8}}{36} + \frac{1}{6r_{E_8}} = \frac{5}{6} + \frac{1}{48}$ (Coxeter principal part plus a rank singlet). [E] (arithmetic) / [C] (typing).

10.2 Level 2 — new exact counting identities ([E])

$$|R(E_8)| = 240 = 16 \cdot 15 = \dim S^+ (\dim S^+ - 1), \quad (45)$$

$$\dim E_8 = 248 = 8 \cdot 31 = (g_{\text{car}} + N_{\text{fam}})(2^{g_{\text{car}}} - 1), \quad (46)$$

$$|R(E_8)| = 8(31 - 1) = \text{rank}(E_8)(2^{g_{\text{car}}} - 2), \quad \text{rank}(E_8) = 8 = g_{\text{car}} + N_{\text{fam}} = 5 + 3, \quad (47)$$

$$10b_1 = 41 = \frac{|R(E_8)|}{6} + 1, \quad I_1^{\text{EW}} = \frac{41}{48} = \gamma_{\text{car}} + \frac{1}{\Omega_{\text{adm}}}, \quad (48)$$

with $31 = 2^{g_{\text{car}}} - 1$ the number of non-empty occupation words of the $[5, 4, 2]$ carrier code.

10.3 Level 2.5 — the even-flip transition atlas (a falsifiable program)

Identify S^+ with the even-weight binary code on five bits. A transition $x \rightarrow y \neq x$ has flip $F = x \oplus y$, even and nonempty, so $|F| \in \{2, 4\}$; per state there are $\binom{5}{2} + \binom{5}{4} = 10 + 5 = 15$ flips, giving

$$240 = 16 \cdot (10 + 5) = \underbrace{160}_{|F|=2} + \underbrace{80}_{|F|=4}, \quad (49)$$

and the 120 undirected flips match the 120 E_8 root *pairs* ([E], counting). This makes the atlas conjecture *testable*: a genuine root map must send each flip to a norm-2 vector in $\mathbb{R}^8$ with inner products in $\{-2, -1, 0, 1, 2\}$ reproducing the E_8 profile (per root: 1 self, 56 at +1, 126 orthogonal, 56 at -1, 1 anti).

Honest obstruction ([O]). The flip atlas splits $160 + 80$ by flip size; the standard E_8 (D_8) split is $112 + 128$. Since $160 + 80 \neq 112 + 128$, no flip-size-preserving map is an E_8 root system; a metrization, if it exists, must mix flip sizes. Until a norm-2, E_8 -profile embedding is exhibited, E_8 stays a *counting* atlas, not a derived root origin.

Sharper two-stage audit (Coxeter before Weyl). A full root-system isomorphism is hard; a much earlier, cheaper test is the Coxeter spectrum. *Stage A*: find a natural order-30 operator C_{TFPT} on the even-flip atlas (or its 8-dimensional primitive-character space) with characteristic

polynomial $\chi_{C_{\text{TFPT}}}(t) = \Phi_{30}(t)$. *Stage B (only if A holds):* the norm-2, E_8 -profile, Weyl-closure metrization. Stage A can fail or carry far sooner than B, so it is the right first falsifier. [O]

Audit only — non-load-bearing speculation: the dim 496 heterotic count [O]. *This box feeds no derivation; it is an arithmetic coincidence kept for completeness only, under the no-free-pattern rule.* The even half S^+ and odd half S^- of the exterior algebra each have dimension $2^{g_{\text{car}}-1} = 16$ and each carry $16 \cdot 15 = 240$ transitions. Together the full carrier ($2^{g_{\text{car}}} = 32$ states) gives

$$\dim S^+(2^{g_{\text{car}}} - 1) = 16 \cdot 31 = 496 = \dim(E_8 \times E_8) = \dim SO(32) = \binom{32}{2} \quad (\text{count}),$$

i.e. the two halves read as two E_8 root atlases, all $\binom{32}{2} = 496$ pairs as $SO(32)$ — the two anomaly-free heterotic gauge dimensions. The *dimension* equality is exact; that the carrier algebra *is* either gauge group is unproven and not used anywhere. [O] (audit-only).

10.4 Level 2.7 — the carrier Coxeter theorem (the strongest new bridge)

Use the universal root-system identity $|R| = \text{rank} \cdot h$ (Coxeter number h). With the carrier values $\text{rank}(E_8) = g_{\text{car}} + N_{\text{fam}} = 8$ and $|R(E_8)| = \dim S^+(\dim S^+ - 1) = 240$,

$$h_{E_8} = \frac{|R(E_8)|}{\text{rank}(E_8)} = \frac{240}{8} = 30 = N_{\text{fam}} \binom{g_{\text{car}}}{2} = N_{\text{fam}} A_{\Lambda} = 2g_{\text{car}} N_{\text{fam}} = 2^{g_{\text{car}}} - 2 \quad [\text{E}]. \quad (50)$$

The Coxeter number of E_8 is exactly three families times the cosmological pair action: each family contributes one Λ -decuple ($A_{\Lambda} = \binom{5}{2} = 10$) to the Coxeter period. Equivalently the action ladder is a Coxeter ladder,

$$A_H = \frac{h_{E_8}}{2N_{\text{fam}}} = 5, \quad A_{\Lambda} = \frac{h_{E_8}}{N_{\text{fam}}} = 10, \quad (51)$$

so Hubble reads half a Coxeter period per family and Λ a full Coxeter period per family. [C]

Rank–Coxeter coupling. Equivalently $\dim S^+ = 2 \text{rank}(E_8) = 16$ and $h_{E_8} = 2(\dim S^+ - 1) = 2 \cdot 15 = 30$, so $\dim E_8 = (g_{\text{car}} + N_{\text{fam}})(h + 1) = 8 \cdot 31$. The two counting forms $16 \cdot 15 = 8 \cdot 30 = 240$ coincide because $\dim S^+ = 2r$. Dropping ν^c sends $\dim S^+ \rightarrow 15$, which breaks *both* $\dim S^+ = 2r$ and $\dim S^+(\dim S^+ - 1) = 240$: ν^c is the state that closes the rank–Coxeter coupling, not an optional singlet. [E]/[C]

10.5 Level 2.8 — positive-root moment: $|R^+(E_8)| = \text{Tr}_{S^+} X^2 = 5!$

The number of positive E_8 roots equals the master hypercharge moment:

$$|R^+(E_8)| = 120 = \text{Tr}_{S^+} X^2 = 5! \quad [\text{E}], \quad (52)$$

so the second hypercharge moment of one family *is* the positive-root half-system. The whole integer chain then reads off $|R^+|$:

$$\sum_{f,j} L_{f,j} = \frac{|R^+(E_8)|}{3} = 40, \quad 10b_1 = 1 + \frac{|R^+(E_8)|}{3} = 41, \quad \delta_2 = 4! |R^+(E_8)| c_3^8. \quad (53)$$

The $\mathbb{Z}_6$ slice is then $|R(E_8)|/6 = \frac{|R^+|}{3} = 40 = \Omega_{\text{adm}}\gamma$: one $\mathbb{Z}_6$ phase slice of the E_8 root atlas is exactly the full flavor transport budget. [E] (arithmetic) / [C] (interpretation).

10.6 Level 2.9 — the 2·3·5 Coxeter cyclotomic compiler (meta-theorem)

The three hard carrier atoms — weak rank 2, color rank / family count 3, carrier rank $g_{\text{car}} = 5$ — generate the E_8 data *cyclotomically*, without taking E_8 as input. Set

$$h := 2 \cdot 3 \cdot g_{\text{car}} = 30, \quad r := \varphi(h) = \varphi(30) = 8 = g_{\text{car}} + N_{\text{fam}}, \quad (54)$$

where h is the Coxeter number and $r = \varphi(2 \cdot 3 \cdot 5) = 8$ the rank. Then the standard $|R| = rh$, $\dim = r(h+1)$ give

$$|R(E_8)| = rh = 8 \cdot 30 = 240, \quad \dim E_8 = r(h+1) = 8 \cdot 31 = 248, \quad h+1 = 2^{g_{\text{car}}} - 1 = 31. \quad (55)$$

Moreover the E_8 exponents are exactly the primitive residues modulo $h = 30$:

$$\text{Exp}(E_8) = (\mathbb{Z}/30\mathbb{Z})^\times = \{1, 7, 11, 13, 17, 19, 23, 29\}, \quad \sum_{m \in (\mathbb{Z}/30\mathbb{Z})^\times} m = 120 = \text{Tr}_{S^+} X^2 = |R^+(E_8)|, \quad (56)$$

so the second hypercharge moment equals the sum of the E_8 exponents and the positive-root count. The $\mathbb{Z}_6 \times \mathbb{Z}_5 \simeq \mathbb{Z}_{30}$ tact has $\varphi(30) = 8$ primitive characters, whose phases $e^{2\pi i m/30}$ are the Coxeter eigenphases of E_8 ; its characteristic polynomial is the 30th cyclotomic polynomial

$$\chi_C(t) = \Phi_{30}(t) = t^8 + t^7 - t^5 - t^4 - t^3 + t + 1. \quad (57)$$

Reading: E_8 here is not primarily a root count but the *Coxeter cyclotomic readout of the 2, 3, 5 carrier*. [E] (arithmetic) / [C] (interpretation).

10.7 Level 3 — conjectural geometric readings ([C])

Conjecture 1 (Carrier transition atlas). $R(E_8) \cong \{(a, b) \in S^+ \times S^+ : a \neq b\}$ as a structure-bearing transition set, since $16 \cdot 15 = 240$. *Open:* a structure-preserving map (roots, lengths, Weyl data), not only the count.

Conjecture 2 (Least common refinement / rank-8 over the exhausted carrier). $240 = \text{lcm}(\Omega_{\text{adm}}, D_{\text{start}})$ reads E_8 as the least common refinement of the occupancy lattice and the $\mathbb{Z}_6$ phase lattice; $248 = (g_{\text{car}} + N_{\text{fam}})(2^{g_{\text{car}}} - 1)$ reads it as a rank-8 structure over the 31 non-empty carrier words.

Two routes to E_8 — keep their status separate. Levels 1–2 are exact arithmetic. There are now *two* distinct routes to E_8 , and they must not be conflated:

- **Glue route (closed, [E]):** $E_8 = (D_5 \oplus A_3) + \mu_4\text{-glue}$ is built explicitly in Section 3 ($\text{disc}(D_5) = \text{disc}(A_3) = \mathbb{Z}_4$, glue index 4, 240 roots of norm 2, simple-root Gram $\det 1$). This closes the former “blind 240-metrisation”.
- **Direct flip-atlas route (open, [O], optional bonus):** the even-flip transition set $240 = 160 + 80$ is *not* a flip-size-preserving E_8 map, since $160 + 80 \neq 112 + 128$. A direct structure-preserving metrisation of the flip graph remains open — a nice-to-have, not load-bearing, because the glue route already delivers E_8 .

So E_8 is closed via the glue; only the direct flip atlas is open.

11 $\mathbb{Z}_6$ monodromy, flavor and CP readouts

Machine checks for this section: [verification/v88_cp_phase_audit.py]

A single $\mathbb{Z}_6$ motif recurs in the quotient $G_{\text{phys}} = (SU(3) \times SU(2) \times U(1)_Y) / \mathbb{Z}_6$, the transport cusps $\{1, (2/3)^6, (1/3)^6\}$, $\delta_{\text{ph}} = z_\star^{1/6}$, and the CP phases

$$\delta_{\text{CKM}}^{(0)} = \frac{\pi}{3}, \quad \delta_{\text{CKM}} = \frac{\pi}{3} + 3\lambda_C^2 = 1.19823 \text{ rad}, \quad \delta_{\text{CP}}^{\text{PMNS}} = \frac{4\pi}{3} = 240^\circ. \quad (58)$$

From the documented inputs $s_{12} = \lambda_C$, $s_{23} = u/(1 + \lambda_C)$, $s_{13} = \lambda_C^3/3$ one gets the new derived readouts $|V_{td}| = 0.00908$, $|V_{ts}| = 0.04264$, $J_{\text{CKM}} = 3.33 \times 10^{-5}$. On the neutrino side, with $\sin^2 \theta_{13} = e^{-5/6}u = 0.02311$, $\sin^2 \theta_{23} = 0.4557$, $\delta_{\text{CP}} = 4\pi/3$ and the solar candidate $\sin^2 \theta_{12} = \frac{1}{3} - \frac{u}{2} = 0.30675$, a near-maximal $|J'_{\text{CP}}| \approx 0.030$ (sharp readout for DUNE/Hyper-Kamiokande). [\[C\]](#)

11.1 CP as triple seed variance; a closed observable quadrilateral

The Cabibbo angle is the Bernoulli width of the seed. The *exact* CKM phase is the family transport holonomy (document 3), $\delta_{\text{CKM}} = \arg \text{Tr } \Pi_{\Delta}(\nabla_F^*)$; on the small-area branch this has the compact readout form (with an $O(\lambda_C^4)$ remainder)

$$\lambda_C = \sqrt{u(1-u)}, \quad \boxed{\delta_{\text{CKM}} - \frac{\pi}{3} = N_{\text{fam}} u(1-u) = 3\lambda_C^2} \quad [\text{E}]/[\text{C}] \text{ (readout; exact = holonomy).} \quad (59)$$

The coefficient 3 is the family number N_{fam} : weak CP is the $\mathbb{Z}_6$ base angle $\pi/3$ plus N_{fam} times the seed (Bernoulli) variance. The $\mathbb{Z}_6$ base angle now has an A_3 root: since $\mathbb{Z}_6 = |R^+(A_3)| = 6$ (the positive-root hexagon of the family algebra), the CP phases are integer steps of the A_3 holonomy quantum $2\pi/|R^+(A_3)|$,

$$\boxed{\delta_{\text{CKM}} = \frac{2\pi}{|R^+(A_3)|} + N_{\text{fam}} u(1-u), \quad \delta_{\text{CP}}^{\text{PMNS}} = 4 \cdot \frac{2\pi}{|R^+(A_3)|} = \frac{4\pi}{3}} \quad [\text{E}]/[\text{C}]. \quad (60)$$

So weak CP = (A_3 positive-root phase step) + (family count $\times$ seed variance), while strong CP is the *determinant* sector, erased. The solar angle reads as a carrier root plus a half-seed, $\sin^2 \theta_{12} = \frac{1}{3} - \frac{u}{2} = -y_- - \frac{u}{2}$ (since $-y_- = \frac{1}{3}$), so PMNS = carrier root + seed + A_3 holonomy phase. [\[C\]](#) Strong CP is killed by the determinant selector ($\arg \det M_u = \arg \det M_d = 0$, $\bar{\theta} = 0$), while weak CP survives as the seed-deformed $\mathbb{Z}_6$ holonomy: CP is not forbidden, only *determinant* CP is. [\[C\]](#) — the strong-CP closure lives on the admissible QFT branch and is *conditional* on the analytic hypotheses of the QFT closure (document 3, Part III: reflection positivity, temperedness, clustering, mass gap, OS reconstruction); its status is therefore weaker than the discrete lattice identities, even though the semantics “determinant CP forbidden, holonomy CP allowed” is robust. Since $u = \Omega_b + \beta_{\text{rad}}$ exactly, eliminating the seed closes the four UV-shadow observables into one cycle:

$$\lambda_C^2 = (\Omega_b + \beta_{\text{rad}})(1 - \Omega_b - \beta_{\text{rad}}), \quad \sin^2 \theta_{13} = e^{-5/6}(\Omega_b + \beta_{\text{rad}}), \quad \Omega_b = \left(1 - \frac{1}{4\pi}\right)e^{5/6} \sin^2 \theta_{13}. \quad (61)$$

One measured member of $\{\lambda_C, \theta_{13}, \Omega_b, \beta_{\text{rad}}\}$ fixes the other three. [\[E\]](#) (the four rows remain operationally distinct readout classes).

12 Mass hierarchy as a hexagon with one extra turn

Machine checks for this section: [\[verification/v18_quark_yukawa.py\]](#) [\[verification/v20_lepton_c_derivation.py\]](#)

The transport length packets are

$$L_u = (7, 3, 0), \quad L_d = (7, 5, 2), \quad L_e = (8, 5, 3), \quad \sum_{f,j} L_{f,j} = 40 = \Omega_{\text{adm}} \gamma_{\text{car}} = 10b_1 - 1. \quad (62)$$

Modulo 6 they lie on related triples of the C_6 hexagon; the first generation carries one extra full turn ($u, d : 1 + 6; e : 2 + 6$), costing $\lambda_Y^6 = (u(1-u))^3 \approx 1.28 \times 10^{-4}$ and explaining why the first generation is so light — “one turn more”, not a free fit. [\[C\]](#) *Flagged proximity ([O], not exact)*: the full-turn cost is close to the topological seam defect, $\lambda_C^6/\delta_{\text{top}} = 1.061$ ($\sim 6\%$), hinting that one

flavor winding $\approx$ one seam-defect unit $\delta_{\text{top}} = \Omega_{\text{adm}} c_3^4$ — a structural smell, not a theorem. Writing the length matrix $L = R + W$ with $R = L \bmod 6$ the residue shadow and W the full-winding part,

$$L = \begin{pmatrix} 7 & 3 & 0 \\ 7 & 5 & 2 \\ 8 & 5 & 3 \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 3 & 0 \\ 1 & 5 & 2 \\ 2 & 5 & 3 \end{pmatrix}}_R + 6 \underbrace{\begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}}_W, \quad \sum R = 22, \sum W = 18 = 3 \cdot 6, \sum L = 40,$$

and the column sums are $(22, 13, 5)$ with $S_1 = 22 = \sum R$ and $S_2 + S_3 = 18 = \sum W$. So the first generation is light because it stores the *entire* residue budget plus the three full C_6 windings; generations 2 and 3 are the winding complement. [E] (arithmetic).

12.1 Lepton flavor decuple: the charged-lepton determinant carries u^{10}

The source masses on the v_{geo} branch are $(\hat{m}_e, \hat{m}_\mu, \hat{m}_\tau) = \frac{v_{\text{geo}}\pi}{\sqrt{2}} (\frac{16}{7}u^5, \frac{4}{3}u^3, \frac{7}{6}u^2)$. Their product is a pure decuple in the seed:

$$\boxed{\frac{\hat{m}_e \hat{m}_\mu \hat{m}_\tau}{(v_{\text{geo}}\pi/\sqrt{2})^3} = \left(\frac{16}{7} \cdot \frac{4}{3} \cdot \frac{7}{6}\right) u^{5+3+2} = \frac{32}{9} u^{10}} \quad [\text{E}], \quad (63)$$

with $5+3+2 = g_{\text{car}} + b + s = 10 = \binom{5}{2}$. So the charged-lepton Yukawa powers $(n_e, n_\mu, n_\tau) = (5, 3, 2)$ are the three carrier numbers (g_{car}, b, s) , and the lepton determinant carries the same decuple (u^{10}) that the cosmological constant carries in the gauge sector ($A_\Lambda = 10$): Λ and the charged-lepton determinant are two realisations of the same degree- $\binom{5}{2}$ sector.

Sum–product reading. The lepton powers $(5, 3, 2)$ encode *both* decuple numbers at once: their sum is the Λ action, their product is the E_8 Coxeter number,

$$5 + 3 + 2 = 10 = A_\Lambda, \quad 5 \cdot 3 \cdot 2 = 30 = h_{E_8} = 2 \cdot 3 \cdot 5, \quad (64)$$

and $(5, 3, 2)$ is the prime factorisation of $h_{E_8} = 30$. So $\det M_\ell$, ρ_Λ and the E_8 Coxeter period read the same carrier core additively ($= 10$) and multiplicatively ($= 30$). [E]/[C]

12.2 Why the top Yukawa is ≈ 1 : the $L = 0$ anchor (Froggatt–Nielsen ladder)

The transport length L_f is a Froggatt–Nielsen-type charge with expansion parameter λ_C :

$$\boxed{y_f \simeq \lambda_C^{L_f} \times \Lambda_f \ (\Lambda_f = \mathcal{O}(1)), \quad y_t \simeq 1 \text{ because } L_t = 0} \quad [\text{E}]/[\text{C}]. \quad (65)$$

The top sits at $L_t = 0$ (the *unwound* carrier direction), so $y_t = \Lambda_t \simeq 1$ ($\sqrt{2} m_t/v = 0.991$); every other charged fermion is suppressed by integer powers of $\lambda_C = 0.2244$. Numerically (with $\mathcal{O}(1)$ sector prefactors $\Lambda_f \in [0.43, 1.07]$):

f	t	b	τ	c	μ	s	u, d	e
L_f	0	2	3	3	5	5	7	8
$y_f/\lambda_C^{L_f}$	0.99	0.48	0.90	0.65	1.07	0.94	$\sim 0.4\text{--}0.9$	0.46

So the entire charged-fermion hierarchy is one λ_C -ladder anchored by the top at $L = 0$; the lightness of the first generation ($L = 7, 8$) and the heaviness of the top ($L = 0$) are the two ends of the same integer winding spectrum. This is the standard Froggatt–Nielsen mechanism with λ_C as the flavon ratio and L_f as the FN charge — a clean bridge between TFPT transport and an established flavor mechanism.

13 Inflation interface

Machine checks for this section: `[verification/v7_gravity_cosmo.py]` `[verification/v86_nstar_reheating.py]`

The documented scalaron is Starobinsky R^2 ; with the pivot $N_\star = 57.1$:

$$\begin{aligned} n_s &= 1 - \frac{2}{N_\star} = 0.96497, & r &= \frac{12}{N_\star^2} = 0.00368, \\ \frac{M_{\text{scal}}}{M_{\text{Pl}}} &= \sqrt{8\pi} c_3^4 = c_3^{7/2} = (8\pi)^{-7/2} \Rightarrow M_{\text{scal}} \approx 3.06 \times 10^{13} \text{ GeV}. \end{aligned} \quad (66)$$

n_s matches Planck; $r = 0.0037$ is the falsifiable target for LiteBIRD/CMB-S4.

Amplitude (status-clean). The scalaron mass is a pure seam power,

$$\boxed{\frac{M_{\text{scal}}^2}{M_{\text{Pl}}^2} = c_3^{r-1} = c_3^{\Omega_{\text{adm}} - 10b_1} = c_3^7} \quad [\text{E}], \quad (67)$$

an *exact* identity (the exponent $7 = 48 - 41$ is the post-flavor, post-Higgs occupancy deficit). The cosmology interface reduces the geometric gravity action to an FRW R^2 /scalaron branch in which M_{scal} is the Starobinsky mass M ; the identification is therefore derived in that sector, not posited. With the standard large- N amplitude $A_s = \frac{N_\star^2}{24\pi^2} (M/\bar{M}_{\text{Pl}})^2$ this gives

$$A_s = \frac{N_\star^2 c_3^7}{24\pi^2} = \frac{N_\star^2 \delta_2}{8640\pi} = 2.17 \times 10^{-9} \quad [\text{C}]/[\text{O}], \quad (68)$$

matching Planck ($\simeq 2.1 \times 10^{-9}$); $\delta_2 = 2880 c_3^8$ is the second seam defect. The seam-power form is exact [\[E\]](#); the amplitude is then determined *within* the R^2 sector, the only residual being the geometric-gravity (Hodge) closure on which that sector rests [\[C\]](#). *Convention note:* $A_s \propto N_\star^2$, so the quoted 2.17×10^{-9} uses the pivot $N_\star = 57.1$; the alternative seed pivot $N_\star = 3/u - 1 = 55.4$ gives 2.05×10^{-9} . A number should only be quoted with $N_\star$ fixed.

14 Further simple bridges

Machine checks for this section: `[verification/v74_compiler_micro_lemmas.py]`

- **Dark-energy equation of state $w = -1$ (exact).** The TFPT Λ is a genuine vacuum constant $\rho_\Lambda = \frac{3}{4\pi^2} e^{-2\alpha_\star^{-1}} \bar{M}_{\text{Pl}}^4$, not a rolling field, so $w = -1$ identically (no quintessence). A robust measurement $w \neq -1$ would falsify the Λ -branch. [\[E\]](#) (structural).
- **Effective neutrino number $N_{\text{eff}} \simeq 3$.** Three families ($N_{\text{fam}} = 3$) give three light neutrinos and the standard $N_{\text{eff}} \simeq 3.04$. [\[E\]](#).
- **Top Yukawa $\simeq 1$** as the $L = 0$ anchor (above): a bridge to the Froggatt–Nielsen flavor mechanism.
- **Flagged (not clean):** $m_H/v = 0.509$ is near $\frac{1}{2}$ (1.7%); $m_W/m_Z = 0.881$ matches the running $\sqrt{1 - \sin^2 \theta_W(M_Z)}$ (not the tree $\sqrt{5/8} = 0.79$). No clean closed bridge for the Higgs/ W/Z masses yet. [\[O\]](#)

15 Closures that the architecture rests on

Machine checks for this section: `[verification/v36_spectral_action_g2.py]` `[verification/v4_flavor_matrix.py]`

Five results close or sharpen the discrete core. They are *proved in this document set* — the φ_0^{ret} -ladder, the H2 realisation and the strong-CP/dynamics gap in document 2 (`tfpt_2_standard_model`), the $E_6 \times A_2$ shadow in document 3 (`tfpt_3_e8_audit_bootstrap`) — and are collected here because the architecture above depends on them.

1. **The φ_0^{ret} -ladder (whole spectrum, one formula).** Every fermion mass is $\hat{m}_{f,j} = \frac{v_{\text{geo}}}{\sqrt{2}} \lambda_Y^{L_{f,j}} \Lambda_{f,j}$ with $\lambda_Y = \sqrt{\varphi_0^{\text{ret}}(1 - \varphi_0^{\text{ret}})}$, and the closed evaluation gives the universal form $\lambda_Y^L \Lambda = \pi c (\varphi_0^{\text{ret}})^k$ — the nine Λ are hexagonal-resolvent outputs (not free numbers); the *charged-lepton* ones are exact, the *quark* ones are reduced to the (U_{wall}) selector. The φ_0^{ret} -order ladder is $t(0) > \{b, c, \tau\}(2) > \{s, \mu\}(3) > \{u, d\}(4) > e(5)$. [E] (leptons exact, quark ratios closed $v49/v71$) / [O] (absolute quark scale)
2. **H2 reduced via the Mehta–Seshadri framework.** Mehta–Seshadri supplies the equivalence (stable parabolic $\leftrightarrow$ irreducible unitary); the TFPT-specific step — “stable = transport-minimal” and the geodesic C_6 word length — is conditional on the unitarity (U) of ∇_F^* (document 3) and the parabolic-degree $\rightarrow C_6$ dictionary. H2 is the sharp realisation statement: the D_4 -equivariant stable structure on $\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$ realises the branch with $\det R = h(D_5) = 8$, principal 2-minors $(2, 3, 5)$. [E] (given (U))
3. **Strong CP / dynamics via the explicit gap.** $\theta_{\text{eff}} = 0$ needs only γ_5 -Hermiticity + polar mass + sheet involution + reflection positivity (no mass gap). The full dynamics closes on the physical sector because the C_6 transport transfer operator has the *explicit* spectrum $\{(k/3)^6\}$, giving the mass gap $\Delta = 6 \log \frac{3}{2} = 2.43$ in closed form $\Rightarrow$ OS reconstruction $\Rightarrow$ measure on the admissible sector. [E] (phys. sector)
4. **$E_6 \times A_2$ flavor shadow.** The residue matrix sits inside an E_8 secondary branching: $248 = \|R\|_F^2 + \det R + 2(1^\top R a) N_{\text{fam}} = 78 + 8 + 2 \cdot 27 \cdot 3$, with $\|R\|_F^2 = 78 = \dim E_6$, $\det R = 8 = \dim A_2$. The seven E_8 slices form an audit raster (atlas note). [O]
5. **SM completeness.** The discrete structure and dimensionless core are closed; the dimensionful EW/QCD masses $(m_W, m_Z, m_H, \sin^2 \theta_W, \alpha_s)$ are schema-level via $\mathcal{R}_{\text{SM}}$, with m_H resting on the closed UV quartic $\lambda_\Phi = 1/(16\pi^2)$ (near-criticality). The *exact* solar angle θ_{12} (leading $1/N_{\text{fam}} = \frac{1}{3}$) is *conditionally derived*, pending the seam-misalignment lemma — the sharpest remaining SM tension point. [C]

16 Status summary and open audit points

Machine checks for this section: [verification/v22_open_gates_audit.py]

Statement	Status	Note
Dual-root dictionary $(Y, \text{cusps}, \gamma_{\text{car}}, \varphi_{\text{base}})$	[E]	exact, from $\{1/2, -1/3\}$
$10b_1 = 5^2 + 4^2 = \Omega_{\text{adm}}\gamma + 1$	[E]	Pythagorean carrier identity
$I_1^{\text{EW}} = 41/48 = \text{Var}(Y) b_1$	[E]	exact
$\delta_2 = 4! \cdot 5! c_3^8, \text{Tr } X^2 = 5!$	[E]	factorial traces
$5/4 = g_{\text{car}}/(g_{\text{car}} - 1)$ universal compression	[E]	exact
$u = \frac{4}{3}c_3 + 48c_3^4$; one-decoder identities	[E]	exact
$\xi = \frac{3}{4}/(1 + 9/128\pi^3)$	[E]	corrected tree value
Ladder $A_{\text{EW}} : \mathcal{A}_H : \mathcal{A}_\Lambda = 1 : g_{\text{car}} : 2g_{\text{car}}$	[E]/[C]	Hubble rung is a corollary
Ladder $= \binom{g_{\text{car}}}{0} : \binom{g_{\text{car}}}{1} : \binom{g_{\text{car}}}{2}$; $2g_{\text{car}} = \binom{g_{\text{car}}}{2} \Leftrightarrow g_{\text{car}} = 5$	[E]	new overdetermination of $g_{\text{car}} = 5$
2D action lattice $S_x = a(\alpha_\star^{-1}/g_{\text{car}}) + bL_0$	[E]/[C]	two integer labels per scale
Decuple bridge $\rho_\Lambda/\bar{M}_{\text{Pl}}^4 = \frac{3}{4\pi^2} e^{2\delta_{\text{ph}}} [v/(5\beta_{\text{rad}}^2 \bar{M}_{\text{Pl}})]^{10}$	[E]	exact, $10 = 2g_{\text{car}} = \dim \Lambda^2 E$
Family = ladder = Pascal row $\{1, 5, 10\}$; $16 = 1 + 5 + 10$	[E]	central new identification
$g_{\text{car}} = 5$ via $2^{g_{\text{car}}-1} = \binom{g_{\text{car}}}{0} + \binom{g_{\text{car}}}{1} + \binom{g_{\text{car}}}{2}$	[E]	Pascal closure (unique)
$\text{Tr}_{S^+} X^2 = 5!$ generates 40, 41, 240, δ_2	[E]	master moment
$ R(E_8) = 16 \cdot 5 \cdot 3$; $\dim E_8 = 240 + 8$	[E]	E_8 compression
Even-flip atlas $240 = 160 + 80$	[E]/[O]	E_8 metrization open ($\neq 112 + 128$)
$\delta_{\text{CKM}} - \pi/3 = 3u(1 - u)$	[E]	CP = triple seed variance

Observable quadrilateral ($u = \Omega_b + \beta_{\text{rad}}$)	[E]	closed UV-shadow cycle
Power tower $v=x^1$, $H=x^5 R_H$, $\Lambda=x^{10} R_\Lambda$	[E]	K_5 vertices/edges
$496 = 16 \cdot 31 = \dim(E_8 \times E_8) = \dim SO(32)$	[E]/[C]	heterotic-dimension count
$A_{\text{EW}} = \alpha_\star^{-1}/\sqrt{\Delta_Y}$, $\Delta_Y = (b+s)^2 = 25$	[E]	discriminant action theorem
$\hat{m}_e \hat{m}_\mu \hat{m}_\tau \propto \frac{32}{9} u^{10}$; $(5, 3, 2) = (g_{\text{car}}, b, s)$	[E]	lepton flavor decuple
$M_{\text{scal}}/\bar{M}_{\text{P1}} = c_3^{7/2}$	[E]	scalaron seam power
$\delta_{\text{CKM}} - \pi/3 = N_{\text{fam}} u(1-u)$	[E]	weak CP = $N_{\text{fam}} \times$ seed variance
$L = R + W$: $\sum R = 22$, $\sum W = 18$	[E]	first generation = residue store
$ R(E_8) /6 = \sum L = 40$ ($\mathbb{Z}_6$ slice)	[E]	one $\mathbb{Z}_6$ slice = flavor budget
$h_{E_8} = R /\text{rank} = 30 = N_{\text{fam}} \binom{5}{2} = N_{\text{fam}} A_\Lambda$	[E]/[C]	carrier Coxeter theorem
$ R^+(E_8) = \text{Tr}_{S^+} X^2 = 120 = 5!$	[E]	positive-root moment
common decuple $\binom{5}{2} = A_\Lambda = u^{5+3+2} = h_{E_8}/N_{\text{fam}} = 10$	[E]/[C]	one carrier pair-decuple
$\lambda_C/\delta_{\text{top}} = 1.061$ (winding $\approx$ seam defect)	[O]	flagged proximity (not exact)
$y_f \simeq \lambda_C^{L_f}$, top $L_t = 0 \Rightarrow y_t \simeq 1$	[E]/[C]	Froggatt–Nielsen ladder
$w = -1$ (true Λ); $N_{\text{eff}} \simeq 3$	[E]	dark energy / neutrino species
3-generator lattice $\{\alpha_\star^{-1}, L_0, L_c\}$; $M_{\text{scal}} = e^{-\frac{7}{2}L_c}$	[E]	seam axis $L_c = \ln(1/c_3)$
$\dim S^+ = 2 \text{rank}(E_8)$, $h_{E_8} = 2(\dim S^+ - 1) = 2^g - 2$	[E]	rank–Coxeter coupling (ν^c closes it)
$\gcd(2h, 6r) = \dim \mathfrak{g}_{\text{SM}} = 12$, $\text{lcm} = 240$	[E]/[C]	SM algebra = gcd; E_8 count = lcm
$I_1^{\text{EW}} = h_{E_8}/36 + 1/(6r) = 41/48$	[E]/[C]	Coxeter part + rank singlet
$M_{\text{scal}}^2/\bar{M}_{\text{P1}}^2 = c_3^{-1} = c_3^7$ exact; $A_s = N_\star^2 c_3^7/24\pi^2$	[E]/[C]/[O]	seam power exact; A_s conditional (Starobinsky)
$h_{E_8} = 2 \cdot 3 \cdot 5 = 30$, $r = \varphi(30) = 8$	[E]/[C]	2, 3, 5 Coxeter cyclotomic compiler
$\text{Exp}(E_8) = (\mathbb{Z}/30)^\times$, $\sum = 120$, $\chi_C = \Phi_{30}$	[E]/[C]	exponents = primitive residues
lepton $(5, 3, 2)$: $\sum = 10 = A_\Lambda$, $\prod = 30 = h_{E_8}$	[E]/[C]	sum=action, product=Coxeter
Hubble $H_0/\bar{M}_{\text{P1}} = e^{-\alpha_\star^{-1}}/(2\pi\sqrt{\Omega_\Lambda})$	[E]	exact corollary
Λ -anchored closure $G_N = \frac{3c^3}{32\pi^3 \hbar \Lambda_{\text{geom}}} e^{-2\alpha_\star^{-1}}$	[E]/[C]	G_N to -0.35%
$g_{\text{car}} = 5$ overdetermined	[C]	monotone $\alpha^{-1}(g_{\text{car}})$
$ R(E_8) = 240 = 16 \cdot 15$, $248 = 8 \cdot 31$	[E]	counting identities
$E_8 \leftrightarrow$ carrier-transition map	[C]	needs structure map
$\delta_{\text{CP}} = 4\pi/3$, $\theta_{12} = \frac{1}{3} - \frac{u}{2}$	[C]	PMNS readouts
$ V_{td} , V_{ts} , J_{\text{CKM}}, J_{\text{CP}}^\nu; n_s, r$	[C]	derived readouts

Open audit points: (1) *discharged* — the EM closure $F_{U(1)}(\alpha) = 0$ is now stated explicitly (Grammar III), with existence/uniqueness and root $\alpha_\star^{-1} = 137.0359992168$ (2.9×10^{-10} , 1.9σ vs CODATA-2022); (2) keep $\gamma_{\text{car}} = 5/6$ vs γ_E distinct; (3) type M_{P1} vs $\bar{M}_{\text{P1}}$ and “pure” ($A_\Lambda = 274.07$) vs “total” ($-\ln(\delta_{\text{top}} e^{-2\alpha_\star^{-1}}) = 283.10$, $-\ln(\rho_\Lambda/\bar{M}_{\text{P1}}^4) = 276.65$) actions consistently; (4) version the Λ prefactor branch; (5) the scalaron-mass $\rightarrow$ Starobinsky- M identification is the *only* condition for $A_s = N_\star^2 c_3^7/(24\pi^2)$ (the seam power $M_{\text{scal}}^2/\bar{M}_{\text{P1}}^2 = c_3^7$ itself is exact); (6) the E_8 *lattice* is now built explicitly as $(D_5 \oplus A_3) + \mu_4$ -glue (Section 3), so the former “blind 240-metrisation” is closed. The flavor status (companion parabolic-weight note) now splits cleanly: the *anchor multiset* $\{1, 1, 2\}$ is derived (Schur–Horn + stability) [E]; the *sector assignment* $(1, 1, 2)$ follows from the D_5 cusp ordering [E]/[C]; the *first-generation* +6 *winding* is the hierarchy convention [C]; the six *non-anchor residues* are then forced (residue sets = D_6 orbit of $\{0, 1, 3\}$, minus anchor, monotone) [E]. The remaining geometric input at the time of writing was (H2), the specific D_6 cusp labelling on $\mathbb{P}^1 \setminus \mu_4$ — sharply characterised by the residue-matrix spectral selector ($\det R = h(D_5)$, principal minors $(2, 3, 5)$), with a naive cusp-position shortcut explicitly ruled out [C]. The explicit Riemann–Hilbert construction confirms the rigid $SU(3)$ system and its A_3 Coxeter rotation. *Status update:* the H2 dictionary has since been made *exact* (hexagon = sign-twisted family spectrum, addresses pinned, margins forced, v118–v122), the deck is *derived* (v141), and the residue is the $R4'$ selector

establishment — two line pairings (v139/v142); what stays geometric is only the parabolic realisation premise (GATE.QGEO, research contracts).

Honest negatives (reported to keep the discipline visible). The TFPT lepton ladder gives a Koide ratio $Q = (\sum m_\ell)/(\sum \sqrt{m_\ell})^2 = 0.6645$, close to but *not* the empirical $2/3 = 0.66667$ (-0.33%); TFPT is Koide-*like* but does not reproduce the precise relation. The baryon asymmetry $\eta_B \approx 6 \times 10^{-10}$ and the proton/electron ratio $m_p/m_e = 1836$ do *not* fall on a clean action charge or L_0 ladder; they remain outside the present grammar. These are stated so the strong items are not read as a uniform success.

17 Consolidated picture

The three nested grammars.

Carrier: $3 + 2 \Rightarrow \{y_-, y_+\} = \{-\frac{1}{3}, \frac{1}{2}\} \Rightarrow Y$, $\dim S^+ = 16$, $N_{\text{fam}} = 3$, $\Omega_{\text{adm}} = 48$, $b_1 = \frac{41}{10}$,

Seed: $u = \frac{4}{3}c_3 + 48c_3^4 \Rightarrow \{\lambda_C, \beta_{\text{rad}}, \Omega_b, \sin^2 \theta_{13}\}$,

Action: $\alpha_\star^{-1} \Rightarrow A_{\text{EW}} = \frac{\alpha_\star^{-1}}{g_{\text{car}}}$, $\mathcal{A}_\Lambda = 2\alpha_\star^{-1}$, $\mathcal{A}_H = \alpha_\star^{-1} \Rightarrow \frac{\rho_\Lambda}{M_{\text{Pl}}^4} = \frac{3}{4\pi^2} e^{2\delta_{\text{ph}}} \left[\frac{v_{\text{geo}}}{5\beta_{\text{rad}}^2 M_{\text{Pl}}} \right]^{10}$.

A small grammar generates many sectors: the Standard-Model packet, flavor, the electroweak hierarchy, the cosmological constant, the metrology, and the inflation/CMB interface. The strongest new statements are exact: the decouple bridge ties Λ to the tenth power of the stripped electroweak scale with $10 = 2g_{\text{car}}$; the Hubble scale is the square root of Λ ; and the Einstein normaliser is the corrected tree value $\frac{3}{4}/(1 + 9/128\pi^3)$. None of this is claimed as a proof of a final theory; it is offered as a tightening of the existing architecture for your assessment.

The compiler in one line: TFPT = even carrier code + seed decoder + 2·3·5 Coxeter compiler. From the three atoms (2, 3, $g_{\text{car}}=5$):

$$h_{E_8} = 2 \cdot 3 \cdot 5 = 30, \quad r_{E_8} = \varphi(30) = 8 = g_{\text{car}} + N_{\text{fam}},$$

$$|R(E_8)| = rh = 240, \quad \dim E_8 = r(h+1) = 8 \cdot 31 = 248,$$

$$\text{Exp}(E_8) = (\mathbb{Z}/30)^\times, \quad \sum \text{Exp} = 120 = \text{Tr}_{S^+} X^2 = |R^+|, \quad \chi_C = \Phi_{30},$$

$$\dim S^+ = 1 + 10 + 5 = 16 = 2r, \quad A_{\text{EW}} : \mathcal{A}_H : \mathcal{A}_\Lambda = 1:5:10, \quad \mathcal{A}_H = \frac{h}{6}, \quad \mathcal{A}_\Lambda = \frac{h}{3}, \quad N_{\text{fam}} = \frac{h}{A_\Lambda} = 3,$$

$$\sum L = \frac{|R^+|}{3} = 40, \quad 10b_1 = 1 + \frac{|R^+|}{3} = 41 = 1 + h \frac{\dim S^+}{\dim \mathfrak{g}_{\text{SM}}},$$

$$\det M_\ell \propto u^{10}, \quad \frac{\rho_\Lambda}{M_{\text{Pl}}^4} \propto \left[\frac{v_{\text{geo}}}{5\beta_{\text{rad}}^2 M_{\text{Pl}}} \right]^{10}, \quad A_{\text{EW}} = \frac{\alpha_\star^{-1}}{\sqrt{\Delta_Y}}, \quad \sqrt{\Delta_Y} = g_{\text{car}} = 5.$$

The common decuple (= 10): ($\frac{g_{\text{car}}}{2}$) (carrier pairs) = A_Λ (cosmological action) = u^{5+3+2} (lepton determinant) = h_{E_8}/N_{fam} (Coxeter per family). Λ , the charged-lepton determinant, the E_8 Coxeter structure and the action ladder all read the *same* carrier pair-decuplet. The open core is one question: can the even-flip atlas be metrised as a genuine E_8 root system (the 160+80 vs 112+128 obstruction)?

Part II

The Coxeter–cyclotomic compiler and the explicit E_8 construction

Abstract

This short companion compresses the TFPT bridge layer to a single discrete object. The claim is that the Standard-Model family, the cosmological action ladder, the E_8 counting data, the flavor budget, inflation and the cosmological constant are *not* independent coincidences but projections of one structure: the even-weight code C^+ on $g_{\text{car}} = 5$ slots, together with its $\mathbb{Z}_{30}$ Coxeter spectrum and the single seam seed $c_3 = 1/(8\pi)$. The decisive new result is that C^+ is precisely the half-spin weight orbit of $D_5 = \mathfrak{so}_{10}$ ($\text{Aut}(C^+) = W(D_5)$, order 1920), that the family geometry $\mathbb{P}^1 \setminus \mu_4$ is an $A_3 = \mathfrak{su}_4$, and that E_8 is then the standard *closure algebra* of $D_5 \times A_3$ — not an input. The fusion is exact: $\text{disc}(D_5) = \text{disc}(A_3) = \mathbb{Z}_4$, so the four family corners μ_4 are precisely the glue code that closes $D_5 \oplus A_3$ into the unimodular E_8 lattice. The glue is metric: the discriminant-form norms $q(D_5) = \frac{5}{4}$ and $q(A_3) = \frac{3}{4}$ sum to the E_8 root norm 2 — and these are exactly the pre-existing TFPT constants $\delta_2/\delta_{\text{top}}^2 = \frac{5}{4}$ and $\xi_{\text{tree}} = \frac{3}{4}$. The A_3 side is explicit: the residue pairing on $H^1(\mathbb{P}^1 \setminus \mu_4)$ is the A_3 Cartan form, and the carrier corner rotation $z \mapsto iz$ acts as the A_3 Coxeter element (order 4, spectrum $\{i, -1, -i\}$). The lift is no longer a count: the 240 E_8 roots are *explicitly* built from $D_5 \oplus A_3$ plus the spinor (μ_4) glue, all of norm 2, Cartan determinant 1. Seven statements carry it; every lattice identity below is checked exactly, and the analytic metric is computed and characterised (a D_4 -equivariant deformation), so no *lattice* structural step is left open in the $D_5 \times A_3$ glue route (the upstream boundary kernel, EM closure and flavor Hecke saturation remain separate open problems). Master statement: *five carrier slots give D_5 , four family corners give A_3 , and μ_4 glues $D_5 \times A_3$ into the E_8 atlas.*

The compression.

$$\begin{aligned} \text{TFPT} &= \underbrace{D_5 \text{ half-spin carrier}}_{C^+} + \underbrace{A_3 \text{ four-point family}}_{\mathbb{P}^1 \setminus \mu_4} + c_3 \text{ seam seed}, \\ E_8 &= \text{closure of } D_5 \times A_3 \end{aligned}$$

with $c_3 = \frac{1}{8\pi}$, $g_{\text{car}} = 5$, $h := 2 \cdot 3 \cdot g_{\text{car}} = 30$, $r := \varphi(h) = \varphi(30) = 8$, $N_{\text{fam}} = 3$, $N_{\Phi} = 1$, $\dim \mathfrak{g}_{\text{SM}} = 12$. Equivalently $C^+ + \mathbb{Z}_{30}$ Coxeter compiler + c_3 seed; the $D_5 \times A_3$ reading is the structural lift of the same object.

Theorem 1 — the even carrier code

Let E be the rank- $g_{\text{car}} = 5$ carrier, split $E = E_3 \oplus E_2$ (color $b = 3$, weak $s = 2$). The one-generation spinor packet is the even exterior algebra, equivalently the even-weight binary code

$$S^+ = \Lambda^0 E \oplus \Lambda^2 E \oplus \Lambda^4 E \cong C^+ = \{x \in \mathbb{F}_2^5 : |x| \text{ even}\}, \quad \dim S^+ = |C^+| = 2^{g_{\text{car}}-1} = 16. \quad (69)$$

The two carrier roots $y_+ = \frac{1}{2}$, $y_- = -\frac{1}{3}$ generate all hypercharges ($Y(Q_L) = y_+ + y_- = \frac{1}{6}$, $Y(u^c) = 2y_-$, $Y(e^c) = 2y_+$, $Y(\nu^c) = 0$) and the transport cusps $\{1, \frac{2}{3}, \frac{1}{3}\} = \{2y_+, -2y_-, 2(y_+ + y_-)\}$, with $\gamma_{\text{car}} = \text{Tr}_E Y^2 = 3y_-^2 + 2y_+^2 = \frac{5}{6}$. [E]

Gauge-rank factorisation ([[verification/v79_review_identities.py](#)]). The carrier polynomial factors into the two gauge ranks directly:

$$\boxed{6Y^2 - Y - 1 = (2Y - 1)(3Y + 1)}, \quad Y = \frac{1}{2} \ (s=2, \ SU(2)), \quad Y = -\frac{1}{3} \ (b=3, \ SU(3)),$$

i.e. $(sY - 1)(bY + 1)$ — the polynomial splits into one linear representative of each Standard-Model non-abelian factor; the 5-slot carrier *forces* the split into $SU(3) \times SU(2)$.

Hypercharge [verification/v79_review_identities.py]	Lucas–Binet resonances	[E]	(mode reading	[C],
The integer carrier charges $(q_+, q_-) = (3, -2)$ (the primitive trace-free pair, $X^2 - X - 6 = (X - 3)(X + 2)$, Lean <code>unique_carrier_pair</code>) generate the Lucas U -sequence				
$D_n = \frac{3^n - (-2)^n}{5} : \quad D_1, \dots, D_6 = 1, 1, 7, 13, 55, 133,$				
which is, <i>in order</i> , $(N_\Phi, \cdot, \text{scalaron exponent } 7, \Delta_Q=13, \text{quark numerator } 55, \dim E_7=133)$. In particular the quark ratio reads $c_u/c_d = D_5/(N_{\text{fam}}^2 D_4) = \frac{55}{117}$ — a single recurrence of the hypercharge generator, not a free integer. [E] for the arithmetic; the per-mode physical identification is a [C] cross-domain reading (only $n \leq 6$, the carrier-slot range, is meaningful).				

Theorem 2 — Exterior–Action Duality

On C^+ the flip $F = x \oplus y$ is even, so $|F| \in \{0, 2, 4\}$. The three relations R_0, R_2, R_4 form a symmetric association scheme (the even Hamming scheme on five bits) with

$$\begin{array}{l} \text{valencies } (1, 10, 5) = (\dim \Lambda^0 E, \dim \Lambda^2 E, \dim \Lambda^4 E), \\ \text{spectral multiplicities } (1, 5, 10) = A_{\text{EW}} : \mathcal{A}_H : \mathcal{A}_\Lambda \end{array} \quad [\text{E}]. \quad (70)$$

The eigenvalues (computed directly) are $A_2 : \{10^{(1)}, 2^{(5)}, (-2)^{(10)}\}$ and $A_4 : \{5^{(1)}, (-3)^{(5)}, 1^{(10)}\}$, with $A_2 + A_4 = K_{16} - \mathbf{1}$. The two graphs are named objects:

$$\begin{array}{l} R_2(C^+) = \text{folded/complement, SRG}(16, 10, 6, 6) \text{ (the Clebsch complement);} \\ R_4(C^+) = \text{Clebsch graph, SRG}(16, 5, 0, 2) \end{array} \quad [\text{E}]. \quad (71)$$

Remark. The Standard-Model family is the *valency* side $(1, 10, 5)$ of one scheme; the cosmological action ladder is its *spectral* side $(1, 5, 10)$. The action ladder is therefore not merely “also binomial” — it is the spectral dual of the even exterior family. [C]

This duality is the MacWilliams transform ([verification/v79_review_identities.py]). C^+ is the even-weight code $[5, 4, 2]$; its dual $(C^+)^{\perp}$ is the repetition code $[5, 1, 5] = \{00000, 11111\}$ (the uniform/vacuum word). The MacWilliams identity sends the weight enumerator of the dual to that of C^+ , namely $(1, 10, 5)$ — so the valency $\leftrightarrow$ spectral duality *is* the coding-theoretic UV/IR duality, and the carrier grading $\Lambda^{\text{even}}(5) = 1+10+5$ is the MacWilliams image of the repetition vacuum. [E] (code identity) / [C] (holographic reading).

The full Bose–Mesner algebra (machine-checkable) fixes the duality beyond eigenvalues:

$$A_2^2 = 10 \mathbf{1} + 6A_2 + 6A_4, \quad A_2 A_4 = 3A_2 + 4A_4, \quad A_4^2 = 5 \mathbf{1} + 2A_2, \quad (72)$$

with eigenmatrix and multiplicity orthogonality

$$P = \begin{pmatrix} 1 & 10 & 5 \\ 1 & 2 & -3 \\ 1 & -2 & 1 \end{pmatrix}, \quad \det P = -64, \quad \boxed{P^{\top} \text{diag}(1, 5, 10) P = \text{diag}(16, 160, 80)} \quad [\text{E}], \quad (73)$$

so the flip split $240 = 160 + 80$ ($|F| = 2, 4$) appears directly in the scheme’s eigenmatrix orthogonality.

Carrier = D_5 (**verified**). The 16 vertices of C^+ are exactly the weights of the $D_5 = \mathfrak{so}_{10}$ half-spinor $\frac{1}{2}(\pm 1, \dots, \pm 1)$ with an even number of minus signs. The flip-2 and flip-4 graphs are translation- and permutation-invariant, so the *affine* group $C^+ \rtimes S_5$ (translations $\rtimes$ coordinate permutations) acts by graph/scheme automorphisms; the group closure gives the automorphism group of the even Hamming scheme (equivalently the Clebsch graph),

$$\boxed{\text{Aut}_{\text{aff}}(C^+) = C^+ \rtimes S_5, \quad |\text{Aut}_{\text{aff}}(C^+)| = 2^{g_{\text{car}}-1} g_{\text{car}}! = 16 \cdot 120 = 1920 = |W(D_5)|} \quad [\text{E}], \quad (74)$$

and $1920 = r |R(E_8)| = 8 \cdot 240$. (The *linear* code automorphism group alone is only the coordinate group S_5 of order 120; the factor $2^{g_{\text{car}}-1} = 16$ is the translation/scheme part. It is the affine/scheme group that equals $W(D_5)$.) Edges of the Clebsch graph: $|E(R_4)| = \frac{16 \cdot 5}{2} = 40 = |R(D_5)| = \sum_{f,j} L_{f,j}$ (the entire flavor transport budget). So the five-slot even code is the half-spin weight structure of $\mathfrak{so}_{10}$, the classical home of one SM family with ν^c .

Theorem 3 — the Coxeter closure equation selects $g_{\text{car}} = 5$

Requiring the even family to equal the action truncation through pair degree, $\dim S^+ = 2^{g-1} = \binom{g}{0} + \binom{g}{1} + \binom{g}{2}$, i.e.

$$\boxed{2^g - 2 = g(g+1) \iff g = 5} \quad [\mathbf{E}], \quad (75)$$

the unique positive-integer solution. At $g = 5$ the two sides equal 30, so *carrier exhaustion* $2^g - 2$ and the *Coxeter period* $g(g+1)$ coincide only at five slots: $2^{g_{\text{car}}} - 2 = 30 = h$ and $g_{\text{car}}(g_{\text{car}} + 1) = 30$.

Theorem 4 — the $\mathbb{Z}_{30}$ cyclotomic Coxeter compiler

From the three atoms $(2, 3, g_{\text{car}})$, with $h = 2 \cdot 3 \cdot g_{\text{car}} = 30$ and $r = \varphi(h) = 8$:

$$\boxed{|R(E_8)| = rh = 240, \quad \dim E_8 = r(h+1) = 8 \cdot 31 = 248, \quad h+1 = 2^{g_{\text{car}}} - 1 = 31} \quad [\mathbf{E}]. \quad (76)$$

The E_8 exponents are the primitive residues modulo 30,

$$\text{Exp}(E_8) = (\mathbb{Z}/30\mathbb{Z})^\times = \{1, 7, 11, 13, 17, 19, 23, 29\}, \quad \chi_C(t) = \Phi_{30}(t) = t^8 + t^7 - t^5 - t^4 - t^3 + t + 1, \quad (77)$$

and $|(\mathbb{Z}/30)^\times| = \varphi(30) = 8 = r$. The local SM algebra and the root atlas are the two arithmetic projections of $(r, h) = (8, 30)$:

$$\Omega_{\text{adm}} = 6r = 48, \quad D_{\text{start}} = 2h = 60, \quad \gcd(6r, 2h) = \dim \mathfrak{g}_{\text{SM}} = 12, \quad \text{lcm}(6r, 2h) = rh = 240. \quad (78)$$

Theorem 5 — the root transition atlas and the positive-edge moment

The directed even-flips of C^+ are the complete graph K_{16} :

$$\boxed{\begin{aligned} |R^+(E_8)| &= \binom{\dim S^+}{2} = \binom{16}{2} = 120 = \text{Tr}_{S^+} X^2 = 5! = \sum_{m \in (\mathbb{Z}/30)^\times} m, \\ |R(E_8)| &= 2 \binom{16}{2} = 240 \end{aligned}} \quad [\mathbf{E}]. \quad (79)$$

($X = 6Y$.) So the positive E_8 roots are the unordered transitions of one family; the second hypercharge moment, the positive-root count and the sum of the E_8 exponents coincide at 120. The CP pairing $m \leftrightarrow 30 - m$ of the exponents gives four pairs, each of sum $h = 30$:

$$\boxed{\text{Tr}_{S^+} X^2 = (N_{\text{fam}} + N_{\Phi}) h = 4 \cdot 30 = 120 = \dim \mathfrak{so}_{16} = \underbrace{45}_{D_5} + \underbrace{15}_{A_3} + \underbrace{60}_{(10,6)}} \quad [\mathbf{E}], \quad (80)$$

so the hypercharge moment is simultaneously four CP-paired Coxeter periods *and* the adjoint half $\mathfrak{so}_{16} = (45, 1) \oplus (1, 15) \oplus (10, 6)$ of the E_8 closure.

How the root *origin* is now obtained — see Theorem 7. A naive flip-size-preserving map $R_{\text{flip}} \rightarrow \mathbb{R}^8$ cannot be E_8 : the flip split $240 = 160 + 80$ ($|F| = 2, 4$) differs from the D_8

split $112 + 128$. The correct route is *not* a direct 240-metrization but the standard branching $E_8 \supset D_5 \times A_3$ (Theorem 7), which builds the root spaces from known D_5 and A_3 data. Until the A_3 lattice is derived from the four punctures (Theorem 7, [O]), E_8 remains a Coxeter atlas with a concrete, falsifiable lift.

Theorem 6 — downstream readouts from (h, r, c_3)

All discrete budgets and the seam seed are functions of $(h, r, c_3) = (30, 8, 1/(8\pi))$:

$$\dim S^+ = 2r = 16, \quad h = 2(\dim S^+ - 1) = 30, \quad (81)$$

$$10b_1 = 5r + 1 = 41 = 1 + h \frac{\dim S^+}{\dim \mathfrak{g}_{\text{SM}}}, \quad I_1^{\text{EW}} = \frac{5r + 1}{6r} = \frac{41}{48}, \quad (82)$$

$$u = \frac{r}{6}c_3 + 6r c_3^4 = \frac{4}{3}c_3 + 48c_3^4, \quad \delta_2 = 4! \binom{16}{2} c_3^8 = 2880 c_3^8, \quad (83)$$

$$\xi_{\text{tree}} = \frac{6}{r} = \frac{\dim \mathfrak{g}_{\text{SM}}}{\dim S^+} = \frac{3}{4}, \quad \frac{M_{\text{scal}}^2}{M_{\text{Pl}}^2} = c_3^{\Omega_{\text{adm}} - 10b_1} = c_3^{r-1} = c_3^7. \quad (84)$$

Two-graph product table. With the carrier-pair graph K_5 ($|E(K_5)| = \binom{5}{2} = 10 = \mathcal{A}_\Lambda$) and the family-corner graph K_4 on μ_4 ($|E(K_4)| = \binom{4}{2} = 6$, $|R(A_3)| = 2|E(K_4)| = 12$), every large budget is a product of the two graphs:

$$\boxed{40 = 10 \cdot 4, \quad 60 = 10 \cdot 6, \quad 120 = 10 \cdot 12, \quad 240 = 10 \cdot 24} \quad [\text{E}], \quad (85)$$

i.e. $(\sum L, D_{\text{start}}, |R^+(E_8)|, |R(E_8)|) = 10 \cdot (|\mu_4|, |E(K_4)|, |R(A_3)|, |S_4|)$. Two occupancy identities follow,

$$\boxed{10b_1 = 1 + |\mu_4| |E(K_5)| = 1 + 4 \cdot 10 = 41}, \quad (86)$$

$$\boxed{\Omega_{\text{adm}} = N_{\text{fam}} \dim S^+ = (N_{\text{fam}} + N_\Phi) |R(A_3)| = 3 \cdot 16 = 4 \cdot 12 = 48} \quad [\text{E}],$$

so the Higgs singleton is the fourth (μ_4) direction that closes $N_{\text{fam}} = 3$ to **4** of A_3 . **Occupancy deficit ladder.** The admissible occupancy splits cleanly,

$$\boxed{\begin{aligned} \Omega_{\text{adm}} &= \underbrace{40}_{\sum L (\text{flavor})} + \underbrace{1}_{\text{Higgs}} + \underbrace{7}_{\text{scalaron}} = 48, & \Omega_{\text{adm}} - \sum L &= 8 = h(D_5) = r(E_8), \\ \Omega_{\text{adm}} - 10b_1 &= 7 = h(D_5) - 1 \end{aligned}} \quad [\text{E}], \quad (87)$$

so the scalaron exponent $r - 1 = 7$ is exactly the post-flavor, post-Higgs occupancy remainder, $M_{\text{scal}}^2/\bar{M}_{\text{Pl}}^2 = c_3^{\Omega_{\text{adm}} - 10b_1} = c_3^7$. With $h(D_5) = 8$ and $|R^+(A_3)| = 6$ the seed itself is a D_5/A_3 ratio-product, $u = \frac{h(D_5)}{|R^+(A_3)|} c_3 + h(D_5) |R^+(A_3)| c_3^4$, and $\xi = \frac{|R^+(A_3)|}{h(D_5)} (1 + |R^+(A_3)|^2 c_3^3)^{-1} = \frac{3}{4} (1 + 36c_3^3)^{-1}$. The seed readouts and the two decuples are

$$\lambda_C^2 = u(1-u), \quad \beta_{\text{rad}} = 2c_3 u, \quad \sin^2 \theta_{13} = e^{-5/6} u, \quad \boxed{\omega_b = \frac{\lambda_C}{\mathcal{A}_\Lambda} = \frac{\lambda_C}{10} = 0.02244} \quad (\text{Planck } 0.02237), \quad (88)$$

$$\boxed{\begin{aligned} \det M_\ell &\propto u^{h/N_{\text{fam}}} = u^{10}, \\ \frac{\rho_\Lambda}{\bar{M}_{\text{Pl}}^4} &= \frac{3}{4\pi^2} e^{2\delta_{\text{ph}}} \left[\frac{v_{\text{geo}}}{5\beta_{\text{rad}}^2 M_{\text{Pl}}} \right]^{h/N_{\text{fam}}}, \end{aligned}} \quad (89)$$

with lepton powers $(5, 3, 2) = (g_{\text{car}}, b, s)$: $\sum = 10 = \mathcal{A}_\Lambda$ (the Λ decuple), $\prod = 30 = h$ (the Coxeter period). The first-generation transport sums read the non-primitive $\mathbb{Z}_{30}$ load,

$$(G_1, G_2, G_3) = (22, 13, 5) = (h - r, \dim \mathfrak{g}_{\text{SM}} + 1, g_{\text{car}}), \quad h - r = 30 - \varphi(30) = 22, \quad (90)$$

Flavor transport is a $D_5 \times A_3$ object. The transport packets $L_u=(7, 3, 0)$, $L_d=(7, 5, 2)$, $L_e=(8, 5, 3)$ split into a D_5 budget and an A_3 geometry:

$$\boxed{\sum_{f,j} L_{f,j} = 40 = |R(D_5)| \text{ (Clebsch edges)}, \quad L_{f,j} \bmod 6 \text{ lives on } C_6 = Z_6 = |R^+(A_3)|} \quad [\mathbf{E}], \quad (91)$$

the residues being the D_6 -orbit of the perfect set $\{0, 1, 3\} \subset C_6$ (cyclic distances $\{1, 2, 3\}$): $u \equiv \{0, 1, 3\}$, $d \equiv 2 - \{0, 1, 3\}$, $e \equiv 2 + \{0, 1, 3\}$, with the universal first-generation $+6$ winding. So the carrier D_5 sets the total budget and the family A_3 (its 6-element positive-root hexagon) sets the per-entry distances. The scalaron mass is a pure seam power $[\mathbf{E}]$; the amplitude is a *conditional* readout, kept $[\mathbf{C}]/[\mathbf{O}]$ until the scalaron variable is formally identified with the Starobinsky M parameter. That identification is now *heat-kernel grounded* (G2): the spectral-action Seeley–DeWitt coefficient $a_4|_{R^2} \sim R^2/72$ delivers exactly the Starobinsky R^2 term, with $M^2/\bar{M}_{\text{Pl}}^2 = 6(4\pi)^2/f_0$ and the TFPT closure c_3^7 fixing the cutoff moment f_0 (`[verification/v36_spectral_action_g2.py]`):

$$\underbrace{\frac{M_{\text{scal}}^2}{\bar{M}_{\text{Pl}}^2} = c_3^{r-1} = c_3^7}_{[\mathbf{E}]}, \quad \underbrace{A_s = \frac{N_\star^2}{24\pi^2} \frac{M_{\text{scal}}^2}{\bar{M}_{\text{Pl}}^2} = \frac{N_\star^2 c_3^7}{24\pi^2} = 2.17 \times 10^{-9}}_{[\mathbf{C}]/[\mathbf{O}] \text{ (if } M_{\text{scal}} = M_{\text{Star}})}. \quad (92)$$

Theorem 7 — the $D_5 \times A_3$ lift of the E_8 atlas

Since $A_3 = \mathfrak{su}_4 \cong \mathfrak{so}_6 = D_3$, one has the standard chain $E_8 \supset \mathfrak{so}_{16} \supset \mathfrak{so}_{10} \oplus \mathfrak{so}_6 = D_5 \times A_3$, with the exact branching (verified dimensions and root/weight counts):

$$\boxed{\mathfrak{e}_8 = (45, 1) \oplus (1, 15) \oplus (10, 6) \oplus (16, 4) \oplus (\overline{16}, \bar{4})}, \quad (93)$$

$$\underbrace{45+15+60+64+64}_{=248} = \dim E_8, \quad \underbrace{40+12+60+64+64}_{=240} = |R(E_8)|.$$

The adjoint **120** of $\mathfrak{so}_{16}$ is $(45, 1) \oplus (1, 15) \oplus (10, 6)$ and its spinor **128** is $(16, 4) \oplus (\overline{16}, \bar{4})$. The TFPT dictionary makes each summand a known object:

piece	value	TFPT reading
(45, 1)	$45 = 40 + 5$	D_5 carrier adjoint: Clebsch edges $40 = R(D_5) $ plus carrier rank $g_{\text{car}} = 5$
(1, 15)	$15 = 12 + 3$	A_3 family+Higgs adjoint: $\dim \mathfrak{g}_{\text{SM}} = 12$ plus family rank $N_{\text{fam}} = 3$
(10, 6)	$10 \cdot 6$	Λ decuple $\mathcal{A}_\Lambda = \binom{5}{2} = 10$ times Z_6 transport $\binom{4}{2} = 6$
$(16, 4) \oplus (\overline{16}, \bar{4})$	$2 \cdot 64$	matter half-spinor S^+ times the four corners μ_4 , plus conjugate sheet

Proof route (replaces the old blind 240-metrization). *Stage 1* (D_5 , $[\mathbf{E}]$): C^+ is the D_5 half-spin orbit, $R_4 = \text{Clebsch}$, $\text{Aut} = W(D_5)$, $|E(R_4)| = 40 = |R(D_5)|$ — Theorem 2. *Stage 2* (A_3 , $[\mathbf{E}]$): the carrier geometry fixes $X_f^\circ = \mathbb{P}^1 \setminus \mu_4$ with $H_1(X_f^\circ, \mathbb{Z}) = \mathbb{Z}^3$ and a rigid $D_4 \subset PGL_2$ generated by $z \mapsto iz$ and $z \mapsto 1/z$. The logarithmic forms $\omega_i = dz/(z - p_i)$ regular at infinity span H^1 by their differences, dual under the residue pairing to the puncture loops γ_i with $\text{Res}(\omega_i, \gamma_j) = \delta_{ij}$; hence the canonical (residue) metric on the puncture classes is I_4 , and the twelve puncture-difference cycles $\gamma_i - \gamma_j$ are exactly the A_3 roots:

$$\boxed{\{\gamma_i - \gamma_j\}_{i \neq j} : \# = 12 = |R(A_3)| = \dim \mathfrak{g}_{\text{SM}}, \quad \text{rank} = 3 = N_{\text{fam}}, \quad \text{Gram} = \text{Cartan}(A_3), \quad \det = 4} \quad [\mathbf{E}], \quad (94)$$

with $\dim A_3 = 15 = \dim S^+ - 1$, and the A_3 positive roots equal the exponent sum $1+2+3 = 6 = Z_6 = |E(K_4)|$. *The corner rotation realises the A_3 Coxeter element.* The carrier generator $z \mapsto iz$ permutes μ_4 as the 4-cycle $(p_1 p_2 p_3 p_4) \in S_4 = W(A_3)$ and acts on $H^1 = \mathbb{C}^3$ as an isometry of the Cartan form with

$$\boxed{\text{ord} = 4 = h(A_3) = |\mu_4|, \quad \text{spec} = \{i, -1, -i\}, \quad \chi(t) = \frac{t^4 - 1}{t - 1} = \Phi_4(t)\Phi_2(t)} \quad [\mathbf{E}], \quad (95)$$

i.e. the eigenvalues are the A_3 exponents $\{1, 2, 3\}$ as fourth roots of unity — the exact analogue of the E_8 Coxeter element reading Φ_{30} . The full $D_4 = \langle z \mapsto iz, z \mapsto 1/z \rangle$ embeds in $W(A_3) = S_4$ (order 8, index 3) by isometries. *Stage 3 (standard, [E]):* assemble E_8 from the $D_5 \times A_3$ branching — no 240-map required.

The μ_4 glue theorem (why the four corners appear everywhere). The two factors have equal discriminant groups, and E_8 is unimodular:

$$\boxed{\text{disc}(D_5) = \text{disc}(A_3) = \mathbb{Z}_4, \quad [E_8 : D_5 \oplus A_3] = \sqrt{\frac{\det D_5 \cdot \det A_3}{\det E_8}} = \sqrt{\frac{4 \cdot 4}{1}} = 4 = |\mu_4|} \quad [\text{E}]. \quad (96)$$

So $E_8 = (D_5 \oplus A_3) + \mu_4\text{-glue}$: the four family corners $\mu_4 = \mathbb{Z}_4$ are precisely the glue code that fuses the carrier and the family into the unimodular E_8 lattice. The glue pairs the D_5 spinor class ($= S^+$, the family) with the A_3 class — exactly the $(16, 4) \oplus (\overline{16}, \overline{4})$ spinor block. Hence μ_4 plays three roles at once: family corners, A_3 origin, and E_8 glue.

Discriminant-norm glue lemma (and the TFPT 5/4, 3/4 identification). The two glue generators carry the discriminant quadratic form $q \in \mathbb{Q}/2\mathbb{Z}$:

$$\boxed{\begin{aligned} q(D_5) &= \frac{n}{4}|_{n=5} = \frac{5}{4} \text{ (spinor class)}, & q(A_3) &= \frac{n}{n+1}|_{n=3} = \frac{3}{4} \text{ (corner class)}, \\ q(D_5) + q(A_3) &= 2 \equiv 0 \pmod{2}, & q(D_5) - q(A_3) &= \frac{1}{2} \end{aligned}} \quad [\text{E}]. \quad (97)$$

The diagonal $\langle (s, g) \rangle \cong \mathbb{Z}_4$ is isotropic ($k^2 \cdot 2 \equiv 0$ for all k), so the glue closes to an *even unimodular* lattice — this is E_8 .

The two glue norms are just $(g_{\text{car}}, N_{\text{fam}})$ over the glue index. Because $\text{rank } D_5 = g_{\text{car}} = 5$ (so $q(D_5) = \frac{n}{4}|_{n=5} = g_{\text{car}}/|\mu_4|$) and $\text{rank } A_3 = N_{\text{fam}} = 3$ (so $q(A_3) = \frac{n}{n+1}|_{n=3} = N_{\text{fam}}/|\mu_4|$, with $n+1 = 4 = |\mu_4|$),

$$\boxed{q(D_5) = \frac{g_{\text{car}}}{|\mu_4|} = \frac{5}{4}, \quad q(A_3) = \frac{N_{\text{fam}}}{|\mu_4|} = \frac{3}{4}}$$

and *all four arithmetic operations* on the pair are then forced compiler atoms:

$$\begin{aligned} \underbrace{q(D_5) + q(A_3) = \frac{g_{\text{car}} + N_{\text{fam}}}{|\mu_4|} = 2}_{E_8 \text{ root norm}}, & \quad \underbrace{q(D_5) - q(A_3) = \frac{|\mathbb{Z}_2|}{|\mu_4|} = \frac{1}{2}}_{\delta \text{ (lepton transport)}}, \\ \underbrace{q(D_5)q(A_3) = \frac{g_{\text{car}}N_{\text{fam}}}{|\mu_4|^2} = \frac{15}{16}}_{\dim \mathfrak{su}(4) / \dim S^+}, & \quad \underbrace{\frac{q(D_5)}{q(A_3)} = \frac{g_{\text{car}}}{N_{\text{fam}}} = \frac{5}{3}}. \end{aligned}$$

So the distinguished lepton transport value $\delta = \frac{1}{2} = (g_{\text{car}} - N_{\text{fam}})/|\mu_4| = |\mathbb{Z}_2|/|\mu_4|$ is the carrier-minus-family count over the glue index — a fully atomic origin, not a chosen rational (it also equals the harmonic-frame holonomy diagonal modulus, [verification/v41_leg_assignment.py]); and (sum, ratio) invert uniquely back to $(\frac{5}{4}, \frac{3}{4})$. [verification/v51_boundary_half_step.py] The sum $\frac{5}{4} + \frac{3}{4} = 2$ is literally the E_8 spinor-root norm $\frac{1}{2}(\pm)^8$ split across $D_5 \times D_3$: $5 \cdot \frac{1}{4} + 3 \cdot \frac{1}{4} = 2$. Strikingly, both norms are pre-existing TFPT constants:

$$\boxed{\frac{\delta_2}{\delta_{\text{top}}^2} = \frac{2880 c_3^8}{(48 c_3^4)^2} = \frac{5}{4} = q(D_5), \quad \xi_{\text{tree}} = \frac{\dim \mathfrak{g}_{\text{SM}}}{\dim S^+} = \frac{12}{16} = \frac{3}{4} = q(A_3)} \quad [\text{E}], \quad (98)$$

so $\delta_2/\delta_{\text{top}}^2 + \xi_{\text{tree}} = 2$: the two-loop/defect ratio is the D_5 spinor norm and the gravitational tree normalization is the A_3 corner norm, and their sum is the E_8 root norm.

The whole integer skeleton from one pair $(g_{\text{car}}, N_{\text{fam}}) = (5, 3)$ **[E]**
`( [verification/v53_compiler_core.py] )`

Every load-bearing integer of the compiler flows from the single boundary pair $(g_{\text{car}}, N_{\text{fam}}) = (5, 3)$ by sum, difference and squares:

$$\text{rank } E_8 = g_{\text{car}} + N_{\text{fam}} = 8, \quad |\mathbb{Z}_2| = g_{\text{car}} - N_{\text{fam}} = 2, \quad |\mu_4| = \frac{g_{\text{car}} + N_{\text{fam}}}{2} = 4,$$

and $(N_{\text{fam}}, |\mu_4|, g_{\text{car}}) = (3, 4, 5)$ is the Pythagorean triple. This is not decorative: the grand-mass-volume exponent $\Delta_Y = g_{\text{car}}^2 = 25$ ($\det M_{\text{SM}} \sim (\varphi_0^{\text{ret}})^{25}$; document 3) splits over the sectors as a *difference of squares*,

$$\Delta_Y = g_{\text{car}}^2 = \underbrace{N_{\text{fam}}^2}_{\text{down}=9} + \underbrace{(g_{\text{car}} - N_{\text{fam}})(g_{\text{car}} + N_{\text{fam}})}_{=|\mathbb{Z}_2| \cdot \text{rank } E_8} = 9 + 16 = \dim S^+$$

i.e. the down sector carries N_{fam}^2 and up+lepton carries $|\mathbb{Z}_2| \cdot \text{rank } E_8 = \dim S^+$ (the K -row sums $(6, 9, 10)$). **The anchor is the Dirac square root of this skeleton:** $a = (1, 1, 2)$ is the *unique* 3-multiset whose elementary symmetric functions are $(|\mu_4|, g_{\text{car}}, |\mathbb{Z}_2|) = (4, 5, 2)$, equivalently its characteristic polynomial is $\chi_a(t) = (t-1)^2(t-2) = t^3 - |\mu_4|t^2 + g_{\text{car}}t - |\mathbb{Z}_2|$ — the compiler atoms *are* its char-poly coefficients. So the theory's only transcendental is π (through $c_3 = \frac{1}{8\pi}$, $8 = \text{rank } E_8$, and the seed $\frac{1}{6\pi}$, $6 = |R^+(A_3)|$); everything else is this $(5, 3)$ -generated integer skeleton.

Boundary Carrier Selection Theorem (Theorem A) [E]/[C]

E_8 is not assumed: it is *selected* as the unique even-unimodular audit hull of the admissible boundary-carrier data. **Boundary side:** $\mu_4 = \{1, i, -1, -i\}$ gives $X_f = \mathbb{P}^1 \setminus \mu_4$ with $H_1 = \mathbb{Z}^3$, hence A_3 and $N_{\text{fam}} = 3$. **Carrier side:** a minimal D -type carrier with a 16-dim half-spinor forces D_5 (only $n = 5$ gives $2^{n-1} = 16$; $S^+ = \Lambda^{\text{even}} \mathbb{C}^5 = 1+10+5$). **Glue:** $\text{disc}(D_5) = \text{disc}(A_3) = \mathbb{Z}_4$ (isomorphic, cyclic, anti-isometric), $q(D_5) + q(A_3) = 2$ (the E_8 root norm), glue index $4 = |\mu_4|$, and E_8 is the unique even unimodular rank-8 lattice. **Exclusion:** the four conditions

$$(i) \dim S^+ = 16, \quad (ii) N_{\text{fam}} = 3, \quad (iii) \text{cyclic } \mathbb{Z}_4 \text{ glue}, \quad (iv) q_D + q_A = 2$$

are met *only* by $D_5 \oplus A_3$; D_8 (non-cyclic $\mathbb{Z}_2 \times \mathbb{Z}_2$, no family geometry), $E_7 \oplus A_1$ (1 family, no 16-spinor), $E_6 \oplus A_2$ (2 families, $\mathbb{Z}_3$), $A_4 \oplus A_4$ (no D -spinor, $\mathbb{Z}_5$) and A_8 (single factor) each fail at least one. **Typing:** the lattice/classification facts are [E]; the physical selection (reflection-positive seam $c_3 = 1/8\pi$ + minimal chiral carrier $\Rightarrow$ exactly these inputs) is the [C]/[O] selection axiom (P1 analytic interface + P2 carrier interface), not a zero-input derivation. `[verification/v47_selection_theorem.py]`

Group-level closure (μ_4 as the common centre). The lattice glue lifts to a clean group statement, because both factors share the *same* centre and μ_4 is the diagonal:

$$Z(\text{Spin}(10)) = \mathbb{Z}_4, \quad Z(SU(4)) = \mathbb{Z}_4, \quad |\mu_4| = 4, \quad G_{\text{closure}} = \frac{\text{Spin}(10) \times SU(4)}{\Delta \mathbb{Z}_4} \subset E_8 \quad \text{[E]}, \quad (99)$$

with $\text{rank} = 5 + 3 = 8 = \text{rank } E_8$. Every adjoint block is diagonal- $\mathbb{Z}_4$ neutral (generator $\omega = i$; charges mod 4): $(45, 1), (1, 15) : 0$; $(10, 6) : 2+2=0$; $(16, 4) : 1+3=0$; $(\overline{16}, \overline{4}) : 3+1=0$, so $45+15+60+64+64 = 248$ are genuine reps of the quotient. This promotes μ_4 from a lattice glue index to the *common centre quotient* — the physically cleanest language for the gluing: the D_5 half-spinor 16 pairs with the $SU(4)$ fundamental 4 so the spinor charges ± 1 cancel the fundamental charges ∓ 1 .

Root-length completion. Every E_8 block reaches norm 2:

block	norm split	reading
$(45, 1), (1, 15)$	2, 2	D_5, A_3 root adjoints
$(10, 6)$	$1 + 1 = 2$	$SO(10)$ vector $\oplus$ $SO(6)$ vector
$(16, 4) \oplus (\overline{16}, \overline{4})$	$\frac{5}{4} + \frac{3}{4} = 2$	D_5 spinor norm + A_3 corner norm

The old question “can the 240 flip atlas be metrized to E_8 ?” is now the sharp question “are the TFPT D_5 and A_3 metrics in standard normalization?” — and the $5/4, 3/4$ match says yes.

Explicit construction (the lift is now built, not counted). Embedding $D_5 \oplus A_3 = D_5 \oplus D_3 \subset D_8$ and adjoining the even spinor coset (the μ_4 glue), the 240 roots are produced explicitly and verified:

$$\{\pm e_i \pm e_j\}_{i < j}^8 (112) \cup \{\frac{1}{2}(\pm)^8 : \text{even}\} (128) = 240 \text{ roots, all norm } 2, \text{ Cartan } \det = 1 \quad [\mathbf{E}]. \quad (100)$$

Splitting each root into its D_5 (first five) and A_3 (last three) parts gives exactly the branching blocks — $(2, 0):40$, $(0, 2):12$, $(1, 1):60$, and $(\frac{5}{4}, \frac{3}{4}):128$ — and the 128 spinor roots realise $\frac{5}{4} + \frac{3}{4} = 2$ on the nose. All 240 vectors are integer combinations of one set of 8 simple roots, so the assembled lattice *is* E_8 . The blind 240-metrization is therefore eliminated: E_8 is explicitly $(D_5 \oplus A_3) + \mu_4$ -glue.

Remark (the metric, computed). This reframes E_8 from origin to closure: E_8 is the closure algebra of the D_5 carrier and the A_3 family geometry, glued by μ_4 . The root system, Cartan form, $W(A_3) = S_4$ and the corner-rotation = A_3 Coxeter element are exact (residue pairing + explicit carrier symmetry). The transcendental geometric metric $G_{ij} = -\log |p_i - p_j|$ on μ_4 has been computed: it is only D_4 -symmetric, with a regularisation-independent anisotropy $-\log 2$ on the root space, hence *not* proportional to $\text{Cartan}(A_3)$. It need not be: the E_8 -relevant A_3 is the canonical *residue* pairing, and the geometric metric is a D_4 -equivariant deformation that preserves the root system, Weyl group and Coxeter element. So the lattice data is exact $[\mathbf{E}]$; the analytic metric is a *characterised* deformation, not an open gap.

Uniform Coxeter law. All three factors obey $|R| = h \cdot \text{rank}$, and the two TFPT Coxeter numbers are themselves carrier data:

$$\begin{aligned} h(D_5) &= 8 = r(E_8) = \varphi(30), & h(A_3) &= 4 = |\mu_4|, \\ h(D_5) h(A_3) &= 2^{g_{\text{car}}} = 32, & 2^{g_{\text{car}}} - 2 &= 30 = h(E_8) \end{aligned} \quad [\mathbf{E}], \quad (101)$$

so the Coxeter closure of Theorem 3 reappears as $h(E_8) = h(D_5)h(A_3) - 2$, and $h(D_5) + h(A_3) = 12 = \dim \mathfrak{g}_{\text{SM}} = |R(A_3)|$.

Theorem 8 — the $(1, 1, 2)$ anchor microcode and the φ_0 -ladder

The parabolic flavor bundle $E_\bullet = \mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$ on $\mathbb{P}^1 \setminus \mu_4$ has splitting type $a = (1, 1, 2)$, whose power sums $p_n(a) = 2 + 2^n$ generate the whole compiler budget:

The anchor power compiler.

$$(p_1, \dots, p_5) = (4, 6, 10, 18, 34) = (|\mu_4|, |R^+(A_3)|, \mathcal{A}_\Lambda, N_{\text{fam}}|R^+(A_3)|, \mathbb{Z}_6 \text{ lift}),$$

$$q(A_3) = \frac{p_2}{2p_1} = \frac{3}{4}, \quad q(D_5) = \frac{p_3}{2p_1} = \frac{5}{4}, \quad \sum L = p_1 p_3 = 40,$$

$$\Omega_{\text{adm}} = 2p_1 p_2 = 48, \quad D_{\text{start}} = p_2 p_3 = 60, \quad |R(E_8)| = 4p_2 p_3 = 240.$$

The elementary symmetric data $\chi_a(t) = (t-1)^2(t-2)$ give $(e_1, e_2, e_3) = (4, 5, 2) = (|\mu_4|, g_{\text{car}}, |\mathbb{Z}_2|)$, so the anchor reconstructs the carrier rank, and $10b_1 = e_1^2 + e_2^2 = 4^2 + 5^2 = 41$. The seed itself is the two-term anchor form $\varphi_0^{\text{ret}} = \frac{2p_1}{p_2} c_3 + 2p_1 p_2 c_3^4 = \frac{4}{3} c_3 + 48c_3^4$.

The φ_0 -ladder. With $\lambda_Y = \sqrt{\varphi_0^{\text{ret}}(1 - \varphi_0^{\text{ret}})}$, every fermion mass is the single closed form $\hat{m}_{f,j} = \frac{v_{\text{geo}}}{\sqrt{2}} \lambda_Y^{L_{f,j}} \Lambda_{f,j}$ with $\lambda_Y^L \Lambda = \pi c (\varphi_0^{\text{ret}})^k$, so all nine holonomies are resolvent outputs and the masses form the φ_0^{ret} -order ladder $t(0) > \{b, c, \tau\}(2) > \{s, \mu\}(3) > \{u, d\}(4) > e(5)$.

Secondary E_8 atlas. Beyond the main $D_5 \times A_3$ split, the residue matrix appears in the $E_6 \times A_2$ branching, $248 = \|R\|_F^2 + \det R + 2(\mathbf{1}^\top R a) N_{\text{fam}} = 78 + 8 + 2 \cdot 27 \cdot 3$, and the scalaron exponent in $E_7 \times A_1$, $112 = 7 \cdot 16$. The seven E_8 slices are charted in the companion note `e8_secondary_branching_atlas` as a falsifiable audit raster.

The $E_6 \times A_2$ block is the $\tau=\omega$ face ([`verification/v407_dn_pairings_omega.py`], **SEAM.EQUIV.02**). A_2 is the Eisenstein lattice of the family CM point $\tau=\omega$ (Cartan $\det = 3 = N_{\text{fam}}$), and the (d, n) selector's three pairings $\{n \cdot \mathbf{1}, n \cdot a, n \cdot \sigma\} = \{2, 8, 121\}$ are exactly its family-slice atoms $\{|Z_2|, \dim A_2 = \det R = \text{rank } E_8, \|Pl(K)\|_1^2\}$, so deriving (d, n) (the R_4' residue, **v139**) folds into the dual keystone (**v405**), leaving only “why $11 = \|Pl(K)\|_1$ ”.

Master statement and status

Five carrier slots give D_5 , four family corners give A_3 , and $D_5 \times A_3$ gives the E_8 atlas.

Equivalently: $\mathcal{A}_\Lambda = \frac{h}{N_{\text{fam}}} = 10 = \binom{5}{2}$, $h = 30 = 2 \cdot 3 \cdot 5$, $(5, 3, 2) : \Sigma = 10, \Pi = 30$; Λ , $\det M_\ell$, the E_8 Coxeter structure and the action ladder read the same five-slot pair-decuple. The Standard model, flavor, Λ , inflation and E_8 are projections of the even carrier code C^+ ($= D_5$ half-spinor) on five slots together with the four-point family A_3 .

Statement	Status	Note
Even code C^+ , $\dim S^+ = 16$, dual-root Y	[E]	Theorem 1
Exterior–Action duality $(1, 10, 5) \leftrightarrow (1, 5, 10)$	[E]	scheme eigenvalues verified
$R_4 = \text{Clebsch } (16, 5, 0, 2)$, $R_2 = (16, 10, 6, 6)$; $\text{Aut}_{\text{aff}} = W(D_5) = 1920$	[E]	carrier = D_5 half-spinor
$2^g - 2 = g(g+1) \Leftrightarrow g = 5$	[E]	Coxeter closure (unique)
$h=30=2 \cdot 3 \cdot 5$, $r=\varphi(30)=8$, $\chi_C = \Phi_{30}$	[E]/[C]	cyclotomic compiler
$ R^+ = \binom{16}{2} = \text{Tr } X^2 = 4h = 120$	[E]	positive-edge moment
$40, 60, 120, 240 = 10 \cdot (4, 6, 12, 24)$; $41=1+4 \cdot 10$	[E]	$K_5 \times K_4$ product table
$10b_1=5r+1$, u totient, $M_{\text{scal}}^2 = c_3^{r-1}$, $\omega_b = \lambda_C/10$	[E]/[C]	downstream readouts
$A_s = N_*^2 c_3^7 / 24\pi^2$	[C]/[O]	if $M_{\text{scal}} = M_{\text{Star}}$
$\mathfrak{e}_8 = (45, 1) \oplus (1, 15) \oplus (10, 6) \oplus (16, 4) \oplus (\overline{16}, \overline{4})$	[E]/[C]	$D_5 \times A_3$ lift
240 roots built from $D_5 \oplus A_3 + \mu_4$ glue, $\det 1$	[E]	explicit E_8 construction
flavor $\sum L=40= R(D_5) $, entries on $C_6= R^+(A_3) $	[E]	transport = $D_5 \times A_3$
A_3 from $\mathbb{P}^1 \setminus \mu_4$: residue pairing = Cartan, $\det 4$	[E]	Stage 2 (Theorem 7)
corner $z \mapsto iz = A_3$ Coxeter el.: $\{i, -1, -i\}$, $\Phi_4 \Phi_2$	[E]	order 4 = $h(A_3) = \mu_4 $
$\text{disc}(D_5) = \text{disc}(A_3) = \mathbb{Z}_4$, glue = μ_4	[E]	$E_8 = (D_5 \oplus A_3) + \mu_4$
$q(D_5) = \frac{5}{4} = \delta_2 / \delta_{\text{top}}^2$, $q(A_3) = \frac{3}{4} = \xi_{\text{tree}}$, sum 2	[E]	discriminant-norm glue
$120 = \dim \mathfrak{so}_{16}$; $\Omega_{\text{adm}} = 40 + 1 + 7$	[E]	adjoint half; deficit ladder
$h(D_5) = 8 = r(E_8)$, $h(A_3) = 4 = \mu_4 $, $ R = h \cdot r$	[E]	uniform Coxeter law
$\text{Exp}(E_8)$ as 41/48 budget deficits	[O]	exponent ledger (fingerprint)
geometric metric D_4 -deform (anisotropy $-\log 2$), residue = A_3	[E]	metric computed/characterised

Scope. This note compresses; it does not re-derive the upstream boundary kernel (the P1 hardening chain) or the carrier rigidity theorem (Lean), and it makes no claim of a final theory. The arithmetic and the $D_5/\text{Clebsch}/W(D_5)$, $A_3/\text{residue-Cartan}/\text{Coxeter}$, and μ_4 -glue identifications are exact, and the E_8 lattice is now built explicitly from $D_5 \oplus A_3$ plus the μ_4 glue (240 roots, $\det 1$). The only residual [C] is the analytic identification of the carrier geometry's L^2 cusp Hodge metric with the canonical residue metric up to scale, which does not affect the explicit root/lattice/Weyl construction. Honest non-fits (Koide $0.6645 \neq 2/3$, η_B , m_p/m_e) stay outside the compiler.

Appendix: machine-checkable $E_8 = (D_5 \oplus A_3) + \mu_4$ construction

The lift is reproducible by a short script (all entries below verified exactly).

Cartan matrices and discriminant groups.

$$C(D_5) = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & 0 \\ 0 & 0 & -1 & 0 & 2 \end{pmatrix}, \quad C(A_3) = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}; \quad \det C(D_5) = \det C(A_3) = 4.$$

Smith normal forms give $\text{disc}(D_5) = \text{disc}(A_3) = \mathbb{Z}_4$; the glue index is $[E_8 : D_5 \oplus A_3] = \sqrt{4 \cdot 4/1} = 4 = |\mu_4|$, with discriminant-form norms $q(D_5) = \frac{5}{4}$, $q(A_3) = \frac{3}{4}$, $q(D_5) + q(A_3) = 2 \equiv 0 \pmod{2}$ (even-glue). Writing $\mu := |\mu_4| = 4$ the norms are $q(D_5) = \frac{\mu+1}{\mu}$, $q(A_3) = \frac{\mu-1}{\mu}$, so $q(D_5)q(A_3) = 1 - \frac{1}{\mu^2} = \frac{15}{16}$ and

$$\boxed{\dim S^+(1 - q(D_5)q(A_3)) = 16 \cdot \frac{1}{16} = 1} \quad [\mathbf{E}], \quad (102)$$

i.e. the glue product fills all $15 = \dim S^+ - 1$ non-trivial half-spin states and the defect is exactly the singleton (the ν^c /Higgs closure). And E_8 has a tidy μ_4 -cardinal form, $\dim E_8 = \binom{|\mu_4|^2}{2} + 2|\mu_4|^3 = \binom{16}{2} + 2 \cdot 64 = 120 + 128 = 248$ [C].

Explicit roots. Split $\mathbb{R}^8 = \mathbb{R}_{(D_5)}^5 \oplus \mathbb{R}_{(A_3)}^3$. The 240 roots are

$$\{\pm e_i \pm e_j : 1 \leq i < j \leq 8\} \text{ (112)} \cup \{\tfrac{1}{2}(\pm 1)^8 : \text{even \# of } -\} \text{ (128)},$$

all of norm 2. Splitting each by $(\|\cdot\|_{D_5}^2, \|\cdot\|_{A_3}^2)$ gives the branching blocks

$$(2, 0): 40 [D_5], \quad (0, 2): 12 [A_3], \quad (1, 1): 60 [(10, 6)], \quad (\tfrac{5}{4}, \tfrac{3}{4}): 128 [(16, 4) \oplus (\overline{16}, \overline{4})],$$

so $\mathfrak{e}_8 = (45, 1) \oplus (1, 15) \oplus (10, 6) \oplus (16, 4) \oplus (\overline{16}, \overline{4})$, $248 = 45 + 15 + 60 + 64 + 64$, $240 = 40 + 12 + 60 + 64 + 64$.

Simple-root Gram (unimodular check). With the Bourbaki simple roots $\alpha_1 = \frac{1}{2}(e_1 - e_2 - \dots - e_7 + e_8)$, $\alpha_2 = e_1 + e_2$, and $\alpha_k = e_{k-1} - e_{k-2}$ for $k = 3, \dots, 8$ (so $\alpha_3 = e_2 - e_1$, $\alpha_4 = e_3 - e_2$, $\dots$, $\alpha_8 = e_7 - e_6$), the Gram matrix $G_{ij} = \langle \alpha_i, \alpha_j \rangle$ is the E_8 Cartan matrix with $\det G = 1$ and all $G_{ii} = 2$; every one of the 240 roots is an integer combination of the α_i . Hence the assembled lattice is E_8 , not merely a 240-count. [E] (The shifted index $e_{k-2} - e_{k-3}$ would duplicate α_3 and make G singular — use $e_{k-1} - e_{k-2}$.)

Part III

Appendix: ablation of the EM fixed point

The single sharpest reviewer test is: *was the closure $F_{U(1)}(\alpha) = 0$ engineered so that α comes out?* The honest answer is an ablation — vary each input and see which actually move the root. Three one-parameter scans (all others held at their compiler values) separate the *load-bearing* integers from the *fine* last-digit selector.

EM fixed point: only $M = 41$ and $N = 8$ land on α^{-1}




Figure 7: EM-closure ablation (data plot, `verification/make_figures.py`). (a) α_*^{-1} vs the integer budget M : only $M = 41$ lands on CODATA-2022. (b) α_*^{-1} vs the seam denominator N in $c_3 = 1/(N\pi)$: only $N = 8$ lands. Both integers are fixed elsewhere (flavor budget; rank/ $h(D_5)$), so the equation is not freely tunable to 137.036.

(A1) The transport budget $M = \sum L + N_\Phi$ (exponent held at $-\frac{5}{4}$):

M	38	39	40	41	42	43	44
α_*^{-1}	140.530	139.326	138.162	137.0359992	135.946	134.890	133.866

Each unit of M shifts α_*^{-1} by ≈ 1.09 , so *only* $M = 41$ lands on 137.036. This is a hard lock — and $41 = 10b_1 = \sum L + N_\Phi = 40 + 1$ is fixed by the compiler (flavor budget 40 + one Higgs). **Load-bearing.**

(A2) The seam “8” in $c_3 = 1/(N\pi)$ (here $\delta_{\text{top}} = 48c_3^4$ tracks c_3):

N	6	7	8	9	10
α_*^{-1}	76.97	104.85	137.036	173.53	214.34

Only $N = 8$ — i.e. $c_3 = 1/(8\pi) = 1/(\text{rank } E_8 \cdot \pi)$ — reproduces 137.036. The “8” is the same one fixed by the bootstrap ($h(D_5) = \text{rank } E_8 = \varphi(30)$) and the Gauss–Bonnet hardening. **Load-bearing.**

(A3) The carrier discriminant exponent (at $M = 41$, $c_3 = 1/(8\pi)$):

exponent	$-\frac{1}{4}$	$-\frac{3}{4}$	$-\frac{5}{4}$	$-\frac{7}{4}$	$-\frac{9}{4}$
α_*^{-1}	137.035995	137.035997	137.0359992	137.036001	137.036003

The exponent moves only the *seventh* digit; all candidates agree to six. Here $-\frac{5}{4} = q(D_5)$ is selected, but this is a *fine* selector, not a hard lock. **Fine (last-digit).**

Ablation verdict — honest

The EM fixed point is a **hard test of two integers** — the budget $M = 41$ and the seam denominator $8 = \text{rank } E_8$, each decisive (a single-unit change misses by ~ 1 in $\alpha_\star^{-1}$) — and a **fine test of one norm**, the carrier discriminant exponent $-\frac{5}{4} = q(D_5)$ (seventh digit). The equation is therefore *not* freely tunable to 137.036: the two integers that hit it are exactly the compiler/bootstrap numbers, fixed elsewhere. What it does *not* do is pin the carrier norm strongly — that is honest, and it is the right level of claim. [E]

Appendix: computational verification

The load-bearing identities of this document (marked [E], [E], [E]) are re-derived from $\{c_3, g_{\text{car}}\}$ alone and machine-checked by the suite in the repository folder `verification/`; the inline tags `[verification/...]` point to the exact script. Relevant here:

- `v1_e8_glue.py` — the μ_4 glue ($q(D_5)+q(A_3)=2$, $\text{disc}=\mathbb{Z}_4$, 240, 248).
- `v2_carrier_pascal.py` — $g_{\text{car}}=5$, $16=1+5+10$, $N_{\text{fam}}, \Omega_{\text{adm}}, b_1$.
- `v3_em_alpha.py` — the EM fixed point $\alpha_\star^{-1}=137.0359992168$ (existence/uniqueness, ablation).
- `v4_flavor_matrix.py` — R with $\det=8$, minors $\{2, 3, 5\}$, $\sum L=40$.
- `v7_gravity_cosmo.py` — scalaron c_3^7 , Λ/A_s readouts.

Run `python verification/run_all.py` (needs `mpmath`, `numpy`, `sympy`); it exits 0 iff all checks pass. Full map: `verification/README.md`.

External works built upon

The TFPT discrete core (the two axioms, the E_8 glue, the integer skeleton) is self-contained and machine-checked. Its *bridges* to continuum/AQFT physics, to gravity and to the lattice/singularity picture build on established mathematics, listed here by topic; the body refers to these by author name.

Modular theory and algebraic quantum field theory.

- M. Takesaki, *Tomita's Theory of Modular Hilbert Algebras and its Applications*, Lecture Notes in Math. **128**, Springer (1970).
- J. J. Bisognano, E. H. Wichmann, *On the duality condition for a Hermitian scalar field / for quantum fields*, J. Math. Phys. **16** (1975) 985–1007; **17** (1976) 303–321.
- K. Osterwalder, R. Schrader, *Axioms for Euclidean Green's functions* I, II, Comm. Math. Phys. **31** (1973) 83–112; **42** (1975) 281–305.
- S. Doplicher, R. Haag, J. E. Roberts, *Local observables and particle statistics* I, II, Comm. Math. Phys. **23** (1971) 199–230; **35** (1974) 49–85.
- R. Haag, *Quantum field theories with composite particles and asymptotic conditions*, Phys. Rev. **112** (1958) 669–673; D. Ruelle, *On the asymptotic condition in quantum field theory*, Helv. Phys. Acta **35** (1962) 147–163.
- R. Longo, *Index of subfactors and statistics of quantum fields* I, II, Comm. Math. Phys. **126** (1989) 217–247; **130** (1990) 285–309.
- Y. Kawahigashi, R. Longo, M. Müger, *Multi-interval subfactors and modularity of representations in conformal field theory*, Comm. Math. Phys. **219** (2001) 631–669.

Conformal nets, vertex operator algebras and lattice constructions.

- I. B. Frenkel, V. G. Kac, *Basic representations of affine Lie algebras and dual resonance models*, Invent. Math. **62** (1980) 23–66; G. Segal, *Unitary representations of some infinite-dimensional groups*, Comm. Math. Phys. **80** (1981) 301–342.
- T. J. Osborne, A. Stottmeister, *Conformal field theory from lattice fermions*, Comm. Math. Phys. **398** (2023) 219–289, [arXiv:2107.13834](https://arxiv.org/abs/2107.13834), [doi:10.1007/s00220-022-04521-8](https://doi.org/10.1007/s00220-022-04521-8). (This is the seam scaling-limit theorem the TFPT continuum leg cites — the Koo–Saleur $\rightarrow$ Virasoro convergence, $c = \frac{1}{2}$, and the smeared WZW-current Theorem 5.1. The internal TFPT code-token “MMST” and the legacy claim-ids `SEAM.MMST.*` / filenames `v*_mmst_*` denote *this* Osborne–Stottmeister theorem.)

- V. Morinelli, G. Morsella, A. Stottmeister, Y. Tanimoto, *Scaling limits of lattice quantum fields by wavelets*, Comm. Math. Phys. **387** (2021) 299–360, [arXiv:2010.11121](#), [doi:10.1007/s00220-021-04152-5](#); and *Operator-algebraic renormalization and wavelets*, Phys. Rev. Lett. **127** (2021) 230601. (The operator-algebraic (wavelet/Wilson–Kadanoff) renormalization *framework* the Osborne–Stottmeister theorem above builds on.)
- M. S. Adamo, Y. Moriwaki, Y. Tanimoto, *Osterwalder–Schrader axioms for unitary full vertex operator algebras*, [arXiv:2407.18222](#) (2024).
- M. S. Adamo, L. Giorgetti, Y. Tanimoto, *Rational and non-rational two-dimensional conformal field theories arising from lattices*, [arXiv:2506.01008](#) (2025); *Representations of conformal nets associated with infinite-dimensional groups*, [arXiv:2508.07109](#) (2025).
- P. Naaijken, D. Penneys, B. Wallick, *(commuting-projector reflection positivity / LTO-RP)*, [arXiv:2605.10693](#).

Lattices and quadratic forms.

- J. H. Conway, N. J. A. Sloane, *Sphere Packings, Lattices and Groups*, 3rd ed., Grundlehren der math. Wiss. **290**, Springer (1999).
- C. L. Siegel, *Über die analytische Theorie der quadratischen Formen* (the Minkowski–Siegel mass formula), Ann. of Math. **36** (1935) 527–606.

Singularities and geometry.

- E. Brieskorn, *Beispiele zur Differentialtopologie von Singularitäten*, Invent. Math. **2** (1966) 1–14.
- J. Milnor, *Singular Points of Complex Hypersurfaces*, Ann. of Math. Studies **61**, Princeton Univ. Press (1968).

Spectral geometry, gravity and thermal time.

- A. H. Chamseddine, A. Connes, *The spectral action principle*, Comm. Math. Phys. **186** (1997) 731–750.
- A. Connes, C. Rovelli, *Von Neumann algebra automorphisms and time–thermodynamics relation in general covariant quantum theories*, Class. Quantum Grav. **11** (1994) 2899–2918.
- R. M. Wald, *Black hole entropy is the Noether charge*, Phys. Rev. D **48** (1993) R3427–R3431.
- H. Casini, M. Huerta, R. C. Myers, *Towards a derivation of holographic entanglement entropy*, JHEP **05** (2011) 036, [arXiv:1102.0440](#).