

TFPT — The Standard Model from the Compiler

The φ_0 -ladder, the flavor block from parabolic transport, and the worked closures (θ_{12} , quark c , the gap, Starobinsky M)

Stefan Hamann

Alessandro Rizzo

June 18, 2026 · v5.1 (rev 188)

What this document is about

The Standard Model as *one* φ_0 -ladder mass formula (all masses, CKM, PMNS, neutrinos), the flavor matrix from parabolic transport on $\mathbb{P}^1 \setminus \mu_4$, the five worked closures (θ_{12} , quark c , the explicit mass gap, Starobinsky M , the H2 splitting), and **gravity/quantum gravity as the seam response** (the Seam Response Grammar $\xi u = c_3$, the heat-kernel grounding of $R + R^2$, and the metric-coupled gap bound that *gap-decouples* the physical QG sector, leaving only the ambient projective limit as a mathematical-completeness item).

The TFPT document set — what is treated where

Plain language: TFPT (Topological Fixed-Point Theory) is a small discrete compiler. Two inputs — the seam constant $c_3 = \frac{1}{8\pi}$ and the carrier rank $g_{\text{car}} = 5$ — build $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ and read off the Standard-Model skeleton, the constants and the scale grammar. The series is **the introduction plus five numbered papers and three companions**, best read in order:

0. **introduction** — reading guide, compiler closure, predictions, the dependency DAG and the single proof ledger (`verification/status_ledger.csv`). (*entry point*)
1. **tfpt_1_architecture_e8** — the two axioms, the derivation map, the EM fixed point α^{-1} , the $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ construction, the QBL theorem chain.
2. **tfpt_2_standard_model** — the SM in one φ_0 -ladder formula, flavor from parabolic transport (sheet diamond, dual anchor, Q_+ cohomology), the worked closures.
3. **tfpt_3_e8_audit_bootstrap** — the seven E_8 slices as an audit raster, the cascade bridge, the Möbius bootstrap.
4. **tfpt_4_frontier** — honest status of η_B , m_p/m_e , Koide, dark matter, full QG.
5. **tfpt_5_redteam** — the adversarial audit: declared attacks, what survives, what each reduces to, and the kill tests.

Companions: **R.** **tfpt_research_contracts** — the open interfaces (v_{geo} , G_{net} , F_{transfer}) as numbered contracts; **H.** **tfpt_horizon_readouts** — Appendix H (unit-system reframe): one seam constant as the universal horizon thermal code, the Nariai/anchor surface, the resummed seam clock; **O.** **origin_theory** — why no free fundamental number remains (exact core + one honestly-typed cyclic interpretation).

You are reading document 2 (Standard Model).


TFPT status at a glance — read this card the same way in every document

Two inputs, one compiler. From the boundary constant $c_3 = \frac{1}{8\pi}$ (axiom P1) and the five-slot carrier $g_{\text{car}} = 5$ (axiom P2) — jointly the elementary symmetric data of one anchor $a = (1, 1, 2)$ — the discrete compiler $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ is *built*, and the Standard-Model skeleton (gauge group, three families, hypercharges, the flavor matrix), the dimensionless constants ($\alpha^{-1} = 137.0359992$) and the scale grammar are read off as fixed points. The algebraic core is machine-checked; physical readouts run through *named bridges* whose status is typed below.

What is closed, and what genuinely remains. The discrete/algebraic layer is closed (**[E]**); the everything-open residual reduces to *three named interface problems*:

Interface	Question	Status
v_{geo}	the one metrology unit ($= 1/\sqrt{G} = m/\mu$); No-Unit Thm: no compiler scale	primitive [O]
G_{net}	simple-current extension $(D_5)_1 \otimes (A_3)_1 \rtimes \langle (1, 1) \rangle \cong (E_8)_1$	[E] / [C]
F_{transfer}	one functor, four typed interfaces (Koide, η_B , axion, m_p/m_e)	[C]

Everything carries a claim ID resolving to a row of `verification/status_ledger.csv` (the single source of truth); **if the text and the ledger ever disagree, the ledger wins.** The historical labels ($U_{\text{wall}}, G_{\text{metric}}, F_{\text{frontier}}$) are retained only for ledger continuity — the live residual is $v_{\text{geo}} \oplus G_{\text{net}} \oplus F_{\text{transfer}}$.
 Marker key: **[E]** exact / machine-proven · **[C]** conditional (holds under named hypotheses) · **[O]** open / axiom · **[X]** falsifiable kill test. Per-claim fine type in `verification/status_ledger.csv`.

[E] exact / machine-proven**[C]** conditional**[O]** open / axiom**[X]** kill test

Contents

I	The Standard Model in one master formula	5
1	The principle: one master formula instead of many Yukawa numbers	5
2	The seed φ_0^{ret} as a decoder	5
3	The flavor residue matrix is the compiler signature	6
4	Charged leptons: completely closed in φ_0^{ret}	8
5	Quarks: hierarchy from the same word-lengths	9

6 The Sheet Diamond and the Winding Line: masses and transport from one surface	11
6.1 Closing the audit gates: is $M=41$, K , Q derived or imported?	20
7 Mixing: CKM and PMNS from holonomy	22
7.1 Downstream flavor readout: the rare kaons $K \rightarrow \pi \nu \bar{\nu}$	22
8 Neutrinos	23
9 How the legacy layers simplify (bridge map)	24
10 Is the entire Standard Model thereby derived?	24
11 Closure status of the five former open questions	25
12 How the original-paper predictions simplify	29
Conclusion	30
 II The flavor block: parabolic weights and the transport theorem	 31
13 The setup: parabolic data on the family curve	31
14 Exact scaffolding (proved)	32
15 The Riemann–Hilbert side (what the monodromy fixes)	32
16 The parabolic-weight computation (resolved this round)	33
17 Global computation: the anchor multiset is derived	34
17.1 Part (ii): the $\mathbb{Z}_6$ structure of the residues, and the frontier	35
17.2 The flavor transport theorem	35
18 The Parabolic Anchor Compiler	36
18.1 Budget master table: everything is $d_p = -\deg E = 4$ times a Coxeter datum	37
18.2 Flavor budget split, row sums, scalaron deficit	37
18.3 The anchor power compiler	37
18.4 Residue-matrix spectral selector — sharpening (H2)	38
19 Closing the flavor block: two theorems	39
Status	41
 III Closing computations and proofs	 43
20 The explicit mass gap closes dynamics and clustering	43
21 The seam elliptic complex is Fredholm (APS)	44
22 The $R^{n \geq 3}$ truncation is $O(1/N_\star^2)$-bounded	44
23 The ambient closure: one spectral-action measure, IR-complete by decoupling	45
23.1 Seam response grammar: gravity is the inverse seed gain	47

24 The solar angle θ_{12}: derived from the seam	48
25 Quark c-rationals: best values at current precision	49
26 H2: the moduli computation, set up explicitly	50
Status	52
Appendix: computational verification	53

Part I

The Standard Model in one master formula

In one sentence. The fermion spectrum — masses, Yukawa structure, CKM, the PMNS skeleton and the neutrinos — follows from *one* master formula with *three* ingredients: the seed φ_0^{ret} , the carrier base $\lambda_Y = \sqrt{\varphi_0^{\text{ret}}(1 - \varphi_0^{\text{ret}})}$ and the *residue matrix of the compiler*. No free Yukawa *exponents* remain; the charged-lepton amplitudes are closed, the quark mass *ratios* are closed too (integer Plücker readouts, [v49/v71](#)), and only the *absolute* quark amplitude scale is a finite (U_{wall}) anchor. **Markers:** [E] exact / machine-proven · [C] conditional (named hypotheses) · [O] open / axiom · [X] kill test — the fine per-claim type is in `verification/status_ledger.csv`.

1 The principle: one master formula instead of many Yukawa numbers

In the source-layer construction each fermion mass arises through a Yukawa coupling $y_{f,j}$ built as a transport word-length $L_{f,j}$ on a hexagon (C_6) times a holonomy amplitude. In the compiler picture this is *one* line:


Mass master formula (closed branch)
$\hat{m}_{f,j} = \frac{v_{\text{geo}}}{\sqrt{2}} y_{f,j}, \quad y_{f,j} = \lambda_Y^{L_{f,j}} \Lambda_{f,j}, \quad \lambda_Y = \sqrt{\varphi_0^{\text{ret}}(1 - \varphi_0^{\text{ret}})}$
with <i>one</i> seed $\varphi_0^{\text{ret}} = \frac{1}{6\pi} + \frac{3}{256\pi^4}$, the integer word-lengths $L_{f,j} = r_{f,j} + 6n_{f,j}$ (residue + winding) and $O(1)$ holonomy amplitudes $\Lambda_{f,j}$.

The point of the simplification: $L_{f,j}$ **is not a fit but the output of the compiler**. The residues $r_{f,j}$ form the matrix whose invariants are pure compiler numbers (Section 3); the windings $n_{f,j}$ are the anchors $\{1, 1, 2\}$. What used to be a separate “flavor transport” theorem is now the readout of a fixed matrix.

The seed base is the halved $(E_8)_1$ Weyl anomaly. The seed base $\frac{1}{6\pi} = \frac{4}{3}c_3$ carries the prefactor $\frac{4}{3} = \dim S^+ / \dim \mathfrak{g}_{\text{SM}} = 16/12$. That same $\frac{4}{3}$ is the conformal trace anomaly of the seam: over the seam sphere S^2 ($\chi = 2$, $\int R\sqrt{g} = 8\pi$) the $(E_8)_1$ ($c = 8$) Weyl anomaly is $\frac{c}{24\pi} \int R\sqrt{g} = \frac{c\chi}{6} = \frac{8}{3}$, halved by the one-sided $|\mathbb{Z}_2|$ Möbius identification to $\frac{4}{3}$. So the Cabibbo-seed prefactor is the halved $(E_8)_1$ conformal anomaly, not a free coefficient (`[verification/v172_trace_anomaly_seed.py]`).

2 The seed φ_0^{ret} as a decoder

A single seed drives four independent low-energy observables. This is the strongest compression of the flavor input:



Numerically: $\varphi_0^{\text{ret}} = 0.0531720$, $\lambda_C = 0.224376$ (PDG 0.2245), $\sin^2 \theta_{13} = 0.02311$ (PDG ≈ 0.0222). The exponent $\gamma = 5/6$ is the carrier trace $\text{tr}_E Y^2$; its complement $1 - \gamma = 1/6$ reappears in the neutrino ratio.

3 The flavor residue matrix is the compiler signature

The core of the simplification: the word-length matrix $L = R + 6W$ splits into the residue matrix R (the hexagon packets of the three sectors) and the winding matrix W (one +6 per first generation = anchor). R carries all the discrete flavor information — and its invariants are *exclusively* compiler numbers:

$$R = \begin{pmatrix} 1 & 3 & 0 \\ 1 & 5 & 2 \\ 2 & 5 & 3 \end{pmatrix} \quad \begin{array}{l} \text{rows} = \text{sectors } u, d, e \\ \text{tr } R = 9 = N_{\text{fam}}^2 \\ \text{minors } 2, 3, 5 \Rightarrow 2 \cdot 3 \cdot 5 = 30 = h(E_8) \\ \text{det } R = 8 = h(D_5) \end{array}$$

Spectral signature of the flavor matrix (verified)

The characteristic polynomial of the residue matrix is built entirely from compiler constants:

$$\chi_R(t) = t^3 - \underbrace{9}_{N_{\text{fam}}^2} t^2 + \underbrace{10}_{\mathcal{A}_L = \binom{5}{2}} t - \underbrace{8}_{h(D_5)},$$

where the linear coefficient $10 = 2 + 3 + 5$ is the sum of the three principal 2×2 minors and their *product* $2 \cdot 3 \cdot 5 = 30$ is the Coxeter number $h(E_8)$. The flavor matrix is therefore not an independent object — it is a *reading of the same compiler* (N_{fam} , $\mathcal{A}_L$, $h(D_5)$, $h(E_8)$). [E] [[verification/v4_flavor_matrix.py](#)]

Concretely:

$$L = \begin{pmatrix} 7 & 3 & 0 \\ 7 & 5 & 2 \\ 8 & 5 & 3 \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 3 & 0 \\ 1 & 5 & 2 \\ 2 & 5 & 3 \end{pmatrix}}_{R \text{ (det 8, minors 2,3,5)}} + 6 \underbrace{\begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}}_{W \text{ (anchor \{1,1,2\})}}, \quad \sum L = 40 = \Omega_{\text{adm}} \cdot \frac{5}{6}.$$

The sum $\sum L = 40$ is the transport budget number; with the Higgs contribution $N_\Phi = 1$ it gives $40 + 1 = 41 = 10 b_1$ — exactly the coefficient the EM fixed point $\alpha_\star^{-1}$ reads.

The winding-deformation theorem: $R \rightarrow L$ in one parameter [E]

Treat the winding as a one-parameter rank-one deformation $R_s = R + s \mathbf{1} e_1^\top$ ($R_0 = R$, $R_6 = L$). Then

the *entire* flavor linear algebra is one line,

$$\chi_{R_s}(t) = t^3 - (9 + s)t^2 + (10 + 5s)t - (8 + 2s),$$

with $\det R_s = 8 + 2s$, $\text{tr } R_s = 9 + s$, $\text{PrinMin}_2(R_s) = (5, 3(s+1), 2(s+1))$, and $R_s^{-1}\mathbf{1} = \frac{1}{4+s}(1, 1, -1)^\top$. The physical winding value is fixed, not chosen:

$$s = |R^+(A_3)| = 6 \text{ is triply locked: } \underbrace{6 + 4s = 30}_{R_s^\top a \rightarrow h(E_8)}, \underbrace{8 + 2s = 20}_{\det \rightarrow 2\mathcal{A}_L}, \underbrace{9 + s = 15}_{\text{tr} \rightarrow \dim A_3} \Rightarrow s = 6.$$

At $s = 6$ this gives $\chi_L = t^3 - 15t^2 + 40t - 20$ with $15 = \dim A_3$, $40 = |R(D_5)|$, $20 = |R(A_4)| = 2\mathcal{A}_L$, and $\text{PrinMin}_2(L) = (5, 21, 14) = (g_{\text{car}}, 3 \cdot 7, 2 \cdot 7)$ — the lift multiplies the sheet and family minors by the successor $s+1 = 7 = |R^+(A_3)|+1$ while the carrier minor 5 stays fixed. So the “+6 winding” is not “add six because hexagon”: it is the unique value selected three independent ways, and the appearance of 7 in the scalaron is $s+1$.

Winding reality, the determinant lift, and the invariant dual anchor (machine-verified)

The rank-one winding $R \rightarrow L = R + 6\mathbf{1}e_1^\top$ is far from a cosmetic offset:

- **Winding makes the flavour system spectrally real.** R has a *complex* eigenpair ($\text{disc } \chi_R = -2^2 \cdot 1999 < 0$, spectrum $7.857, 0.572 \pm 0.832i$), while L has three positive reals, with a fully compiler-factorised discriminant (two equivalent readings)

$$\text{disc } \chi_L = 39200 = 2^5 \cdot g_{\text{car}}^2 \cdot 7^2 = 2(\mathcal{A}_L \cdot \dim G_2)^2 = 2(10 \cdot 14)^2.$$

- **The anchor Coxeter descent.** The anchor projects onto a clean three-scale ladder,

$$R^\top a = (6, 18, 8), \quad L^\top a = (30, 18, 8) = (h(E_8), N_{\text{fam}}|R^+(A_3)|, h(D_5)),$$

so the winding lifts $|R^+(A_3)| = 6 \rightarrow h(E_8) = 30$ (a $+24 = |W(A_3)|$ shift), with descent gaps $30 - 18 = 12 = |R(A_3)| = \dim \mathfrak{g}_{\text{SM}}$ and $18 - 8 = 10 = \mathcal{A}_L$. Column-wise the lift is $\mathbf{1}^\top R = (4, 13, 5) \rightarrow \mathbf{1}^\top L = (22, 13, 5)$, adding 18 only in the first generation — which is thus literally the store of the family-hexagon winding.

- **The determinant lift is the A_3 winding over μ_4 .** Since $e_1^\top R^{-1}\mathbf{1} = \frac{1}{4} = \frac{1}{|\mu_4|}$,

$$\det L = \det R \left(1 + \frac{|R^+(A_3)|}{|\mu_4|}\right) = 8 \left(1 + \frac{6}{4}\right) = 20 = 2\mathcal{A}_L,$$

so the D_5 selector 8 is lifted to the decuple-doublet 20 by the A_3 winding. The cokernel moves $\mathbb{Z}_8 \rightarrow \mathbb{Z}_{20}$, i.e. the winding injects the carrier factor 5 into the torsion.

- **The dual anchor is winding-invariant.** The first column is the anchor, $Re_1 = a = (1, 1, 2)$, and the dual covector is preserved by the lift,

$$a^\top R^{-1} = a^\top L^{-1} = \left(-\frac{1}{2}, -\frac{1}{2}, 1\right), \quad \left(-\frac{1}{2}, -\frac{1}{2}, 1\right) \cdot \mathbf{1} = 0,$$

so winding changes the absolute hierarchy but *not* the anchor-dual sheet balance. The uniform response keeps the sheet value fixed: $\mathbf{1}^\top \text{adj}(R)\mathbf{1} = \mathbf{1}^\top \text{adj}(L)\mathbf{1} = 2 = |\mathbb{Z}_2|$, while the inverse response denominator runs $|\mu_4| = 4 \rightarrow \mathcal{A}_L = 10$. [E]

Audit-level only: flavor linear-algebra fingerprints (not load-bearing). The following are *audit fingerprints* (exact linear algebra of R, L), kept here, not in any derivation, under the no-free-pattern rule. [E]

- **Cofactor seam normal.** Generations 2, 3 share columns, $c_2 = (3, 5, 5)^\top$, $c_3 = (0, 2, 3)^\top$; their cross product $n = c_2 \times c_3 = (5, -9, 6) = (g_{\text{car}}, -N_{\text{fam}}^2, |R^+(A_3)|)$ gives the determinant geometrically: $n \cdot a = 8 = \det R$, $n \cdot (a+6\mathbf{1}) = 20 = \det L$, $n \cdot \mathbf{1} = 2 = |\mathbb{Z}_2|$.
- **Adjugate selector ($\Rightarrow$ why only gen. 1 winds).** n is also the first row of both adjugates, $e_1^\top \text{adj}(R) = e_1^\top \text{adj}(L) = (5, -9, 6)$, so $(5, -9, 6)R = (8, 0, 0)$ and $(5, -9, 6)L = (20, 0, 0)$ — it annihilates the unchanged c_2, c_3 plane.

- **Length polynomial as an action polynomial.** $\chi_L(t) = t^3 - \dim(A_3)t^2 + |R(D_5)|t - |R(A_4)| = t^3 - 15t^2 + 40t - 20$.
- **Principal minors.** $\text{PrinMin}_2(L) = (5, 21, 14) = (g_{\text{car}}, 3 \cdot 7, 2 \cdot 7)$ with $\Sigma = 40 = |R(D_5)|$, $\Pi = 1470 = 7^2 h(E_8)$.
- **Commutator is a pure winding rotation.** $[R, L]$ has rank 2, trace 0, and char. poly $\lambda(\lambda^2 + 432)$ — a single 2D rotation in the non-anchor plane with a null direction.

4 Charged leptons: completely closed in φ_0^{ret}

For the leptons the hexagonal resolvent solves the amplitudes Λ_ℓ in closed form — only pure φ_0^{ret} powers with rational carrier prefactors remain:

Absolute lepton source masses

$$(\hat{m}_e, \hat{m}_\mu, \hat{m}_\tau) = \frac{v_{\text{geo}} \pi}{\sqrt{2}} \left(\frac{16}{7} (\varphi_0^{\text{ret}})^5, \frac{4}{3} (\varphi_0^{\text{ret}})^3, \frac{7}{6} (\varphi_0^{\text{ret}})^2 \right),$$

and hence the *parameter-free* ratios

$$\frac{\hat{m}_\mu}{\hat{m}_\tau} = \frac{8}{7} \varphi_0^{\text{ret}}, \quad \frac{\hat{m}_e}{\hat{m}_\mu} = \frac{12}{7} (\varphi_0^{\text{ret}})^2, \quad \frac{\hat{m}_e \hat{m}_\tau}{\hat{m}_\mu^2} = \frac{3}{2} \varphi_0^{\text{ret}}.$$

Where the lepton rationals come from: $\delta = \frac{1}{2}$ resolvent + product rule [E]

The coefficients $(\frac{16}{7}, \frac{4}{3}, \frac{7}{6})$ are not fitted — they are *derived*, and the key is that they are *rational*, so they cannot come from the irrational pole δ_{ph} . They come from the *distinguished* transport value $\delta = \frac{1}{2}$ (rational), evaluated at the lepton leg $y = 1$ ($= |Y(e^c)|$). **The value $\delta = \frac{1}{2}$ itself is a glue-norm readout:** the two E_8 -glue norms are $q(D_5) = g_{\text{car}}/|\mu_4| = \frac{5}{4}$ and $q(A_3) = N_{\text{fam}}/|\mu_4| = \frac{3}{4}$, so $\delta = q(D_5) - q(A_3) = (g_{\text{car}} - N_{\text{fam}})/|\mu_4| = |\mathbb{Z}_2|/|\mu_4| = \frac{1}{2}$ — the carrier-minus-family count over the glue index (the same pair whose *sum* is the E_8 root norm 2, architecture document), matching the harmonic-frame holonomy diagonal modulus $\frac{1}{2}$. So δ is fixed, not chosen ([[verification/v51_boundary_half_step.py](#)]). With that value,

$$\text{amplitude}(r) = \left| \frac{1}{1 - \frac{1}{2}\zeta^r} \right|^2 = \frac{1}{\frac{5}{4} - \cos(r\pi/3)}, \quad \zeta = e^{i\pi/3}, \quad r = L \bmod 6.$$

Non-anchor leptons (dressed by the winding factor $|\mu_4| = 4$, $w = \lfloor L/6 \rfloor$):

$$c = |\mu_4|^w \frac{1}{\frac{5}{4} - \cos(r\pi/3)} : \quad c_e = 4^1 \cdot \frac{4}{7} = \frac{16}{7}, \quad c_\mu = 4^0 \cdot \frac{4}{3} = \frac{4}{3} \quad (\text{exact}).$$

Anchor (heaviest lepton τ) fixed by the *sector product rule*

$$\boxed{c_e c_\mu c_\tau = \frac{2^{g_{\text{car}}}}{N_{\text{fam}}^2} = \frac{32}{9}} \implies c_\tau = \frac{32/9}{(16/7)(4/3)} = \frac{7}{6} \quad (\text{exact}).$$

Every ingredient is a compiler datum ($\delta = \frac{1}{2}$, $|\mu_4|$, hexagon angles, g_{car} , N_{fam}). The product constant reads $2^{g_{\text{car}}} = 32 = \dim(S^+ \oplus S^-)$ (the full carrier Clifford space) over N_{fam}^2 , and the lepton word-lengths obey $\sum_j L_j = 16 = \dim S^+$, $\sum_j (L_j - K_j) = 6 = |R^+(A_3)|$. *Edge-centre ring:* the two boundary generations close on the middle one, $c_e c_\tau = \frac{16}{7} \cdot \frac{7}{6} = \frac{8}{3} = |\mathbb{Z}_2| c_\mu$, so the lepton amplitudes form a closed algebraic ring with sheet-parity $|\mathbb{Z}_2| = 2$. [[verification/v20_lepton_c_derivation.py](#)] [[verification/v21_solar_product_quark.py](#)] [[verification/v37_plucker_anchor.py](#)]

Update (the hexagon is the family monodromy, [[verification/v118_hexagon_family_dictionary.py](#)]): with the flavor-wall monodromy now exact ($\widetilde{M}_0 \in W(A_3)$, research contracts), the hexagon constants stop being separate inputs [E]: $\text{spec}(\widetilde{M}_0) \cup \text{spec}(-\widetilde{M}_0) = \mu_6$ (the six sites = family spectrum + sheet twist), $\det(\mathbf{1} - t\widetilde{M}_0) = 1 - t^3$, and

$$c_e = \frac{|\mu_4|}{2 \det(\mathbf{1} - \frac{1}{2}\widetilde{M}_0)} = \frac{16}{7}, \quad c_\mu = \frac{N_{\text{fam}}}{2 \det(\mathbf{1} + \frac{1}{2}\widetilde{M}_0)} = \frac{4}{3}, \quad c_\tau = \frac{|\mu_4| \det(\mathbf{1} - \frac{1}{2}\widetilde{M}_0)}{N_{\text{fam}}} = \frac{7}{6},$$

with the product rule $= |\mu_4|/\det(\mathbf{1} + \frac{1}{2}\widetilde{M}_0) = \frac{32}{9}$ — the lepton 7 and 9 *are* the two family-resolvent determinants ($\times 2^3$), and $\delta = \frac{1}{2} = |(\widetilde{M}_0)_{22}|$ is supplied by the monodromy itself. The $e/\mu/\tau$ *leg assignment* remains the established input above; nothing is re-fitted.

The lepton ring is an étale Frobenius algebra [\[E\]/\[C\]](#) ([\[verification/v229_lepton_frobenius_algebra.py\]](#)). The three coefficients are the spectrum of the commutative algebra $A = \mathbb{Q}[t]/(m)$, $m = (t-c_e)(t-c_\mu)(t-c_\tau)$: A is étale (separable), so the trace form $(x,y) \mapsto \text{Tr}(xy)$ is nondegenerate ($\det \text{Gram} = \text{disc}(m) \neq 0$) — the defining property of a (commutative) Frobenius algebra — and it lives over the μ_6 resolvent (C_6 has characteristic polynomial t^6-1 , spectrum $\mu_6 = \mu_3 \text{ family} \times \mu_2 \text{ sheet}$). So the charged-lepton boundary is a genuine finite Frobenius object [\[E\]](#); the PMNS extension (its automorphisms plus the hexagonal CM phase $j=0$, [\[verification/v220_cp_hexagonal_modulus.py\]](#)) is the natural [\[C\]](#) candidate, not a closure of the matrix.

Scope (honest). This law is lepton-specific. Applied to the down sector it fails provably: $r_d=1, r_s=5$ share the same amplitude, so the law forces $c_d/c_s=|\mu_4|=4$, whereas the source ratio gives 0.94. Quarks (colour, different cusp legs) are structurally different and their exact c 's stay open (cf. the U_f^* reduction above).

Ratio	Formula	Prediction	PDG
$\hat{m}_\mu/\hat{m}_\tau$	$\frac{8}{7}\varphi_0^{\text{ret}}$	0.06077	0.05946
$\hat{m}_e/\hat{m}_\mu$	$\frac{12}{7}(\varphi_0^{\text{ret}})^2$	0.004847	0.004836
$\hat{m}_e\hat{m}_\tau/\hat{m}_\mu^2$	$\frac{3}{2}\varphi_0^{\text{ret}}$	0.07976	0.08133

The factor $B = \frac{3}{2}$ in the Koide-type product is the rank ratio $\text{rank } E_3/\text{rank } E_2$ of the carrier — again a compiler number, not a fit. The e/μ relation hits the PDG to 0.2%.

5 Quarks: hierarchy from the same word-lengths

The quark sectors use the same master formula with $L_u = (7, 3, 0)$, $L_d = (7, 5, 2)$. Where two generations share the same word-length the mass ratio is pure transport amplitude; where they differ by 2 a factor λ_Y^{-2} appears:

Ratio	Origin	Prediction	PDG order
$\hat{m}_u/\hat{m}_d$	Λ_u/Λ_d (same $L=7$)	0.470	~ 0.5
$\hat{m}_c/\hat{m}_s$	$\lambda_Y^{-2} \Lambda_c/\Lambda_s$	13.6	$\sim 11-14$
$\hat{m}_t/\hat{m}_b$	$\lambda_Y^{-2} \Lambda_t/\Lambda_b$	40.8	~ 41

The residue classes mod 6 link quarks and leptons: $L_c = L_\tau = 3$, $L_s = L_\mu = 5$ — the same hexagon position, different sectors. This is the geometric root of quark–lepton complementarity.

The φ_0^{ret} -ladder: *all* nine masses in one closed form

The quark Λ are *not* free numbers: they are evaluations of the *same* finite hexagonal resolvent D_y^{-1} (proven in closed form, [\[verification/v17_hexagonal_resolvent.py\]](#)) at the *same* pole δ_{ph} as the leptons. For every fermion the *universal* closed form holds

$$\lambda_Y^{L_{f,j}} \Lambda_{f,j} = \pi c_{f,j} (\varphi_0^{\text{ret}})^{k_{f,j}} \implies \hat{m}_{f,j} = \frac{v_{\text{geo}} \pi}{\sqrt{2}} c_{f,j} (\varphi_0^{\text{ret}})^{k_{f,j}}$$

with carrier rationals $c_{f,j}$ and integer φ_0^{ret} -order $k_{f,j}$. The leptons give $c = (\frac{16}{7}, \frac{4}{3}, \frac{7}{6})$ *exactly*; the quark c are fixed by the same resolvent (the published Λ table is only three-digit). The φ_0^{ret} -orders form the **mass ladder**

$$\underbrace{t}_{k=0} > \underbrace{b, c, \tau}_{k=2} > \underbrace{s, \mu}_{k=3} > \underbrace{u, d}_{k=4} > \underbrace{e}_{k=5},$$

each rung a factor $\varphi_0^{\text{ret}} \approx 1/19$. The whole fermion hierarchy is a φ_0^{ret} power ladder with carrier-rational rungs — quarks *and* leptons in *one* family. [\[E\]](#) (leptons exact) / [\[C\]](#) (quark rationals from the resolvent)

Quark *ratio* closure: the invariant readouts of the character sector [E]

The right invariants are not the six absolute c_q (gauge-dependent through the external legs and the sector normalisation) but the three intra-sector *ratios*, which close on pure compiler building blocks. With $\Delta_Q := |R(A_3)|+1 = 13$, $\Omega_{\text{adm}} = 48$, $N_{\text{fam}} = 3$ and the anchor power sum $p_5(a) = 1^5+1^5+2^5 = 34$,

$$\frac{c_u}{c_d} = \frac{g_{\text{car}} \cdot 11}{N_{\text{fam}}^2 \Delta_Q} = \frac{55}{117}, \quad \frac{c_c}{c_s} = \frac{p_5(a)}{\Omega_{\text{adm}} - 1} = \frac{34}{47}, \quad \frac{c_t}{c_b} = \frac{N_{\text{fam}}}{2\Delta_Q} = \frac{3}{26}$$

With the φ_0^{ret} -orders $(k_u, k_d, k_c, k_s, k_t, k_b) = (4, 4, 2, 3, 0, 2)$ the mass ratios are

$$\frac{\hat{m}_u}{\hat{m}_d} = \frac{55}{117} = 0.4700855, \quad \frac{\hat{m}_c}{\hat{m}_s} = \frac{34}{47}(\varphi_0^{\text{ret}})^{-1} = 13.6050, \quad \frac{\hat{m}_t}{\hat{m}_b} = \frac{3}{26}(\varphi_0^{\text{ret}})^{-2} = 40.812,$$

matching the closed-branch targets (0.470085, 13.607, 40.80) to $\lesssim 0.03\%$. A rational normalisation gauge compatible with these ratios is

$$(c_u, c_d, c_s, c_c, c_b, c_t) = \left(\frac{1}{2}, \frac{117}{110}, \frac{47}{41}, \frac{34}{41}, \frac{8}{3}, \frac{4}{13}\right),$$

whose $\Lambda_q = \pi c_q(\varphi_0^{\text{ret}})^k / \lambda_Y^L$ all land in the $O(1)$ band ($\Lambda_{u,d,s,c,b,t} \approx 0.44, 0.93, 0.95, 0.65, 0.47, 0.97$). The denominators carry $41 = 4^2+5^2 = 10b_1$. The individual c_q are a gauge; the three ratios are the invariant readouts. [verification/v24_quark_ratio_closure.py]

Honest note on the “11”. Two of the three ratios have *clean single-reading* building blocks: $c_c/c_s = p_5(a)/(\Omega_{\text{adm}}-1)$ (the anchor power sum $p_5=34$) and $c_t/c_b = N_{\text{fam}}/(2\Delta_Q)$. The first ratio is the odd one out: its numerator factor 11 admits *many* TFPT readings ($\dim S^+ - g_{\text{car}}, g_{\text{car}} + |R^+(A_3)|, h(D_5) + N_{\text{fam}}, 2g_{\text{car}}+1, A_{\Lambda}+1, \Delta_Q-2$, the E_8 exponent 11, ...), so it is *not uniquely forced* as a loose integer. **This worry is now retired** (Plücker box below): $11 = \|\text{Pl}(K)\|_1$ is *generated* as the L^1 norm of the anchor-plane minors of K (v42), and by Readout Rigidity (v49, with the selector stratum now *derived* — $\det R = 8$ lattice and $\text{Spec}(Q_+) = \{1, 2, 3\}$ from D_4 , v69/v71) the *ratio* $c_u/c_d = \frac{55}{117}$ is *constant* on that stratum, i.e. an integer Plücker readout that needs *no* transcendental solve. The wall-selection (U) enters only the *absolute* amplitude normalisation (an anchor), not the ratio. [verification/v26_flavor_frontier_unification.py] Quark masses are scheme-sensitive (esp. light quarks); we state dimensionless source-scale ratios, not pole masses (cf. FLAG 2024).

Anchor-plane Plücker localisation of the “11” (and 26) [E]

The cleanest way to retire the “many readings” worry is to stop reading loose integers and read *orientation invariants* of the one canonical plane in play — the *anchor plane* spanned by the two TFPT generators $\mathbf{1} = (1, 1, 1)$ and $a = (1, 1, 2)$. For $M \in \{R, Q, K, L\}$ form the left anchor block and its 2×2 minors (Plücker coordinates)

$$B_M = \begin{pmatrix} \mathbf{1}^\top M \\ a^\top M \end{pmatrix}, \quad \text{Pl}(M) = ([12], [13], [23]).$$

Then $\text{Pl}(K) = (-N_\Phi, |R^+(A_3)|, |\mu_4|) = (-1, 6, 4)$ and

$$\|\text{Pl}(K)\|_1 = 1 + 6 + 4 = 11$$

is the *Plücker norm of the mass-power anchor plane* — a single coordinate-free object, not a free integer. The right block $C_M = (M \mathbf{1} \ Ma)$ gives $\text{Pl}_R(K) = (12, 12, -2)$ with $\|\text{Pl}_R(K)\|_1 = 26$, and $\sum \text{Pl}_R(L) = 20+30+10 = 60$ is the E_8 cascade start. **Canonical readings** ([verification/v94_sheet_diamond.py]): both quark-ratio integers live in the *same* operator K — the left block reads flavor *orientation*, the right block reads transport *propagation*:

$$\frac{c_u}{c_d} = \frac{g_{\text{car}} \|\text{Pl}(K)\|_1}{N_{\text{fam}}^2 \Delta_Q}, \quad \frac{c_t}{c_b} = \frac{N_{\text{fam}}}{\|\text{Pl}_R(K)\|_1}, \quad \frac{c_c}{c_s} = \frac{\sum \text{Pl}(L) - \sum \text{Pl}(Q)}{1 + \sum \text{Pl}(L)},$$

with the older 26-readings ($\text{Pl}(L)_{13} = 26$, $2\Delta_Q = 26$) now *corollaries* of the canonical right-Plücker norm, not independent choices. **What this does and does not buy.** The Plücker values are exact identities [E], and the “11” now *lives in the mass-power matrix* K via the canonical plane rather than

in a free integer pool — the best-motivated single reading. Combined with Readout Rigidity (below), this *closes the ratio*: $c_u/c_d = \frac{55}{117}$ is fixed combinatorially, no U_f^* evaluation required. What stays open is the *absolute* amplitude scale (an anchor) and a mild choice of invariant in the c_c/c_s block. [verification/v37_plucker_anchor.py]

Why the leg is exterior, not scalar (Exterior Leg Lemma). u and d share the hexagon-radial data $(r, w) = (1, 1)$, and the C_6 resolvent reads only (r, w) — so *any scalar* leg forces $c_u/c_d \equiv 1$. A scalar sees radius, not orientation; the u/d separation is intrinsically the oriented anchor-plane *area* $\text{Pl}_{1,a}(K)$ (the exterior square $\Lambda^2 F$ of the $SU(3)_F$ composition), whose L^1 norm *generates* the “11”. [verification/v42_exterior_leg.py] The norm $\|\text{Pl}(K)\|_1 = 11$ also carries a topological reading: it is the dimension of the QBL-truncated ($\deg \leq 2$) Fock space of the four μ_4 -corner fermions, $\sum_{k \leq 2} \binom{4}{k} = 1+4+6 = 11 = \dim S^+ - g_{\text{car}}$ (a Witten index; the discarded high-degree complement is $\binom{4}{3} + \binom{4}{4} = 5 = g_{\text{car}}$, the full corner Fock space $2^4 = 16 = \dim S^+$), so $c_u/c_d = 55/117$ reads the QBL-protected boundary state-space dimension ([verification/v174_seam_fock_readings.py]).

Readout Rigidity — the quark *ratios* are closed combinatorially [E] (residual: absolute scale [O])

The earlier “reduce c_u/c_d to the transcendental wall-selection” framing is *superseded*. The four flavor operators are integer lattice maps on $H_1(\mathbb{P}^1 \setminus \mu_4) = \mathbb{Z}^3$ with compiler-atom determinants $(\det Q, \det K, \det R, \det L) = (3, 4, 8, 20)$, product $1920 = |W(D_5)|$, and the *discrete selector stratum*

$$\mathcal{S}_{8,Q} = \{\det R = 8, \text{SNF}(R) = (1, 1, 8), \text{Spec}(Q_+) = \{1, 2, 3\}\}$$

is now *fully derived*: $\det R = 8 = \text{rank } E_8$ and the SNF are the lattice invariant (v70, $\det Q = 3 = N_{\text{fam}}$), and $\text{Spec}(Q_+) = \{1, 2, 3\}$ is the D_4 -equivariant result (v69). By **Readout Rigidity** (Theorem U2, [verification/v49_readout_rigidity.py]) the anchor-plane readout is *constant* on $\mathcal{S}_{8,Q}$, so the quark *ratios*

$$\frac{c_u}{c_d} = \frac{g_{\text{car}} \|\text{Pl}(K)\|_1}{N_{\text{fam}}^2 \Delta_Q} = \frac{5 \cdot 11}{9 \cdot 13} = \frac{55}{117}, \quad \frac{c_c}{c_s}, \frac{c_t}{c_b}$$

are *pure integer Plücker readouts* — independent of the continuous D_4 position, no transcendental monodromy / Hitchin solve ([verification/v71_simple_r_bridge.py]). The single genuinely transcendental piece, U_{point} (full uniqueness of ρ^*), enters *only* the absolute amplitude scale and is an anchor of the same nature as the one dimensional scale — *not* a missing computation. The monodromy machinery (§Riemann–Hilbert below) was over-engineering for the ratios; it remains the route to the *absolute* normalisation only.

6 The Sheet Diamond and the Winding Line: masses and transport from one surface

The mass-power matrix and the transport word-length matrix are *not* two independent machines: they are the two sheet-readouts of *one* parabolic projection Q acting on the residue matrix R . Collect the φ_0^{ret} -orders $k_{f,j}$ of the nine charged fermions into the **mass-power matrix**

$$K = \begin{pmatrix} 4 & 2 & 0 \\ 4 & 3 & 2 \\ 5 & 3 & 2 \end{pmatrix} \quad (\text{rows } u, d, e), \quad Q := L - K = \begin{pmatrix} 3 & 1 & 0 \\ 3 & 2 & 0 \\ 3 & 2 & 1 \end{pmatrix}, \quad \Sigma := \text{diag}(1, -1, -1), \quad (1)$$

with Σ the *sheet* ($\mathbb{Z}_2$) involution. Then, exactly,

$$\boxed{K = R + Q\Sigma, \quad L = R + Q(I + \Sigma)} \quad [\text{E}] \quad [\text{verification/v10_projection_involution.py}], \quad (2)$$

and since $I + \Sigma = \text{diag}(2, 0, 0)$ and Q ’s first column is $(3, 3, 3)^\top$, the “hand-added” first-generation winding is *forced*: $Q(I + \Sigma) = 6W = 2N_{\text{fam}} \cdot \mathbf{1}e_1^\top$, i.e. sheet ($\mathbb{Z}_2$) $\times$ families ($\mathbb{Z}_3$) = the C_6 transport. So the φ_0^{ret} -mass powers are the Σ -odd readout, and the transport lengths are the $(I + \Sigma)$ readout, of the *same* projection Q on R .




Figure 1: Mass ladder (data plot, `verification/make_figures.py`). The nine charged fermions (PDG masses) plotted against their word-length L on the φ_0^{ret} -ladder, coloured by sector; the dotted line is the geometric reference $\propto \lambda_Y^L$ with $\lambda_Y = \sqrt{\varphi_0^{\text{ret}}(1 - \varphi_0^{\text{ret}})}$. The whole five-order hierarchy is one ladder.

The Sheet Diamond: all four operators are points of *one* surface [E]
(`[verification/v94_sheet_diamond.py]`)

The whole flavor algebra lives on a single two-parameter operator surface,

$$M(s, t) = R + Q \operatorname{diag}(s, t, t) : \\ R = M(0, 0), \quad K = M(1, -1), \quad L = M(2, 0), \quad F := R + Q = M(1, 1),$$

with the two global invariants

$$\det M(s, t) = 3st^2 + 9st + 6s + t^2 + 5t + 8, \quad \det B_{M(s, t)} = 6st + 12s + 3t^2 + 15t + 16.$$

The mass-transport pencil $K + xQ$ is the *cut* $(s, t) = (1+x, x-1)$, on which the anchor-block determinant factorises as $(3x+2)(3x+5)$ — the master cover of the operator-pencil sections below (unique up to Möbius reparametrisation, `[verification/v85_master_cover.py]`). The factorisation is **not generic**: the surface carries its own negative controls — the diagonal cut $t=s$ gives $9s^2+27s+16$ with discriminant 153 (non-square; the *same* control discriminant as the R -direction pencils), and the $R \rightarrow L$ cut $t=0$ is linear (degenerate direction, no cover).

Centered normal form (`[verification/v95_centered_diamond.py]`). The diamond is a *centered cross*: with the center $C := M(1, 0) = \frac{R+L}{2} = \frac{K+F}{2}$ and the two axes $U := Q \operatorname{diag}(1, 0, 0) = N_{\text{fam}} \mathbf{1} e_1^\top$ (winding, rank 1, spectrum $\{3, 0, 0\}$) and $V := Q \operatorname{diag}(0, 1, 1)$ (sheet, rank 2),

$$\boxed{Q = U + V, \quad R = C - U, \quad L = C + U, \quad K = C - V, \quad F = C + V} \quad [\mathbf{E}],$$

so five objects collapse to three, and the winding line below *is* the U -axis ($L = R + 2U$). The sheet axis carries exactly the cusp class: $\operatorname{Spec}(V) = \{0, 1, 2\} = N_{\text{fam}}\{0, \frac{1}{3}, \frac{2}{3}\}$ (cf. $Q_+ = 3 \text{ cusp} + 1$ with $\{1, 2, 3\}$, `v69/v72`). The center is tightly typed: $\det C = 14$, $\operatorname{SNF}(C) = (1, 1, 14)$, $\sum_{ij} C_{ij} = 31 = 2^{g_{\text{car}}} - 1 = 1 + h^\vee(E_8)$ (the same 31 as the IR gap-bound numerator $\|V_{\text{metric}}\| \leq 31/(8\pi^2)$), $\det B_C = 28 = 2 \cdot 14$, $\operatorname{Pl}(C) = (0, 14, 14)$ and $\operatorname{Pl}_R(C) = 7(2, 3, 1)$ — the same ray $(2, 3, 1)$ as $\operatorname{Pl}_R(L) = 10(2, 3, 1)$, scalaron-7 vs decuple-10 scaling. *Typing*: the normal form and all integers are exact **[E]**; the “ G_2 center” reading ($14 = \dim G_2$, also the constant term of $\chi_{[K, Q]} = t^3 + 5t + 14$ and $\sqrt{\sum B_F}$) is an *audit* Lie-dimension match, recorded, not promoted.

Axis curvature and two ladders (`[verification/v218_diamond_axis_geometry.py]`). The cross has a *discrete differential geometry*: the determinant is *linear* along the winding axis and *quadratic* along the sheet axis,

$$\det(C + xU) = 14 + 6x, \quad \det(C + yV) = 14 + 14y + 4y^2, \quad \det B_{C+xU} = 2 \det(C + xU),$$

so winding flows flat (A_3 -driven, slope $6 = |R^+(A_3)|$) while the sheet split *curves* with second differences $8 = \text{rank } E_8$ (determinant) and $6 = |R^+(A_3)|$ (anchor block) — the two curvatures are exactly the triply-forced 8 and the A_3 positive roots. Two exact ladders ride the same sheet axis. The **Plücker transfer ladder** $\text{Pl}(K) = (-1, 6, 4) \rightarrow \text{Pl}(C) = (0, 14, 14) \rightarrow \text{Pl}(F) = (1, 22, 30)$ lifts in steps $(1, 8, 10) = (N_\Phi, \text{rank } E_8, A_\Lambda)$ then $(1, 8, 16) = (N_\Phi, \text{rank } E_8, \dim S^+)$ — first the decuple, then the full spinor generation, so the F_{transfer} source→pole step *is* this two-stage sheet lift [C]. The **spectral ramification ladder** factors the cubic discriminant of Q, K, C, F as $q(r)^2 \text{Disc}(q)$ with squares $(1, 3, 4, 6) = (N_\Phi, N_{\text{fam}}, |\mu_4|, |R^+(A_3)|)$ and kernels $(13, 48, 65, 105) = (\Delta_Q, \Omega_{\text{adm}}, g_{\text{car}} \Delta_Q, N_{\text{fam}} g_{\text{car}} \cdot 7)$, the corner F carrying $105 = 3 \cdot 5 \cdot 7$ (family · carrier · scalaron). *Audit:* the anchor-defect spectrum $\Delta_a(Q, K, R, C, L, F) = (8, 16, 18, 21, 24, 26)$ and the corner-determinant pair-sum atlas (total $64 = \dim S^+ |\mu_4|$, staircase sum $81 = N_{\text{fam}}^4$) are recorded fingerprints — the $2 \dim G_2$ and $\dim F_4$ pair sums stay audit-only, like the G_2 center above.

F_{transfer} is a path, not four interfaces ([verification/v224_diamond_ftransfer_path.py]). The corners $K=M(1, -1)$, $C=M(1, 0)$, $F=M(1, 1)$ lie on the *sheet axis* $s=1$: the determinant there is the curved $\det M(1, t) = 4t^2 + 14t + 14$ (values $(4, 14, 32)$, 2nd difference $8 = \text{rank } E_8$), while the winding axis $\det M(s, 0) = 6s + 8$ is flat. So F_{transfer} is read as *one discrete geodesic* $K \rightarrow C \rightarrow F$ on the diamond (the Plücker steps $(1, 8, 10)$, $(1, 8, 16)$ above), of which the Koide source→pole Möbius relaxation is the 1-D projection (v37/v183); the external QFT solver evaluates continuously *along* this known path [C]. The diagonal cut $t=s$ mixes the axes (negative control), so only the $s=1$ sheet axis is the branch-preserving transfer direction.

The center collects the three local norms ([verification/v230_center_budget_norms.py]). The center's row sums are $(7, 11, 13)$, and these are the three *local norms* of the theory: $7 = N_{\mathbb{Z}[\omega]}(3+2\omega)$ (the Eisenstein/hexagonal CM norm = scalaron exponent), $13 = N_{\mathbb{Z}[i]}(3+2i)$ (the Gaussian/square CM norm = Δ_Q), and $11 = \binom{4}{0} + \binom{4}{1} + \binom{4}{2} = 1+4+6$ (the QBL boundary Fock count on the four μ_4 corners = $\dim S^+ - g_{\text{car}} = 16-5$). So $C = (\text{hex norm, boundary Fock count, square norm})$: the two exceptional CM rings ([verification/v222_cm_norm_duality.py]) and the seam boundary count meet in one operator center, the ring assignment rigid [E].

The Winding Line. On $R_s = R + s \mathbf{1} e_1^\top$ (the $R \rightarrow L$ channel) anchor area and torsion volume run *synchronously* for all s :

$$\boxed{\det R_s = 8 + 2s, \quad \det B_{R_s} = 16 + 4s = 2 \det R_s},$$

and the physical winding $s = 6 = |R^+(A_3)|$ is fixed by *each* of three closure conditions alone: $\text{tr } R_s = 15 = \dim A_3$, $\det R_s = 20 = \det L$, and the Coxeter lift $(R_s^\top a)_1 = 30 = h(E_8)$. The mechanism is the **cofactor seam normal** $n = c_2 \times c_3 = (5, -9, 6)$ (cross product of the second and third *columns* of R) — the common first adjugate row of R and L (columns 2, 3 are winding-invariant) — with

$$n \cdot \mathbf{1} = 2 = |\mathbb{Z}_2| \text{ (per-winding det increment)}, \quad n^\top R = (8, 0, 0), \quad n^\top L = (20, 0, 0) :$$

the normal annihilates the stable columns, so *generation one is the only open torsion channel* — the +6 winding is not sorted by hand. *Honest scope:* the winding line crosses the spectral *reality* threshold at $s^* \approx 2.83$ (not at 6), so reality is a *property* of the physical point — $\text{disc } \chi_L = 39200 = 2(10 \cdot 14)^2$, spectrum real positive — not its selector; and the corner $F = R+Q$ is an *audit* operator (the $F_4 \times G_2$ shadow: $\det B_F = 52 = \dim F_4$, $\sum B_F = 196 = (\dim G_2)^2$), not a physical transport.

The dual anchor ([verification/v134_dual_anchor.py]). The anchor's *inverse* response is the same on both ends of the winding line, and it is the Nariai root of the horizon paper:

$$\boxed{d := a^\top R^{-1} = a^\top L^{-1} = \left(-\frac{1}{2}, -\frac{1}{2}, 1\right)}, \quad d \cdot \mathbf{1} = 0, \quad d \cdot a = 1, \quad (1, 1, -2) = -2d.$$

The invariance is structural (Sherman–Morrison): $L = R + 6 \mathbf{1} e_1^\top$ makes a covector winding-invariant iff it kills $R^{-1} \mathbf{1} = \frac{1}{4}(1, 1, -1)$; the anchor does, while $\mathbf{1}$, e_1 and n do not ($\frac{1}{4}, \frac{1}{4}, -\frac{5}{2}$ — the membership is special). (d, n) is the dual normal pair of the flavor boundary: d the *sheet* normal (kills $\mathbf{1}$, dual to a , ∞ Nariai), n the *torsion* normal (first-generation channel). [E]; bridge reading [C].

Walls of the surface ([verification/v135_det_surface.py]). The surface determinant factorises,

$$\boxed{\det M(s, t) = 3s(t+1)(t+2) + t^2 + 5t + 8},$$

exposing two *iso-volume walls independent of s*: $t = -2 \Rightarrow \det = 2 = |\mathbb{Z}_2|$ and $t = -1 \Rightarrow \det = 4 = |\mu_4|$ — so $\det K = 4$ is a whole *layer* of the surface ($K = M(1, -1)$ lies on the μ_4 wall), not an isolated

hit — while the winding line $t = 0$ carries $\det = 6s + 8$, i.e. $(\det R, \det C, \det L) = (8, 14, 20)$, and the fourth corner closes the pattern with $\det F = \det M(1, 1) = 32 = 2^{g_{\text{car}}}$. The anchor block obeys $\det B_{M(s,0)} = 2 \det M(s, 0)$ *exactly* on the winding line and has its own $\mathbb{Z}_2$ wall, $\det B_{M(s,-2)} = -2$ ($t = -1$ is *not* a B -wall: $6s + 4$, recorded honestly). Row-budget axes: $\text{rowsum}(C) = (7, 11, 13)$, $\text{rowsum}(U) = (3, 3, 3)$ (democratic winding), $\text{rowsum}(V) = (1, 2, 3) = \text{Spec}(Q_+)$ verbatim; the centre budget $(7, 11, 13) = (\text{scalaron}, \|\text{Pl}(K)\|_1, \Delta_Q)$ is an audit match. Two anchor micro-identities: $e_1^2 + e_2^2 = 41 = 10b_1$ (the anchor Pythagoras for the abelian coefficient, companion to $\Delta_Y = 25 = 9+16$) and $p_1^2 + p_2^2 = 52 =$ the carrier coset root count $(40+12, [\text{verification/v128_graded_hull.py}])$. All [E]; the “2d surface with atom walls” reading [C].

The Q_+ grading is cohomology: one spectrum in four readouts [E]
 ([verification/v137_qplus_cohomology.py])

The sheet axis is not just numerology on V : it is the cohomology of the seam curve. $b_1(\mathbb{P} \setminus \mu_4) = 3 = N_{\text{fam}}$, with holomorphic basis $\omega_k = z^{k-1} dz / (z^4 - 1)$, $k = 1, 2, 3$ (regular at infinity; the trivial character is absent), and under the μ_4 deck $z \mapsto iz$

$$\omega_k \longmapsto i^k \omega_k, \quad \text{res}_\zeta \omega_k = \frac{\zeta^k}{4} :$$

$H^1 = \chi_1 \oplus \chi_2 \oplus \chi_3$ — each nontrivial μ_4 character exactly once, and the residue vector over the punctures is the character vector. The character degrees $(1, 2, 3)$ are exactly $\text{Spec}(Q_+)$ ($Q_+ = 3 \text{ cusp} + 1$, v69/v72), and the cusp weights $\{0, \frac{1}{3}, \frac{2}{3}\}$ underneath are $\text{spec}(A_3^*)$ (v115):

$$\text{MS weights} = \text{cusp classes} = H^1 \text{ characters} = \text{Spec}(Q_+) - 1 \text{ over } N_{\text{fam}}$$

— one spectrum in four readouts; and $(1, 2, 3)$ are the *exponents* of A_3 ($W(A_3) = S_4$, v117; $h(A_3) = 4 = |\mu_4|$): the Q_+ generation grading is the A_3 exponent grading, realised on the seam curve’s cohomology. *Typing*: all spectrum statements [E]; what remains of GATE.QGEO is the *naturality* of the identification — the canonical map from the H^1 character basis to the V -axis generation basis ($\text{rowsum}(V) = (1, 2, 3)$ verbatim, [verification/v135_det_surface.py]) — typed [C].

The canonical map exists and is rigid ([verification/v140_canonical_map.py]). The naturality residue collapses. Of the established μ_4 actions on generation space, the plain deck $U = \text{diag}(1, i, -i)$ has characters $(0, 1, 3)$ — an invariant line, missing 2 — and *cannot* match H^1 ; but the **sheet-twisted** deck $-U$ (characters $(2, 3, 1)$) and the cusp exponential i^{Q_+} (characters $(1, 2, 3)$) both carry the character set $\{1, 2, 3\}$ — exactly the H^1 set, by the same sign twist that builds the v118 hexagon. Both sides are multiplicity-free, so by Schur the equivariant isomorphism *exists and is unique* up to one scalar per character: the canonical map is well-defined — existence and uniqueness are no longer open. Under the i^{Q_+} -map the H^1 Euler degree pulls back to Q_+ itself; under the $-U$ -map to $Q_+ \circ \rho$ with ρ the (123) family rotation: the two candidates differ by *exactly one* $\mathbb{Z}_3$ rotation. *Typing*: character algebra, rigidity and grading transport [E].

The deck selection theorem: the $\mathbb{Z}_3$ choice is derived ([verification/v141_deck_selection.py]). The discrete choice does not survive contact with the integer model the suite already owns: the derived integer deck $G = T_A \Sigma$ ([verification/v97_sheet_conjugation_bridge.py]/[verification/v98_discriminant_dictionary.py]) has the $Q_+=1$ cusp line as an *exact* eigenline with eigenvalue $-1 = i^2$ — the self-conjugate character 2 — and the conjugate pair $\{1, 3\}$ on the E -plane. Of the three $\mathbb{Z}_3$ rotations only the sheet-twisted assignment $(2, 3, 1) = \text{chars}(-U)$ puts character 2 on the $Q_+=1$ line: **the geometric boundary deck is the sheet-twisted class; the cusp exponential i^{Q_+} is excluded** (same eigenvalue multiset $\{-1, \pm i\}$, wrong grading pairing — the (-1) -eigenlines sit on Q_+ degrees 1 vs 2). An explicit $Q_+=1$ -line-preserving conjugacy $SGS^{-1} = -U$ exists; under the relabeling $k \rightarrow -k$ only the plane orientation flips = the established sheet $\mathbb{Z}_2$ (the two Lagrangian glues, v92) — no new residual. Consequence: the H^1 Euler degree pulls back to $Q_+ \circ \rho$; cusp grading and Euler grading differ on generation space by exactly the one family rotation ρ = the coker Q triality (v72). *Typing*: selection, conjugacy and obstruction [E] (lattice-level theorem); what remains of GATE.QGEO is only the realisation premise the gate always carried (the integer D_4 model is the parabolic geometry, v69/v98) [C] — with *no* discrete freedom left.

The Möbius D_4 realisation: the premise at its floor ([verification/v146_moebius_d4.py]). The realisation premise itself now has every checkable part checked. The *full* dihedral symmetry group of the

seam curve, $\langle \delta: z \mapsto iz, \iota: z \mapsto 1/z \rangle = D_4$, acts on the explicit cohomology basis $\omega_k = z^{k-1}dz/(z^4-1)$ exactly as the integer model: $\delta^*\omega_k = i^k\omega_k$ (the deck, v137), and

$$\iota^*\omega_1 = \omega_3, \quad \iota^*\omega_2 = \omega_2, \quad \iota^*\omega_3 = \omega_1$$

— the exponent duality with parity +1 on the self-conjugate line and $\det = -1$, *no extra phases*. In the cusp basis the anchor-forced conjugation T_A is this permutation (fixes the character-2 line with +1, $\det = -1$ = the glue-swap parity), and Σ is the $\delta\iota$ class (−1 on the character-2 line, $\det = +1$ = glue-fix): the v98 reflection-class split lands exactly on the geometric reflection classes. *Typing*: the dihedral match [E]; with spectra (v137), cyclic part (v141) and reflections now all matched, the premise reduces to the module identification itself — “generation space is H^1 of the seam curve with its parabolic weights” [C] — with no remaining checkable mismatch.

Spectra and the determinant ladder $\rightarrow |W(D_5)|$ [E]

The two operators carry the compiler atoms in their characteristic polynomials,

$$\chi_Q = (t-1)(t^2-5t+3) [g_{\text{car}}, N_{\text{fam}}], \quad \chi_K = (t-1)(t^2-8t+4) [h(D_5), |\mu_4|], \quad (3)$$

with discriminants $\Delta_Q = 13 = |R(A_3)|+1$ and $\Delta_K = 48 = \Omega_{\text{adm}}$. Their determinants form a torsion ladder ($\text{coker} = \mathbb{Z}_3, \mathbb{Z}_4, \mathbb{Z}_8, \mathbb{Z}_{20}$),

$$(\det Q, \det K, \det R, \det L) = (3, 4, 8, 20), \quad (4)$$

$$\det Q \cdot \det K \cdot \det R \cdot \det L = 1920 = |W(D_5)| \quad [\text{E}],$$

tying flavor residues, mass powers, transport and the D_5 Weyl group (the same 1920 that appears in the horizon Hawking-power denominator). The sheet decomposition is sharp: $\text{Spec}(Q_+) = \{1, 2, 3\}$ are the A_3 exponents ($\det Q_+ = 6 = |R^+(A_3)|$), $\chi_{Q_-} = t(t^2-3)$ (a square root of N_{fam}), and $\chi_{K_-} = t(t^2-8)$ (a square root of $h(D_5)$). Finally the EM budget sits in the K anchor block: $a^\top K a = 41 = 10b_1$ with $a = (1, 1, 2)$, and $\det \begin{pmatrix} \mathbf{1}^\top K \mathbf{1} & \mathbf{1}^\top K a \\ a^\top K \mathbf{1} & a^\top K a \end{pmatrix} = \det \begin{pmatrix} 25 & 29 \\ 35 & 41 \end{pmatrix} = 10 = \mathcal{A}_\Lambda$. [verification/v10_projection_involution.py], [verification/v12_mass_generation_polynomials.py]

Mass volume and the EM budget are one anchor operation ([verification/v74_compiler_micro_lemmas.py]). The two diagonal entries of B_K are the two physical readouts: $\mathbf{1}^\top K \mathbf{1} = 25 = g_{\text{car}}^2$ is the grand *mass-volume* exponent ($\det M_{\text{SM}} \sim (\varphi_0^{\text{ret}})^{25}$), and $a^\top K a = 41 = 10b_1$ is the *EM budget*. Their difference is one generation,

$$a^\top K a - \mathbf{1}^\top K \mathbf{1} = 41 - 25 = 16 = \dim S^+$$

i.e. *EM budget = mass volume + one generation*. So the number 41 that the EM fixed point reads is not “40+1” but the anchor quadratic form of the mass-power matrix, tied to the mass volume g_{car}^2 by a single half-spinor $\dim S^+$.

Mass and transport are one pencil: $\det(K + xQ)$ carries the scalaron sequence [E]

Because $L = K + Q$, the mass-power matrix K and the transport matrix L are the two endpoints ($x = 0, 1$) of a *single* one-parameter compiler pencil $P(x) = K + xQ$, whose determinant is

$$\det(K + xQ) = 3x^3 + 7x^2 + 6x + 4, \quad (3, 7, 6, 4) = (N_{\text{fam}}, \Omega_{\text{adm}} - 10b_1, |R^+(A_3)|, |\mu_4|),$$

i.e. the coefficient sequence is N_{fam} , the *scalaron exponent* 7, the A_3 positive roots and the glue index — one curve linking flavor, the Starobinsky scalaron, the family and the glue. The pencil is anchored at four integer nodes by compiler atoms: $P(-1) = 2 = |\mathbb{Z}_2|$ (the $K-Q$ sheet endpoint, with $\det(K-Q) = |\mathbb{Z}_2|$, $\text{tr}(K-Q) = N_{\text{fam}}$), $P(0) = 4 = \det K = |\mu_4|$, $P(1) = 20 = \det L$, and $P(2) = 68 = 2p_5(a)$ (twice the anchor power sum $p_5 = 34$). [verification/v52_pencil_endpoints.py]

And the consecutive differences are the sheet→matter chain ([verification/v74_compiler_micro_lemmas.py]). The node values (2, 4, 20, 68) step by

$$P(0)-P(-1) = 2 = |\mathbb{Z}_2|, \quad P(1)-P(0) = 16 = \dim S^+, \quad P(2)-P(1) = 48 = \Omega_{\text{adm}}$$

i.e. $2 \rightarrow 16 \rightarrow 48$ — *sheet* $\rightarrow$ *one generation* $\rightarrow$ *three generations* — so the single determinant pencil already encodes the passage from the $\mathbb{Z}_2$ sheet to the one-family half-spinor to the full admissible matter packet. The (K, Q, L) operators are therefore not separate artefacts but one curve. The non-commutativity of the mass $\rightarrow$ transport step is a fixed G_2 -sized obstruction: $[K, L] = [L, Q] = [K, Q]$ (forced by $L = K + Q$) with $\chi_{[K, Q]}(t) = t^3 + 5t + 14$, reading $g_{\text{car}} = 5$ and $\dim G_2 = 14$ (audit-level [O]).
[verification/v37_plucker_anchor.py]

Operator-pencil geometry: the anchor plane as a small area in operator space [E]
([verification/v80_operator_pencil_geometry.py])

Read the four operators through the *anchor block* $B_M = \begin{pmatrix} \mathbf{1}^\top M \mathbf{1} & \mathbf{1}^\top M a \\ a^\top M \mathbf{1} & a^\top M a \end{pmatrix}$. **(i) Anchor Singularity Theorem.** On the pencil $P(x) = K + xQ$,

$$\det B_{K+xQ} = 9x^2 + 21x + 10 = (N_{\text{fam}}x + |\mathbb{Z}_2|)(N_{\text{fam}}x + g_{\text{car}}) = (3x + 2)(3x + 5) \quad [\text{E}],$$

so the oriented anchor area *degenerates* at exactly $x = -|\mathbb{Z}_2|/N_{\text{fam}} = -\frac{2}{3}$ and $x = -g_{\text{car}}/N_{\text{fam}} = -\frac{5}{3}$. The first zero is the Koide target / transport-gap base ($\lambda_2 = (\frac{2}{3})^6$); the second is the D_5/A_3 glue asymmetry. **So $\frac{2}{3}$ is not just a recurring fraction — it is the first singular point of the mass $\leftrightarrow$ transport anchor plane**, a structural *location* where an oriented area collapses. **(ii) Type checker.** The anchor-block determinants classify the operators: $\det B_Q = 9 = N_{\text{fam}}^2$ (family square), $\det B_K = 10 = \mathcal{A}_\Lambda$ (pair sector), $\det B_R = 16 = \dim S^+$ (one generation), $\det B_L = 40 = |R(D_5)|$ (carrier root budget) — each a determinant of a fixed block, not a loose integer. **(iii) Buildup chain.** The pencil coefficients $(3, 7, 6, 4)$ have partial sums $(3, 10, 16, 20) = (N_{\text{fam}}, \mathcal{A}_\Lambda, \dim S^+, \det L)$, so the scalaron exponent is the internal flavor transition $7 = \mathcal{A}_\Lambda - N_{\text{fam}}$.

Audit layer (exact values [E], decorative reading [C]). $D(x) = \det(K+xQ)$ and its derivatives at $x = -1, 0, 1, 2$ give $(2, 4, 20, 68), (1, 6, 29, 70), (-4, 14, 32, 50)$ (29 the top E_8 exponent, $32 = 2^{g_{\text{car}}}$, 50 the $A_4 \times A_4$ off-block); and the missing corner $F = R + Q$ has $\det B_F = 52 = \dim F_4$ with $248 = 52 + 14 + 26 \cdot 7$ — the genuine $E_8 \supset F_4 \times G_2$ branching $(52, 1) \oplus (1, 14) \oplus (26, 7)$ as an operator *shadow*. These are exact identities; their atlas/embedding significance is kept audit-level ([C]), since the coefficients are themselves atoms.

The Koide point is one branch of a double cover [E]
([verification/v81_singular_pencil_matrices.py])

The anchor-block determinant $\det B_{K+xQ} = (3x + 2)(3x + 5)$ is *quadratic*, so

$$y^2 = \det B_{K+xQ} = (3x + 2)(3x + 5)$$

is a 2:1 cover of the pencil line, *ramified exactly over its two zeros*. Hence the Koide/gap point $x = -|\mathbb{Z}_2|/N_{\text{fam}} = -\frac{2}{3}$ and the carrier point $x = -g_{\text{car}}/N_{\text{fam}} = -\frac{5}{3}$ are **the two branch points of one double cover**: *Koide is literally “the other side” of the carrier point*. The deck involution is $y \mapsto -y$ of degree 2 = $|\mathbb{Z}_2|$ — the sheet itself — which is the line-bundle reflection of the $\text{Spin}(10) \rightarrow \text{SO}(10)$ half-spinor cover the carrier lives on. Three rigid readings:

- **Branch locus = spine ends / family.** The ramification is at $-2/3$ and $-5/3$, i.e. $-(|\mathbb{Z}_2|, g_{\text{car}})/N_{\text{fam}}$ — the *two ends* of the spine $(2, 3, 4, 5)$ over N_{fam} .
- **Discriminant** $= N_{\text{fam}}^4 = 81$ (a perfect square $\Rightarrow$ rational, split branch points); $\sqrt{81} = 9 = N_{\text{fam}}^2 = \det B_Q$.
- **Separation** $= 1 = \text{one transport period}$ (the $K \rightarrow L$ step $x : 0 \rightarrow 1$); the deck-translation $x \mapsto x + 1$ carries the carrier point onto the Koide point.
- **Branch divisor = (scalaron, $\mathcal{A}_\Lambda$).** Sum of branch points $= -7/3 = -7/N_{\text{fam}}$, product $= 10/9 = \mathcal{A}_\Lambda/N_{\text{fam}}^2$; the integer labels $\{|\mathbb{Z}_2|, g_{\text{car}}\} = \{2, 5\}$ have sum $= 7$, prod $= 10 = \mathcal{A}_\Lambda$, diff $= 3 = N_{\text{fam}}$. **So the scalaron is the trace of the branch divisor** — its several readouts ($\Omega_{\text{adm}} - 10b_1, \mathcal{A}_\Lambda - N_{\text{fam}}, g_{\text{car}} + |\mathbb{Z}_2|$, the mixed anchor area) collapse to this one geometric origin.
- **The sheet lives on the cut.** The fibre $y^2 = \det B$ is $\mathcal{A}_\Lambda$ at K and $4\mathcal{A}_\Lambda$ at L , so transport multiplies y by $|\mathbb{Z}_2|$; the $\mathbb{Z}_2$ sheet endpoint $x = -1$ sits *between* the branch points where $y^2 = \det B_{K-Q} = -|\mathbb{Z}_2| < 0$ — on the imaginary branch cut joining Koide and the carrier.

At each branch point the pencil clears its 3-denominator into an integer compiler matrix. *Koide*

$C_{2/3} = 3K - 2Q$ reproduces the carrier glue: $\text{tr} = 15 = \dim A_3$, $\sum = 45 = \dim D_5$, $\det = 60 = D_{\text{start}}$, and the (now forced rank-one) anchor block $B_{C_{2/3}} = \begin{pmatrix} 45 & 55 \\ 63 & 77 \end{pmatrix} = \begin{pmatrix} g_{\text{car}} \\ 7 \end{pmatrix} \begin{pmatrix} N_{\text{fam}}^2 & 11 \end{pmatrix}$ with $\sum = 240 = |R(E_8)|$. Carrier $C_{5/3} = 3K - 5Q$ is the charge-neutral side ($\sum_{ij} = 0$) with $\chi(\lambda) = (\lambda + |\mathbb{Z}_2|)(\lambda^2 + \lambda + |R^+(A_3)|) = (\lambda + 2)(\lambda^2 + \lambda + 6)$, splitting off the sheet eigenvalue $-|\mathbb{Z}_2|$. So the recurring $\frac{2}{3}$ is not a quotient that “appears everywhere” — it is one ramification point of the mass $\leftrightarrow$ transport double cover. The sharper open question is then *why the leptonic source-to-pole transfer renormalises onto a branch point, and which sheet ($y \rightarrow \pm$) the parity selects* — a cleaner target than “why $2/3$ ”. (The D_5/A_3 and E_8 labels on the clearing matrices are exact; their per-entry atlas reading stays audit-level [C].) *Gravitational twin*: the same split N_{fam} -linear double-cover structure reappears in classical gravity — the Schwarzschild–de Sitter mass line has $\text{disc} \propto (1 - 3m)(1 + 3m)$ (branch points $\pm 1/N_{\text{fam}}$, separation $2/3$, deck = horizon swap), the Nariai roots are the traceless anchor $(1, 1, -2)$, and $2/3$ is the exact Nariai/de-Sitter entropy bound (Appendix H; [verification/v101_horizon_anchor.py]).

The Koide branch is a forced attractor, and the clean split is non-generic [E]/[C]
([verification/v82_koide_attractor_splitting.py])

This sharpens the open question above. Work in the branch coordinate $q = -3x$ (Koide $q = 2$, carrier $q = 5$). **(A) The source-to-pole map is forced, not postulated.** Any branch-divisor-preserving Möbius map of the pencil line that fixes *both* branch points is, in the cross-ratio coordinate $\rho = (q - 2)/(5 - q)$, exactly $\rho \mapsto \mu\rho$ — hence *unique* once μ is fixed. The multiplier is not a new input: $\mu = (2/3)^6$ is the subleading eigenvalue λ_2 of the *already-established* unique gapped boundary transfer operator (spectrum $\{1, (2/3)^6, (1/3)^6\}$, gap $6 \log \frac{3}{2} > 0$; [verification/v54_seam_horizon_keystones.py], [verification/v56_unique_attractor.py]), and the Koide branch $x = -2/3 = -|\mathbb{Z}_2|/N_{\text{fam}}$ is precisely its cusp weight $2/3 = \{0, 1, |\mathbb{Z}_2|\}/N_{\text{fam}}$ (the carrier maps to Koide under the deck period $q \mapsto q - N_{\text{fam}}$, the same cusp class). With $\mu = (2/3)^6$ the unique map is the integer Möbius map

$$F(q) = \frac{2(569q - 3325)}{665q - 3517}, \quad F(2) = 2, \quad F(5) = 5, \quad F'(2) = (2/3)^6 < 1, \quad F'(5) = (3/2)^6 > 1,$$

so Koide is the **attractor** and the carrier the **repeller**, and $q_n \rightarrow 2$ i.e. $x \rightarrow -2/3$ at the *same* rate that governs the SM flavor gap and horizon recovery. The three structural postulates needed for a Koide derivation thus collapse to *one* physical identification — *source-to-pole transfer = branch-preserving relaxation of the established transfer operator* — after which rate, uniqueness and direction all follow. That single identification stays [C]; the rest is [E]. **(B) The clean rational split is non-generic.** The splitting-type exponents $(2, 1, 1)$ (the anchor $a = (1, 1, 2)$) placed across the three family slots give three different covers: $(1, 1, 2)$ gives $\det B_{K+Q} = (3x + 2)(3x + 5)$, $\text{disc} = 81 = N_{\text{fam}}^4$ (*split*, rational labels $\{2, 5\}$); $(2, 1, 1)$ gives $(x + 2)(9x + 11)$, $\text{disc} = 49 = 7^2$ (*split*); $(1, 2, 1)$ gives $5x^2 + 10x + 3$, $\text{disc} = 40 = |R(D_5)| = \det B_L$, which is *not* a square — a **non-split** cover with Galois-conjugate branch points $-1 \pm \sqrt{10}/5$ centred exactly on the sheet $x = -1$ ($q = 3$), separation $2\sqrt{A_\Lambda}$. (Negative control, as for the cover itself: the R -pencil has $\text{disc} = 153$, also not a square.) So Koide and the carrier appearing as *separate rational* singularities is a special feature of the canonical anchor placement, not a generic one — a hardening of anchor-first. The physical fermion-sector identification of the three placements is not claimed and stays [C].

Branch kernels select the sectors; one deck involution [E]
([verification/v96_branch_kernel_selection.py], [verification/v97_sheet_conjugation_bridge.py])

The “which sheet does parity select” question above had a blocker: no *independently motivated* sector $\rightarrow$ sheet map. The cover supplies one itself. At each branch point the anchor block is **rank one**, with canonical *integer* kernels:

$$\begin{aligned} \text{Koide } x = -\frac{2}{3}: \quad w &= 11 \mathbf{1} - 9a = (2, 2, -7), \quad v = 7 \mathbf{1} - 5a = (2, 2, -3); \\ \text{carrier } x = -\frac{5}{3}: \quad w &= \mathbf{1}, \quad v = 8 \mathbf{1} - 7a = (1, 1, -6), \end{aligned}$$

— at the carrier point the block annihilates the *democratic vector itself*. **Collapse lemma [E]**: rank one forces $P(x_0)w \perp \text{span}\{\mathbf{1}, a\}$, hence proportional to the antisymmetric direction $(-1, 1, 0)$ of the anchor’s two *equal* slots — on both sides:

$$P(-\frac{2}{3})w = \frac{20}{3}(1, -1, 0), \quad P(-\frac{5}{3})w = \frac{2}{3}(-1, 1, 0), \quad v^\top P(x_0) \propto (-1, 1, 0) \text{ at both points.}$$

Reading the rows as the sectors (u, d, e) : *up and down are the deck-odd pair* ($f_u = -f_d$; convention-free, $f_u f_d < 0$ at both points), and the *lepton pairing vanishes* — the leptons sit *on* the ramification, consistent with Koide being a charged-lepton relation. Indeed the lepton pairing forms *are* the two $\det B$ factors: the **sector zero ladder** reads lepton $\rightarrow \{-\frac{2}{3}, -\frac{5}{3}\}$ (the branch points), up $\rightarrow -\frac{3}{2} = -N_{\text{fam}}/|\mathbb{Z}_2|$ *doubly* (the whole up-row of $P(-\frac{3}{2})$ is the odd direction $-\frac{1}{2}(1, -1, 0)$ — the previously unexplained $\text{GL}(2)$ rung), down $\rightarrow \{0 (= K), -\frac{9}{5} = -N_{\text{fam}}^2/g_{\text{car}}\}$. The deck translation flips the orientation $(+, -) \rightarrow (-, +)$ with magnitude ratio $10 = \mathcal{A}_\Lambda$. *Controls*: anchor $(2, 1, 1)$ moves the collapse to $(0, 1, -1)$; $(1, 2, 1)$ is non-split (disc 40), no rational kernels at all.

The bridge to the glue chiralities ([[verification/v97_sheet_conjugation_bridge.py](#)]). The two E_8 glues are swapped by *either single* spinor conjugation and fixed by the simultaneous one ([[verification/v92_glue_uniqueness.py](#)]); on the family side, conjugation negates the cusp classes (0 fixed, $\frac{1}{3} \leftrightarrow \frac{2}{3}$ on the Q_+ eigenspaces $e_1 = (0, 0, 1)$, $e_2 = (0, 1, 2)$, $e_3 = (1, 0, 0)$). That conjugation has a *unique anchor-compatible* integer realisation: $T\mathbf{1} = \mathbf{1}$ and $Ta = a$ both force the same scale, giving

$$T_A = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 2 & -2 & 1 \end{pmatrix}, \quad T_A^2 = I, \quad \det T_A = -1, \quad \text{and} \quad a = e_2 + e_3$$

— *the anchor is the conjugation-symmetrised vector of the swapped pair* (the two alternative pairings admit *no* anchor-compatible scale). On the quotient by the anchor plane, $T_A = -1$ exactly like the sector transposition: **one deck action where the cover lives**. The parity matches the glue arithmetic ($\det = -1$ single/swapping vs $\det \Sigma = +1$ simultaneous/fixing); $\langle T_A, \Sigma \rangle$ closes the *same* $D_4 = \mathbb{Z}_4 \times \mathbb{Z}_2$ as the Q -geometry, with $\chi_{T_A \Sigma} = (t+1)(t^2+1)$ identical to the μ_4 homology matrix; and the **sheet-index lemma**: the generation- μ_4 and the homology- μ_4 are rationally but *not* $\text{GL}(3, \mathbb{Z})$ -conjugate — every intertwiner has *even* determinant, minimum $|\det| = 2 = |\mathbb{Z}_2|$ (one sheet doubling; independent of $\det Q = N_{\text{fam}}$).

The dictionary is derived, not assumed ([[verification/v98_discriminant_dictionary.py](#)]). The generation space carries an *integer* μ_4 : $G := T_A \Sigma$ acts on the cusp basis as

$$G e_1 = -e_1, \quad G e_3 = e_2, \quad G e_2 = -e_3 \quad (G^4 = I, \quad \chi_G = (t+1)(t^2+1)),$$

i.e. the $B_1 \oplus E$ decomposition of the D_4 -equivariant Q -geometry realised *integrally*: e_1 spans the B_1 line (G -eigenvalue -1 = the self-conjugate $\mathbb{Z}_4$ class 2), and $\text{span}\{e_2, e_3\}$ is the E -plane (eigenvalues $\pm i$ = the conjugate class pair $\{1, 3\}$). **So the cusp-0 line is the self-conjugate discriminant line, and the swapped cusp pair is the conjugate class pair** — the dictionary “ Q_+ grading = A_3 discriminant grading” holds inside the model. Consistently: $T_A G T_A^{-1} = G^{-1}$ (the discriminant conjugation $k \mapsto -k$), $q_{A_3}(1) = q_{A_3}(3) = \frac{3}{8}$ (equal on the swapped pair) with $q_{A_3}(2) = \frac{1}{2}$ fixed, and the two D_4 reflection classes separate exactly the glue arithmetic: $T_A e_1 = +e_1$, $\det = -1$ (glue-swapping) vs $\Sigma e_1 = -e_1$, $\det = +1$ (glue-fixing), with $\Sigma = T_A G$. *Honest scope*: the sheet question therefore carries **no separate [C] residual any more** — what remains is exactly the already-open Q -geometry realisation gate (GATE.QGEO, the parabolic construction that realises the documented Q on $\mathbb{P}^1 \setminus \mu_4$); conditional on that established typing, the sector $\rightarrow$ sheet selection is *closed*: up/down are the deck-odd pair, the leptons sit on the ramification, and the deck involution is the A_3 -side spinor conjugation realised as T_A .

Grand Mass Volume: the absolute mass-scaling normalisation [E]

The individual c_q are a gauge, but the *absolute* normalisation of the mass spectrum is fixed by the sector mass-matrix *determinants* — each a clean φ_0^{ret} -power whose exponent is the K -row sum:

$$\det M_{\text{up}} \sim (\varphi_0^{\text{ret}})^6 = (\varphi_0^{\text{ret}})^{|R^+(A_3)|}, \quad \det M_{\text{down}} \sim (\varphi_0^{\text{ret}})^9 = (\varphi_0^{\text{ret}})^{N_{\text{fam}}^2}, \quad \det M_{\text{lep}} \sim (\varphi_0^{\text{ret}})^{10} = (\varphi_0^{\text{ret}})^{\mathcal{A}_\Lambda},$$

$$\det M_{\text{SM}} \sim (\varphi_0^{\text{ret}})^{6+9+10} = (\varphi_0^{\text{ret}})^{25} = (\varphi_0^{\text{ret}})^{g_{\text{car}}^2} = (\varphi_0^{\text{ret}})^{\Delta_Y}$$

the *Grand Mass Volume*. The Q -row sums $(4, 5, 6) = (|\mu_4|, g_{\text{car}}, |R^+(A_3)|)$ total $15 = \dim A_3$, and $25 + 15 = 40 = \sum L = |R(D_5)|$ (mass volume + shift volume = transport budget). So the absolute scaling is an [E] determinant identity (each sector = its K -row sum); the only genuine scale input is the overall v_{geo} , not a per-coefficient freedom. [[verification/v46_grand_mass_volume.py](#)]

Uniqueness (not fitted). K and Q are the *unique* nonnegative-integer 3×3 matrices with their row sums, column sums, characteristic polynomial and per-sector monotone hierarchy — verified by

exhaustive enumeration [verification/v11_unique_KQ.py]. They are forced by small compiler budgets plus spectrum, not chosen.

Theorem Q — the Σ -split as a parabolic projection (sharper uniqueness) [E]/[C]

The sheet involution $\Sigma = \text{diag}(1, -1, -1)$ splits Q into two parabolic pieces:

$$Q_+ = \frac{1}{2}(Q + \Sigma Q \Sigma) = \begin{pmatrix} 3 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 2 & 1 \end{pmatrix}, \quad Q_- = \frac{1}{2}(Q - \Sigma Q \Sigma) = \begin{pmatrix} 0 & 1 & 0 \\ 3 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

$$\chi_{Q_+}(t) = (t-1)(t-2)(t-3) \Rightarrow \text{Spec}(Q_+) = \{1, 2, 3\} \text{ (} A_3 \text{ exponents),}$$

$$\chi_{Q_-}(t) = t(t^2 - 3) \Rightarrow Q_-^2|_{\text{supp}} = 3 = N_{\text{fam}}.$$

Uniqueness: Q is the *unique* nonnegative-integer matrix with rows = (4, 5, 6), cols = (9, 5, 1), $\chi_{Q_+} = (t-1)(t-2)(t-3)$ and $\chi_{Q_-} = t(t^2 - 3)$ (exhaustive enumeration, [verification/v50_q_geometry.py]). So Q_+ is the A_3 exponent grading, Q_- the μ_4 sheet coupling, and $K = R + Q\Sigma$ is the *second Σ -shadow of the same parabolic flavor operator*, not an imported mass trick.

D_4 -equivariant derivation ([verification/v69_d4_q_geometry.py]) — **gate advanced** [C]/[E]. The four punctures $\mu_4 = \{1, i, -1, -i\}$ are the vertices of a square with symmetry $D_4 = \mathbb{Z}_4 \rtimes \mathbb{Z}_2$ (the 90° rotation $z \mapsto iz$ is μ_4 ; the reflection is the sheet parity Σ). The $\mathbb{Z}_4$ 4-cycle on the punctures acts on $H_1(\mathbb{P}^1 \setminus \mu_4) = \mathbb{Z}^4 / (\text{sum}=0) = \mathbb{Z}^3$ (the three families) with eigenvalues $\{i, -1, -i\}$, and the D_4 permutation rep decomposes as $A_1 \oplus B_1 \oplus E$, so $H_1 = B_1 \oplus E$. On this,

$$Q_+ = 3(\text{cusp weights } \{0, \frac{1}{3}, \frac{2}{3}\}) + 1 = \text{diag}(1, 2, 3),$$

$$Q_- = E\text{-block coupling } \sqrt{N_{\text{fam}}} \text{ (} B_1 \text{ in the kernel),}$$

which reproduce $\chi_{Q_+} = (t-1)(t-2)(t-3)$ and $\chi_{Q_-} = t(t^2 - 3)$ exactly. So the Σ -split is the $D_4 = \mathbb{Z}_4 \rtimes \mathbb{Z}_2$ structure ($\mathbb{Z}_4 = \mu_4$ gives the three eigenspaces and Q_+ ; the $\mathbb{Z}_2$ reflection is Σ , giving the even/odd split). **Typing:** the spectra and the Σ -split are now *derived* from the D_4 -equivariant geometry [E]; the residual is only that the cusp weights $\{0, \frac{1}{3}, \frac{2}{3}\}$ are the established monodromy exponents (v61) and the explicit integer Q still needs the lattice datum — now *sharpened to a single named invariant* ([verification/v70_q_integer_lift.py]): in the μ_4 -puncture homology basis the $\mathbb{Z}_4$ rotation is the *unimodular* $R = \begin{pmatrix} 0 & 0 & -1 \\ 1 & 0 & -1 \\ 0 & 1 & -1 \end{pmatrix}$ (det = -1, eigenvalues $\{-1, i, -i\}$), while the transport has $\boxed{\det Q = 3 = N_{\text{fam}}}$ (SNF $\text{diag}(1, 1, 3)$). So the integer lift is exactly the parabolic-degree-0 embedding carrying the extra $\det = N_{\text{fam}}$ (the family multiplicity) that D_4 alone (unimodular) does not supply — one named lattice invariant, no longer a diffuse “derive Q ”.

And that invariant is the cusp order itself ([verification/v72_q_det_from_cusp.py]). The extra $\det Q = N_{\text{fam}}$ is *not* an independent lattice input: $\det Q = |\text{coker } Q| = 3$ equals the *order of the cusp monodromy class* (eigenvalues $\{1, \omega, \omega^2\}$, $\omega^3=1$) = the common *denominator* of the very cusp weights $\{0, \frac{1}{3}, \frac{2}{3}\}$ that fix $\text{Spec}(Q_+)$. So $\det Q = N_{\text{fam}}$ comes from the *same* cusp datum as the spectrum, and $\text{coker } Q = \mathbb{Z}/N_{\text{fam}}$ is the deck group of the degree- N_{fam} triality cover defining the family space — the integer lift is derived from the geometry, not posited ([E] chain; the $\text{coker}=\text{deck-group}$ identification is the standard non-abelian Hodge reading [E]).

Status: Q -geometry derived (spectra, Σ -split and integer lift) [E]

What is *proved* ([E]): the algebraic identities $K = R + Q\Sigma$, $L = R + Q(I + \Sigma)$, the spectra, the 1920 determinant ladder, the anchor block — all machine-checked. **Newly advanced** ([verification/v69_d4_q_geometry.py]): the D_4 -equivariant origin of the Q -spectra and the Σ -split is now derived — Q_+ is the $3w+1$ image of the cusp weights $w \in \{0, \frac{1}{3}, \frac{2}{3}\}$ on the three $\mu_4 = \mathbb{Z}_4$ eigenspaces of $H_1(\mathbb{P}^1 \setminus \mu_4)$, and Q_- is the E -block coupling $\sqrt{N_{\text{fam}}}$ on $H_1 = B_1 \oplus E$, with Σ the $\mathbb{Z}_2$ reflection of $D_4 = \mathbb{Z}_4 \rtimes \mathbb{Z}_2$. So the φ_0^{ret} -mass ladder and the transport are two Σ -shadows of one D_4 -equivariant parabolic operator. **Integer lift closed to the cusp order** ([verification/v72_q_det_from_cusp.py]): the one remaining lattice datum $\det Q = N_{\text{fam}}$ is the order of the cusp class = the denominator of the same cusp weights, i.e. the *same* geometry that fixes the spectrum ($\text{coker } Q = \mathbb{Z}/N_{\text{fam}} = \text{triality deck group}$). So the Q, Σ algebra is now an [E] layer whose spectra, Σ -split and integer lift are all read from the order- N_{fam} cusp structure; the only standing input is the established fact $N_{\text{fam}} = 3 = \text{rank } A_3 = \dim H_1$.




Figure 2: Gauge running from the carrier content (two-loop SM RGEs; the β -functions are confirmed by the external generator PyR@TE, `[verification/v159_pyrate_gauge_crosscheck.py]`). The $U(1)_Y$ slope is the abelian index $b_1 = 41/10 = 10^{-1}(g_{\text{car}}2^{g_{\text{car}}-2}+1)$; the strong coupling reaches confinement, $\alpha_3^{-1} \rightarrow 0$ at $\Lambda_{\text{QCD}} \approx 0.4$ GeV (`[verification/v164_qcd_scale.py]`); and the Standard Model does not unify (the E_8 here is the audit hull, not a gauge group).

6.1 Closing the audit gates: is $M=41$, K , Q derived or imported?

A reviewer asks whether the compiler *derives* its objects or silently *imports* them. We answer the three flavor/EM gates explicitly (`[verification/v13_open_gates.py]`); each is either closed or reduced to a named geometric statement, with the honest residual flagged.

Gate 1 (closed): the EM budget $M=41$ is the $U(1)$ index, not a fit [E]

The transport budget that enters the EM closure is *forced*:

$$M = \sum_{f,j} L_{f,j} + N_{\Phi} = 40 + 1 = 10b_1 = 41, \quad \frac{4}{5}M c_3^6 = 8b_1 c_3^6, \quad (5)$$

so the log-coefficient of $F_{U(1)}$ is $8b_1 c_3^6$ with b_1 the *same* $U(1)$ hypercharge index derived from g_{car} ($10b_1 = g_{\text{car}}2^{g_{\text{car}}-2}+1 = 41$). M is therefore not an independent EM knob: it is the flavor transport budget = the abelian index, both equal to 41 from the carrier alone. An *independent* third-party cross-check confirms this: feeding the carrier field content into the textbook one-loop formula gives the full gauge vector $(b_1, b_2, b_3) = (\frac{41}{10}, -\frac{19}{6}, -7)$, and the external RGE generator PyR@TE 3 (`arXiv:2007.12700`) writes $\beta_{g_1} = (\frac{41}{10})g_1^3$ verbatim, with $10b_1 = 40_{(\text{ferm. } U(1) \text{ index})} + 1_{(\text{Higgs})} = 41$ reproducing this box exactly — so the F_{transfer} gauge input is not a free knob (`[verification/v159_pyrate_gauge_crosscheck.py]`).

Gate 2 (forced): K and Q are not read off the masses [E]

K (the φ_0^{ret} -mass-exponent matrix) and Q are the *unique* nonnegative integer matrices with their sector/generation budgets (themselves compiler numbers: rows $(6, 9, 10) = (|R^+(A_3)|, N_{\text{fam}}^2, \mathcal{A}_{\Lambda})$, columns $(13, 8, 4) = (|R(A_3)|+N_{\Phi}, h(D_5), |\mu_4|)$) and their characteristic polynomials, subject to per-sector monotonicity — proved by exhaustive enumeration (`[verification/v11_unique_KQ.py]`). The mass exponents are thus *fixed by compiler data*, not fitted to the measured masses.

Gate 3 (sharpened): the geometric origin of Q reduces to two named pieces [C]

Under the sheet involution Σ , $Q = Q_+ + Q_-$ splits exactly, and the two parts carry pure A_3/μ_4 data:

$$\text{Spec}(Q_+) = \{1, 2, 3\} = A_3 \text{ exponents}, \quad \det Q_+ = 6 = |R^+(A_3)|; \quad Q_-^2|_{\text{supp}} = N_{\text{fam}} = 3. \quad (6)$$

So Q_+ is the A_3 exponent grading and Q_- a sheet square-root of N_{fam} . The open “derive Q from geometry” target therefore reduces to two *specific, finite* identifications on $\mathbb{P}^1 \setminus \mu_4$: (i) Q_+ = the A_3 Hodge/exponent grading of H^1 , (ii) Q_- = the μ_4 sheet-monodromy coupling. With those, $K = R + Q\Sigma$ makes the mass ladder a *consequence* of the flavor compiler. *Honest residual*: the sector/generation budget *assignment* and the exact realisation of $Q_{\pm}$ on the parabolic bundle are the remaining geometric steps; the cosmology pivot $N_{\star}$ stays a *reheating input* ([C]), not a compiler output.

Gate 4 (new): the solar angle from a shared dual anchor [E]/[C]

The residue matrix R and the word-length matrix $L = R + 6\mathbf{1}e_1^{\top}$ (a *rank-one* winding deformation, $L - R = [[6, 0, 0], [6, 0, 0], [6, 0, 0]]$) act *identically* on the dual of the anchor $a = (1, 1, 2)$:

$$a^{\top} R^{-1} = a^{\top} L^{-1} = (-\tfrac{1}{2}, -\tfrac{1}{2}, 1) \quad [\text{E}].$$

Mechanism ([verification/v74_compiler_micro_lemmas.py]). This is *not* a coincidence: since $a^{\top} R^{-1}\mathbf{1} = 0$, the Sherman–Morrison correction for the rank-one update $L = R + 6\mathbf{1}e_1^{\top}$ has numerator $a^{\top} R^{-1}(6\mathbf{1}) = 0$, so $a^{\top} L^{-1} = a^{\top} R^{-1}$ *exactly* — the winding deformation is *invisible* to the dual anchor. So the gen $1 \leftrightarrow 2$ TBM symmetry vector (the two equal entries; the 1 marks the third-generation anchor) is *stable* under transport, which is the structural reason the solar texture survives. This ties the solar misalignment to the seam. **New: the coefficient is the A_3 family-lattice discriminant norm.** The misalignment is

$$\varepsilon = q(A_3)\varphi_0 = \tfrac{3}{4}\varphi_0, \quad q(A_3) = \tfrac{3}{4} = \frac{N_{\text{fam}}}{|\mu_4|},$$

the *same* $q(A_3)$ that satisfies the E_8 glue condition $q(D_5) + q(A_3) = \frac{5}{4} + \frac{3}{4} = 2$. Its *leading* term is exactly $q(A_3) \cdot \frac{1}{6\pi} = \frac{1}{8\pi} = c_3$, and the TBM sum-rule gives

$$\sin^2 \theta_{12} = \tfrac{1}{3} - \tfrac{2}{3}\varepsilon = \tfrac{1}{3} - \tfrac{\varphi_0}{2} = 0.30675 \quad (\text{NuFIT 6.0: } 0.307),$$

with $\sin^2 \theta_{13} = e^{-5/6}\varphi_0$, $\delta_{CP} = 4\pi/3$, and a $\mu\tau$ -symmetric atmospheric sector $\theta_{23} \rightarrow 45^\circ$ (the octant is *not* selected; current global fits leave it open). The solar angle is thus a *reduced* readout whose coefficient is now *geometric*: $\varepsilon = q(A_3)\varphi_0$, the A_3 glue-norm. The residual is only *why* the misalignment equals the A_3 discriminant norm ([C]). [verification/v16_solar_dual_anchor.py] [verification/v21_solar_product_quark.py]

Scale tag. These angles are stated as dimensionless seeds; the F_{transfer} running between the seam scale and M_Z is governed only by the charged-lepton term $\frac{3}{2}Y_e Y_e^{\dagger} Y_e$ (PyR@TE SM RGEs), i.e. by $y_{\tau}^2 \sim 10^{-4}$, so $|\Delta \sin^2 \theta_{ij}| \lesssim 10^{-4}$ — two to three orders below current precision. The lepton-mixing seeds are therefore *scale-independent* F_{transfer} outputs (seam value = measured value), unlike the y_t^2 -driven quark mass ratios, which carry a scale tag ([E]/[C], [verification/v163_rg_stability_flavor.py]).

Inverse Anchor Normalisation Theorem [E] ([verification/v79_review_identities.py])

The dual-anchor identity is one row of a sharper invariance. On the *inverses* of the flavor operators the all-ones vector reads the reciprocal compiler atoms,

$$\mathbf{1}^{\top} Q^{-1} \mathbf{1} = \tfrac{1}{3} = \tfrac{1}{N_{\text{fam}}}, \quad \mathbf{1}^{\top} R^{-1} \mathbf{1} = \mathbf{1}^{\top} K^{-1} \mathbf{1} = \tfrac{1}{4} = \tfrac{1}{|\mu_4|}, \quad \mathbf{1}^{\top} L^{-1} \mathbf{1} = \tfrac{1}{10} = \tfrac{1}{\mathcal{A}_{\Lambda}},$$

and the anchor is *self-dual* under the physical (residue/mass/transport) operators:

$$a^{\top} R^{-1} a = a^{\top} K^{-1} a = a^{\top} L^{-1} a = 1 \quad (\text{while } a^{\top} Q^{-1} a = \tfrac{7}{3}).$$

So the passage mass-powers $K \rightarrow$ residues $R \rightarrow$ transport L leaves the anchor plane *exactly* normalised — the anchor is its own dual under physical evolution — and the exception $a^{\top} Q^{-1} a = \frac{7}{3}$ pins the self-duality to precisely the residue/mass/transport triple, not the pure Σ -difference Q . [E]

7 Mixing: CKM and PMNS from holonomy

Quantity	Formula	Prediction	PDG
s_{12}^{CKM}	$\lambda_C = \sqrt{\varphi_0^{\text{ret}}(1 - \varphi_0^{\text{ret}})}$	0.2244	0.2245
s_{23}^{CKM}	$\varphi_0^{\text{ret}}/(1 + \lambda_C)$	0.0434	~ 0.041
s_{13}^{CKM}	$\lambda_C^3/3$	0.00377	~ 0.0038
δ_{CKM}	$\frac{\pi}{3} + 3\lambda_C^2$ (leading, [C])	1.198 rad	~ 1.14
$\sin^2 \theta_{13}^{\text{PMNS}}$	$\varphi_0^{\text{ret}} e^{-5/6}$ (UV shadow)	0.0231	~ 0.0222

The cubic rule $|V_{ub}| = |V_{us}|^3/3$ and the CP phase $\delta = \arg \text{tr} \Pi_\Delta$ (oriented holonomy triangle) are direct consequences of the *one* family connection ∇_F^* — the same one that generates the word-lengths in the flavor theorem.

New: the CP sector falls out of the closed CKM (not separately fitted)

The four closed entries $s_{12}, s_{23}, s_{13}, \delta$ already *determine* the whole CP sector — quantities not previously written down:

$$J = c_{12}c_{23}c_{13}^2 s_{12}s_{23}s_{13} \sin \delta = 3.33 \times 10^{-5} \text{ (PDG 3.08)}, \quad \beta = 22.1^\circ \text{ (PDG 22.2)},$$

$$\gamma = 68.6^\circ \text{ (PDG 65.7)}, \quad \alpha = 89.2^\circ \text{ (PDG } \sim 85),$$

together with the full magnitudes $|V_{cb}| = 0.0434$, $|V_{td}| = 0.0091$, $|V_{ts}| = 0.0426$. The Jarlskog invariant and the unitarity-triangle angles are thus *leading-order readouts* of the closed CKM skeleton: the CP phase is the leading closed form $\delta = \frac{\pi}{3} + 3\lambda_C^2$ (its higher-order C_6 -transport correction is *not* in closed form), so the few-percent offsets ($J=3.33$ vs 3.08, $\gamma=68.6^\circ$ vs 65.7°) are expected and the CP sector is typed **[E]/[C]**, *not* **[E]** — only β happens to be near-exact. (To upgrade to **[E]** the higher-order δ correction Δ_Δ must be given explicitly.)

7.1 Downstream flavor readout: the rare kaons $K \rightarrow \pi \nu \bar{\nu}$

The closed CKM point is not an end state — it feeds the theoretically cleanest flavor-changing neutral-current probe, the rare decays $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and $K_L \rightarrow \pi^0 \nu \bar{\nu}$. These are *not* a new compiler power: the only TFPT-specific content is the CKM point above; the short-distance loop functions (X_t , P_c , $\delta P_{c,u}$, $\kappa_\pm$, Brod–Gorbahn–Stamou) are standard external electroweak/QCD input. Inserting the exact TFPT products $\lambda_t = V_{ts}^* V_{td}$, $\lambda_c = V_{cs}^* V_{cd}$ (equivalently the exact apex $(\bar{\rho}, \bar{\eta}) = (0.1374, 0.3509)$) gives [\[verification/v202_rare_kaon.py\]](#)

$$\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = 9.45 \times 10^{-11}, \quad \text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) = 3.33 \times 10^{-11},$$

typed **[C]** (the branching ratios are not parameter-free). The neutral mode is purely CP-violating, $\text{BR}(K_L) \propto (\Im \lambda_t)^2 \propto \bar{\eta}^2$, so it is a direct probe of the holonomy phase δ_{CKM} ; the charged mode carries a sizeable charm admixture and tests the overall CKM norm. The model-independent Grossman–Nir ratio is $\text{BR}(K_L)/\text{BR}(K^+) = 0.352 \ll 4.3$, safely inside the isospin corridor. **Dated kill test [X]**: a stable NA62 $\text{BR}(K^+)$ *outside* $[7, 12] \times 10^{-11}$, or a KOTO-II $\text{BR}(K_L)$ off the predicted Grossman–Nir point, breaks the TFPT flavor bridge for this sector (the compiler core is untouched). The current NA62 value $(13.0_{-3.0}^{+3.3}) \times 10^{-11}$ sits $\sim 1.2\sigma$ above the prediction — consistent today; the $\sim 2\sigma$ δ_{CKM} tension (**v88**) makes the corridor a genuinely discriminating, live falsification channel.

Weak mixing from D_5 . Because the carrier C^+ is the $D_5 = \mathfrak{so}(10)$ half-spinor, the weak mixing inherits the canonical $\mathfrak{so}(10)$ unification value $\sin^2 \theta_W = \frac{3}{8} = N_{\text{fam}}/h(D_5)$ at the carrier scale, running to 0.231 at M_Z (standard). **[E]** (structural) / **[C]** (running).

Honest note on the search. A generic “spine-ratio” scan ($\sin^2 \theta_W$, α_s , the Ω ratios, N_{eff} , m_H/v) finds $\leq 2\%$ hits abundantly — which is *weak* evidence, since $O(1)$ numbers are densely covered

by ratios of $\{8, \dots, 60\}$. We therefore count only *named* invariants (like $3/8 = N_{\text{fam}}/h(D_5)$ or the exact endpoints), not generic ratios.

8 Neutrinos

The neutrino sector is the Majorana branch of the same construction: $U_{\text{PMNS}} = U_{e,L}^\dagger U_{\nu,L}$.

Quantity	Origin	Value
$\sin^2 \theta_{13}$	$\varphi_0^{\text{ret}} e^{-5/6}$ (UV), closure 0.02311	≈ 0.0222
$\sin^2 \theta_{12}$	primary (freeze): $\frac{1}{3} - \frac{\varphi_0^{\text{ret}}}{2}$ (seed, $\varepsilon = q(A_3)\varphi_0^{\text{ret}}$); two derived variants in the named box	0.306747 (NuFIT 6.0 0.307)
m_2/m_3	$\sim \pi\varphi_0^{\text{ret}} = (1 - \gamma) + \frac{3}{256\pi^3}$	≈ 0.167
Σm_ν	closure readout (normal ordering)	5.88×10^{-2} eV
$m_{\beta\beta}$	Majorana effective mass ($0\nu\beta\beta$)	1.52×10^{-3} eV
δ_{CP}^ν	phase lattice	240°
M_R	heavy scale (type-I seesaw)	$\approx 1.3 \times 10^{15}$ GeV

The solar angle θ_{12} is fixed by the neutrino transport matrix $U_{\text{PMNS}} = U_{e,L}^\dagger U_{\nu,L}$. The **single frozen prediction** is the seed value $\sin^2 \theta_{12}^{\text{seed}} = \frac{1}{3} - \frac{\varphi_0^{\text{ret}}}{2} = 0.306747$ (misalignment $\varepsilon = q(A_3)\varphi_0^{\text{ret}} = \frac{3}{4}\varphi_0^{\text{ret}} = c_3 + 36c_3^4$ — leading term c_3 , fourth-order puncture correction exact — conditional on the seam-misalignment lemma); the seam-leading 0.306808 and the full non-linear 0.307020 are *derived variants* of the same texture, *not* alternative predictions (named box below). The older democratic $\frac{1}{3}$ and QLC $\frac{\pi}{4} - \theta_{12}^{\text{CKM}}$ readings are superseded by this seed closure. The ratio $m_2/m_3 \sim \pi\varphi_0^{\text{ret}}$ carries the carrier complement $1 - \gamma = 1/6$ — neutrino and charged sectors read the same seed. The heavy scale $M_R \sim 10^{15}$ GeV links the seesaw bridge to the inflation/reheating scale (the scalaron-reheating chain, **v86**).

M_ν as the type-I seesaw of D_F (`[verification/v263_mnu_from_df.py]`). The texture above is no longer only a posited matrix: the light mass matrix is the seesaw of the finite Dirac operator's own blocks, $M_\nu = -m_D M_R^{-1} m_D^\top$ (Dirac m_D , Majorana M_R from D_F , **v252**). **[E]** it is symmetric and seesaw-suppressed, and a $\mu\tau$ -symmetric (m_D, M_R) gives a $\mu\tau$ -symmetric M_ν (the texture is in the image of the seesaw map); **[C]** diagonalising it returns $\theta_{23} = 45^\circ$ and $\sin^2 \theta_{12} = \frac{1}{3} - \frac{\varphi_0^{\text{ret}}}{2}$ under the seam misalignment, reproducing **v9**. Honestly open: the leading texture gives $\theta_{13} = 0$ (the physical $\sin^2 \theta_{13} \sim \varphi_0^{\text{ret}} e^{-5/6}$ is a separate readout, **[O]**), the CP *magnitude* $\delta_{\text{CP}} = 240^\circ$ is the μ_6 /triality phase (**v231/v233**, not from M_ν), and the absolute ν -mass scale (M_R) is the free seesaw scale — so this is an *operator* advance (PMNS is now a seesaw object), *not* a closure of full PMNS dynamics.

The reactor-angle exponent is the carrier trace (`[verification/v268_theta13_carrier_trace.py]`). The reactor angle is a *separate* channel from the seesaw M_ν : its value $\sin^2 \theta_{13} = \varphi_0^{\text{ret}} e^{-5/6} = 0.0231$ (**v16**) now has its *exponent* closed exactly — $\gamma = 5/6 = \text{tr}_E Y^2$, the carrier hypercharge-squared trace ($Y = \text{diag}(-\frac{1}{3}, -\frac{1}{3}, -\frac{1}{3}, \frac{1}{2}, \frac{1}{2})$, the roots of $6Y^2 - Y - 1 = 0$; complement $1 - \gamma = \frac{1}{6}$ the neutrino ratio). **[E]** this is its own carrier-trace channel: the $\mu\tau$ -breaking $\varepsilon = \frac{3}{4}\varphi_0^{\text{ret}}$ that fixes θ_{12} induces only $\sin^2 \theta_{13} \sim \varepsilon^2 \approx 1.6 \times 10^{-3} \ll 0.023$, so θ_{13} is *not* the seesaw $\mu\tau$ breaking. **[O]** the CP *magnitude* and the absolute ν -mass scale remain the open PMNS items.

The full PMNS matrix and the CP strength (`[verification/v270_pmns_jarlskog_assembly.py]`). The four channels — $\sin^2 \theta_{12} = \frac{1}{3} - \frac{\varphi_0^{\text{ret}}}{2}$ (**v9/v263**), $\sin^2 \theta_{23} = \frac{1}{2}$ (**v9/v263**), $\sin^2 \theta_{13} = \varphi_0^{\text{ret}} e^{-5/6}$ (**v16/v268**) and $\delta = 4\pi/3$ (**v231/v233**) — assemble into *one* exactly-unitary complex matrix $U = R_{23} U_{13}(\delta) R_{12}$. **[E]** the assembly is unitary; **[C]** the leptonic Jarlskog $J_{\text{PMNS}} = \Im(U_{e1} U_{\mu 2} U_{e2}^* U_{\mu 1}^*)$, i.e. $s_{12} c_{12} s_{23} c_{23} s_{13} c_{13}^2 \sin \delta = -0.02965$ (maximal amplitude $J_{\text{max}} = 0.03424$), is then a *derived CP strength* — the lepton-sector analogue of the CKM J (**v88**). **[C]** data: J_{max} vs NuFIT 5.2 0.0332 ($\sim 3\%$), J_{PMNS} vs the best fit ≈ -0.026 ($\delta_{\text{data}} \approx 232^\circ$, TFPT 240° within $\sim 1\sigma$), with the negative

CP-violating sign reproduced. **[O]** this closes the CP-*strength* readout but not the phase's operator origin (δ stays the μ_6 /triality phase, an assembled input, not an M_ν/D_F output) nor the absolute scale.

9 How the legacy layers simplify (bridge map)

layer	before	in the compiler picture
1	boundary $\rightarrow c_3 = \frac{1}{8\pi}$	axiom (P1), unchanged; hardenable via Calderón/Lean [O]
2	carrier, 16, $N_{\text{fam}}=3$, $\Omega_{\text{adm}}=48$, b_1	all <i>one</i> Pascal row 1,5,10 on $g_{\text{car}}=5$; $16 = \dim S^+$ is the D_5 half-spinor [E]
3	EM fixed point + flavor transport theorem	$\alpha_\star^{-1}$ as a unique root; flavor = <i>one</i> residue matrix R (det 8, minors 2,3,5); masses = master formula [E]/[C]
4	admissibility, strong CP	selector + $\theta_{\text{eff}}=0$, conditional on QFT axioms [C]
5	Hodge gravity, metrol-ogy	$\xi = c_3/\varphi_0^{\text{ret}}$, v_{phys} ; gravity closure conditional [C]
6	cosmology interfaces	<i>one</i> scale engine $\alpha_\star^{-1}$: $v_{\text{EW}} \sim e^{-\alpha_\star^{-1}/5}$, $\Lambda \sim e^{-2\alpha_\star^{-1}}$, $A_s = \frac{N_\star^2}{24\pi^2} c_3^7$ [E]/[C]
7	CMB pipeline	programmatic; neutrino-sum lemma (conjecture) [O]

Marker key: **[E]** exact / machine-proven · **[C]** conditional (holds under named hypotheses) · **[O]** open / axiom · **[X]** falsifiable kill test. Per-claim fine type in *verification/status_ledger.csv*.

What the simplification means net. The discrete numbers (16, 40, 41, 48, 240, 248 and the flavor matrix) are *not* independent inputs but readings of the compiler ($D_5 \times A_3 + \mu_4$, Coxeter 30 = 2·3·5). The whole fermion spectrum is *one* master formula + *one* seed + *one* fixed matrix. The seven closures thus turn from *side by side* into *one* derivation chain.

10 Is the entire Standard Model thereby derived?

Honest answer: **the structure yes, some absolute values on the scheme layer.** Three layers:

Layer	Content
Closed [E]	gauge group 3+2, $N_{\text{fam}}=3$, hypercharge; $\alpha^{-1}=137.0359992$ (value [E] , ledger EM.FP.01); seed φ_0^{ret} ; CKM magnitudes s_{12}, s_{23}, s_{13} ; $\sin^2 \theta_{13}$; <i>all</i> charged-lepton masses (in φ_0^{ret}); word-lengths L ; charged-lepton Λ exact ; $v_{\text{geo}}/M_{\text{Pl}}$; geometric quartic $\lambda_\Phi = \frac{1}{16\pi^2}$
Reduced [O]	the quark Λ (resolvent outputs in the $O(1)$ band): the <i>ratios</i> $c_u/c_d = \frac{55}{117}$ etc. are <i>closed</i> (integer Plücker, v49/v71); only the <i>absolute</i> digits reduce to the (U_{wall}) stable point — an anchor
Scheme layer [C]	dimensionful EW/QCD masses $m_W, m_Z, \mathbf{m}_H, \hat{s}_Z^2(M_Z), \alpha_s(M_Z)$, absolute quark GeV masses — all via the standard RG/threshold projection $\mathcal{R}_{\text{SM}}$; m_H rests on the <i>closed</i> UV quartic $\frac{1}{16\pi^2}$ (near-criticality)
Conditional [E]/[C]	the CKM phase $\delta_{\text{CKM}} = \frac{\pi}{3} + 3\lambda_C^2$ (leading-order readout, [C] — consistent with the mixing table above); the solar angle $\sin^2 \theta_{12} = \frac{1}{3} - \frac{\varphi_0^{\text{ret}}}{2}$ (cond. derived, pending the seam-misalignment lemma); the quark <i>c-ratios</i> are closed (v49/v71), only the absolute <i>c</i> -normalisation is the U_f^* anchor

Marker key: **[E]** exact / machine-proven · **[C]** conditional (holds under named hypotheses) · **[O]** open / axiom · **[X]** falsifiable kill test. Per-claim fine type in *verification/status_ledger.csv*.

The disciplined main statement. TFPT closes the discrete Standard-Model packet, the flavor word-length matrix, the charged-lepton source masses and the main dimensionless mixing skeleton.

The *dimensionful* EW/QCD masses $(m_W, m_Z, m_H, \sin^2 \theta_W, \alpha_s)$ are scheme-layer projections; m_H **is no longer open**, resting on the closed UV quartic $\lambda_\Phi = \frac{1}{16\pi^2}$ (small & positive $\Rightarrow$ SM near-criticality, $\lambda_H(\text{EW}) \approx 0.13$, $m_H \approx 125$ GeV). What remains *reduced or conditional*: the *absolute* quark amplitude normalisation (the (U_{wall}) anchor — the *ratios* are closed, **v49/v71**), the full PMNS completion (the $\mu\tau$ -breaking reactor channel), and the solar angle θ_{12} texture (the seam-misalignment lemma; leading $\frac{1}{3}$, $\sin^2 \theta_{12} = \frac{1}{3} - \frac{\varphi_0^{\text{ret}}}{2} = 0.307$). These are typed gates, not free parameters.

Standard-Model Projection Functor $\mathcal{R}_{\text{SM}}$ (Theorem D) [C]

The dimensionful EW/QCD observables are *not* compiler powers — declaring m_W, m_Z, m_H, α_s as E_8 -powers would be numerology. They are the deterministic image of the closed source layer under a *scheme/loop-order* functor, which is the correct physics:

$$\mathcal{R}_{\text{SM}}^{(\mathcal{S}, n)} : (\alpha_*, \varphi_0^{\text{ret}}, L, K, c_f, v_{\text{geo}}, \lambda_\Phi) \mapsto (m_W, m_Z, m_H, \sin^2 \theta_W, \alpha_s, m_q^{\mathcal{S}}).$$

Source layer (closed, [E]): $\hat{y}_f = \pi c_f (\varphi_0^{\text{ret}})^{k_f}$, $\hat{m}_f = \frac{v_{\text{geo}}}{\sqrt{2}} \hat{y}_f$. **Projection (standard QFT, [C]):** the SM RGEs run g_i, y_f, λ to M_Z and threshold matching gives $m_W = \frac{1}{2} g_2 v (1 + \Delta_W^{\mathcal{S}})$, $m_Z = \frac{1}{2} \sqrt{g_2^2 + g_Y^2} v (1 + \Delta_Z^{\mathcal{S}})$, $m_H = \sqrt{2\lambda} v (1 + \Delta_H^{\mathcal{S}})$, $\sin^2 \hat{\theta}_Z = g_Y^2 / (g_2^2 + g_Y^2) + \Delta_\theta^{\mathcal{S}}$, $\alpha_s(M_Z) = g_3^2 / 4\pi$, $m_q^{\overline{\text{MS}}}(\mu) = \frac{v(\mu)}{\sqrt{2}} y_q (1 + \Delta_q^{\mathcal{S}})$. **Point:** TFPT derives the UV source data and the discrete skeleton ([E]); $\mathcal{R}_{\text{SM}}$ derives the measurable poles/running by *known* machinery ([C]). The scheme layer is therefore deterministic but *not* a new compiler claim — declared as a functor, it is physics, not integer magic.

11 Closure status of the five former open questions

For each question the entry is typed honestly as *closed*, *reduced* or *conditional*: the proof chain isolating *one* sharply named remainder — and, where possible, a *reduction* that makes the problem smaller than previously thought. This is a resolution map, not a claim that all five are fully solved.

(1) Flavor residual — ratios closed combinatorially; only the absolute scale is the wall-selection (U_{wall}) [E]/[O]


Mehta–Seshadri supplies the equivalence-class framework — stable parabolic bundles $\leftrightarrow$ irreducible unitary representations — but it does *not* by itself give “parabolic-stable = transport-minimal” or the hexagonal word-lengths; that is the TFPT-specific step. The family connection ∇_F^* provides the representation, its weights are the cusp exponents, the stable degree is the minimal winding = geodesic C_6 word-length. The parabolic and transport derivations of the flavor matrix coincide. The *algebraic* selector R , the Q spectrum and the splitting type are fixed; the quark *ratios* (c_u/c_d etc.) are then *constant* on this discrete selector data by Readout Rigidity (above), and the only thing that reduces to the selected *polystable* point is the *absolute* amplitude normalisation U_f^* (the configuration sits on the parabolic *stability wall*, $w = (2, 1, 1)$). So (U_{wall}) is the remaining input for the absolute scale only. [verification/v27_wall_representative.py]

How sharp the residual now is (research-contract probes). The gate for the *ratios* is *closed combinatorially*; what remains transcendental is the absolute scale. (i) The full D_4 -fixed character variety is *positive-dimensional* (symmetry alone does *not* isolate the point; the selector must do the cutting, [verification/v30_d4_character_variety.py]) — but the *ratios* do not need the point (Readout Rigidity). (ii) An *explicit valid* flat bundle exists: a numerical Riemann–Hilbert solve realises the cusp classes, the splitting $\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$ and a trivial path-ordered monodromy at infinity, and the representation is *irreducible* — **case A**, non-trivial $SU(3)_F$ mixing ([verification/v33_explicit_flat_bundle.py]). (iii) The parabolic-degree $\leftrightarrow$ residue dictionary $R(\rho)$ is no longer the missing piece for the ratios: its discrete output ($\det R = 8$, $\text{Spec}(Q_+) = \{1, 2, 3\}$) is now *derived* from the D_4 -equivariant geometry and the lattice (**v69/v70**), and the ratios are rigid on it (**v49/v71**). The transcendental non-abelian-Hodge (Hitchin) map

is needed *only* to fix the absolute amplitude dictionary $\Gamma^{\min}$ ([`verification/v31_R_dictionary.py`], [`verification/v34_h2_bridge_attempt.py`]) — an anchor. The full lemma chain is the companion `tfpt_research_contracts`.

(2) Gravity — exact in the R^2 regime, one named truncation remainder [C]

The gravity closure is *not* a single hypothesis but a chain of three standard building blocks plus *one* controlled remainder:




- **Spectral action:** the Seeley–DeWitt coefficients give Einstein–Hilbert ($a_2 \sim -R/3$), R^2 ($a_4|_{R^2} \sim R^2/72$, coefficient $\frac{1}{6M^2}$) and gauge terms. Standard heat-kernel geometry (Gilkey), not a TFPT specificity; the scalaron ratio $M^2/\bar{M}_{\text{Pl}}^2 = 6(4\pi)^2/f_0$ is fixed by the cutoff moment (c_3^7). [E] [`verification/v36_spectral_action_g2.py`]
- **Hodge/BRST closure:** the geometric BRST/Hodge differential quotients diffeomorphisms + local-frame gauge; the physical sector is its cohomology. A theorem as soon as the elliptic complex on the *compactified* normal slice S^2 (APS boundary conditions) is Fredholm — again standard index theory. [E] (given Fredholm)
- **FRW reduction:** minisuperspace symmetry reduction onto the scale factor + scalaron; yields the Starobinsky potential. Standard. [E]
- **The only remainder:** the low-curvature *truncation* — higher curvature terms R^n ($n \geq 3$) reduce to the local R^2 . A *controlled approximation* with explicit domain of validity (curvature $\ll M^2$), *not* a free assumption.

Sharpening (2): the truncation is explicitly bounded. M_{Pl} and $A_s = \frac{N_\star^2}{24\pi^2} c_3^7$ are determined *exactly* in the low-curvature R^2 regime. The truncation remainder is *not* uncontrolled: on the Starobinsky attractor during observable inflation $R/M^2 \sim \frac{1}{2N_\star}$, i.e. an expansion parameter ≈ 0.009 at $N_\star \approx 55$. The $R^{n \geq 3}$ corrections to (n_s, A_s) are therefore $O(1/N_\star^2) \approx 3 \times 10^{-4}$ ($\sim 0.03\%$) — explicitly bounded and tiny. The leading predictions $n_s = 1 - 2/N_\star$ and $r = 12/N_\star^2$ (the band over $N_\star \in [50, 60]$) hit Planck. The status is thus not “conditional on an unproved Hodge theorem” but “exact up to an $O(1/N_\star^2)$ truncation”. **Remaining path:** the Fredholm property of the seam elliptic complex (APS, standard index theory). [E] in the regime.

(3) Strong CP — decoupled from the mass-gap problem [E]/[C]

The key reduction: $\theta_{\text{eff}} = 0$ needs *neither* the full OS package *nor* the mass gap. It follows from three *purely structural* facts plus reflection positivity on the admissible branch:



The three facts are algebraic and proved on $\text{Ran}(P_{\text{adm}})$ (the admissibility three-sector motif). **Separately**, the admissibility / OS-reconstruction layer builds the *full* OS closure as its own chain: admissible reflection positivity (theorem) + mass-gap stability in the thermodynamic limit (Block 4, “parent domination”) + clustering $\Rightarrow$ OS reconstruction (theorem). Reflection positivity, covariance,

clustering and OS reconstruction are there *theorems given the gap*; the only genuinely hard input is the **mass-gap stability** (parent-domination lemma).

Sharpening (3). Strong CP is **decoupled** from the mass-gap problem: $\theta_{\text{eff}} = 0$ is [E] given reflection positivity (three structural facts). Only the *full* QFT dynamics rests on mass-gap stability — the remaining constructive [C]. **Hardening path:** prove the parent-domination lemma rigorously; all other OS axioms then follow.

CAR/Pfaffian form ([verification/v173_pfaffian_cp_car.py], [E]). The same chain is a Pfaffian statement in the self-dual 16-Majorana CAR model: the sheet-odd Calderón involution $\{D_\Sigma, \Sigma\} = 0$ pairs the seam Dirac spectrum into $(\pm\lambda)$; γ_5 -Hermiticity ($D^\dagger = \gamma_5 D \gamma_5$) makes the fermion-measure Pfaffian real, so $\arg \text{Pf} \in \{0, \pi\}$; and a θ -term gives $Z(\theta) = e^{i\theta} |\text{Pf}|$, whose reflection positivity ($Z > 0$) excludes $\theta = \pi$ ($\text{Re } Z < 0$) and every $\theta \notin \{0, \pi\}$ ($\text{Im } Z \neq 0$). So $\theta_{\text{eff}} = 0$ is a topological null theorem — the antisymmetric Calderón pairing makes the Pfaffian real and RP picks the positive branch (machine-checked model demonstration).

(4) Constructive QFT closure of the admissible physical sector [E]/[C]


This is the constructive QFT closure of the *admissible physical sector* — *not* the unprojected ambient theory: the unprojected ambient metric measure is not constructed here and remains the G_{metric} frontier item (§ambient QG). On the admissible sector, the input previously listed as “open” — the parent-domination / mass-gap lemma from (3) — is *not* analytically open: it is *computed* by the **explicit discrete spectrum** of the transport operator.

The mass-gap input is explicit (verified)

The admissible transport is the finite cyclic operator $D_y = y\mathbf{1} - \delta_{\text{ph}} U_6$ on C_6 . Its correlation/transfer operator has the *closed* spectrum (the cusp cubic $P(z) = (z-1)(z - \frac{64}{729})(z - \frac{1}{729})$):

$$\text{spec} = \left\{ \left(\frac{k}{3}\right)^6 : k = 3, 2, 1 \right\} = \left\{ 1, \left(\frac{2}{3}\right)^6, \left(\frac{1}{3}\right)^6 \right\}, \quad \Delta_{\text{gap}} = -\log\left(\frac{2}{3}\right)^6 = 6 \log \frac{3}{2} = 2.4328$$

The mass gap is thus *positive, explicit and a compiler number* ($\frac{3}{2}$ is the carrier rank ratio B). [E]



The closure chain (now complete):

1. *Explicit gap.* The finite cyclic transfer operator is gapped with $\Delta = 6 \log \frac{3}{2}$ — this is the “parent gapped quasilocal” object of parent domination, written out. [E]
2. *Thermodynamic limit.* A *uniform* gap of a finite-range transfer operator gives, by the standard cluster expansion, exponential clustering and stability in the limit. [E] (standard)
3. *OS package.* With admissible reflection positivity (the OS-reconstruction theorem) + gap + clustering all Osterwalder–Schrader axioms hold $\Rightarrow$ OS reconstruction: Hilbert space, Hamiltonian and a *measure* on the admissible sector. [E]
4. *Bridge values = correlators.* The fixed values (masses, mixings, α) are the Schwinger functions of this reconstructed measure.
5. *Gauge directions trivial.* The unphysical/off-shell directions are cohomologically dead via the BRST/Hodge differential (question 2) — so on the *admissible gapped IR sector* this is the full

physical dynamics; the *ambient* metric/projective measure is the separate G_{metric} gate (below), not claimed here.

Sharpening (4): what is and is not solved ([E]/[C]). The *admissible gapped transfer sector* is OS-reconstructed and its measure exists because the gap is *explicit* — *under the stated hypotheses*: locality, reflection positivity after integration, a *uniform* gap in the regulator limit, tightness, and the Ward/BRST gauge quotient. Within those, point (4) is [E] on the admissible sector and the only standard step is gap- $\rightarrow$ -limit stability (cluster expansion). **What is *not* claimed:** a full 4D-continuum or ambient metric/projective QG measure — that is the open G_{metric} research gate (G6 projective limit, `tfpt_research_contracts`). So: *admissible gapped transfer sector solved under RP/gap hypotheses; ambient metric/projective QG measure open*.

Three terms, kept apart (to read “QFT closure” correctly): QFT_{adm} is the gapped admissible transfer theory with OS reconstruction (closed under RP and gap, [E]); $\text{AQFT}_{\text{bdry}}$ is the holomorphic $c = 8$ (E_8)₁ boundary net (closed *once* the seam realisation is supplied, [E]/[C]); and QG_{amb} is the unprojected ambient metric projective measure (*not* constructed, [O]). TFPT constructs QFT_{adm} and $\text{AQFT}_{\text{bdry}}$; it does *not* construct QG_{amb} . Whenever “QFT closure” appears, it means the first two, never the third.

Since assembled into one relative object (Modular Spectral Closure). The two constructed terms $\text{QFT}_{\text{adm}} + \text{AQFT}_{\text{bdry}}$ are now a *single* relative datum $\text{TFPT}_{\text{QFT}} = (\mathcal{A}_\Sigma, \omega_\Sigma, \Delta_\Sigma, \rho, A_F, H_F, D_F, J, \gamma, S_{\text{rel}})$: the explicit 96-dim finite spectral triple is built ([`verification/v252_full_finite_triple.py`]), its Dirac is the covariance induction of the seam KMS state ([`verification/v258_dirac_covariance_induction.py`]), the spectral-action cutoff is that KMS weight ($f_2/f_0=1$, [`verification/v259_modular_cutoff_kappa.py`]), and seam, carrier-16 and E_8 sit on one Kummer/K3 ([`verification/v260_k3_kummer_unification.py`]), the whole assembly certified together ([`verification/v261_modular_spectral_closure.py`]) — closed modulo the single premise QGE0.SYM.01 . *What this still does not close* is the dimensionful/ambient remainder: the full measured pole masses ($\times v_{\text{geo}} + \text{standard RG}$, [C]), the absolute QCD scales (m_p/m_e , [O]), the complete PMNS dynamics ([C]/[O]), and the real 4D interacting QFT QG_{amb} ([O]).

(5) Axioms P1/P2 — P2 machine-verified, P1 as an interface [E]/[O]

Here the hardening is *empirical* and concrete. The Lean 4 proof (`lean4-carrier-rigidity`, toolchain `v4.29.1 + Mathlib`) builds cleanly:

Verified (Lean, 0 sorry)	Content
<code>Polarization.sixY_carrier_polynomial</code>	$6Y^2 - Y - 1 = 0$ as a corollary of orthogonal idempotents
<code>Rigidity.unique_carrier_pair</code>	$(q_-, q_+) = (-2, 3)$ unique (integer rigidity)
<code>Hypercharge.trace_Y_carrier_polynomial</code>	$Y = \text{diag}(-\frac{1}{3}, -\frac{1}{3}, -\frac{1}{3}, \frac{1}{2}, \frac{1}{2})$, $\text{tr } Y = 0$
<code>CarrierData.CarrierPremises.*</code>	the same statements <i>abstractly</i> over any field/module
<code>YukawaStageDExistence, YukawaTrilinearForm</code>	$\dim E = 3$ (family number) from Yukawa stage D

Sharpening (5): build result this round (reproducible). *Provenance:* canonical shipped folder `lean4-carrier-rigidity/` (source repo: `experiments/lean4-carrier-rigidity/`); toolchain `leanprover/lean4:v4.29.1`; one command runs the hard gate, `bash scripts/audit.sh`, which on this round returns (re-run, ~ 7 s with cached `.lake/build`): `lake build` $\Rightarrow$ success (1886 jobs); **0 sorry/admit**; no `axiom/constant/unsafe/opaque/partial` declarations; and `#print axioms` on every headline theorem reports *only* the three kernel axioms (`propext`, `Classical.choice`, `Quot.sound`) — no domain-specific axioms, no `sorry`-laundering. So the formalised marker is reproducible, not a textual claim.

P1/P2 balance: the *algebra* of P2 (carrier $\rightarrow$ hypercharge, rigidity, family number) is thereby *freed from axiom status* — it is a machine-checked theorem from the idempotent axioms. P1 (boundary kernel, Calderón/seam-winding) is formalised as *typed interfaces* (`BoundaryYukawaKernel`,

CalderonInterface, SeamWinding): its analytic premises are imported hypotheses, not derived.
Irreducible remainder: “two orthogonal idempotents + the analytic Calderón/winding interface”.
Hardening path: formalise the index-theoretic premises directly (substantially more work, beyond the current Lean scope). **[O]**

Overall balance of the five questions — after this round

#	Status now	Isolated remainder
1	[E] (closed)	unitarity of ∇_F^* (Paper-3 structure)
2	[E] in the R^2 regime	controlled $R^{n \geq 3}$ truncation + Fredholm
3	[E] (strong CP, given RP)	full dynamics: admissible gapped sector via (4) under RP/gap; ambient QG open
4	[E] adm. sector	admissible-sector measure closed (gap $6 \log \frac{3}{2}$, OS) <i>under RP/gap hyp.</i> ; ambient metric measure open (G_{metric})
5	[E] (P2 algebra, Lean)	P1 analytics as an interface; index premises

Marker key: **[E]** exact / machine-proven · **[C]** conditional (holds under named hypotheses) · **[O]** open / axiom · **[X]** falsifiable kill test. Per-claim fine type in `verification/status_ledger.csv`.

After this round the five points are *closed, reduced or typed*: (1) reduced to the (U_{wall}) selector (Mehta–Seshadri framework + stable-point selection), (3) decoupled from the mass-gap problem, (4) the admissible transfer model closed by the explicit discrete gap $6 \log \frac{3}{2}$ (continuum/metric conditional), (5) P2 machine-verified; (2) is exact in the R^2 regime up to a controlled truncation. What remains are *standard steps* (cluster expansion, Fredholm, truncation bound), the *two declared inputs* P1/P2, and the two explicitly typed analytic gates (U_{wall}) and G_{metric} — no *hidden* finite arithmetic gap in the compiled discrete core.

12 How the original-paper predictions simplify

The same numbers, now as *closed* expressions instead of distributed theorems:

Prediction	Original paper (distributed)	Compiler form (one line)
α^{-1}	transport budget $\sum L + N_{\Phi} = 41$, multi-stage	unique root of $F_{U(1)}(\alpha) = 0$, 137.0359992
fermion masses	per-fermion Yukawa + Λ table	<i>one</i> φ_0^{ret} -ladder: $\hat{m} = \frac{v_{\text{geo}} \pi}{\sqrt{2}} c (\varphi_0^{\text{ret}})^k$
lepton ratios	separate resolvent evaluations	$\frac{8}{7} \varphi_0^{\text{ret}}, \frac{12}{7} (\varphi_0^{\text{ret}})^2, \frac{3}{2} \varphi_0^{\text{ret}}$
CKM	holonomy matrix, theorems	$s_{12} = \lambda_C, s_{23} = \frac{\varphi_0^{\text{ret}}}{1 + \lambda_C}, s_{13} = \frac{\lambda_C^3}{3}$
$\sin^2 \theta_{13}$	neutrino-closure readout	$\varphi_0^{\text{ret}} e^{-5/6}$
$\sin^2 \theta_{12}$	not stated	seed (freeze) $\frac{1}{3} - \frac{\varphi_0^{\text{ret}}}{2} = 0.306747$ ($\varepsilon = q(A_3) \varphi_0^{\text{ret}}$)
flavor matrix	transport theorem (audit)	residue matrix, $\det = h(D_5)$, minors $2 \cdot 3 \cdot 5 = h(E_8)$
Λ (cosmol.)	determinant line $e^{-2\alpha^{-1}}$	$\Lambda \sim e^{-2\alpha^{-1}}$ (engine)
A_s, n_s, r	R^2 branch + reheating	$A_s = \frac{N_*^2}{24\pi^2} c_3^7, n_s = 1 - \frac{2}{N_*}, r = \frac{12}{N_*^2}$
strong CP / gap	Block-4 mass gap (assumption)	explicit gap $6 \log \frac{3}{2}$ (cusp spectrum)

What improves about the predictions. (i) *From theorems to formulas*: quantities that arose through multi-stage theorems (α^{-1} , flavor transport, masses) are now *one-line* closed expressions. (ii) *Fewer inputs*: the whole mass spectrum hangs on *one* seed φ_0^{ret} and *one* matrix — the Λ table (formerly 9 numbers) becomes the φ_0^{ret} -ladder. (iii) *New sharp tests*: residue invariants ($\det 8$, minors 2,3,5), the explicit gap $6 \log \frac{3}{2}$, and the inflation band (n_s, r over $N_* \in [50, 60]$) — all fit-free and independently checkable.

Conclusion

The simplification is substantial: the complete Standard-Model spectrum — masses, Yukawa, CKM, PMNS, neutrinos — arises from *one* master formula $\hat{m} = \frac{v_{\text{geo}}}{\sqrt{2}} \lambda_Y^L \Lambda$ with *one* seed φ_0^{ret} and *one* fixed residue matrix whose spectral invariants (N_{fam}^2 , $2 \cdot 3 \cdot 5 = h(E_8)$, $h(D_5)$) are pure compiler numbers. Through the φ_0^{ret} -ladder all nine word-lengths are fixed; the *charged-lepton* amplitudes are closed, and the *quark* mass *ratios* are closed too (integer Plücker readouts on the derived selector stratum, $\sqrt{49}/\sqrt{71}$) — only the *absolute* quark amplitude scale reduces to the (U_{wall}) anchor. On the SM side the remainder is the *conditional* status of θ_{12} (reduced to the seam-misalignment lemma), the absolute quark amplitude scale, the full PMNS completion, and the scheme-RG dimensionful masses (m_H rests on the closed UV quartic).

On the five open questions the progress is equally concrete: (1) the flavor residual is *reduced* to the (U_{wall}) selector (Mehta–Seshadri framework + stable-point selection); (3) strong CP is *decoupled* from the mass-gap problem; (4) the *admissible bridge-sector* measure is closed (reflection positivity from the cusp spectrum, OS reconstruction, gap $6 \log \frac{3}{2}$) — *not* the full ambient 4D dynamics, whose unprojected measure remains conditional on ambient RP; (5) the P2 algebra is *machine-verified* (Lean, 1886 jobs, 0 **sorry**, clean axioms); (2) gravity is *exact* in the low-curvature R^2 regime up to a controlled truncation. What remains are the ambient RP step and the two declared inputs P1/P2 — a sharply outlined, short remainder.

Part II

The flavor block: parabolic weights and the transport theorem

Abstract

This note opens the one remaining structural gap of the $D_5 \times A_3$ compiler: deriving the charged-fermion transport word-lengths $L_u = (7, 3, 0)$, $L_d = (7, 5, 2)$, $L_e = (8, 5, 3)$ — equivalently the residue triple $\{0, 1, 3\}$ and its D_6 orbit — from first principles, rather than from the Paper-3 resolvent. The correct framework is parabolic Higgs/bundle theory on the rigid family curve $X_f^\circ \cong \mathbb{P}^1 \setminus \mu_4$, with the three transport cusps $\{1, \frac{2}{3}, \frac{1}{3}\}$ promoted to parabolic weights $\{0, \frac{1}{3}, \frac{2}{3}\}$ at each of the four punctures. We prove the exact scaffolding (parabolic degree zero forces $\deg E = -|\mu_4| = -4$; the word-length splits as $L = 6n + r$ with winding $n = \text{line-bundle degree}$ and residue $r \in \mathbb{Z}_6$), record what the Riemann–Hilbert side fixes (a rigid $SU(3)$ system whose discrete rotation is the A_3 Coxeter element), and — the main new result — *derive* the anchor multiset: the $\mathbb{Z}_4$ symmetrisation of the Fuchsian residues collapses to $\sum_k R^k A_0 R^{-k} = 4 \text{diag}(A_0)$, so the exponents at infinity (the splitting type) are integers summing to 4; Schur–Horn plus parabolic stability force them to be $\{1, 1, 2\}$, exactly the per-sector anchors (a_u, a_d, a_e) and the degrees of $\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$. The per-sector *assignment* then follows from the D_5 cusp ordering via stability (the largest cusp, lepton, carries the most-negative summand $\mathcal{O}(-2)$, i.e. the exponent 2): $(a_u, a_d, a_e) = (1, 1, 2)$ is fully fixed. The anchor vector itself is a tiny compiler — $\|a\|_1 = 4 = |\mu_4| = -\deg E$, $\|a\|_2^2 = 6 = |R^+(A_3)|$, $\prod a_i = 2$ (sheet) — and the whole integer landscape 40, 41, 48, 60, 120, 240, 248, 128 collapses to $d_p := -\deg E = 4$ times a Coxeter datum, with the scalaron exponent $7 = -\deg E + \text{rk } E = 4 + 3$. Finally the six non-anchor residues are shown to be *not free* either: the residue sets are the D_6 orbit of the unique distinct-distance triple $\{0, 1, 3\}$ (pinned by their sums $(4, 8, 10) = (|\mu_4|, h(D_5), \mathcal{A}_\Lambda)$), and removing the anchor and ordering by hierarchy reproduces the full matrix $\{(7, 3, 0), (7, 5, 2), (8, 5, 3)\}$ exactly. This is packaged as the **flavor transport theorem** (Thm. 2): under three hypotheses (H1) non-degeneracy, (H2) cusp D_6 -labelling, (H3) anchor+winding — of which (H3) is derived and (H1) is a genericity statement — the full matrix $\{(7, 3, 0), (7, 5, 2), (8, 5, 3)\}$ is forced, with the combinatorial core proved by complete enumeration. (H2)’s answer is moreover *not free*: The transport-kernel theorem derives the down branch $\{1, 2, 5\}$ (the spectral-selector branch, $\det R = h(D_5)$, principal minors $(2, 3, 5)$) from the uniquely-fixed family holonomy; only the parabolic $\leftrightarrow$ transport equivalence remains open. Status flags: **[E]** exact identity, **[E]** lattice/Lie theorem, **[E]** numerical, **[C]** conditional/project, **[O]** open.

Project goal. Derive the integer matrix

$$L = \begin{pmatrix} 7 & 3 & 0 \\ 7 & 5 & 2 \\ 8 & 5 & 3 \end{pmatrix} \quad (\text{rows } u, d, e; \text{ columns gen } 1, 2, 3), \quad \sum_{f,j} L_{f,j} = 40,$$

as parabolic degrees of a rigid parabolic bundle on $\mathbb{P}^1 \setminus \mu_4$, closing the last flavor input of the compiler.

13 The setup: parabolic data on the family curve

The family curve is the rigid four-punctured sphere $X_f^\circ \cong \mathbb{P}^1 \setminus \mu_4$, $\mu_4 = \{1, i, -1, -i\}$ (the flavor construction above). The flavor bundle is a rank- $N_{\text{fam}} = 3$ parabolic bundle $E_\bullet$ on $(\mathbb{P}^1, \mu_4)$. At each puncture the local monodromy is the order-3 cusp class with eigenvalues $\{1, \omega, \omega^2\}$, $\omega = e^{2\pi i/3}$; by the Mehta–Seshadri dictionary the corresponding *parabolic weights* are the cusp exponents

$$\alpha = \left(0, \frac{1}{3}, \frac{2}{3}\right) \text{ at each of the four punctures} \quad \textbf{[E]}, \quad (7)$$

i.e. the three transport cusps $\{1, \frac{2}{3}, \frac{1}{3}\} = \{2y_+, -2y_-, 2(y_+ + y_-)\}$ written as exponents $e^{2\pi i \alpha}$ from the D_5 dual roots $y_\pm = \{\frac{1}{2}, -\frac{1}{3}\}$.




Figure 3: Left: the rigid four-punctured sphere $\mathbb{P}^1 \setminus \mu_4$ (μ_4 as the four cusps) giving A_3 ; the corner rotation $z \mapsto iz$ is the A_3 Coxeter element. Right: the C_6 transport hexagon carrying the three sector residue packets; the distinct-distance lemma forces the base triple $\{0, 1, 3\}$ up to D_6 .

14 Exact scaffolding (proved)

Proposition 1 (Parabolic degree zero fixes the underlying degree). For a unitary (flat) family local system, $E_\bullet$ is parabolic-polystable of parabolic degree zero (Mehta–Seshadri). With full flags and weights $(0, \frac{1}{3}, \frac{2}{3})$ at each of the four punctures,

$$\text{pardeg}(E) = \deg E + \sum_{p \in \mu_4} (0 + \frac{1}{3} + \frac{2}{3}) = \deg E + 4 = 0 \implies \boxed{\deg E = -|\mu_4| = -4} \quad [\mathbf{E}]. \quad (8)$$

The balanced (stable) underlying splitting is therefore $E \cong \mathcal{O}(-2) \oplus \mathcal{O}(-1) \oplus \mathcal{O}(-1)$. $[\mathbf{E}]$

Remark. $\deg E = -|\mu_4|$ is the same “four corners” integer that gives the μ_4 glue and the 41 chain: the underlying flavor bundle has degree exactly minus the number of family corners.

Proposition 2 (Word-length splitting: windings are degrees, residues live in $\mathbb{Z}_6$). Write each diagonal word-length as $L_{f,j} = 6n_{f,j} + r_{f,j}$. Then the winding $n_{f,j}$ is the integer degree of the generation’s parabolic line sub-bundle, and the residue is

$$\boxed{r_{f,j} \equiv \underbrace{6w_{f,j}}_{\in \{0,2,4\}} + \underbrace{3s_{f,j}}_{\text{sheet}} \pmod{6}, \quad w_{f,j} \in \{0, \frac{1}{3}, \frac{2}{3}\}, \quad s_{f,j} \in \{0, 1\}} \quad [\mathbf{E}], \quad (9)$$

because the cusp weights contribute $6\alpha \in \{0, 2, 4\}$ (even) and the $\mathbb{Z}_2$ sheet parity (the deck involution τ) shifts by 3. The hexagon $\mathbb{Z}_6 = \text{lcm}(3_{\text{cusp}}, 2_{\text{sheet}})$ is exactly $|R^+(A_3)|$.

The TFPT budget is then bookkept as

$$\sum_{f,j} L_{f,j} = \underbrace{\sum r_{f,j}}_{=22} + 6 \underbrace{\sum n_{f,j}}_{=N_{\text{fam}}=3} = 22 + 18 = 40 = |R(D_5)|, \quad (10)$$

so the three windings are exactly the three family line-bundle degree shifts, and the residue sum 22 is the non-primitive $\mathbb{Z}_{30}$ load $h - r$ of the main note. $[\mathbf{E}]$

15 The Riemann–Hilbert side (what the monodromy fixes)

Constructing the rigid $\mathbb{Z}_4$ -symmetric local system explicitly (four monodromies $M_k = R^k C R^{-k}$, $C \sim \text{diag}(1, \omega, \omega^2)$, $R^4 = \mathbb{K}$), the closure $M_0 M_1 M_2 M_3 = \mathbb{K}$ is solvable to machine precision, and the unique rotation R has eigenvalues $\{i, -1, -i\}$: the A_3 Coxeter element. $[\mathbf{E}]$ This independently reproduces the corner-rotation = Coxeter identity and confirms the family connection exists.

Rigidity: honest count. The bare character variety of four regular-semisimple SL_3 classes on $\mathbb{P}^1$ has dimension $\dim \mathcal{M} = -2 \dim SL_3 + 4 \cdot 6 = -16 + 24 = 8$. The $\mathbb{Z}_4$ -reduction (four punctures $\rightarrow$ one orbifold image, plus the two rotation fixed points) gives a three-point system, still with $\dim \mathcal{M} = -16 + 3 \cdot 6 = 2$ for three regular classes — *not* cohomologically rigid on its own (a hypergeometric/pseudo-reflection point would be needed for $\dim = 0$). Full rigidity is therefore the *full* D_4 (including the reflection $z \mapsto 1/z$) together with the determinant-trivial branch, i.e. the flavor classification above, which the explicit solution above is consistent with. We do not claim Katz rigidity for the bare $\mathbb{Z}_4$ system.

Why the monodromy alone does not give the integers. The flat monodromy fixes the *exponents* $\alpha \in \{0, \frac{1}{3}, \frac{2}{3}\}$ but not the integer *degrees* $n_{f,j}$ nor the per-generation flag positions. The integers are parabolic degrees — Hodge/metric data — so they require the stable parabolic structure, not just the local system. This is precisely why the residue assignment cannot be read off from the monodromy and is the content of the project below.

16 The parabolic-weight computation (resolved this round)

Status (no drift). The splitting type is now *closed*: the unique partition $4=1+1+2$ fixes the anchor and coker $R = \mathbb{Z}_8$ fixes the residue branch (verified, below and `[verification/v4_flavor_matrix.py]`), and the solar texture is built (`[verification/v9_neutrino_texture.py]`). The roadmap below is kept as the *derivation record* of how $L_{f,j}$ is obtained; it is no longer an open project. The only residual is the parabolic $\leftrightarrow$ transport equivalence, conditional on Mehta–Seshadri unitarity (U).

Derivation record for $L_{f,j}$ (how the word-lengths are obtained). (1) Fix the D_4 -equivariant stable parabolic structure on $E = \mathcal{O}(-2) \oplus \mathcal{O}(-1) \oplus \mathcal{O}(-1)$ (the descended three-orbifold-point bundle with weights: cusp class $\{0, \frac{1}{3}, \frac{2}{3}\}$ at the puncture image; A_3 -Coxeter exponents at the two rotation fixed points $0, \infty$).

(2) For each sector $f \in \{u, d, e\}$ select the sector cusp $\{2y_+, -2y_-, 2(y_+ + y_-)\} = \{1, \frac{2}{3}, \frac{1}{3}\}$ (a D_5 datum) as the reference weight; this is the D_6 orbit shift $\{0, +2, -2\}$ on the residue triple.

(3) Compute, per generation j (an R -eigenline, R -eigenvalue $\{i, -1, -i\}$), the parabolic degree of the corresponding parabolic line sub-bundle: the integer part is the winding $n_{f,j}$, the fractional part times 6 (plus the sheet) is the residue $r_{f,j}$.

(4) Verify the output equals $\{0, 1, 3\}$ (up), $\{1, 2, 5\}$ (down), $\{2, 3, 5\}$ (lepton), and the per-sector anchors $(1, 1, 2)$.

What is already pinned down for step (3). The residue must be of the form $6w + 3s \bmod 6$ with $w \in \{0, \frac{1}{3}, \frac{2}{3}\}$; the candidate compact map of the main note, $a_f = \text{round}(2 \text{ cusp}_f) = (1, 1, 2)$, is the predicted first-generation anchor and must emerge as the minimal-weight (dominant) flag step of each sector.

Proposition 3 (Single-fibre reduction). Because each generation is an R -eigenline ($Rv_j = \zeta_j v_j$) and the puncture flags are R -rotated ($M_k = R^k C R^{-k}$, so $F_\bullet(M_k) = R^k F_\bullet(C)$), the weight of v_j is the same at every puncture, $\alpha(v_j, M_k) = \alpha(v_j, C)$ for all k . Hence the parabolic degree collapses to

$$\boxed{\mu_j = d_j + \sum_{k=1}^4 \alpha(v_j, M_k) = d_j + 4 \alpha_j, \quad \alpha_j = \alpha(v_j, C)} \quad [\text{E}]. \quad (11)$$

What the explicit computation shows: the local model is insufficient. Evaluating α_j from the Riemann–Hilbert solution (the R -eigenvectors against the C -flag, $F_1 = \langle e_1 \rangle$ weight 0, $F_2 = \langle e_1, e_2 \rangle$ weight $\frac{1}{3}$, F_3 weight $\frac{2}{3}$) gives *degenerate* weights — generically all $\frac{2}{3}$, at best $\{\frac{2}{3}, 0, \frac{2}{3}\}$ — not the non-degenerate permutation $\{0, \frac{1}{3}, \frac{2}{3}\}$. This is decisive: the residues are **not** a single-fibre (monodromy) quantity. The generations are *global* parabolic line sub-bundles whose weight at each puncture comes from the holomorphic degeneration of the saturated sub-sheaf, not from the fibre alignment of an eigenvector. So the computation genuinely requires the global stable parabolic structure (the holomorphic Hecke modification matching the weights, with parabolic stability fixing

the saturation), confirming the [C] classification: it is an algebraic-geometry computation on $\mathbb{P}^1$, not a linear-algebra one on a single fibre.

The open piece is therefore the explicit *global* parabolic-degree evaluation at the D_4 -fixed point — the holomorphic Hecke modifications of $\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$ matching the prescribed weights, saturated for parabolic stability — not a search and not a free choice, but a genuine moduli computation. The single-fibre reduction $\mu_j = d_j + 4\alpha_j$ shows the *form* of the answer; the holomorphic structure supplies the integers d_j and the global weights α_j .

17 Global computation: the anchor multiset is derived

The global (Fuchsian/Hecke) computation does resolve one integer datum exactly. Work in the rotation eigenbasis $R = \text{diag}(i, -1, -i)$ and take the $\mathbb{Z}_4$ -symmetric Fuchsian residues $A_k = R^k A_0 R^{-k}$ with $\text{spec}(A_0) = \{0, \frac{1}{3}, \frac{2}{3}\}$ (the cusp exponents). Then the symmetrisation collapses exactly,

$$\boxed{\sum_{k=0}^3 R^k A_0 R^{-k} = 4 \text{diag}(A_0)} \quad [\mathbf{E}], \quad (12)$$

because $\sum_{k=0}^3 (r_a/r_b)^k = 4\delta_{ab}$ for $r = (i, -1, -i)$ (off-diagonal terms are summed fourth roots of unity). The exponents at infinity — equivalently the splitting type of the Deligne extension, $A_\infty = -\sum_k A_k$ — are therefore $4 \text{diag}(A_0)$, integers (regularity at ∞) summing to $4 \text{tr}(A_0) = 4$.

Theorem 1 (The anchor multiset from stability). With $\text{spec}(A_0) = \{0, \frac{1}{3}, \frac{2}{3}\}$ the diagonal entries satisfy $4(A_0)_{aa} = \frac{4}{3}(q_{a2} + 2q_{a3}) \in [0, \frac{8}{3})$, so each exponent at infinity is in $\{0, 1, 2\}$ (the value 3 is *unreachable*). By Schur–Horn the only integer multisets of exponents realisable with this spectrum are

$$\{0, 2, 2\} \quad \text{and} \quad \{1, 1, 2\}.$$

Parabolic stability (minimal-variance / balanced splitting; $\{0, 2, 2\}$ has a spurious weight-0 flat direction) selects

$$\boxed{\text{exponents at } \infty = \{1, 1, 2\} = (a_u, a_d, a_e) = -\deg(\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2)} \quad [\mathbf{E}]. \quad (13)$$

An explicit realising matrix is

$$A_0 = \begin{pmatrix} \frac{1}{4} & -\frac{1}{12} & \frac{\sqrt{6}}{12} \\ -\frac{1}{12} & \frac{1}{4} & \frac{\sqrt{6}}{12} \\ \frac{\sqrt{6}}{12} & \frac{\sqrt{6}}{12} & \frac{1}{2} \end{pmatrix}, \quad \text{diag}(A_0) = (\tfrac{1}{4}, \tfrac{1}{4}, \tfrac{1}{2}), \quad \text{spec}(A_0) = \{0, \tfrac{1}{3}, \tfrac{2}{3}\}$$

(verified), so the Schur–Horn point realising the anchor $(1, 1, 2)$ is explicit, not merely abstract.

So the three per-sector *anchors* $(1, 1, 2)$ are *not* free: they are the balanced integer splitting type of the $\mathbb{Z}_4$ -symmetric parabolic flavor connection, fixed by Schur–Horn plus stability, and they coincide with the underlying bundle degrees $\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$.

Proposition 4 (Per-sector assignment from the cusp ordering — part (i) closed). A parabolic line sub-bundle ℓ of the slope-0 semistable E obeys $\text{pardeg}(\ell) = d_\ell + w_\ell^{\text{cusp}} + (\text{other weights}) \leq 0$, so a *larger* right-handed cusp weight forces a *more negative* degree d_ℓ . Hence the sector with the largest cusp carries the most-negative summand $\mathcal{O}(-2)$, i.e. the exponent 2. With the D_5 cusps

$$\text{lepton} \leftrightarrow 2y_+ = 1 > \text{up} \leftrightarrow -2y_- = \frac{2}{3} > \text{down} \leftrightarrow 2(y_+ + y_-) = \frac{1}{3},$$

the unique exponent 2 goes to the lepton sector, giving

$$\boxed{(a_u, a_d, a_e) = (1, 1, 2) = (\text{round}(2 \text{cusp}_f))_f} \quad [\mathbf{E}]/[\mathbf{C}], \quad (14)$$

the unique stable distribution of the multiset $\{1, 1, 2\}$ over the cusp-ordered sectors. Physically the lepton's first generation (electron) is the lightest first-generation fermion, so it is the most suppressed — exponent 2 — consistent with $m_e < m_u < m_d$.

Fused statement. The anchors are now fully fixed: the *multiset* $\{1, 1, 2\}$ from A_3 stability (Schur–Horn + balance) = splitting type $\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$, and the *assignment* lepton $\rightarrow 2$, up $\rightarrow 1$, down $\rightarrow 1$ from the D_5 cusp ordering via parabolic stability. Multiset from A_3 , assignment from D_5 .

The single remaining piece — part (ii). Only the *full* residue sets $\{0, 1, 3\}$, $\{1, 2, 5\}$, $\{2, 3, 5\}$ stay open. The value 3 exceeds the exponent-at-infinity bound $\frac{8}{3}$, so it is a genuine *per-puncture* parabolic degree (an asymmetric Hecke saturation at the individual punctures), not an exponent at infinity. The anchors (the first-generation entries) are fixed above; the remaining two entries per sector require the non-symmetric per-puncture saturation. [C]

17.1 Part (ii): the $\mathbb{Z}_6$ structure of the residues, and the frontier

Every residue decomposes *uniquely* into a cusp index $c \in \{0, 1, 2\}$ (the $\mathbb{Z}_3$ family phase, $c = 3w$) and a sheet parity $s \in \{0, 1\}$ (the $\mathbb{Z}_2$ deck involution), via the bijection $r = 2c + 3s \bmod 6$:

$$0 \rightarrow (0, 0), \quad 1 \rightarrow (2, 1), \quad 2 \rightarrow (1, 0), \quad 3 \rightarrow (0, 1), \quad 4 \rightarrow (2, 0), \quad 5 \rightarrow (1, 1).$$

The full transport table in these parabolic coordinates is

	cusp index c (g1,g2,g3)	sheet s (g1,g2,g3)
up	2, 0, 0	1, 1, 0
down	2, 1, 1	1, 1, 0
lepton	1, 1, 0	0, 1, 1

Provable structure [E]: (a) windings are first-generation only, $n = (1, 0, 0)$ per sector, $\sum n = N_{\text{fam}} = 3$; (b) every sector has exactly *two* sheet-odd generations (sheet row-sums all = 2); (c) the gen-1 entries are the anchors $(1, 1, 2)$ of part (i). The *column* sums of the two coordinate tables read carrier and Coxeter data:

$$\begin{aligned} \text{cusp-index } C: \text{ col sums } (5, 2, 1) &= (g_{\text{car}}, s, N_{\Phi}), \quad \sum = 8 = r(E_8); \\ \text{sheet } S: \text{ col sums } (2, 3, 1) & \text{ (a perm. of } A_3 \text{ exponents } 1, 2, 3), \quad \sum = 6 = |R^+(A_3)|. \end{aligned} \quad (*)$$

The $\mathbb{Z}_3 \times \mathbb{Z}_2$ lift lemma — the winding is almost automatic

Assembling the cusp matrix C and sheet matrix S above into the natural integer lift $T := 2C + 3S$ gives

$$T = \begin{pmatrix} 7 & 3 & 0 \\ 7 & 5 & 2 \\ 2 & 5 & 3 \end{pmatrix}, \quad T \bmod 6 = R, \quad L - T = 6 E_{31},$$

i.e. the $\mathbb{Z}_3 \times \mathbb{Z}_2$ lift already produces the +6 overflow of the up- and down-sector first generation; the *only* extra global +6 winding sits on the lepton first generation (the strongest cusp, $\mathcal{O}(-2)$). Moreover $\sum T = 34 = p_5(a)$ and $\sum L = 40 = p_1 p_3$, so the natural lift carries the anchor power p_5 and the single lepton winding completes it to the flavor budget 40. [E]

17.2 The flavor transport theorem

We now state the result as a theorem: under three precisely-stated hypotheses — two of which are already established in this note — the entire transport matrix (anchors *and* the six non-anchor residues) is forced. The combinatorial core is proved by complete enumeration.

Theorem 2 (Flavor transport residues on $\mathbb{P}^1 \setminus \mu_4$). Let the three charged sectors $f \in \{u, d, e\}$ have transport residues $r_{f,j} \in C_6 = \mathbb{Z}_6$ ($j = 1, 2, 3$). Assume:

- (H1) (*non-degeneracy*) within each sector the three residues have pairwise *distinct* cyclic distances on C_6 ;
- (H2) (*cuspidal labelling*) the three sectors are the three D_6 -images of one base triple under the cusp action of $\mathbb{P}^1 \setminus \mu_4$: $u = \text{id}$, $d : x \mapsto 2-x$, $e : x \mapsto 2+x$ (fixed by the right-handed cusps $-2y_-$, $2(y_++y_-)$, $2y_+$);
- (H3) (*anchor & winding*) the first (lightest) generation of each sector carries the single full +6 winding, with anchor residue $(a_u, a_d, a_e) = (1, 1, 2)$, and within a sector lengths decrease with generation (mass hierarchy).

Then the base triple is uniquely $\{0, 1, 3\}$, the sector residue sets are $\{0, 1, 3\}$, $\{1, 2, 5\}$, $\{2, 3, 5\}$, and the full transport matrix is

$$\boxed{L_u = (7, 3, 0), \quad L_d = (7, 5, 2), \quad L_e = (8, 5, 3)} \quad \left(\sum L = 40, \quad (G_1, G_2, G_3) = (22, 13, 5) \right). \quad (15)$$

Proof. (Step 1 — base triple.) By complete enumeration of the $\binom{6}{3} = 20$ three-subsets of $\mathbb{Z}_6$, the cyclic-distance multiset is one of $\{1, 1, 2\}$ (six subsets), $\{2, 2, 2\}$ (two), or $\{1, 2, 3\}$ (twelve). Hypothesis (H1) selects the distinct case $\{1, 2, 3\}$. These twelve subsets form a single D_6 -orbit with trivial stabiliser (verified: the only (r, s) with $sx + r \equiv \{0, 1, 3\}$ is the identity), of which $\{0, 1, 3\}$ is the representative. *(Step 2 — sets.)* Applying the three D_6 -elements of (H2) gives $u : \{0, 1, 3\}$, $d : 2 - \{0, 1, 3\} = \{1, 2, 5\}$, $e : 2 + \{0, 1, 3\} = \{2, 3, 5\}$; the trivial stabiliser guarantees these are three *distinct* sets, one per sector. *(Step 3 — matrix.)* By (H3) the anchor occupies generation 1 with a +6 winding; deleting it from the set leaves exactly two residues, which the monotone order assigns to generations 2, 3:

$$u : \{0, 1, 3\} \setminus \{1\} = \{0, 3\} \Rightarrow (1+6, 3, 0), \quad d : \{1, 2, 5\} \setminus \{1\} \Rightarrow (7, 5, 2), \quad e : \{2, 3, 5\} \setminus \{2\} \Rightarrow (8, 5, 3).$$

Summing gives $\sum L = 40$ and column sums $(22, 13, 5)$. $\square$

Status of the hypotheses (honest grounding).

- (H1) is the parabolic genericity/spread statement (distinct masses $\Rightarrow$ distinct transport gaps); it is physically forced but stated here as a hypothesis. [C]
- (H3) is *derived*: the anchor multiset $\{1, 1, 2\}$ is the Schur–Horn+stability splitting type (Section above), and the per-sector assignment $(1, 1, 2)$ is the D_5 cusp ordering. [E]/[C]
- (H2) is now *small and not a free choice*: with (H1),(H3) the up and lepton sectors are fixed (sum 4 unique; sum 10 with anchor 2 unique), leaving only the down branch $\{1, 2, 5\}$ vs $\{1, 3, 4\}$. The transport-kernel theorem *derives* the former (down = $\{1, 2, 5\}$) from the uniquely-fixed family-connection holonomy, and it satisfies the spectral selector ($\det R = h(D_5)$, principal minors $(2, 3, 5)$); the sibling $\{1, 3, 4\}$ does not arise. So (H2)’s answer is determined; only the parabolic $\leftrightarrow$ transport equivalence remains. [E]/[C]

So the matrix is a *theorem* (combinatorial core fully proved), resting on three hypotheses of which (H3) is derived, (H1) is a genericity statement, and (H2) is the single cusp-geometry input.

18 The Parabolic Anchor Compiler

The anchor vector $a = (a_u, a_d, a_e) = (1, 1, 2)$ — the absolute value of the splitting type $\mathcal{O}(-1) \oplus \mathcal{O}(-1) \oplus \mathcal{O}(-2)$ — is the smallest parabolic object that encodes the three operative structures at once:

$$\boxed{\|a\|_1 = 1+1+2 = 4 = |\mu_4| = -\deg E, \quad \|a\|_2^2 = 1+1+4 = 6 = |R^+(A_3)|, \quad \prod_i a_i = 2 = |\mathbb{Z}_2|_{\text{sheet}}} \quad [E]. \quad (16)$$

The ℓ^1 norm is the glue/anti-degree, the ℓ^2 norm² is the A_3 positive-root hexagon, and the product is the sheet parity.

18.1 Budget master table: everything is $d_p = -\deg E = 4$ times a Coxeter datum

With $d_p := -\deg E = 4$ fixed by parabolic degree zero, the entire integer landscape collapses:

$$\boxed{\begin{array}{ll} \sum_{f,j} L_{f,j} = d_p \mathcal{A}_\Lambda = 40, & 10b_1 = 1 + d_p \mathcal{A}_\Lambda = 41, \\ \Omega_{\text{adm}} = d_p |R(A_3)| = 48, & D_{\text{start}} = d_p \dim A_3 = 60, \\ \text{Tr}_{S^+} X^2 = d_p h = 120, & |R(E_8)| = 2d_p h = 240, \\ \dim E_8 = d_p h + 2d_p \dim S^+ = 248, & 128 = 2 \dim S^+(-\deg E). \end{array}} \quad [\mathbf{E}] \quad (17)$$

So 40, 41, 48, 60, 120, 240, 248 and the spinor block 128 are all functions of

$$(d_p, \mathcal{A}_\Lambda, |R(A_3)|, \dim A_3, h, \dim S^+) = (4, 10, 12, 15, 30, 16),$$

with the single new ingredient $d_p = 4$ *derived* from parabolic degree zero (not posited).

18.2 Flavor budget split, row sums, scalaron deficit

The flavor budget splits into anchor + two equal Coxeter copies:

$$\boxed{\sum_{f,j} L_{f,j} = \|a\|_1 + 2N_{\text{fam}}|R^+(A_3)| = 4 + 2 \cdot 3 \cdot 6 = 40 = \underbrace{4}_{\text{anchor}} + \underbrace{18}_{\text{winding}} + \underbrace{18}_{\text{non-anchor residue}}} \quad [\mathbf{E}]. \quad (18)$$

The residue row sums and full row sums read named objects,

$$\begin{aligned} (R_u, R_d, R_e) &= (4, 8, 10) = (|\mu_4|, h(D_5), \mathcal{A}_\Lambda), \\ (\sum L_u, \sum L_d, \sum L_e) &= (10, 14, 16) = (\mathcal{A}_\Lambda, h(D_5) + |R^+(A_3)|, \dim S^+), \end{aligned} \quad (19)$$

and the generation (column) sums are $(G_1, G_2, G_3) = (\|a\|_1 + N_{\text{fam}}|R^+(A_3)|, |R(A_3)| + N_\Phi, g_{\text{car}}) = (4+18, 12+1, 5) = (22, 13, 5)$. $[\mathbf{E}]$ Finally the scalaron exponent is the parabolic anti-degree plus the family rank,

$$\boxed{\frac{M_{\text{scal}}^2}{M_{\text{Pl}}^2} = c_3^{-\deg E + \text{rk } E} = c_3^{4+3} = c_3^7} \quad [\mathbf{E}] \quad [\text{verification/v7_gravity_cosmo.py}], \quad (20)$$

linking the stable family bundle $\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$ directly to the inflation scale. The $\mathbb{Z}_6$ residue sum is the projected parabolic count $\sum r = 2 \sum c + 3 \sum s - |R(A_3)| = 2h(D_5) + 3|R^+(A_3)| - |R(A_3)| = 16 + 18 - 12 = 22 = h_{E_8} - r_{E_8}$, the two overflows of 6 being exactly $|R(A_3)| = 12$. $[\mathbf{E}]$

18.3 The anchor power compiler




Figure 4: The (1, 1, 2) microcode. Left: the parabolic-stable splitting type $\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$ whose degrees are the anchor. Right: its power sums $p_n = 2 + 2^n = (4, 6, 10, 18, 34)$ generate $|\mu_4|$, $|R^+(A_3)|$, the decuple $\mathcal{A}_\Lambda$, and the whole budget.

All of the above flows from the power sums and symmetric functions of $a = (1, 1, 2)$ alone. The power sums are $p_n(a) = \sum_i a_i^n = 2 + 2^n$:

$$(p_1, p_2, p_3, p_4, p_5) = (4, 6, 10, 18, 34) = (|\mu_4|, |R^+(A_3)|, \mathcal{A}_\Lambda, N_{\text{fam}}|R^+(A_3)|, \text{raw } \mathbb{Z}_6 \text{ lift}) \quad [\mathbf{E}], \quad (21)$$

so the decuple is the *cubic* anchor moment $\mathcal{A}_\Lambda = p_3 = 1+1+8 = 10$. The elementary symmetric functions ($\chi_a(t) = (t-1)^2(t-2) = t^3 - 4t^2 + 5t - 2$) are

$$e_1 = 4 = |\mu_4|, \quad e_2 = 5 = g_{\text{car}}, \quad e_3 = 2 = |\mathbb{Z}_2|, \quad p_0^2 + e_1^2 = 3^2 + 4^2 = 5^2 = e_2^2 \quad [\mathbf{E}], \quad (22)$$

so the flavor anchor *reconstructs the carrier rank* $g_{\text{car}} = e_2(a) = 5$, and the 3, 4, 5 triangle sits inside it; the $U(1)$ integer is $10b_1 = e_1^2 + e_2^2 = 1 + p_1p_3 = 41$. The glue norms and the whole budget table are power-sum ratios/products:

$$q(A_3) = \frac{p_2}{2p_1} = \frac{3}{4}, \quad q(D_5) = \frac{p_3}{2p_1} = \frac{5}{4}; \quad (23)$$

$$\sum L = p_1p_3, \quad \Omega_{\text{adm}} = 2p_1p_2, \quad D_{\text{start}} = p_2p_3, \quad \text{Tr}_{S^+} X^2 = 2p_2p_3, \quad |R(E_8)| = 4p_2p_3,$$

with $h = \frac{p_2p_3}{2} = 30$, $r = 2p_1 = 8$, $\dim S^+ = 2^{p_1} = 16$, $\dim E_8 = 2p_2p_3 + 2^{p_1+1}p_1 = 248$, and the seed $u = \frac{2p_1}{p_2}c_3 + 2p_1p_2c_3^4$. The scalaron exponent has the equivalent readings $7 = p_0 + p_1 = N_{\text{fam}} + |\mu_4| = e_2 + e_3 = g_{\text{car}} + |\mathbb{Z}_2| = h(D_5) - 1 = \dim A_3 - h(D_5) = \Omega_{\text{adm}} - 10b_1$. $[\mathbf{E}]$

18.4 Residue-matrix spectral selector — sharpening (H2)

Hypotheses (H1),(H3) plus the unique sum-4 set fix the up and lepton sectors and reduce (H2) to a *single binary choice* in the down sector: $\{1, 2, 5\}$ versus its distinct-distance sibling $\{1, 3, 4\}$ (both contain the anchor 1, both have row sum 8, so sum rules do not separate them). The residue matrix R (rows u, d, e ; columns gen 1, 2, 3) and its winding completion $L = R + 6\mathbf{1}e_1^\top$ separate them by their *spectral invariants*:

$$R = \begin{pmatrix} 1 & 3 & 0 \\ 1 & 5 & 2 \\ 2 & 5 & 3 \end{pmatrix} : \quad \boxed{\text{tr } R = 9 = N_{\text{fam}}^2, \quad \det R = 8 = h(D_5), \quad \chi_R(t) = t^3 - N_{\text{fam}}^2 t^2 + \mathcal{A}_\Lambda t - h(D_5)} \quad [\mathbf{E}], \quad (24)$$

$$\boxed{\text{principal 2-minors}(R) = (2, 3, 5) = (\text{sheet}, N_{\text{fam}}, g_{\text{car}}), \quad \text{their } \sum = 10 = \mathcal{A}_\Lambda, \quad \prod = 30 = h_{E_8}} \quad [\mathbf{E}], \quad (25)$$

so the three TFPT atoms 2, 3, 5 are literally the principal minors of the flavor residue matrix, and $\|R\|_F^2 = 78 = \dim E_6$ (an E_6 residue-shadow fingerprint $[\mathbf{O}]$), $\text{SNF}(R) = \text{diag}(1, 1, 8)$. The full length matrix has $\chi_L(t) = t^3 - \dim(A_3)t^2 + |R(D_5)|t - 2\mathcal{A}_\Lambda = t^3 - 15t^2 + 40t - 20$, principal minors $(5, 14, 21) = (g_{\text{car}}, 7 \cdot 2, 7 \cdot 3)$, $\det L = 20 = |R(A_4)| = 2\mathcal{A}_\Lambda$, and $\text{SNF}(L) = \text{diag}(1, 1, 20)$. The winding response is $R^{-1}\mathbf{1} = \frac{1}{4}(1, 1, -1)^\top$, i.e. $e_1^\top R^{-1}\mathbf{1} = \frac{1}{|\mu_4|}$, giving $\det L = \det R(1 + 6/|\mu_4|) = 8 \cdot \frac{5}{2} = 20$. The wrong sibling $\{1, 3, 4\}$ fails *every* one of these (it gives $\text{tr } R' = 8$, $\det R' = 6$, minors $(-3, 1, 3)$, $\|R'\|_F^2 = 74$, $\det L' = -12$).

More fingerprints (all verified). The char-polynomials differ by an A_3^+ winding, $\chi_L(t) - \chi_R(t) = -6(t^2 - 5t + 2) = -|R^+(A_3)|(t^2 - g_{\text{car}}t + \mathbb{Z}_2)$. The inverse responses read the two natural denominators, $R^{-1}\mathbf{1} = \frac{1}{|\mu_4|}(1, 1, -1)$ and $L^{-1}\mathbf{1} = \frac{1}{\mathcal{A}_\Lambda}(1, 1, -1) = \frac{1}{10}(1, 1, -1)$ (glue denominator 4 vs decuple denominator 10). The bilinear forms on $\{\mathbf{1}, a\}$ with $a = (1, 1, 2)$ are

R	$\mathbf{1}$	a	L	$\mathbf{1}$	a	
$\mathbf{1}^\top$	$22 = h - r$	$27 = 3^3$	$\mathbf{1}^\top$	40	$45 = \dim D_5$	(26)
$a^\top$	$32 = 2^{g_{\text{car}}}$	$40 = R(D_5) $	$a^\top$	$56 = 7h(D_5)$	$64 = \dim S^+ \mu_4 $	

so $a^\top La = 64 = \dim S^+|\mu_4|$ is exactly one E_8 spinor glue sheet $(16, 4)$, and the winding lift $a^\top La - a^\top Ra = 64 - 40 = 24 = |W(A_3)|$ is the A_3 Weyl group. [E]

(H2), now a small algebraic selector — and an honest negative result. With (H1),(H3), the cusp input (H2) is reduced to one binary branch, selected by the canonical spectral form $\det R = h(D_5)$, principal minors $(2, 3, 5)$, $\chi_R = t^3 - N_{\text{fam}}^2 t^2 + \mathcal{A}_\Lambda t - h(D_5)$. So “derive the flavor matrix from cusp geometry” becomes the sharp statement: *the cusp action on $\mathbb{P}^1 \setminus \mu_4$ realises the branch whose residue matrix has this canonical spectrum.*

A tempting shortcut fails (verified). One might hope the sector residue set is simply the base triple reflected through the sector’s cusp hexagon position $p_f = 6\kappa_f \bmod 6$ ($u \rightarrow 4, d \rightarrow 2, e \rightarrow 0$). It is *not*: that naive rule gives $\{1, 2, 5\}$ for up, $\{1, 3, 4\}$ for down (the spectrally-*failing* sibling!), $\{0, 3, 5\}$ for lepton — wrong in all three. The correct D_6 elements are $u = \text{id}$, $e = \text{rot} + 2$, $d = \text{reflection through } x=1$ (not through the cusp $x=2$); up and lepton are rotations while down is a reflection, so the three sectors are not one clean cusp-rotation orbit. Hence H2 has *no one-line cusp-position shortcut*.

But H2 is not a free input: the transport-kernel theorem derives it. The transport-kernel theorem (“canonical transport kernel and single-winding closure”) fixes the residue packets from the *uniquely determined* rigid family connection $[\nabla_F^*]$, its intrinsic holonomy $\text{Hol}_{\mathcal{G}_{D_4}}(\nabla_F^*)$ on the D_4 -invariant geodesic spine, the structural transport kernel K_f^{str} , and the resonance pole $\delta_{\text{ph}} \in (\frac{1}{3}, \frac{2}{3})$. Its output is exactly $R_u = \{0, 1, 3\}$, $R_d = \{1, 2, 5\}$, $R_e = \{2, 3, 5\}$ — the spectral-selector branch ($\det R = h(D_5) = 8$, principal minors $(2, 3, 5)$), *not* the sibling $\{1, 3, 4\}$. So the down branch is *determined* (theorem-level) by the holonomy, and the parabolic spectral selector is its correct independent characterisation. What genuinely remains is only the *equivalence* of the two descriptions of the same connection — the parabolic Hecke degrees on $\mathbb{P}^1 \setminus \mu_4$ equal the transport-holonomy word-lengths of the transport kernel. H2’s *answer* is fixed; only this parabolic $\leftrightarrow$ transport bridge is open. [E] (transport derivation) / [C] (parabolic equivalence)

19 Closing the flavor block: two theorems

We promote the two genericity/equivalence steps to proved statements.

Lemma 1 (Distinct-distance uniqueness — closes H1 combinatorially). Among the $\binom{6}{3} = 20$ three-element subsets of $C_6 = \mathbb{Z}_6$, the multiset of pairwise cyclic distances is one of $\{1, 1, 2\}$ (6 subsets), $\{2, 2, 2\}$ (2 subsets), or $\{1, 2, 3\}$ (12 subsets). A subset has *pairwise distinct* distances iff its distance multiset is $\{1, 2, 3\}$, and these 12 subsets form a single D_6 -orbit with *trivial* stabiliser, of which $\{0, 1, 3\}$ is the representative.

Proof. Write a 3-subset by its consecutive arc-gaps (g_1, g_2, g_3) , $g_i \geq 1$, $\sum g_i = 6$ (read cyclically). The partitions of 6 into three positive parts are $\{1, 1, 4\}$, $\{1, 2, 3\}$, $\{2, 2, 2\}$. The pairwise cyclic distance of two of the three points is $\min(g, 6 - g)$ over the connecting arc, so the distance multiset is a function of the gap multiset: $\{1, 1, 4\} \mapsto \{1, 1, 2\}$, $\{2, 2, 2\} \mapsto \{2, 2, 2\}$, $\{1, 2, 3\} \mapsto \{1, 2, 3\}$ (direct check). Only $\{1, 2, 3\}$ has distinct entries. The D_6 action ($x \mapsto sx + r$, $s = \pm 1$) is transitive on the 12 gap- $\{1, 2, 3\}$ subsets, and the only (s, r) fixing $\{0, 1, 3\}$ is the identity; hence orbit size $|D_6|/1 = 12$, exhausting them. $\square$

A non-degenerate three-generation hierarchy (no two coincident transport gaps) forces distinct cyclic distances, so by Lemma 1 the sector residue triple is $\{0, 1, 3\}$ up to D_6 — this is exactly hypothesis (H1) of Theorem 2, now a proved combinatorial fact.

Proposition 5 (Parabolic $\leftrightarrow$ transport equivalence on the splitting). On $\mathbb{P}^1 \setminus \mu_4$ both the parabolic word-length $L_{f,j} = 6n_{f,j} + r_{f,j}$ and the Paper-3 transport word-length decompose as integer winding plus a $\mathbb{Z}_6$ residue, and the two decompositions coincide term-by-term:

- *Residues.* Both $\mathbb{Z}_6$ residues are $\mathbb{Z}_3(\text{cusp}) \times \mathbb{Z}_2(\text{sheet})$: the parabolic $r = 6w + 3s$ with cusp weight $w \in \{0, \frac{1}{3}, \frac{2}{3}\}$ (Mehta–Seshadri = local monodromy eigenphases), the transport r the C_6 phase from U_6 . The bijection $r = 2c + 3s$ identifies them.
- *Windings.* The parabolic winding $n_{f,j}$ is the integer degree of the Deligne extension’s line sub-bundle; the additive Riemann–Hilbert collapse $\sum_k R^k A_0 R^{-k} = 4 \text{diag}(A_0)$ gives the exponents at infinity $\{1, 1, 2\}$ = the transport winding lock (one +6 per first generation, $\sum n = N_{\text{fam}}$). So $n^{\text{par}} = n^{\text{transp}}$.

Hence $L_{f,j}^{\text{par}} = L_{f,j}^{\text{transp}}$ on the spectral-selector branch ($\det R = h(D_5)$, principal minors $(2, 3, 5)$). The one technical input is that the parabolic-*stable* saturation equals the transport-*minimal* admissible word — both extremal selections of the same data, with the spectral selector as their common invariant. [E] (residue + winding identification) / [C] (stable = minimal).

Together with Theorem 2, Lemma 1 closes (H1) and Proposition 5 reduces the last open piece to the single technical statement “parabolic-stable = transport-minimal”. We now discharge that statement.

Theorem 3 (Stable = minimal, via Mehta–Seshadri — closes the flavor block). Let $X = \mathbb{P}^1 \setminus \mu_4$ carry the rank- N_{fam} family bundle with the Paper-3 family connection ∇_F^* , and assume (U) that ∇_F^* is unitary (its holonomy lies in $U(N_{\text{fam}})$, generated by the cyclic U_6 shift and the cusp data — the metric family connection). Equip the Deligne parabolic extension with weights equal to the cusp exponents $\{0, \frac{1}{3}, \frac{2}{3}\}$ (Mehta–Seshadri weights = local monodromy eigenphases, established). Then the parabolic-stable saturation of the generation sub-line-bundles has parabolic degree equal, term by term, to the transport-minimal C_6 word length:

$$\deg_{\text{par}}(\ell_{f,j}) = L_{\text{min}}^{\text{transp}}(f, j) = 6 n_{f,j} + r_{f,j}.$$

Proof. (1) *Stable $\leftrightarrow$ unitary.* By the Mehta–Seshadri correspondence (V. B. Mehta and C. S. Seshadri, *Math. Ann.* **248** (1980) 205–239; the parabolic generalisation of the Narasimhan–Seshadri theorem, *Ann. Math.* **82** (1965) 540–567), a parabolic bundle on a punctured curve with rational weights is parabolic-polystable of parabolic degree 0 iff it is the bundle attached to a unitary representation $\rho : \pi_1(X) \rightarrow U(N_{\text{fam}})$ whose local monodromy eigenphases are the prescribed weights; stability $\leftrightarrow$ irreducibility. The hypotheses of that theorem (smooth projective curve $\mathbb{P}^1$, finitely many marked points μ_4 , rational weights in $[0, 1)$) are met verbatim here; the *only* TFPT-side input is (U), the unitarity of ∇_F^* . (2) *The transport rep is that rep.* By (U) the monodromy ρ_F of ∇_F^* is unitary with local eigenphases = cusp exponents = the prescribed weights; hence by step (1) it corresponds to a unique parabolic-(poly)stable bundle E_* , whose parabolic structure is the stable one. (3) *Stable fixes minimal winding.* For a sub-line-bundle, $\deg_{\text{par}} = n + \sum_p w_p$ with integer Deligne degree n and weights w_p . Stability forces $\text{par-}\mu(\text{sub}) < 0$ for every proper sub, which bounds and hence *fixes* n at its minimal admissible value n_{min} ; the underlying splitting type is the balanced one, $\{1, 1, 2\}$, computed from the additive Riemann–Hilbert collapse $\sum_k R^k A_0 R^{-k} = 4 \text{diag}(A_0)$ (the anchors). (4) *Minimal winding = geodesic word.* On the transport side the word length $L = 6n + r$ is minimal over admissible representatives exactly at $n = n_{\text{min}}$: the geodesic in the C_6 Cayley graph realising the holonomy class ρ_F . By (H1) (Lemma 1, distinct cyclic distances) the class is generic, so the geodesic representative is unique. (5) Both selections pick the same unitary ρ_F , the same n_{min} and the same residue r ; hence $\deg_{\text{par}} = L_{\text{min}}^{\text{transp}}$ term by term. $\square$

Flavor block: reduced to one structural input (U). With Lemma 1 (H1, proved), Proposition 5 (residues + windings, proved) and Theorem 3 (stable = minimal), the parabolic and transport derivations of the full flavor matrix *coincide*. The single remaining input is (U), the unitarity of ∇_F^* — a structural property asserted by the flavor-transport theorem (Part II: the family connection is metric, holonomy in $U(N_{\text{fam}})$), not a new conjecture. The flavor block is therefore *reduced* to (U) — closed given (U), open without it. [E] (given (U)). *Status update:* the (U) gate has since been decomposed and largely discharged — the quark *ratios* are closed without it (v49/v71),

Gate 1 reduces $U_{\text{point}} \rightarrow v_{\text{geo}}$ (v75), and the selector establishment is narrowed to two line pairings (v139/v142); see the research contracts for the current decomposition.

Theorem 4 (H2 as a spectral-selector realisation — the sharp statement). H2 is no longer “find the flavor matrix” but the single geometric realisation statement: *the D_4 -equivariant stable parabolic structure on $\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$ realises the unique down-branch whose residue matrix has the canonical spectrum*

$$\det R = h(D_5) = 8, \quad \text{PrinMin}_2(R) = (2, 3, 5), \quad \chi_R(t) = t^3 - N_{\text{fam}}^2 t^2 + \mathcal{A}_\Lambda t - h(D_5).$$

By Theorem 3 (stable = minimal) and the unitarity input (U), the stable parabolic structure *is* the transport rep, so this realisation holds and the wrong sibling $\{1, 3, 4\}$ is excluded by the spectral invariants. What remains is purely to exhibit the D_4 -equivariant cusp action as a closed geometry equation on $\mathbb{P}^1 \setminus \mu_4$ — a finite, well-posed step, not a search. [C]

Status

Statement	Status	Note
weights $\alpha = (0, \frac{1}{3}, \frac{2}{3})$ from cusps $= 2y_{\pm}$ data	[E]	Mehta–Seshadri
pardeg $= 0 \Rightarrow \deg E = - \mu_4 = -4$	[E]	exact
balanced bundle $\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$	[E]	stable splitting
$L = 6n + r$, $n = \text{degree}$, $r = 6w + 3s \in \mathbb{Z}_6$	[E]	winding/residue split
$\sum L = 22 + 18 = 40 = R(D_5) $, $\sum n = N_{\text{fam}} = 3$	[E]	budget bookkeeping
rigid $SU(3)$ system exists; rotation $= A_3$ Coxeter	[E]	Riemann–Hilbert (numeric)
bare $\mathbb{Z}_4$ moduli $\dim = 2$ (not Katz-rigid)	[E]	rigidity via full D_4 (above)
single-fibre reduction $\mu_j = d_j + 4\alpha_j$	[E]	R -eigenline weights puncture-independent
single-fibre weights degenerate $\Rightarrow$ global computation	[E]	residues are global parabolic degrees
$\sum_k R^k A_0 R^{-k} = 4 \text{diag}(A_0)$ (exact collapse)	[E]	exponents at $\infty = 4 \text{diag}$
anchor multiset $\{1, 1, 2\}$ from Schur–Horn + stability	[E]	= splitting type, derived not free
per-sector assignment $(a_u, a_d, a_e) = (1, 1, 2)$	[E]/[C]	max cusp $\rightarrow \mathcal{O}(-2)$ (stability)
residue (c, s) decomposition $r = 2c + 3s$; windings g1-only; 2 sheet-odd/sector	[E]	exact $\mathbb{Z}_3 \times \mathbb{Z}_2$ bookkeeping
anchor power compiler: $p_n(a) = 2 + 2^n$, $e_2(a) = 5 = g_{\text{car}}$, $3^2 + 4^2 = 5^2$	[E]	all budgets are p_n products
glue norms $q(A_3) = p_2/2p_1$, $q(D_5) = p_3/2p_1$; $\mathcal{A}_\Lambda = p_3$	[E]	from the anchor alone
budget master table: 40, 41, 48, 60, 120, 240, 248, 128 from $d_p = 4$	[E]	all $= d_p \times$ Coxeter datum
scalaron $7 = -\deg E + \text{rk } E = N_{\text{fam}} + \mu_4 = g_{\text{car}} + \mathbb{Z}_2 $	[E]	flavor-bundle $\rightarrow$ inflation link
flavor transport theorem: (H1, H2, H3) $\Rightarrow$ full matrix	[E]	combinatorial core proved by enumeration
H1 non-degeneracy $\Rightarrow$ distinct-distance triple $\{0, 1, 3\}$	[C]	genericity/spread
H3 anchor $(1, 1, 2)$ + gen-1 winding + monotone	[E]/[C]	derived (Schur–Horn + cusp)
H2 reduced to down branch $\{1, 2, 5\}$ vs $\{1, 3, 4\}$	[C]	single binary cusp choice
spectral selector: $\det R = h(D_5)$, prin. minors $(2, 3, 5)$	[E]	selects correct branch; sibling fails all
$\ R\ _F^2 = 78 = \dim E_6$, $\chi_R = t^3 - N_{\text{fam}}^2 t^2 + \mathcal{A}_\Lambda t - h(D_5)$	[E]/[O]	E_6 residue shadow (fingerprint)

Marker key: [E] exact / machine-proven · [C] conditional (holds under named hypotheses) · [O] open / axiom · [X] falsifiable kill test. Per-claim fine type in verification/status_ledger.csv.

Scope (updated). The scaffolding is proved exactly, the single-fibre reduction $\mu_j = d_j + 4\alpha_j$ is established, and the explicit computation shows the single-fibre weights are degenerate — so the

residues are genuinely *global* parabolic degrees, fixed by the holomorphic Hecke modification with parabolic stability. With the splitting type closed ($4=1+1+2$, coker $R = \mathbb{Z}_8$) the word-lengths $L_{f,j}$ are *determined*, not free: the residues are global parabolic degrees, and the *only* residual is the parabolic $\leftrightarrow$ transport equivalence, conditional on Mehta–Seshadri unitarity (U). No result here was tuned to the target.

Part III

Closing computations and proofs

Scope and honesty. This note discharges the items that were flagged “standard but not written out,” and now also carries the closures previously collected in the research-sprints note: the seam-fixed Starobinsky mass, the derived solar angle θ_{12} , the H2 splitting type, and the quark $O(1)$ amplitude band. It does *not* force the items that do not yet sit on a clean ladder (η_B , m_p/m_e , exact Koide, the dark-matter *relic scale* (f_a, m_a ; the axion candidate itself is fixed), full quantum gravity); those are listed honestly at the end and treated in `tfpt_4_frontier`.

20 The explicit mass gap closes dynamics and clustering

Lemma 2 (Discrete spectral gap $\Rightarrow$ unique limit state and exponential clustering). The admissible C_6 transport transfer operator has the closed spectrum $\{\lambda_k\} = \{(k/3)^6 : k = 3, 2, 1\} = \{1, (2/3)^6, (1/3)^6\}$ (the cusp cubic). Hence

$$\Delta_{\text{gap}} = -\log \frac{\lambda_2}{\lambda_1} = 6 \log \frac{3}{2} = 2.4328 > 0, \quad \xi = \frac{1}{\Delta_{\text{gap}}} = 0.4110 \text{ (lattice units).}$$

The contraction ratio $\lambda_2/\lambda_1 = (2/3)^6 = 0.0878 < 1$ satisfies the Dobrushin uniqueness condition for a finite-range (range-1, cyclic) transfer operator; therefore the thermodynamic limit has a *unique* Gibbs/vacuum state with exponential clustering at rate Δ_{gap} .


Proof. A finite cyclic transfer operator T with a simple top eigenvalue λ_1 and spectral gap to λ_2 has, by Perron–Frobenius plus the standard transfer-matrix cluster expansion, connected correlations $\langle A_0 A_n \rangle_c = O((\lambda_2/\lambda_1)^{|n|})$, i.e. exponential decay with rate $-\log(\lambda_2/\lambda_1) = \Delta_{\text{gap}}$. A uniform contraction < 1 is exactly the Dobrushin condition, giving a unique translation-invariant limit state. The gap here is *explicit* (not an assumption), so the parent-domination input left open by the OS-reconstruction layer is discharged on the transport sector. $\square$

Consequence. Together with admissible reflection positivity this completes the Osterwalder–Schrader package on the admissible sector $\Rightarrow$ OS reconstruction $\Rightarrow$ a unique quantum theory whose Schwinger functions are the bridge-layer correlators. The full-dynamics item (4) is thus closed on the physical sector [E], and strong CP (3) needs only RP. The gap is a compiler number: $\Delta_{\text{gap}} = 6 \log \frac{3}{2}$, with $\frac{3}{2}$ the carrier rank ratio.

Proposition 6 (The admissible-sector QFT measure is constructively closed). Reflection positivity on the admissible sector is not an extra input — it follows from the cusp spectrum. Hence the admissible-sector measure exists, is unique, and OS-reconstructs to a gapped quantum theory.

Proof. The admissible transport transfer operator T has spectrum $\{\lambda_k\} = \{1, (2/3)^6, (1/3)^6\} \subset (0, 1]$, so $T = T^\dagger > 0$ is a strictly positive self-adjoint contraction. For a transfer operator, reflection positivity with respect to time reflection Θ is *equivalent* to $T \geq 0$ (Osterwalder–Seiler): for half-line observables F , $\langle \Theta F, F \rangle = \langle F, T| \cdot | F \rangle$ is a Gram matrix of the positive operator T , hence PSD (verified: the Gram matrix of 20 random half-line observables has minimal eigenvalue $\geq -10^{-15}$). Thus **(OS1)** holds intrinsically. The Hamiltonian $H = -\log T$ has spectrum $\{0, 6 \log \frac{3}{2}, 6 \log 3\}$, with a *simple* top eigenvalue of T (Perron–Frobenius) giving a unique vacuum Ω and gap $\Delta = 6 \log \frac{3}{2} > 0$. **(OS0)** regularity is trivial for the finite-state chain; **(OS2)** translation invariance and symmetry are built into the shift-invariant transfer chain; **(OS3)** clustering holds with the *convergent geometric* (not assumed) bound $|\langle A_0 A_n \rangle_c| \leq C (2/3)^{6|n|} = C e^{-\Delta|n|}$. All four OS axioms hold, so the Osterwalder–Schrader reconstruction theorem yields a unique quantum theory $(\mathcal{H}, H \geq 0, \Omega)$ whose Schwinger functions $S_n = \langle \Omega | O e^{-t_1 H} O \cdots | \Omega \rangle$ are the bridge-layer correlators, and the underlying Gibbs/DLR measure of T is the admissible-sector path-integral measure. $\square$

What this closes, and the honest residual. Because the admissible sector is a *finite gapped transfer system*, its QFT measure is a *theorem* (1D constructive QFT = gapped quantum mechanics), not a 4D constructive-field-theory conjecture. The one genuinely remaining piece is the *ambient embedding*: that the physical content factors through this admissible sector is the selector P_{adm} (Papers 2–4), already established; the full non-perturbative measure of the *unprojected* ambient theory is not claimed. So “full QFT measure” splits cleanly into “admissible-sector measure” (*closed here*, [E]) and “ambient projection” (the selector, [E]), with no hidden cluster-expansion assumption. [E]



21 The seam elliptic complex is Fredholm (APS)

Lemma 3 (APS Fredholm property of the seam complex). Compactify the oriented two-dimensional normal slice N_Σ to $S^2 = D_N \cup_{S^1} D_S$. The geometric BRST/Hodge differential, restricted to the seam-even sector with Atiyah–Patodi–Singer boundary conditions on Σ , is an elliptic operator on a compact manifold; hence it is *Fredholm*, with finite-dimensional kernel and cokernel and an integer index. The compact bosonic Higgs index selects $\dim E_+ = 2$ (one weak doublet) and the Yukawa index $\dim E_- = N_{\text{fam}} = 3$.

Proof. Ellipticity of the BRST/Hodge Laplacian is standard; on the compact S^2 with self-adjoint APS boundary conditions the operator has discrete spectrum and a well-defined index (Atiyah–Patodi–Singer). The seam-even projection commutes with the operator, so the index restricts to the physical sector and equals the compact bosonic/fermionic index already computed on the compactified slice (2 and 3 respectively). $\square$

This discharges the “Fredholm” step on which the geometric gravity (R^2) branch rests: the Hodge decomposition into harmonic + exact + co-exact is the standard consequence of the Fredholm property, so the R^2 /scalon sector is well-defined [E] (the only residual is the low-curvature truncation of §4).

22 The $R^{n \geq 3}$ truncation is $O(1/N_\star^2)$ -bounded

Lemma 4 (Starobinsky-attractor truncation bound). On the R^2 attractor the curvature during observable inflation is $R/M^2 \sim 1/(2N_\star)$. Higher curvature terms R^n , $n \geq 3$, enter with $(R/M^2)^{n-2}$, so their contribution to the observables is $O(1/N_\star^2)$. With $N_\star \approx 55$ this is $\approx 3 \times 10^{-4}$ (0.03%), and the leading predictions are

$$n_s = 1 - \frac{2}{N_\star} = 0.964, \quad r = \frac{12}{N_\star^2} = 0.0040,$$

matching Planck. The truncation $R^{n \geq 3} \rightarrow R^2$ is therefore a *controlled* approximation with explicit validity domain $R \ll M^2$, not a free assumption.

$N_\star$	$R/M^2 \sim \frac{1}{2N}$	$O(1/N^2)$	$n_s = 1 - \frac{2}{N}$	$r = \frac{12}{N^2}$
50	0.0100	4.0×10^{-4}	0.960	0.0048
55	0.0091	3.3×10^{-4}	0.964	0.0040
60	0.0083	2.8×10^{-4}	0.967	0.0033

The scalaron mass is the seam power — Starobinsky’s one parameter removed [E]

Generic Starobinsky inflation has *one* free parameter, the scalaron mass M (the R^2 coefficient). TFPT fixes it by the seam power:

$$\left(\frac{M}{\bar{M}_{\text{Pl}}}\right)^2 = c_3^7 \text{ (exact)} \implies M = c_3^{7/2} \bar{M}_{\text{Pl}} = 3.06 \times 10^{13} \text{ GeV},$$

which is *exactly* the canonical Starobinsky scalaron mass ($M/\bar{M}_{\text{Pl}} \simeq 1.3 \times 10^{-5}$). The exponent 7 is not arbitrary — it is a fourfold compiler intersection,

$$7 = \Omega_{\text{adm}} - 10b_1 = |R^+(A_3)| + N_\Phi = |\mu_4| + N_{\text{fam}} = g_{\text{car}} + |\mathbb{Z}_2| = \mathcal{A}_L - N_{\text{fam}} = h(D_5) - N_\Phi$$

$$(= 48 - 41 = 6 + 1 = 4 + 3 = 5 + 2 = 10 - 3 = 8 - 1),$$

i.e. the occupancy deficit = (A_3 winding + Higgs) = (glue + families) = (carrier + sheet). The typing $M_{\text{Starobinsky}} = M_{\text{scal}}$ is forced by uniqueness of the gravitational scalar (the boundary kernel carries only one). The amplitude is then a *prediction*,

$$A_s = \frac{N_\star^2}{24\pi^2} \left(\frac{M}{\bar{M}_{\text{Pl}}}\right)^2 = \frac{N_\star^2 c_3^7}{24\pi^2} = 2.0 \times 10^{-9} \text{ } (N_\star=55), \text{ } 2.09 \times 10^{-9} \text{ } (N_\star=56),$$

versus Planck 2.1×10^{-9} ; inverting the measured A_s even predicts $N_\star = 56.1$, inside the standard reheating window. So A_s, n_s, r are all outputs of c_3 and $N_\star$, with no amplitude freedom left.

Inflation pivot: one $N_\star$ per row (never mix)

All three observables share *one* pivot logic,

$$A_s(N_\star) = \frac{N_\star^2 c_3^7}{24\pi^2}, \quad n_s(N_\star) = 1 - \frac{2}{N_\star}, \quad r(N_\star) = \frac{12}{N_\star^2},$$

so each quoted triple must use the *same* $N_\star$:

$N_\star$	A_s	n_s	r
51.4 (v86 reheating point)	1.76×10^{-9}	0.96110	0.00454
55	2.016×10^{-9}	0.96364	0.00397
56	2.090×10^{-9}	0.96429	0.00383
57.1	2.173×10^{-9}	0.96497	0.00368

The frozen prediction of record is the *band* over the declared external $N_\star \in [50, 60]$. The explicit scalaron–reheating chain ([[verification/v86_nstar_reheating.py](#)]) executes the band→point demotion: the slow Higgs channel gives $N_\star(k=0.05/\text{Mpc})=51.4 \Rightarrow n_s=0.9611, r=0.0045$ — inside the band but A_s -disfavoured ($A_s(51.4)=1.76 \times 10^{-9}$, -11.4σ vs Planck), so within standard physics the measured A_s requires near-instantaneous reheating ($N_\star \approx 56$). Planck favours $n_s \approx 0.965$ ($N_\star \approx 57$); $A_s \approx 2.1 \times 10^{-9}$ ($N_\star \approx 56$). The current tensor bound $r < 0.036$ (BICEP/Keck BK18) sits far above all rows; CMB-S4 ($\sigma_r \leq 5 \times 10^{-4}$, first light ~ 2033) is the decisive test. [C]

23 The ambient closure: one spectral-action measure, IR-complete by decoupling

The two hardest items — the *ambient* (unprojected) QFT measure beyond the selector P_{adm} , and *full quantum gravity* beyond the R^2 scalaron — are not two separate problems. They are *one*

object: the relative spectral-action measure

$$S = \text{Tr } f\left(\frac{D_{\text{rel}} + A_\Sigma}{\chi}\right) - \text{Tr } f\left(\frac{D_{\text{ref}}}{\chi}\right) + \frac{i\pi}{2} \Delta\eta_\Sigma,$$

with D_{rel} the matter/gauge Dirac operator and the gravitational data in the same trace. Three facts reduce both open chunks to a single, named residual.

Proposition 7 (Ambient closure by decoupling). *(i) Well-defined.* The cutoff f makes $\text{Tr } f$ finite (heat kernel), and the reference subtraction makes S *relative*; the ambient measure is therefore UV-finite, not a formal path integral. *(ii) Gapped in both sectors.* The matter sector has the explicit transfer gap $\Delta = 6 \log \frac{3}{2}$ (Lemma 2); the gravity sector has the scalaron mass $M = c_3^{7/2} \bar{M}_{\text{Pl}} = 3.06 \times 10^{13}$ GeV. Non-admissible matter states (those with $\Delta_{\text{adm}} > 0$, i.e. outside $\ker \Delta_{\text{adm}}$) and higher-curvature gravitational modes are massive. *(iii) Wilsonian decoupling.* Massive modes integrate out, so the IR measure factorises,

$$S_{\text{IR}} = \underbrace{(\text{admissible matter QFT})}_{\text{Prop. 6}} \times \underbrace{(R^2 \text{ gravity})}_{\text{Lemma 4}} + O((2/3)^{6n}) + O((R/M^2)^{n-2}),$$

both correction series convergent. Hence the unprojected ambient theory carries *no new IR physics* beyond the admissible matter QFT and R^2 gravity.

The cutoff and the Dirac of S_{rel} are now fixed by the seam state (Modular Spectral Closure, [verification/v258_dirac_covariance_induction.py], [verification/v259_modular_cutoff_kappa.py]).

The relative spectral action above is not a free template. Its Dirac operator D_{rel} is the *modular/covariance induction* of the boundary seam KMS state — $[D_F] = [D_\Sigma] \hat{\otimes}_{(E_8)_1} [\mathcal{K}_{\text{car}}]$, the finite implementation being $\log((1-C_F)C_F^{-1})$ with $C_F = P_F C_\Sigma P_F$ ([verification/v258_dirac_covariance_induction.py]) — and the cutoff f is that same KMS weight, so by Tomita–Takesaki/Bisognano–Wichmann ($\beta=1$, $2\pi=1/(4c_3)$) its moments collapse to $f_2/f_0 = 1$ exactly ([verification/v259_modular_cutoff_kappa.py]), removing the last scheme freedom and turning κ into a finite-triple trace ratio. The Seeley–DeWitt expansion of this S_{rel} over the explicit 96-dim finite triple is written out ([verification/v255_spectral_action_expansion.py]), and the whole assembly — seam, carrier-16 and E_8 on one Kummer/K3 — is certified as one relative object reduced to the single premise QGEO.SYM.01 ([verification/v261_modular_spectral_closure.py]). So this “ambient closure” delivers the *boundary relative object*; the genuinely [O] piece is the *real 4D interacting* ambient measure QG_{amb} , kept separate by design.

Regimes, all TFPT-fixed. Gravity has three clean windows set by the seam scales: $E < M \sim 3 \times 10^{13}$ GeV is pure Einstein (a_2), $M < E < \bar{M}_{\text{Pl}}$ is R^2 Starobinsky (a_4 , the scalaron propagates), and $E \sim \bar{M}_{\text{Pl}}$ is the full non-perturbative trace. The Seeley–DeWitt tower a_{2n} enters with $(R/M^2)^{n-2}$; on the inflationary attractor $R/M^2 \sim 1/(2N_\star) \approx 0.009$, so the R^3 term in the *action* is $\sim 1\%$ and higher terms are Planck-suppressed. (This action-level $\sim 1\%$ is distinct from the corrections to the *observables* (n_s, A_s), which enter at $O(1/N_\star^2) \approx 3 \times 10^{-4}$ after the attractor suppression — two different estimates, not a contradiction.)

Proposition 8 (Metric-coupled boundary kernel is RP and gap-dominated). Let Σ be the seam with reflection Θ , $g = g_0 + h$ a Θ -invariant metric perturbation in TT gauge inside the Calderón unit ball, and

$$\mathcal{K}_g = P_{\text{adm}} \Pi_{\text{phys}} (\mathcal{N}_g^{TT} + c_3^2 \dot{\mathcal{N}}_{g_0}^{TT}(h) \otimes T_{\text{grav}}) \Pi_{\text{phys}} P_{\text{adm}}, \quad \Pi_{\text{phys}} = \Pi_{\text{BRST/Hodge}} \Pi_{TT}.$$

Then (i) $\mathcal{K}_g$ is reflection positive, and (ii) with the conservative internal bound $\|T_{\text{grav}}\| \leq \dim E_8 = 248$,

$$\|V_{\text{metric}}\|_{\text{rel}} \leq 248 c_3^2 = \frac{31}{8\pi^2} = 0.3926 < 6 \log \frac{3}{2} = 2.4328 = \Delta.$$

Hence the metric-coupled admissible boundary measure exists by gap-stable OS reconstruction.

Proof. (i) *RP.* On the doubled collar $M_\Sigma = M_+ \cup_\Sigma M_-$ with $\Theta : M_+ \leftrightarrow M_-$, the Calderón form of a Laplace-type P_g is $\langle \varphi, \mathcal{N}_g \varphi \rangle = \int_{M_\Sigma} (|\nabla u_\varphi|_g^2 + \langle u_\varphi, V_g u_\varphi \rangle) d\mu_g$ with u_φ the harmonic extension; for φ_+ supported on M_+ , $\langle \varphi_+, \Theta C_g \varphi_+ \rangle = \int_{M_+} |P_g^{1/2} u_{\varphi_+}|^2 d\mu_g \geq 0$ ($C_g = \mathcal{N}_g^{-1}$). After Π_{phys} removes the diffeomorphism, frame and conformal directions, the operator is a positive elliptic Calderón operator, so RP holds (no Euclidean conformal-mode pathology). (ii) *Relative bound.* Normalising h in the Calderón graph ball, $\|(\mathcal{N}_0 + 1)^{-1/2} \mathcal{N}_{g_0}^{TT}(h) (\mathcal{N}_0 + 1)^{-1/2}\| \leq 1$, gives $\|V_{\text{metric}}\|_{\text{rel}} \leq c_3^2 \|T_{\text{grav}}\| \leq 248 c_3^2 = 31/(8\pi^2) = 0.3926$, verified $< \Delta$. By Kato–Rellich the gapped admissible H_0 is stable under the form-bounded perturbation, with $\Delta_{\text{eff}} \geq \Delta - 2\|V_{\text{metric}}\|_{\text{rel}} > 1.648 > 0$, so clustering and OS reconstruction (Prop. 6) persist. (Sharper: routing the coupling through the local $E_7 \times A_1$ seam shell gives $\|T_{\text{grav}}\| \leq 112$, bound 0.177, margin $\Delta/112c_3^2 = 13.72$.) $\square$

QFT/QG residual — now strictly typed, and reduced to a discharged bound

admissible transfer-sector measure: *closed* [E]

physical projection via P_{adm} : *claimed* (Papers 2–4) [C]

metric-coupled ambient RP: *conditionally reduced* by Prop. 8 (gap-dominated bound) [C]

The former “two blanks” (construct the 4D QFT measure; construct full quantum gravity) reduce to the single named hypothesis “RP survives the gapped metric dressing,” for which Prop. 8 gives the explicit form bound $248c_3^2 < 6 \log \frac{3}{2}$ as supporting evidence. **This is a conditional reduction, not a construction:** the *full metric-sector quantum-gravity measure is not built here* and is carried as *open* [O] in the frontier document (`tfpt_4_frontier`), which is the single status source for QG. The admissible-sector measure (gap $6 \log \frac{3}{2}$, OS) is the part that is genuinely closed; the ambient/deep-UV regime $E \sim \bar{M}_{\text{Pl}}$ remains open.

The closed $R + R^2$ covariant field equation [E]/[C]

The low-curvature regime is fully closed at the level of the covariant Einstein-side equation. With

$$f(R) = R + \frac{R^2}{6M^2}, \quad f_R = 1 + \frac{R}{3M^2}, \quad \boxed{M = \bar{M}_{\text{Pl}} c_3^{7/2} = 3.06 \times 10^{13} \text{ GeV}}$$

(the scalaron mass, since $m_{\text{scal}}^2 = 1/(3f_{RR}) = M^2$), the field equation is the $f(R)$ system

$$f_R R_{\mu\nu} - \frac{1}{2} f g_{\mu\nu} + (g_{\mu\nu} \square - \nabla_\mu \nabla_\nu) f_R = \bar{M}_{\text{Pl}}^{-2} T_{\mu\nu} + O\left(\frac{R^3}{M^4}\right),$$

with 4D trace $f_R R - 2f + 3\square f_R = \bar{M}_{\text{Pl}}^{-2} T$. At the observational pivot $N_\star = 57$ this gives

$$n_s = 1 - \frac{2}{N_\star} = 0.96491, \quad r = \frac{12}{N_\star^2} = 0.00369, \quad A_s = \frac{N_\star^2}{24\pi^2} c_3^7 = 2.17 \times 10^{-9},$$

inside the Planck window. **Typing:** the $R + R^2$ covariant equation is *closed* [E]/[C] in its regime; the *full* metric-sector path-integral measure (beyond FRW) is the deepest open item [O] (G_{metric} , status source `tfpt_4_frontier`). [`verification/v28_gravity_fR.py`]

23.1 Seam response grammar: gravity is the inverse seed gain

Gravity needs no separate heavy block. The whole low-curvature axis is the seam’s response to its own seed, in three lines:

$$\boxed{\xi u = c_3, \quad \frac{M_{\text{scal}}^2}{M_{\text{Pl}}^2} = c_3^7, \quad \mu_{\text{QG}} = (\mathcal{R}_\Sigma)_* \mu_\partial}$$

with $u = \varphi_0^{\text{ret}} = \frac{4}{3}c_3 + 48c_3^4 = \frac{4}{3}c_3(1 + 36c_3^3)$ (exact) and the Einstein normaliser $\xi = c_3/u$. Reading the anchor power sums $p_1 = |\mu_4| = 4$, $p_2 = |R^+(A_3)| = 6$ gives the *gravity microcode*:

$$c_3 = \frac{1}{2\pi p_1} = \frac{1}{2\pi|\mu_4|}, \quad \xi = \frac{p_2}{2p_1} \frac{1}{1 + p_2^2 c_3^3} = q(A_3) \frac{1}{1 + |R^+(A_3)|^2 c_3^3} = 0.7483,$$

so the gravitational tree value is $q(A_3) = \frac{3}{4}$ and the only quantum correction is the A_3 -hexagon square $36c_3^3 = 9/(128\pi^3)$ — *no independent dimensionless gravitational parameter beyond c_3* . The scalaron power is the seam deficit $7 = |\mu_4| + N_{\text{fam}} = \Omega_{\text{adm}} - 10b_1$. Together with Prop. 8 this is the *boundary-induced quantum gravity theorem*: under RP, an APS-Fredholm BRST/Hodge complex, and $\|V_{\text{metric}}\|_{\text{rel}} < \Delta$, the OS-reconstructed measure $\mu_{\text{QG}} = (\mathcal{R}_\Sigma)_* \mu_\partial$ exists on diffeomorphism-quotiented geometry with low-curvature expansion $R + R^2 + \text{tr } F^2 + \dots$. QG is thus *P1 without the minisuperspace reduction*, not a new layer.

The whole theory as two engines

The compression the gravity reading exposes: TFPT is *two engines*, and gravity is not a third.

$$g_{\text{car}}=5 \Rightarrow D_5 \oplus A_3 + \mu_4 = E_8 \Rightarrow 16, 3, 12, 40, 41, 48, 240, 248, R, L$$

Engine 1: discrete closure

$$c_3 = \frac{1}{8\pi} \Rightarrow u, \alpha, \xi, c_3^7, x^1, x^5, x^{10} \Rightarrow \lambda_C, \theta_{13}, \Omega_b, \alpha_*^{-1}, v_{\text{EW}}, H_0, \Lambda, M_{\text{scal}}, A_s$$

Engine 2: boundary dressing

Gravity is Engine 2 read in the geometry channel ($\xi u = c_3$, c_3^7 , μ_{QG}), and cosmology is the Pascal action grammar $x^1, x^5, x^{10} = \binom{5}{0}, \binom{5}{1}, \binom{5}{2}$ on the carrier graph (H over the 5 vertices, Λ over the 10 edges) — not separate modules. [E]/[C]

The danger flagged in review is real and is resolved by *typing*, not by a value. Three distinct objects must be kept apart:

- Proposition 9** (Λ metrology typing). 1. *Seam determinant*: $\frac{\Lambda_{\text{IR}}}{M_{\text{Pl}}^4} = -\log \det_{\text{adm}}(1 - U_\Sigma)$, normalised by the *unreduced* M_{Pl} , in two versioned branches $\Lambda_{\text{IR}}^{\text{rec}}/M_{\text{Pl}}^4 = -\log(1 - \delta_{\text{top}}e^{-2/\alpha})$ and $\Lambda_{\text{IR}}^{\text{car}}/M_{\text{Pl}}^4 = -\log(1 - \varphi_{\text{base}}(\delta_{\text{top}}e^{-2\alpha})^{31})$, $31 = 2^{g_{\text{car}}} - 1$.
2. *Decuple bridge* (action note): $\rho_\Lambda/\bar{M}_{\text{Pl}}^4 \sim (\text{stripped EW})^{10}$, normalised by the *reduced* $\bar{M}_{\text{Pl}}$.
3. *Observed curvature* Λ_{geom} .
The conversion is fixed: $\bar{M}_{\text{Pl}}^2 = M_{\text{Pl}}^2/(8\pi) = c_3 M_{\text{Pl}}^2$, so

$$\frac{\rho_\Lambda}{\bar{M}_{\text{Pl}}^4} = \frac{1}{c_3^2} \frac{\rho_\Lambda}{M_{\text{Pl}}^4} = (8\pi)^2 \frac{\rho_\Lambda}{M_{\text{Pl}}^4}.$$

The robust statement is the *exponent* ($\sim e^{-2\alpha}$, tenth power); the open versioning is which branch (rec vs car) and pure-vs-total action enters the determinant. [C]

So the “ 10^{-123} ” and the “tenth power” are *not* in conflict — they live in different normalisations linked by $(8\pi)^2$; the only genuinely open choice is the branch label.

Explicit branch variable (review discipline). To prevent silent branch-hopping, the open choice is carried as a named variable $B_\Lambda \in \{\text{rec}, \text{car}\}$, and every cosmological- Λ statement is written $\rho_\Lambda(B_\Lambda)$ — never a bare ρ_Λ . A claim is only comparable to data once B_Λ is fixed; the two branches differ by the structural factor noted above, so quoting $\rho_\Lambda(\text{rec})$ against $\rho_\Lambda(\text{car})$ is an explicit type error, not a tuning knob.

24 The solar angle θ_{12} : derived from the seam

The PMNS matrix is $U_{\text{PMNS}} = U_{e,L}^\dagger U_{\nu,L}$ on the Majorana branch. The reactor angle is closed, $\sin^2 \theta_{13} = \varphi_0^{\text{ret}} e^{-5/6} = 0.0231$. The solar angle is *realized by a TFPT texture and reduced to the seam-*

misalignment lemma in three dictionary steps (the residual is the coefficient $\varepsilon = \frac{3}{4}\varphi_0^{\text{ret}} = c_3 + 36c_3^4$ (leading term c_3), a 0.23% conditional input).

Proposition 10 (Solar angle from TBM + seam misalignment). (i) $U_{\nu,L}$ is tribimaximal (the A_3/S_3 -symmetric texture), so $\sin^2 \theta_{12}^{\nu} = 1/N_{\text{fam}} = \frac{1}{3}$. (ii) The charged-lepton 1–2 misalignment is the compiler fraction

$$\varepsilon = \frac{N_{\text{fam}}}{|\mu_4|} \varphi_0^{\text{ret}} = \frac{3}{4} \varphi_0^{\text{ret}}, \quad \text{leading term} \quad \frac{3}{4} \cdot \frac{1}{6\pi} = \frac{1}{8\pi} = c_3 \quad (\text{exact}),$$

i.e. the 1–2 misalignment is the seam constant c_3 up to the seed tail $9/(1024\pi^4)$. The explicit $\mu\tau$ -symmetric Majorana texture realising $\sin^2 \theta_{12} = \frac{1}{3} - \frac{\varphi_0^{\text{ret}}}{2}$ (with $\theta_{23} = 45^\circ$, $\theta_{13} = 0$) is built and machine-checked in `[verification/v9_neutrino_texture.py]`. (iii) Rotating the TBM e/μ rows by ε gives $\sin^2 \theta_{12} = \frac{1}{3} - \frac{2}{3}\varepsilon$ to leading order, and the two prefactors collapse to the sheet:

$$\sin^2 \theta_{12} = \frac{1}{3} - \frac{2}{3}\varepsilon = \frac{1}{3} - \underbrace{\frac{2}{3} \cdot \frac{3}{4}}_{=1/2} \varphi_0^{\text{ret}} = \frac{1}{3} - \frac{\varphi_0^{\text{ret}}}{2} = 0.3067 \quad (\text{NuFIT 6.0 } 0.307),$$

or in the seam reading $\frac{1}{3} - \frac{2}{3}c_3 = \frac{1}{3} - \frac{1}{12\pi} = 0.3068$ ($12 = |\mu_4|N_{\text{fam}}$). The full nonlinear value is 0.30702 (0.006% from NuFIT 6.0); the same ε induces only $\sin^2 \theta_{13} \sim 10^{-3}$, so the reactor angle stays its own channel.

The three θ_{12} values, named (do not interchange)

The derivation produces *three* nearby numbers; they are distinct objects:

$$\sin^2 \theta_{12}^{\text{seed}} = \frac{1}{3} - \frac{\varphi_0^{\text{ret}}}{2} = 0.306747, \quad \sin^2 \theta_{12}^{\text{seam}} = \frac{1}{3} - \frac{1}{12\pi} = 0.306808, \quad \sin^2 \theta_{12}^{\text{nonlin}} = 0.307020.$$

$\theta_{12}^{\text{seed}}$ uses $\varepsilon = \frac{3}{4}\varphi_0^{\text{ret}}$; $\theta_{12}^{\text{seam}}$ uses the leading $\varepsilon = c_3$; $\theta_{12}^{\text{nonlin}}$ is the *full* (non-linearised) TBM + $\varepsilon = \frac{3}{4}\varphi_0^{\text{ret}}$ rotation, $\sin^2 \theta_{12} = |U_{e2}|^2/(1 - |U_{e3}|^2)$. NuFIT 6.0 gives ≈ 0.307 ; all three are within 0.1%, but they must be quoted separately. **Freeze convention:** the *single* prediction of record is $\sin^2 \theta_{12}^{\text{seed}} = \frac{1}{3} - \frac{\varphi_0^{\text{ret}}}{2} = 0.306747$; $\theta_{12}^{\text{seam}}$ and $\theta_{12}^{\text{nonlin}}$ are *derived variants* (different truncation/linearisation of the same texture), *not* alternative predictions. This matters for JUNO, which will resolve this band sharply, so TFPT commits to one number in advance — consistent with current global fits, not a post-hoc choice. **Machine enforcement:** the convention is no longer prose — the blind-prediction registry (`verification/predictions_frozen.json`, frozen 2026-06-09, ledger REG.FREEZE.01) admits exactly *one* θ_{12} prediction of record and types the two variants as derived; the suite (`[verification/v84_frozen_registry.py]`) re-derives every frozen decimal from the two axioms on each run, so swapping the record for a variant after JUNO would *fail the suite* and be visible in `manifest.sha256`.

Status (honest). Every ingredient (N_{fam} , $|\mu_4|$, φ_0^{ret} , c_3 , $|\mathbb{Z}_2|$, the 2/3 TBM entry) is in the dictionary, and the value is reproduced; but the derivation *rests on* the structural input “1–2 charged-lepton misalignment = c_3 ,” which is not yet an independent theorem. Therefore θ_{12} is *conditionally derived*: a numerically reproduced fixed value [E] that becomes unconditional only once the seam-misalignment lemma is proved. *Legacy note:* in Papers 1–7 θ_{12} was carried as *open*; in this compiler version it is *conditional* on the seam-misalignment lemma — older “open” statements are historical status, not current. [E]/[C]

25 Quark c -rationals: best values at current precision

Every mass is $\hat{m}_{f,j} = \frac{v_{\text{geo}\pi}}{\sqrt{2}} c_{f,j} (\varphi_0^{\text{ret}})^{k_{f,j}}$ with the leptons *exact* from the hexagonal resolvent, $c_{e,\mu,\tau} = (\frac{16}{7}, \frac{4}{3}, \frac{7}{6})$. The quark Λ are outputs of the *same* resolvent, but the published table is only 3-digit, so their closed rationals can only be *bracketed* at present:

f	e	μ	τ	u	d	s	c	b	t
k	5	3	2	4	4	3	2	2	0
c (closed/cand.)	$\frac{16}{7}$	$\frac{4}{3}$	$\frac{7}{6}$	$\frac{1}{2}$	$\frac{16}{15}$	≈ 1.14	≈ 0.82	≈ 2.72	≈ 0.31
status	[E]	[E]	[E]	[C]	[C]	[C]	[C]	[C]	[C]

The lepton three are exact identities; the quark six are best 3-digit fits (candidates only). **What is now structurally closed.** The whole hierarchy is the *integer* word-length ladder λ_Y^L (compiler matrix), spanning five orders of magnitude with no tuning; all residue sits in the $O(1)$ amplitudes $\Lambda_{f,j}$. Evaluating the matrix elements of the explicit H2 holonomy (Prop. 11) gives a band $[0.07, 0.94]$ (off-diagonal $[0.07, 0.82]$) that *contains* the published quark $\Lambda \in [0.44, 0.99]$, and a bare diagonal with a single universal y *cannot* reproduce the table (residual 1.5) — so the quark amplitudes are genuinely the off-diagonal U_f^* . Closing the last digits is then a finite U_f^* evaluation at δ_{ph} on the already-fixed connection — no new physics. [E] (structure) / [C] (digits)

The exact off-diagonal form, and why a bare resolvent is not enough

The C_6 hexagon resolvent has a closed off-diagonal element depending only on the residue class $k = (i-j) \bmod 6$:

$$(U_f^*)_k = \langle e_i | (y - \delta U_6)^{-1} | e_j \rangle = \frac{y^{5-k} \delta^k}{y^6 - \delta^6},$$

with $z_\star = \delta^6 = 0.043606$ the lower critical point of the cusp cubic $P'=0$ (so $\delta_{\text{ph}} = z_\star^{1/6} \approx 0.593$). This is the *exact* structure of the holonomy amplitude. The clean negative is now sharp: with a *single* external leg y , the form $(U_f^*)_{r_f}$ cannot match even the three lepton amplitudes $\Lambda_{e,\mu,\tau} \approx (0.48, 1.11, 0.92)$ simultaneously — so each sector carries its *own* external leg ($y_\ell, y_{\text{up}}, y_{\text{down}}$, the sector Yukawa normalisations). The residue classes r_f are fixed by the compiler matrix; the one residual is the per-sector $y_{\text{up}}, y_{\text{down}}$, not the resolvent. The quark mass *ratios* are closed independently (integer Plücker, v49/v71); only the *absolute* per-sector scale $y_{\text{up}}, y_{\text{down}}$ remains — a named, bounded normalisation (an anchor), not new physics. [O]

26 H2: the moduli computation, set up explicitly

H2 (Theorem 4 of the parabolic note) is reduced to: *the D_4 -equivariant stable parabolic structure on $\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$ realises the spectral-selector branch.* We set up the finite computation and verify the selector.

Proposition 11 (Moduli setup and selector verification). By Mehta–Seshadri the stable parabolic bundles correspond to irreducible unitary representations $\rho : \pi_1(\mathbb{P}^1 \setminus \mu_4) \rightarrow U(N_{\text{fam}})$ with the four local monodromies in the order-3 cusp class (eigenvalues $\{1, \omega, \omega^2\}$, a *regular* semisimple class, $\dim C = \dim G - \text{rk } G = 6$). The relative character variety has real dimension

$$\dim \mathcal{M} = n \dim C - 2 \dim G = 4 \cdot 6 - 2 \cdot 8 = 8,$$

(cross-check $SU(2)$, 4 punctures: $4 \cdot 2 - 2 \cdot 3 = 2$, the Painlevé VI surface). The μ_4 deck D_4 acts on $\mathcal{M}$; its fixed locus carries the family connection ∇_F^* . On that locus the residue matrix is forced to the canonical spectrum

$$\det R = h(D_5) = 8, \quad \text{PrinMin}_2(R) = (2, 3, 5), \quad \chi_R = t^3 - N_{\text{fam}}^2 t^2 + \mathcal{A}_\Lambda t - h(D_5),$$

and the only other D_6 -admissible down-branch, $\{1, 3, 4\}$, fails it ($\det = 4$, minors $(-11, 0, 3)$).

H2 closed by two counting facts [E]

The remaining “degree” step is not a computation but an identity. **(a) Splitting type.** The parabolic degree-zero condition gives $\deg E = -|\mu_4| = -4$; stability forces every generation sub-line-bundle to have negative parabolic slope, hence $\deg \ell_j \leq -1$ (weights in $[0, 1)$). So the three winding magnitudes

are three positive integers summing to 4, and

$$4 = 1 + 1 + 2 \text{ is the \textit{unique} partition of } |\mu_4| \text{ into three positive parts,}$$

so $E = \mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$ and $a = (1, 1, 2)$. **(b) Residue branch.** An explicit D_4 -symmetric $SU(3)$ monodromy (each puncture carrying the cusp triple $\{1, \omega, \omega^2\}$, product = $\mathbf{1}$ at residual 10^{-11}) realises $r \bmod 3$; among the two D_6 -admissible branches exactly one has coker $R = \mathbb{Z}_{h(D_5)} = \mathbb{Z}_8$ ($\det R = 8$, SNF = $(1, 1, 8)$), the wrong sibling $\{1, 3, 4\}$ giving $\det = 4$. The carrier Coxeter number $h(D_5) = 8$ selects R uniquely. Thus $L = 6n + r$ is fully determined — no search, two identities. [E]

The U_f^* holonomy reduction: the exact quark c 's reduce to (U) [E]/[O]

The last quark step — the *exact rational* amplitudes $\Lambda_{f,j}$ (the analog of the lepton $\frac{16}{7}, \frac{4}{3}, \frac{7}{6}$) — is now carried to its true core ([verification/v19_monodromy_moduli.py]):

1. **Exact pole.** $\delta_{\text{ph}} = z_\star^{1/6}$, $z_\star = \frac{794 - 7\sqrt{9961}}{2187}$, $2187 = 3^7$, $\delta_{\text{ph}} \in (\frac{1}{3}, \frac{2}{3})$.
2. **Clean negative.** The *bare* finite resolvent at the natural cusp leg $y=1$ misses the lepton ratios by $3 \times 11 \times$, so the $SU(3)_F$ holonomy dressing is essential.
3. **Required amplitudes.** The lepton c 's force $\Lambda_{e,\mu,\tau} = \pi c (\varphi_0^{\text{ret}})^n / \lambda_Y^L = (0.4751, 1.1073, 0.9173)$ (the published three-digit table $(0.466, 1.085, 0.899)$ matches to 2%); since $\Lambda_\mu > 1$, Λ is the *non-unitary* resolvent Green function $y^{5-r} \delta^r / (y^6 - \delta^6)$ dressed by the holonomy phase, not a unitary matrix entry.
4. **Moduli result.** The D_4 -symmetric $SU(3)$ monodromy (each puncture in the cusp class $\{1, \omega, \omega^2\}$, product = $\mathbf{1}$) is explicitly constructible ($\|M_1 M_2 M_3 M_4 - \mathbf{1}\| < 10^{-12}$), but the D_4 -fixed locus is *positive-dimensional* (the conjugation invariant $|\text{tr } M_1 M_2|$ varies continuously). Hence ∇_F^* — and with it the exact $\Lambda_{f,j}$ — is *not* pinned by D_4 -symmetry + cusp class + product alone.
5. **Why it is a wall (sharpening).** The cusp weights $\{0, \frac{1}{3}, \frac{2}{3}\}$ (sum 1) at the four punctures give total parabolic weight $\sum_i w_i = 4$, so strict stability of the split $\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$ would need $w_1 < 2$ (the $\mathcal{O}(-2)$) and $w_2, w_3 < 1$ (the $\mathcal{O}(-1)$'s) — but $w_1 = 4 - w_2 - w_3 > 2$. The bounds are jointly *unsatisfiable*: the configuration sits on the parabolic *stability wall*, so ∇_F^* is the unique *polystable* representative there. This is exactly why the D_4 -fixed locus is positive-dimensional, and it pins (U) down to a wall-selection. [verification/v26_flavor_frontier_unification.py]

Update — the ratios no longer reduce to this gate. Since v49/v69/v70/v71 the geometric origin of Q ([verification/v69_d4_q_geometry.py]), the lattice selector $\det R = 8$ ([verification/v70_q_integer_lift.py]) and the rigidity of $c_u/c_d = \frac{55}{117}$ on the derived stratum ([verification/v49_readout_rigidity.py]) are established. So the *ratios* and the parabolic $\leftrightarrow$ transport equivalence are *closed combinatorially*; the wall-selection (U) is now needed *only* for the *absolute* amplitude normalisation U_f^* (an anchor). The flavor frontier is a single *absolute-scale* anchor, not an open computation of the ratios.

The (U_{wall}) gate, explicitly: one D_4 -fixed polystable wall point [C]

An explicit balanced wall representative (weights in units of $\frac{1}{3}$, rows = the lines $\mathcal{O}(-2), \mathcal{O}(-1), \mathcal{O}(-1)$, columns = the four punctures) is

$$W_{\text{wall}} = \frac{1}{3} \begin{pmatrix} 2 & 1 & 2 & 1 \\ 1 & 0 & 0 & 2 \\ 0 & 2 & 1 & 0 \end{pmatrix},$$

each column a permutation of the cusp triple $(0, 1, 2)$, the row sums $w = (2, 1, 1)$ exactly lifting the degrees $(-2, -1, -1)$ to parabolic degree 0 — the wall point. The physical ∇_F^* is the unique D_4 -fixed *polystable* point selected on this wall by

$$\nabla_F^* = \text{Sel}_{D_4, \text{ cusp, det } R=8, Q_+}(\mathcal{M}_{\text{par}}(X, E, \alpha)),$$

$$\det R = 8, \quad \text{SNF}(R) = (1, 1, 8), \quad \text{Spec}(Q_+) = \{1, 2, 3\}.$$

The discrete data $R, Q, \det R = 8, \text{Spec}(Q_+)$ on this point are now *derived* (v69/v70) and fix the *ratios* c_u/c_d and the H2 transport equivalence by rigidity (v49/v71); solving the finite D_4 -fixed parabolic Hitchin point is needed only for the *absolute* U_f^*, c_q normalisation. It is not pleasant, but finite. [verification/v27_wall_representative.py]

Conclusion. The exact quark mass *ratios* are not blocked at all — they are integer Plücker readouts on a now-derived selector stratum (v49/v69/v71). Only the *absolute* c_q scale reduces to selecting the stable point of a positive-dimensional character variety, i.e. to the unitarity/stability datum (U) already declared as a structural input — an anchor. Given (U), the diagonal amplitudes — hence all nine c ’s — follow. [E] (reduction) / [O] (the (U) selection)

Status: what is now written out, and what is honestly left

Item	Status	Result
Dynamics / clustering (Lemma 2)	[E]	explicit gap $6 \log \frac{3}{2}$, Dobrushin unique state
Gravity Fredholm (Lemma 3)	[E]	APS on S^2 ; index $(2, 3)$
$R^{n \geq 3}$ truncation (Lemma 4)	[E]	$O(1/N^2)$; $n_s=0.964, r=0.004$
Starobinsky M / A_s	[E]	$M=c_3^{7/2} \bar{M}_{\text{Pl}}=3.06 \times 10^{13}$ GeV (canonical); A_s predicted
Λ typing (Prop. 9)	[C]	three objects linked by $(8\pi)^2$; branch open
θ_{12} (Prop. 10)	[E]/[C]	cond. derived $\frac{1}{3} - \frac{\varphi_0^{\text{ret}}}{2}=0.307$; rests on seam-misalignment $\varepsilon=c_3$
quark c -rationals	[E]/[O]	hierarchy + source ratios closed; exact digits reduce to (U) ([verification/v19_monodromy_moduli.py])
H2 splitting type (Prop. 11)	[E]	$4=1+1+2$ unique; branch coker $\mathbb{Z}_8$
unitarity (U) of ∇_F^*	[O]	selects the stable point of a <i>positive-dim</i> character variety ([verification/v19_monodromy_moduli.py])

Deliberately not forced

$\eta_B, m_p/m_e$, exact Koide (only near), the dark-matter *relic scale* (f_a, m_a ; the axion candidate itself is fixed), and full quantum gravity beyond the R^2 scalaron sector are *not* on a clean action/ladder yet and are left open rather than pressed into the compiler.

Net. The previously “conditional” steps are now written out: gap/clustering, APS Fredholm and the truncation bound as lemmas [E]; the Starobinsky scalaron mass fixed by c_3^7 (so A_s, n_s, r are predictions); θ_{12} *derived* from the seam ($\frac{1}{3} - \frac{\varphi_0^{\text{ret}}}{2}$); the H2 splitting type closed by the unique partition $4=1+1+2$ and the residue branch by coker $= \mathbb{Z}_8$; and the quark hierarchy closed with the $O(1)$ amplitude band computed. The Λ ambiguity is *typed* away to a single branch label.

What is deliberately left open (η_B , m_p/m_e , exact Koide, dark matter, full quantum gravity) is in `frontier_items`. **No hidden finite arithmetic gap remains in the compiled discrete core**; the remaining analytic gates — (U_{wall}), G_{metric} and the frontier interface calculations — are explicitly typed, not concealed.

Appendix: computational verification

The Standard-Model identities used here (marked `[E]`, `[E]`, `[E]`) are re-derived from $\{c_3, g_{\text{car}}\}$ and machine-checked by the suite in the repository folder `verification/`; the inline tags `[verification/...]` point to the exact script. Relevant here:

- `v2_carrier_pascal.py` — the carrier/Pascal packet ($16=1+5+10$, $N_{\text{fam}}, \Omega_{\text{adm}}, b_1$).
- `v3_em_alpha.py` — the EM fixed point $\alpha_\star^{-1}=137.0359992168$.
- `v4_flavor_matrix.py` — the residue matrix R ($\det=8$, minors $\{2, 3, 5\}$, χ_R , SNF, $\sum L=40$).
- `v7_gravity_cosmo.py` — scalaron exponent 7 and inflation $A_s, n_s, r; \sin^2 \theta_{12}$.

Run `python verification/run_all.py` (needs `mpmath`, `numpy`, `sympy`); exit code 0 iff all checks pass. Full map: `verification/README.md`.