

TFPT — Research Contracts for the Remaining Interfaces:

$$v_{\text{geo}}, G_{\text{net}}, F_{\text{transfer}}$$

(U_{wall}): the parabolic flavor wall-selection and (G_{metric}): the full quantum-gravity measure

Stefan Hamann

Alessandro Rizzo

June 18, 2026 · v5.2 (rev 215)

The TFPT document set — what is treated where

Plain language: TFPT (Topological Fixed-Point Theory) is a small discrete compiler. Two inputs — the seam constant $c_3 = \frac{1}{8\pi}$ and the carrier rank $g_{\text{car}} = 5$ — build $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ and read off the Standard-Model skeleton, the constants and the scale grammar. The series is **the introduction plus five numbered papers and three companions**, best read in order:

0. **introduction** — reading guide, compiler closure, predictions, the dependency DAG and the single proof ledger (`verification/status_ledger.csv`). (*entry point*)
1. **tfpt_1_architecture_e8** — the two axioms, the derivation map, the EM fixed point α^{-1} , the $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ construction, the QBL theorem chain.
2. **tfpt_2_standard_model** — the SM in one φ_0 -ladder formula, flavor from parabolic transport (sheet diamond, dual anchor, Q_+ cohomology), the worked closures.
3. **tfpt_3_e8_audit_bootstrap** — the seven E_8 slices as an audit raster, the cascade bridge, the Möbius bootstrap.
4. **tfpt_4_frontier** — honest status of η_B , m_p/m_e , Koide, dark matter, full QG.
5. **tfpt_5_redteam** — the adversarial audit: declared attacks, what survives, what each reduces to, and the kill tests.

Companions: **R.** **tfpt_research_contracts** — the open interfaces ($v_{\text{geo}}, G_{\text{net}}, F_{\text{transfer}}$) as numbered contracts; **H.** **tfpt_horizon_readouts** — Appendix H (unit-system reframe): one seam constant as the universal horizon thermal code, the Nariai/anchor surface, the resummed seam clock; **O.** **origin_theory** — why no free fundamental number remains (exact core + one honestly-typed cyclic interpretation).

You are reading companion R (Research Contracts).

TFPT status at a glance — read this card the same way in every document

Two inputs, one compiler. From the boundary constant $c_3 = \frac{1}{8\pi}$ (axiom P1) and the five-slot carrier $g_{\text{car}} = 5$ (axiom P2) — jointly the elementary symmetric data of one anchor $a = (1, 1, 2)$ — the discrete compiler $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ is *built*, and the Standard-Model skeleton (gauge group, three families, hypercharges, the flavor matrix), the dimensionless constants ($\alpha^{-1} = 137.0359992$) and the scale grammar are read off as fixed points. The algebraic core is machine-checked; physical readouts run through *named bridges* whose status is typed below.

What is closed, and what genuinely remains. The discrete/algebraic layer is closed ($[E]$); the everything-open residual reduces to *three named interface problems*:

Interface	Question	Status
v_{geo}	the one metrology unit ($= 1/\sqrt{G} = m/\mu$); No-Unit Thm: no compiler scale	primitive $[O]$
G_{net}	simple-current extension $(D_5)_1 \otimes (A_3)_1 \rtimes \langle (1, 1) \rangle \cong (E_8)_1$	$[E]/[C]$
F_{transfer}	one functor, four typed interfaces (Koide, η_B , axion, m_p/m_e)	$[C]$


Everything carries a claim ID resolving to a row of `verification/status_ledger.csv` (the single source of truth); **if the text and the ledger ever disagree, the ledger wins.** The historical labels ($U_{\text{wall}}, G_{\text{metric}}, F_{\text{frontier}}$) are retained only for ledger continuity — the live residual is $v_{\text{geo}} \oplus G_{\text{net}} \oplus F_{\text{transfer}}$.
 Marker key: $[E]$ exact / machine-proven · $[C]$ conditional (holds under named hypotheses) · $[O]$ open / axiom · $[X]$ falsifiable kill test. Per-claim fine type in `verification/status_ledger.csv`.

$[E]$ exact / machine-proven

$[C]$ conditional

$[O]$ open / axiom

$[X]$ kill test



What this document is about

After the compiler closure the entire residual is

$$\text{Rest} = (U_{\text{wall}}) \oplus (G_{\text{metric}}) \oplus (F_{\text{frontier}})$$

This note turns the two genuine research gates into *contracts*: a numbered chain of lemmas, the single theorem that closes each gate, and — for every step — whether it is machine-certifiable today. (F_{frontier}) (Koide, η_B , axion relic scale, m_p/m_e) is *not* a gate: those are deliberately typed QFT/cosmology interfaces (see `tfpt_4_frontier`). **Priority:** (U_{wall}) *first* (finite, algebraic, falsifiable), then (G_{metric}) (deep analytic programme).

Operative aliases (current state of the reductions; the historical gate names above are kept for ledger continuity). The lemma chains below have since *discharged* large parts of both gates, so the operative residual reads sharper:

$$\text{Rest}_{\text{operativ}} = v_{\text{geo}} \oplus G_{\text{net}} \oplus F_{\text{transfer}}$$

— $(U_{\text{wall}}) \rightarrow v_{\text{geo}}$: the quark *ratios* are closed combinatorially (Readout Rigidity + Plücker, U2/v49/v71), U_{point} collapses to the *one* dimensionful unit v_{geo} shared with $1/G$ (v75/v68) — what remains is a scale anchor, not a flavor gap (the *full* amplitude matrix beyond the ratios is the only place where U_{point} still appears); $(G_{\text{metric}}) \rightarrow G_{\text{net}}$: holographically reduced to the single boundary-net statement “*the seam-Calderón inclusion has Jones index 4 = $|\mu_4|$* ” — holomorphy, $(E_8)_1$ and the unique 2D bulk then follow (v89/v92/GATE.METRIC.05-07); and the carrier-side QBL input now merges into the *same* statement: the tower carrier $\rightarrow SO(16)_1 \rightarrow (E_8)_1$ carries one and the same 16-fermion content, whose single quasi-free kernel (rank = c ; Wick/Pfaffian exact) is the certified Calderón datum — closing G_{net} therefore also closes the carrier choice [E] ([verification/v113_quasifree_kernel.py]); $(F_{\text{frontier}}) \rightarrow F_{\text{transfer}}$: named QFT/cosmology *transfer* interfaces, never compiler powers — and *not* four unrelated interfaces but *one* discrete geodesic on the Sheet Diamond (the $K \rightarrow C \rightarrow F$ sheet axis $M(1, t)$, with the Koide source $\rightarrow$ pole as its 1-D Möbius projection), along which the external continuous solver merely evaluates ([verification/v224_diamond_ftransfer_path.py]).

QFT layer (this round). The boundary QFT is now *one relative object* $\text{TFPT}_{\text{QFT}} = (\mathcal{A}_{\Sigma}, \omega_{\Sigma}, \Delta_{\Sigma}, \rho, A_F, H_F, D_F, J, \gamma, S_{\text{rel}})$, closed modulo the *single* premise **QGeo.SYM.01**: its finite Dirac is a covariance induction of the seam KMS state (v258), its spectral-action cutoff *is* that KMS weight (v259), its seam, carrier-16 and E_8 live on one Kummer/K3 surface (v260), and the whole assembly is certified as the *Modular Spectral Closure* (v261). The ambient quantum-gravity measure (G_{metric}) is gap-decoupled and kept separate by design — the QFT layer does not wait on it.

Research Contract 1 — (U_{wall})

Goal. Prove

$$\nabla_F^* = \text{Sel}_{D_4, \text{ cusp}, \det R=8, \text{ Spec}(Q_+)=\{1,2,3\}}(\mathcal{M}_{\text{par}}(\mathbb{P}^1, \mu_4, E, \alpha)),$$

with $E = \mathcal{O}(-2) \oplus \mathcal{O}(-1) \oplus \mathcal{O}(-1)$ and $\alpha = \{0, \frac{1}{3}, \frac{2}{3}\}$ at the four points of μ_4 . The substrate is standard: degree-zero stable parabolic bundles correspond to unitary representations with prescribed local monodromies (Mehta-Seshadri); tame non-abelian Hodge gives the Higgs / flat-connection side. The TFPT content is to pin the *one* D_4 -symmetric, rank-3, four-point point that realises the selectors.

Lemma 1 (U0 — normalisation). *The gate data are equivalent to the $SU(3)$ character variety $\mathcal{X}_{\text{cusp}} = \{(M_1, \dots, M_4) \in C_{\text{cusp}}^4 : M_1 M_2 M_3 M_4 = \mathbf{1}\} / SU(3)$ with $C_{\text{cusp}} = \text{Ad}_{SU(3)} \text{diag}(1, \omega, \omega^2)$, $\omega = e^{2\pi i/3}$, and $\pi_1(\mathbb{P}^1 \setminus \mu_4) = \langle \gamma_1, \dots, \gamma_4 \mid \prod \gamma_i = 1 \rangle$. To show: $\mathcal{M}_{\text{par}}^{D_4} \simeq \mathcal{X}_{\text{cusp}}^{D_4}$ as polystable Mehta-Seshadri / tame Hodge realisation. Machine: the group presentation and D_4 -action are finite and codable (Sage/Wolfram).*

Finite rigidity — now proven [E]. The finite half of U0 is closed exactly: μ_4 has cross-ratio 2 and Möbius stabiliser $\langle z \mapsto iz, z \mapsto 1/z \rangle$ of order $8 = |D_4|$ (a 4-cycle of the marks + a reflection, faithful); $H^1(\mathbb{P}^1 \setminus \mu_4)$ has rank $4 - 1 = N_{\text{fam}} = 3$; and the forms $\omega_k = z^{k-1} dz / (z^4 - 1)$, $k = 1, 2, 3$, are μ_4 -eigenforms ($z \mapsto iz : \omega_k \mapsto i^k \omega_k$) carrying the three nontrivial characters χ_1, χ_2, χ_3 with weights $(1, 2, 3) = \text{Spec}(Q_+)$. Hence a genus-zero boundary with faithful D_4 , four parabolic marks and this μ_4 -decomposition is Möbius-equivalent to $\mathbb{P}^1 \setminus \mu_4$ with the $(1, 2, 3)$ grading (QGeo.RIGID.01, [E], [verification/v168_qgeo_rigidity.py]). The single residual is the seam-side *realisation* — that these hypotheses follow from the seam collar itself — which stays **QGeo.REALIZE.01** ([C]).

Seam Collar Realisation Theorem — the one central proof obligation [E]/[O]
([verification/v176_seam_collar_realisation.py])

Stated as one theorem and certified as a reduction — not a fabricated proof.

Theorem (target). *Given* (1) an oriented one-sided RP seam collar Σ ; (2) a sheet-odd Calderón involution; (3) quasi-free one-particle seam data with gap $(2/3)^6$; (4) a faithful D_4 action on the boundary marks; (5) four parabolic marked points with μ_4 -character decomposition; (6) an admissible $c=8$ boundary net — *then* the conformal boundary of Σ is Möbius-equivalent to $\mathbb{P}^1 \setminus \mu_4$, its H^1 character basis is canonically the generation basis with grading $(1, 2, 3)$, the seam Calderón contraction is the rank-8 K_{Σ} polarisation, and the index-4 simple-current extension is $(E_8)_1$.

The proof decomposes into six steps, five already [E]:

- 1–2 *Geometry / uniformisation* [E] (v168): μ_4 has cross-ratio 2 and a faithful Möbius D_4 stabiliser of order 8; four genus-0 marks with faithful D_4 force the μ_4 square up to Möbius; $H^1(\mathbb{P}^1 \setminus \mu_4)$ has rank $4 - 1 = N_{\text{fam}} = 3$.
- 3 *Calderón data* [E] (v156): the Dirichlet-to-Neumann operator of the 2d Laplacian on the half-space has symbol $|k|$ (harmonic extension $e^{ikx - |k|y}$), the free chiral dispersion — bulk-detail-independent.
- 4 *H^1 character = generation grading* [E] (v137/v168): $\omega_k = z^{k-1}dz/(z^4 - 1)$ carry μ_4 characters i^k with weights $(1, 2, 3) =$ the A_3 exponents $= \text{Spec}(Q_+)$ — natural, not merely isomorphic.
- 5 *One-particle contraction* [E] (v162): a μ_4 -equivariant operator is diagonal in the cusp basis $(\frac{1}{|\mu_4|} \sum_j U^j X U^{-j} = \text{diag } X \text{ for } U = \text{diag}(1, i, -1, -i))$; its spectrum is the cusp weights $\{0, \frac{1}{3}, \frac{2}{3}\}$, so the transfer eigenvalues $(1 - \alpha)^{p_2}$ ($p_2=6$) are $\{1, (2/3)^6, (1/3)^6\}$, gap $(2/3)^6$ — forced, no free knob.
- 6 *AQFT closure* [E] (v154/v175): $(D_5)_1 \otimes (A_3)_1$ with the isotropic μ_4 glue has index $4 = |\mu_4|$, $c = 5+3 = 8$, μ -index $16/16 = 1 \Rightarrow$ holomorphic $\Rightarrow (E_8)_1$ ($248 = 120+128$); full-cone RP holds for all m via the CAR second-quantisation functor.

The proof tree — and the two genuine open doors [O] ([verification/v177_seam_marking_kernel.py]). Taking the marks/ D_4 as a *hypothesis* (as the six-step form above does) is a near-circular trap; the honest decomposition factors the theorem as

$$\text{REALIZE} = \text{MARKS} \cdot \text{UNIFORM} \cdot \text{COHOM} \cdot \text{MODULE} \cdot \text{KERNEL} \cdot \text{NET},$$

of which *four nodes are now closed constructively*: UNIFORM [E] (an order-4 Möbius map conjugates to $z \mapsto iz$, a non-fixed orbit scales to μ_4 , and $\sigma\rho\sigma = \rho^{-1}$ for $\sigma: z \mapsto 1/z$ gives a faithful D_4 — so a genus-0 four-marked faithful- D_4 boundary is $(\mathbb{P}^1, \mu_4)$, strengthening v168's cross-ratio to the full uniformisation — and now **Lean-formalised** as FORM.QGEO.02: $\rho^4 = \text{id}$, $\rho^2 \neq \text{id}$, $\sigma^2 = \text{id}$, $\sigma\rho\sigma = \rho^{-1}$, the orbit scaling to μ_4 , the μ_4 permutation, and the multiplier classification “order-4 $\Leftrightarrow \zeta = \pm i$ ” are machine-proved in TftptCarrier/MobiusUniformisation.lean, AUDIT: PASS); COHOM [E] ($\rho^* \omega_k = i^k \omega_k$, grading $(1, 2, 3)$ — also **Lean-formalised** as FORM.QGEO.03: the pullback characters i^1, i^2, i^3 and the reflection parity $\sigma^* \omega_1 = \omega_3$, $\sigma^* \omega_2 = \omega_2$, $\sigma^* \omega_3 = \omega_1$ are machine-proved in TftptCarrier/CohomologyGrading.lean, AUDIT: PASS); MODULE [E] (a *unique* μ_4 -equivariant iso — multiplicity-free + residue normalisation, reflection $\omega_1 \leftrightarrow \omega_3$, ω_2 fixed; the parity is the Lean FORM.QGEO.03 above, the *uniqueness* stays symbolic); and NET [E] (v154/v175). The residual is therefore *exactly two* named obligations, neither a finite computation: QGEO.MARKS.01 — the *raw* RP seam collar canonically produces the genus-0 four-parabolic-marked D_4 boundary (the four marks from the anchor atom $e_1(a)=4=|\mu_4|$, collisions excluded, $\tau\rho\tau = \rho^{-1}$) — and QGEO.KERNEL.01 — the *raw* seam Calderón operator equals the μ_4 -equivariant free $c=8$ gapped one-particle contraction *as an operator* ($C_\Sigma = U^{-1}C_{\mu_4}U$), not merely the same spectral list. **The whole structural residual of the theory is these two doors**; everything between them is [E]. They need a human constructive-geometry/AQFT argument (or a proof assistant), not more arithmetic.

How far each door opens by computation ([verification/v178_marks_kernel_attempt.py]). Neither door is closed, but each splits into a finite [E] core + one *sharper* premise. For MARKS: given a genus-0 double with an order-4 clock ρ (fixed points $\{0, \infty\}$) and reflection τ , the marks are forced to be a generic ρ -orbit $\{1, i, -1, -i\} = \mu_4$ (not the fixed points), distinct iff $b_1 = 3 = N_{\text{fam}}$ (any collision drops b_1), with $\tau\rho\tau = \rho^{-1}$ giving faithful D_4 — so MARKS reduces to the *single* topological premise “the seam double is genus-0 with an order-4 clock whose marks form one orbit”. For KERNEL: on the 3-dim multiplicity-free H^1 the μ_4 characters $(1, 2, 3)$ are distinct, so by Schur a μ_4 -equivariant operator is forced *diagonal* and is *completely determined by its eigenvalue per character* — two such with the forced transfer spectrum $\{1, (2/3)^6, (1/3)^6\}$ and the unique residue-normalised basis are *equal as operators*, not merely isospectral; the “spectrum vs operator” distinction *vanishes* on the finite block. So KERNEL reduces to the *single* premise “the full DtN on L^2 reduces to this 3-dim gapped transfer block + the fixed rank-8 polarisation” (principal symbol $|k|$, Lee–Uhlmann v156; no drift v158). The two doors are thereby *milder*, but still open.

Marks without a geometric posit: a Lefschetz/character argument ([verification/v195_marks_lefschetz_character.py]). The marks need not be *assumed* as D_4 boundary points: they follow from the H^1 character alone. The order-4 clock $\rho: z \mapsto iz$ fixes exactly $\{0, \infty\}$, so a puncture at a fixed point would contribute a *trivial* character to H^1 ; but the carrier datum is $H^1 = \chi_1 \oplus \chi_2 \oplus \chi_3$ (the A_3 exponents $(1, 2, 3)$, with $\text{Tr}(\rho|H^1) = i + i^2 + i^3 = -1$), which has *no* trivial component — so no puncture sits at a fixed point. With $\text{rank } H^1 = n - 1 = 3 \Rightarrow n = 4$

punctures, all off the fixed locus and each in a free order-4 orbit, the only possibility is a *single* free μ_4 orbit (cross-ratio 2), and the closed-surface Lefschetz number $L(\rho) = 1 + 1 = 2 = |\{0, \infty\}|$ is consistent. So MARKS reduces from the geometric posit “ D_4 marks” to the milder *representation* premise “the raw seam H^1 carries the A_3 -exponent rep” — a carrier-algebra statement, strictly less circular, still [O].

And the two doors are in fact ONE ([verification/v179_conformal_realisation.py]). MARKS’s residual is not even three independent assumptions: (1) genus-0 is P1 — the one-sided Gauss–Bonnet $c_3 = 1/(\mathbb{Z}_2 |2\pi \chi(S^2)|) = 1/(8\pi)$ uses $\chi(S^2) = 2$, and $\chi = 2 - 2g \Rightarrow g = 0$, so “genus-0” is the *same* $\chi = 2$ that supplies the “8” in c_3 (v58/v73); (2) the order-4 clock *is* the carrier — it is the Coxeter element of $W(A_3) = S_4$ (v117), order $h(A_3) = 4 = |\mu_4|$; (3) marks = one orbit is *forced* — the parabolic marks are not the two elliptic fixed points $\{0, \infty\}$, and a free order-4 orbit has exactly $4 = e_1(a)$ points. So MARKS reduces to “the carrier clock acts *conformally* on the genus-0 double”, and KERNEL then *follows* from that same geometry (Lee–Uhlmann $|k|$ + the $c = 8$ rigidity v158 + the cusped-surface spectral split, residual = $H^1 = \mathbb{C}^3 = N_{\text{fam}}$). **Hence the entire structural residual of the theory is a single geometric premise** QGEO.CONF.01: *the raw RP seam normal double is conformally* ($\mathbb{P}^1, \mu_4$) *with the carrier clock as the order-4 Möbius automorphism* $z \mapsto iz$. Given it, everything — genus-0, the four D_4 marks, the (1, 2, 3) grading, the DtN block, the rank-8 polarisation and the $(E_8)_1$ net — follows. This single premise is a constructive-geometry statement (the conformal class of the raw seam *is* the cusped sphere), *not* a finite computation, and is left honestly [O].

And even that conformal premise reduces to a milder isometry premise ([verification/v180_clock_is_mobius.py]). QGEO.CONF.01 need not be assumed as a conformal-class identity: by *uniformisation* (Poincaré–Koebe) the genus-0 double (P1) with its RP-metric conformal structure *is* $\mathbb{P}^1$ — the target is *produced*, not assumed — and by Kerékjártó (1919; Constantin–Kolev 2003) a finite-order orientation-preserving homeomorphism of S^2 is conjugate to a rotation, so the order-4 clock ($h(A_3)=4$) is topologically $z \mapsto iz$. An orientation-preserving *isometry* of a Riemannian surface is holomorphic (angle-preserving), hence a Möbius map of $\mathbb{P}^1$, and an order-4 Möbius map is $z \mapsto iz$ (elliptic, multiplier i , free orbit μ_4). So QGEO.CONF.01 is *implied* by the strictly milder, non-circular premise QGEO.ISO.01: “the carrier clock is an order-4 orientation-preserving *isometry* of the RP seam metric” — which does *not* name $\mathbb{P}^1/\mu_4$ and is the natural statement that the seam transport is metric-compatible (a holonomy preserving the RP inner product). **That isometry premise is the single, now milder, structural residual of the whole theory**; its surface realisation from the raw seam stays honestly [O].

And one step further reaches bedrock ([verification/v181_clock_is_conformal_symmetry.py]). By *equivariant uniformisation* (Nielsen realisation; for cyclic groups Kerékjártó + uniformisation) a finite-order orientation-preserving action on a genus-0 surface preserves a complex structure for which it acts by Möbius maps — so a *conformal automorphism* (strictly weaker than an isometry: every isometry is conformal, conformal allows any positive scale) already suffices to put the order-4 clock in the form $z \mapsto iz$. Since the clock is the Coxeter element of $W(A_3) = S_4$ (v117), i.e. the μ_4 deck, and a deck transformation preserves the pulled-back conformal structure *by definition*, the residual collapses to the *definitional* identification QGEO.SYM.01: *the seam’s conformal structure is the μ_4 -deck structure of the carrier clock* (the carrier μ_4 glue *is* the conformal monodromy of the seam). This is a physical/definitional statement tying P2 to the seam’s conformal geometry — *not* a finite computation and *not* reducible further without relabeling. **This is bedrock**: the chain REALIZE $\rightarrow$ theorem $\rightarrow$ MARKS+KERNEL $\rightarrow$ CONF $\rightarrow$ ISO $\rightarrow$ SYM (v175–v181) ends at a single definitional premise, left honestly [O]; everything above it is a theorem or an established citation.

A sharper, operator-checkable restatement: QGEO.ENERGY.01 ([verification/v192_qgeo_energy_clock.py]). The bedrock can be re-expressed as an energy identity rather than a conformal-class statement: *the carrier clock ρ preserves the Calderón / Dirichlet-to-Neumann energy form of the RP seam double*, $\rho^* \Lambda_\Sigma \rho = \Lambda_\Sigma$. In two dimensions Dirichlet-energy invariance *is* conformal invariance, so this feeds the same chain (conformality $\rightarrow$ Möbius $\rightarrow z \mapsto iz \rightarrow \mu_4$). Its appeal is checkability: $\Lambda_\Sigma = |k|$ is the rank-8 Calderón polarisation already in the net construction, so the premise becomes an operator identity rather than an abstract identification. **Honest scope, however**: “ ρ preserves Λ_Σ ” is plausibly *equivalent* to “ ρ is the conformal deck” (QGEO.SYM.01), so QGEO.ENERGY.01 *relocates* the bedrock to a more falsifiable surface but does *not* eliminate it — it becomes a theorem only if ρ is defined *independently* (from the carrier algebra) and the invariance is then *proven*, which is open. The bedrock therefore stays [O]; the gain is a sharper operator-analytic target, not a closure.

The non-circular sharpening: QGEO.ENERGY.02 ([verification/v193_qgeo_energy_commutator.py]).

The way to make “ ρ defined independently” precise is the *energy commutator* $[\rho, \Lambda_\Sigma] = 0$ on the rank-8 Calderón polarisation, with three sub-claims: (1) ρ is the Coxeter element of $W(A_3) = S_4$ (order $4 = |\mu_4|$, v117), defined from the *carrier* algebra, not the seam; (2) *the non-circularity hinge* — Λ_Σ must be the DtN of the *raw* RP seam (definable from the RP data alone), not the μ_4 normal-form operator; (3) $[\rho, \Lambda_\Sigma] = 0$ forces the μ_4 character grading $(1, 2, 3)$ on H^1 , already established (QGEO.COHO.M.01). An exact consistency rider pins the direction: the induced EH coefficient $k = (\ln m)/(12\pi)$ (v150) reproduces $c_3/2$ iff $\ln m = q(A_3) = \frac{3}{4}$ (the *family* glue norm), while $q(D_5) = \frac{5}{4}$ (the carrier norm) misses by $g_{\text{car}}/N_{\text{fam}} = \frac{5}{3}$ — so the seam energy form must reproduce the *family* norm (gravity is family-geometry-induced). If sub-claim (2) holds, $[\rho, \Lambda_\Sigma] = 0$ turns QGEO.SYM.01 from definitional bedrock into an *energy-rigidity theorem*; until then it is a sharp, falsifiable *proof target*, [O].

Subclaim (2) tackled — one notch closer, not closed ([verification/v194_raw_seam_dtn_rp.py]). The non-circularity hinge is essentially *met* at the canonical level: Osterwalder–Schrader reconstruction yields a canonical transfer operator $T = e^{-H}$ from the RP state and reflection alone — no coordinates (v54; §(4) of tfpt_2) — on a *quasi-free* seam state (v155: the boundary marginal of a Gaussian bulk is the compression $P\Gamma P$, a contraction). So Λ_Σ is RP-definable without ever naming $\mathbb{P}^1 \setminus \mu_4$. The commutator $[\rho, \Lambda_\Sigma] = 0$ then *closes on the finite cohomology* H^1 [E]: $\rho: z \mapsto iz$ acts on $\langle w_1, w_2, w_3 \rangle$ ($w_k = z^{k-1} dz / (z^4 - 1)$) as $\rho^* w_k = i^k w_k$, distinct characters $i, -1, -i$, so any ρ -equivariant operator is diagonal there (v177). What does *not* close is the *full- L^2* commutation beyond H^1 — it is exactly the statement that the quasi-free seam state’s modular flow is *geometric* (Bisognano–Wichmann). So the bedrock *reduces* from a bare definitional postulate to “intrinsic BW geometric modular covariance of the quasi-free seam state” — a named, standard (hard) AQFT property, one notch sharper, but still [O] (and the BW step must be established *intrinsically*, else it re-presupposes the conformal covariance it should produce).

A concrete variational handle on the residual ([verification/v196_seam_energy_functional.py]). The open BW statement can be made a discretisable extremum problem via the failure functional $E_{\text{fail}}(\Sigma, \rho) = \|[\rho, \Lambda_\Sigma]\|_{HS}^2 + \|\rho^4 - 1\|^2 + \|\Theta \rho \Theta - \rho^{-1}\|^2$, whose zero locus is exactly the QGEO.SYM.01 conditions (the energy commutator, the order-4 clock, and the RP-reflection twist). On the finite H^1 block $E_{\text{fail}} = 0$ at the μ_4 configuration ($\rho = \text{diag}(i, -1, -i)$, Λ μ_4 -equivariant, $\Theta = \text{conjugation}$) and > 0 for any μ_4 -breaking perturbation, so μ_4 is a genuine minimiser [E]; minimising E_{fail} over the *full* raw-seam DtN is precisely the BW residual above. So the bedrock is now a sharply-posed *variational target*, not a bare definition — but still [O].

The decisive crack: principal symbol exact + Tomita–Takesaki ([verification/v198_modular_commutator_reduction.py]). Two observations collapse the residual to a single non-circular state-invariance. *First*, the DtN principal symbol is $|k| = \sqrt{-\partial_\theta^2}$ (Lee–Uhlmann, v156); in the Fourier basis $e^{in\theta}$ it is $\text{diag}(|n|)$ and the clock $\rho: z \mapsto iz$ (rotation by $\pi/2$) is $\text{diag}(i^n)$, both *diagonal*, so $[\rho, |k|] = 0$ *exactly* on all of L^2 — the leading-order commutation is free on the whole boundary, never the open part. *Second*, by Tomita–Takesaki the modular operator Δ_ω is intrinsic to (M, Ω) , so a state-preserving symmetry ($\rho\Omega = \Omega$) commutes with Δ and hence with $\log \Delta \sim \Lambda_\Sigma$: thus $[\rho, \Lambda_\Sigma] = 0$ *follows from* $\omega \circ \rho = \omega$ — using only the state and reflection (which RP supplies) and *no* conformal covariance, which **removes the BW circularity worry above**. Since the finite H^1 block is already μ_4 -invariant (v177), the entire residual is now the single *state-invariance* “the raw quasi-free seam state is μ_4 -invariant” ($\omega \circ \rho = \omega$, equivalently the Gaussian bulk measure is deck-invariant) — the maximally-operational, checkable form of QGEO.SYM.01. A foundational symmetry postulate cannot be derived from nothing (the role $c = \text{const}$ plays in relativity); it is now in its sharpest, non-circular, falsifiable form, [O].

The residual, fully localised and numerically probed ([verification/v199_seam_state_invariance.py], [verification/v200_seam_variational_scan.py]). Attacking $\omega \circ \rho = \omega$ on the *raw* seam (not the μ_4 normal form) confirms it is non-circular — ρ is the Coxeter element of $W(A_3) = S_4$ ($\rho^4 = 1$), an automorphism of the raw CAR algebra, defined entirely carrier-side — and reduces it, via the quasi-free covariance $C_\Sigma = \frac{1}{2}(1 + \text{sgn } H_1)$, to $[\rho, H_1] = 0$, i.e. H_1 is *block-diagonal in the four μ_4 -character classes* $\{n \equiv r \pmod{4}\}$. The principal symbol $|k| = \text{diag}(|n|)$ passes automatically, so the entire residual is the single bounded statement “the sub-principal symbol of the raw DtN has no off-character matrix elements”. The companion variational scan minimises $E_{\text{fail}}(\theta) = \|\rho(\theta)^4 - 1\|^2 + (\text{faithfulness})$ over a free clock angle and finds its *unique* minimum at $\theta = \pi/2$ — the μ_4 clock $z \mapsto iz$ (the only faithful order-4 grading $i, -1, -i$) — a numerical sign that the variational principle selects μ_4 . This is the “next executable theorem” a working mathematician can take up; it does not close the postulate, [O].

The bounded sub-principal residual is the image of the μ_4 marks, not a free postulate ([[verification/v201_seam_subprincipal_marks.py](#)]). One genuine step further: the seam DtN is $\Lambda = |k| + M_f$ with M_f the bounded sub-principal piece = multiplication by a curvature function $f(\theta)$ on the seam circle (the standard Steklov/DtN structure), so $\langle n|M_f|n' \rangle = f_{n-n'}$. Then M_f is μ_4 -character-block-diagonal iff f has Fourier support only on modes $\equiv 0 \pmod{4}$, i.e. iff f is $\mathbb{Z}_4$ -invariant (the principal $|k|$ already commutes, v198). But a curvature sourced by the four marks, $f(\theta) = \sum_{j=0}^3 g(\theta - 2\pi j/4)$ for any profile g , has $f_m = 4g_m$ [$m \equiv 0 \pmod{4}$] — automatically $\mathbb{Z}_4$ -invariant. So the μ_4 -mark orbit (forced by v195) forces the sub-principal symbol block-diagonal; 3 equally-spaced marks ($\mathbb{Z}_3$) or 4 generic marks both break it (negative controls). The residual of $\omega \circ \rho = \omega$ thus collapses to “the DtN sub-principal symbol is *mark-local* (the seam is flat away from the μ_4 marks = the conformal-deck structure, v177)” — a locality/flatness property, structurally definitional, the narrowest form yet of QGEO.SYM.01, [O].

Certified on realistic Steklov profiles, and at the state level ([[verification/v210_mark_local_dtn.py](#)]). The mode-level argument above is carried to concrete, smooth boundary curvature: a real von-Mises bump $g(\theta)$ localised at a puncture, summed over the four marks $f(\theta) = \sum_{j=0}^3 g(\theta - j\pi/2)$, has every off-(mod 4) Fourier coefficient vanishing to $< 10^{-16}$ (exact 4-fold cancellation), the assembled DtN $\Lambda = |D_\theta| + M_f$ has $\|[\rho, \Lambda]\| < 10^{-15}$, and — the genuinely new step — the *quasi-free state* it defines, with covariance $C = \frac{1}{2}(1 + \text{sgn } H_1)$, is clock-invariant: $\rho C \rho^\dagger = C$ to $< 10^{-13}$, i.e. the actual $\omega \circ \rho = \omega$, not merely $[\rho, \Lambda] = 0$. A single off-centre bump, a $\mathbb{Z}_3$ (three-mark) source and four unequally-spaced marks each break both by $O(1)$, and the mark-local commutator stays $< 10^{-10}$ as the truncation grows ($N \in \{16, 32, 64\}$) — not an artefact. This *numerically certifies the implication* “mark-local DtN $\Rightarrow \omega \circ \rho = \omega$ ” for real profiles; the premise that the physical seam *is* mark-local stays [O] (it does not close QGEO.SYM.01).

The same residual, read dynamically: modular flow + a recovery Lindblad generator ([[verification/v238_modular_lindblad_dynamics.py](#)]). The whole QGEO.SYM.01 discussion above is *static* — a commutator condition $[\rho, \Lambda_\Sigma] = 0$. Read as a *dynamics* it is the statement that the modular flow $\sigma_t = \Delta^{it}$ (the Tomita–Takesaki one-parameter group generated by the modular Hamiltonian $\Lambda_\Sigma = \log \Delta_\omega$) *conserves* the μ_4 clock: $\rho \sigma_t \rho^{-1} = \sigma_t \forall t \Leftrightarrow [\rho, \Lambda_\Sigma] = 0$ — so the open premise is exactly “modular time conserves the carrier charge”. The dissipative half is the recovery channel T of v221 read as a generator: $L = \log T$ is a doubly-stochastic (CPTP-classical) Markov generator ($e^L = T$, $e^{L\tau} \rightarrow$ the democratic projector = the arrow of time), and its dissipative gap is $-\log((2/3)^6) = 6 \log(3/2) = \Delta$, the OS mass gap of v64 — the static recovery bound and the dynamical mass gap are *one* number, one exponentiated. Together $G = i\Lambda_\Sigma + L$ is the canonical Hamiltonian + Lindbladian (GKSL) split, the reversible (imaginary) part conserving and the dissipative (real ≤ 0) part setting the arrow. This is a finite-block exhibit of where the static compiler becomes a dynamics; the residual it leaves — that σ_t is the *geometric* modular flow (Connes–Rovelli “thermal time = physical time”) on the full seam algebra — is the same QGEO.SYM.01 bedrock, [O].

From the dynamics to a QFT skeleton on the seam ([[verification/v239_kms_thermal_time.py](#)], [[verification/v240_gns_os_reconstruction.py](#)], [[verification/v241_dhr_sectors.py](#)], [[verification/v242_spin_statistics_modular.py](#)], [[verification/v243_haag_ruelle_braiding.py](#)], [[verification/v244_spectral_action_contract.py](#)]). With the static $\rightarrow$ dynamic flip in hand, the standard AQFT machinery assembles a finite-block QFT skeleton, each step typed honestly. *Thermal time* (v239): the modular flow is KMS at $\beta = 1$, and the seam unit $2\pi = 1/(4c_3)$ fixes the modular temperature as the horizon temperature $T_H = c_3/M$. *Reconstruction* (v240): GNS gives the Hilbert space and a cyclic vector from $(\mathcal{M}, \omega)$, and the OS Hamiltonian $H_{\text{OS}} = -\log T = -L$ is positive with gap Δ — the v64 “ $H \geq 0$ + mass gap” is the recovery generator read as Euclidean time. *Spectrum* (v241/v242): the superselection sectors are the discriminant anyons of the Lean glue form (FORM.GLUE.01); fusion is recovered from the modular S by Verlinde, the Gauss–Milgram sum returns $c = 8$, and the condensation tower $16 \rightarrow 4 \rightarrow 1$ (v235/v237) selects the holomorphic $(E_8)_1$. *Scattering* (v243): the 2-particle S-matrix is the braiding monodromy, factorised (Yang–Baxter) with crossing, trivial on the condensed $(E_8)_1$. The one genuinely 4d step is typed as a contract, CONTRACT.QFT4D.01 (v244): the Connes spectral action of the seam Dirac operator, whose gravity half is v36 and whose matter half (the SM gauge group sits in the carrier, the half-spinor 16 is one generation) is the open [O] target. Its representation-theory core is now discharged ([[verification/v245_spectral_matter_content.py](#)]): the **16** is exactly one *anomaly-free* SM generation ($\sum Y = \sum Y^3 = [SU(2)]^2 U(1) = [SU(3)]^2 U(1) = 0$) with the forced SM hypercharges and the spectral-action value $\sin^2 \theta_W = 3/8$ ($g_3 = g_2 = \sqrt{5/3} g_1$), so only the seam $\rightarrow$ Dirac realisation and the run-down couplings stay open. So the residual to a full 4d QFT is *one premise* (QGEO.SYM.01) plus *one contract* (CONTRACT.QFT4D.01), not a diffuse list.

The perturbative 4d S-matrix is constructible ([[verification/v269_spert_paqft_skeleton.py](#)]). The 4d scattering layer is typed precisely as three *distinct* objects: S_{top} (the 2d DHR braiding monodromy, $|M|=1$ statistics, v243), S_{pert} (the 4d Epstein–Glaser / BV S-matrix of the spectral action), and S_{phys} (LSZ on the OS-reconstructed Wightman functions, v240) — the 2d braiding is *not* the 4d cross-section S-matrix. [E] the spectral-action a_4 interaction is power-counting renormalizable (every operator mass dimension ≤ 4 : 7 marginal + 3 relevant, a *finite* counterterm basis), so [C] by the Epstein–Glaser theorem (causal perturbation theory; Brunetti–Fredenhagen pAQFT) the perturbative S-matrix $S(g) = T \exp(i \int g L_{\text{int}})$ is built order by order by causal factorisation + finite local extension — no path integral, *no* UV divergences. [C] the gap $\Delta = 6 \log(3/2)$ gives an IR-safe adiabatic limit. [O] this is *perturbative* only: the nonperturbative ambient measure QG.AMB.01 stays open, and the canonical 4d reading remains boundary-only (v265); S_{pert} is the perturbative cross-section layer on top (QFT4D.SPERT.01).

A concrete one-loop EG number ([[verification/v271_eg_oneloop_quartic.py](#)]). The skeleton is taken from “exists” to an actual Epstein–Glaser renormalization: the order-1 product $T_1 = :L_{\text{int}}:$ needs no extension, and the first one is the order-2 T_2 , whose UV behaviour is the Steinmann *scaling degree*. [E] the massless propagator has $\text{sd} = 2$ in $d = 4$, the one-loop bubble (two propagators) has $\text{sd} = 4 = d$, so the singular order $\omega = \text{sd} - d = 0 \Rightarrow$ *exactly one* logarithmic local counterterm ($C(\omega+d, d) = 1$ — finite, renormalizable, no infinite tower); the one-loop factor is $\Omega_3/(2(2\pi)^4) = 1/(16\pi^2)$ exactly, the *same* $1/(16\pi^2)$ that normalises the spectral-action a_4 geometric quartic. [C] the ϕ^4 one-loop running is then $\beta_\lambda = 3\lambda^2/(16\pi^2)$ (symmetry factor 3 = the s, t, u channels), with the EG initial condition fixed by the spectral action, $\lambda_\Phi(\text{UV}) = 1/(16\pi^2)$. [O] this is order-2/one-loop, one diagram class; the all-order construction and the nonperturbative measure stay open (QFT4D.SPERT.02).

Red-team caveat ([[verification/redteam/rt_F_qft4d.py](#)]). Power-counting renormalizability is *not* unitarity for the gravity sub-sector: the R^2/Weyl^2 terms give a four-derivative propagator $1/(p^2(p^2+M^2))$ whose massive spin-2 mode has a negative residue (the Stelle ghost). So S_{pert} is a unitary perturbative S-matrix for the *matter+gauge* sector only; the gravity perturbative S-matrix carries the ghost — which is precisely why the *nonperturbative* QG.AMB.01 (asymptotic safety v207 / the boundary $(E_8)_1$ net) is the real frontier, not a new gap.

The gauge running is an EG order-2 number ([[verification/v273_eg_gauge_running.py](#)]). The same scaling-degree-4 mechanism applied to the gauge self-energy gives the SM one-loop coefficients from the carrier/SM content: $b_i = -\frac{11}{3}C_2(G) + \frac{2}{3}\sum_f T(R) + \frac{1}{3}\sum_s T(R)$, i.e. [E] $b_3 = -7$, $b_2 = -\frac{19}{6}$, $b_1 = \frac{41}{10}$ (GUT-normalised; the same 41 as the carrier algebra $10b_1 = g_{\text{car}}^{2g_{\text{car}}-2} + 1$, v159). [C] the physical RG directions ($b_3 < 0$ asymptotic freedom, $b_1 > 0$ the $U(1)$ Landau pole) are EG-derived and feed the unification cross-check (v246/v249); [O] the two-loop matrix (v159, PyR@TE) and the all-order construction stay open (QFT4D.SPERT.03).

The bridge to the physical S-matrix and one-loop unitarity ([[verification/v278_lsz_bridge_unitarity.py](#)]). The perturbative layer connects to the physical asymptotic S-matrix: [E] the EG time-ordered products of S_{pert} are the OS Wick-rotated correlators (v240), so amputating external legs and taking the on-shell residue gives $S_{\text{pert}} \xrightarrow{\text{LSZ}} S_{\text{phys}}$ (the three-layer S-matrix $S_{\text{top}}|S_{\text{pert}} \xrightarrow{\text{LSZ}} S_{\text{phys}}$), valid because the gap $\Delta=6 \log(3/2)$ gives Haag–Ruelle asymptotic states. [E] *one-loop unitarity* is the optical theorem: the s -channel bubble discontinuity $\text{Im} B(s)/\pi = x_+ - x_- = \sqrt{1-4m^2/s}$ is *exactly* the two-body phase space (turning on at the physical threshold $s=4m^2$), so $2 \text{Im} M = \sum \int d\Pi |M|^2$ holds and the matter+gauge sector is unitary order by order. [O] the gravity sector’s Stelle ghost is the lone non-unitary piece $\rightarrow$ QG.AMB.01 (QFT4D.SPERT.04).

The first hard data confrontation ([[verification/v246_unification_data_crosscheck.py](#)]). TFPT’s gauge-charged content *is* exactly the Standard Model (v159) and the theory adds no new states (§ “Dark matter = no new state”); the E_8 cascade is an arithmetic even-integer spine ($D_n=60-2n$, v5), *not* a particle/threshold tower. So a “TFPT-adapted” RGE run *is* the SM run — and the SM does not unify. Tested against the *measured* couplings (v159 betas, plus the standard two-loop gauge matrix): at one loop the pairwise crossings spread over $\sim 10^{13}$ – 10^{17} GeV with residual $\Delta\alpha^{-1} \approx 9$ at 2×10^{16} GeV; at two loops (RK4) the minimal spread is $\Delta\alpha^{-1} \approx 3.2$ at $\mu \approx 1.7 \times 10^{14}$ GeV — the three couplings never meet (Fig. 2). This is a falsifiable *tension*, [X]: with “no new state” there is *no* admissible gauge-threshold source, so closing the spread would require either new gauge-charged matter (a new postulate) or a non-GUT/boundary reading of the cutoff. The spectral-action 4d-GUT route is thus in genuine tension — the same status as the Connes–Chamseddine NCG Standard Model — and is likely falsified unless the theory is extended; the AQFT skeleton, DHR structure and SM rep-core survive

regardless as an emergent boundary QFT. Recorded as a kill-test, never as a confirmation.

The carrier-native Pati–Salam program ([[verification/v247_e8_branching_noi26.py](#)], [[verification/v248_ps_algebra.py](#)], [[verification/v249_ps_unification.py](#)], [[verification/v250_dirac_triple_contract.py](#)]). The unification tension above is resolved *natively* if the carrier $SO(10)$ is gauged (the carrier $D_5 \oplus A_3 = SO(10) \times SU(4)$ is Pati–Salam). *First*, the E_8 audit hull *forbids* the **126**: the **248** branches as $(45, 1) + (1, 15) + (10, 6) + (16, 4) + (\overline{16}, \overline{4})$, so the available $SO(10)$ reps are exactly $\{1, 10, 16, 45\}$ — the renormalisable 126_H is algebraically absent, making the “ 16_H over 126_H ” selection a theorem of the hull, not a fit (v247). *Second*, the Pati–Salam NCG algebra $A_F = \mathbb{H}_L \oplus \mathbb{H}_R \oplus M_4(\mathbb{C})$ gives the unimodular group $SU(2)_L \times SU(2)_R \times SU(4)_c$, the exact SM hypercharges $Y = T_{3R} + (B-L)/2$ and the matching $\alpha_1^{-1} = \frac{3}{5}\alpha_{2R}^{-1} + \frac{2}{5}\alpha_4^{-1}$, $\sin^2 \theta_W = 3/8$ (v248). *Third*, the two-step run unifies with the breaking scale on the scalaron scale, $M_{PS} = \kappa M_s$, $\kappa \in [1.0, 1.7]$ at one loop ($\rightarrow [1.0, 1.2]$ at two loops), and proton decay selecting the content (minimal excluded, the E_8 -allowed single 45 marginal); the negative controls (SM-only fails, 126 breaks the match, no-45 proton-unsafe) all fire (v249). *Fourth*, the residual “carrier is gauged” reduces to ONE constructive object, the spectral triple (A, H, D_F, J, γ) with A_F Pati–Salam and $H_F = 3 \times 16$, whose NCG axioms (the first-order condition above all) and the σ = scalaron identification are the open contract **CONTRACT.QFT4D.DIRAC.01** (v250). *Fifth*, a first concrete step on that triple is taken ([[verification/v251_first_order_sigma.py](#)]): on the lepton block the first-order condition is verified to *hold* for the Yukawa operator and to be *violated exactly* by the Majorana/ σ term — the Chamseddine–Connes–van Suijlekom “beyond first order” mechanism by which the inner fluctuations promote the SM to Pati–Salam and generate σ (honest correction: σ does *not* rescue the first-order condition; its controlled violation *is* the mechanism). *Sixth*, the full 96-dimensional finite spectral triple is then built explicitly ([[verification/v252_full_finite_triple.py](#)]) — H_F the particle+antiparticle doubling of three $SO(10)$ **16**-plets — and the NCG axioms are verified by direct computation: reality J , grading γ , KO-dimension 6, order-zero, the first-order condition (*holding* for the SM algebra including the Majorana, *violated* for the Pati–Salam algebra exactly by the Majorana on $\{\nu_R, e_R\}$), and the spectrum of D_F reproducing the fermion masses and the type-I seesaw $m_{\text{light}} = m_D^2/M_R$ — thereby discharging five of the seven obligations of **CONTRACT.QFT4D.DIRAC.01** to **[E]** (orientability, the no-junk condition and the spectral-action expansion remain open). *Seventh*, the scalaron–Pati–Salam ratio κ in $M_{PS} = \kappa M_s$ is pinned to a tight $O(1)$ interval (≈ 1.3) by the RG and shown to be a scale-free ratio of heat-kernel coefficients, $\kappa = \sqrt{(f_2/f_0) c_{PS}/c_{\text{grav}}}$, over the same 96-dim H_F ([[verification/v253_kappa_scalaron_ps.py](#)]). *Eighth*, the remaining NCG obligations are then discharged or honestly typed: the inner fluctuations Ω_D^1 give exactly the one SM Higgs doublet $(1, 2)_{1/2}$ with no junk and no **126**, while σ (the ν_R Majorana scalar, $B-L=2$) appears only as the beyond-first-order Pati–Salam field ([[verification/v254_inner_fluctuations.py](#)]); the spectral-action Seeley–DeWitt expansion a_0, a_2, a_4 yields the cosmological, Einstein–Hilbert and Yang–Mills+ R^2 +Higgs terms with $b/a^2 = 1/3$, $\sin^2 \theta_W = 3/8$ and the R^2 scalaron, making κ an explicit moment ratio ([[verification/v255_spectral_action_expansion.py](#)]); only strict orientability and strict Poincaré duality genuinely *fail* for this triple — a published feature of the NCG-Standard-Model class (KO-dimension 6 with three ν_R forces a skew intersection form, corank 1; arXiv:0902.2068, hep-th/0601192), not a TFPT defect; the physically-correct weakened versions, *quasi-orientability* (left/right irrep boxes disjoint by chirality) and *modified Poincaré duality*, do hold ([[verification/v256_orientability_poincare.py](#)], [[verification/v257_quasiorient_modpd.py](#)]). *Ninth*, the built D_F is no independent posit but the *modular/covariance induction* of the boundary seam state ([[verification/v258_dirac_covariance_induction.py](#)]): a quasi-free (KMS, $\beta=1$) state is fixed by its covariance C (a positive contraction), with one-particle modular Hamiltonian $H = \log((1-C)C^{-1})$ (Araki; Casini–Huerta) inverse to the KMS occupation $C = (1+e^H)^{-1}$; inducing the carrier covariance $C_F = P_F C_\Sigma P_F$ returns the **v252** operator *exactly* — same spectrum, chirality-odd, the Majorana an off-diagonal covariance entry with the seesaw m_D^2/M_R recovered — so D_F is the Kasparov shadow $[D_\Sigma] \otimes_{(E_8)_1} [K_{\text{car}}]$ of the boundary geometry and the **v18** Yukawas are the *readout* of C_Σ , **[E]** modulo the one **[O]** seam realisation. *Tenth*, the last open spectral-action input — the cutoff function f — is fixed by the same seam state ([[verification/v259_modular_cutoff_kappa.py](#)]): by Tomita–Takesaki/Bisognano–Wichmann the seam is KMS at $\beta=1$ (v239) and the seam unit $2\pi=1/(4c_3)$ (v58) fixes that β , so the heat-kernel cutoff $f(u) = e^{-u}$ is *derived*, not chosen, with moments $f_2/f_0 = f_4/f_2 = 1$ *exactly* (a generic Gaussian cutoff gives $\sqrt{\pi}/2 \neq 1$, so the choice is non-vacuous); hence $\kappa = \sqrt{(f_2/f_0) c_{PS}/c_{\text{grav}}} = \sqrt{c_{PS}/c_{\text{grav}}}$ loses its scheme factor and the **v253/v255** open **[O]** cutoff freedom becomes a **[C]** finite trace ratio over the 96-dim H_F (the RG $\kappa \approx 1.3$ pinning $c_{PS}/c_{\text{grav}} \approx 1.7$). So unification is now *one* theory decision (gauge $SO(10)$?) plus *one* constructive object, both falsifiable —

not a fog. The colour $SU(4)_c$ here sits inside $SO(10)=D_5$ and is distinct from the carrier family factor A_3 .

The implication is now Lean-formalised (FORM.QGEO.01). The exact algebraic core — the 4th-root character orthogonality, the order-4 clock, and `markLocal_blockDiagonal` ($f_{n-n'} \neq 0 \Rightarrow (n \equiv n' \bmod 4)$, i.e. $[\rho, M_f] = 0$) — is machine-proved in the carrier-rigidity Lean project (`TfptCarrier/SeamDeckClosure.lean`; AUDIT: PASS, axioms `propext`, `Classical.choice`, `Quot.sound` only). The premise itself is a typed structure `SeamDeckPremise` — consumed, not proved, exactly as `CalderonProjector` encodes the Paper-1 analytic input — so the conditional theorem “`SeamDeckPremise  $\Rightarrow \omega \circ \rho = \omega$` ” is now [E] while QGEO.SYM.01 stays the one open [O] postulate.

The two residuals are one canonical metric: the pillowcase (`[verification/v214_seam_pillowcase.py]`). The isometry residual QGEO.ISO.01 (v180) and the mark-locality residual QGEO.SUBPRIN.01 (v201) are not independent: both follow from a single canonical choice — *the seam carries the uniformising flat orbifold metric of its conformal class*. The four marks make the orbifold sphere $S^2(2, 2, 2, 2)$, whose orbifold Euler characteristic $\chi_{\text{orb}} = 2 - 4(1 - \frac{1}{2}) = 0$ is *Euclidean*: by Troyanov’s cone-metric uniformisation it carries a flat metric, unique up to scale, with curvature concentrated at the four cone points (Gauss–Bonnet: four deficits π sum to $4\pi = 2\pi\chi(S^2)$) — this is the v201 “flat away from the marks”. And the order-4 clock is no longer an extra assumption but a *consequence* of the cross-ratio 2 that v168 already proved: $j(\lambda) = 256(\lambda^2 - \lambda + 1)^3/(\lambda^2(\lambda - 1)^2)$ gives $j(2) = 1728$, so the modulus is the *square* torus $\tau = i$, whose complex multiplication by $\mathbb{Z}[i]$ supplies the order-4 isometry $z \mapsto iz$ (explicitly $(x, y) \mapsto (-x, iy)$ on $y^2 = x^3 - x$, order 4); a generic cross-ratio gives $j \notin \{0, 1728\}$ and only $\mathbb{Z}/2$, so the clock is specific to μ_4 . Thus the two open QGEO residuals *collapse to one* canonical-metric statement and QGEO.ISO.01 ceases to be an imposed symmetry. A genuine reduction and unification, *not* a closure: the premise “the RP seam metric is the uniformising constant-curvature representative of its conformal class” stays [O] — milder and more canonical than the bare isometry premise, and now *shared* by both residuals (QGEO.PILLOW.01).

The converse, on the canonical metric: raw seam $\Rightarrow$ mark-locality (`[verification/v264_raw_seam_marklocal.py]`). Composing the chain explicitly closes the gap between v216, v201 and v198 on the uniformising metric: the flat pillowcase has its curvature *only* at the four cone points (v216), so the DtN sub-principal symbol M_f is a μ_4 -orbit sum, hence $\mathbb{Z}_4$ -invariant (Fourier support on $4\mathbb{Z}$, verified by FFT), hence block-diagonal in the clock-character basis, hence $[\rho, M_f] = 0$; with the principal symbol $|k|$ already commuting (v198) this gives $[\rho, \Lambda] = 0$ and $\omega \circ \rho = \omega$ *on this metric*. A negative control (curvature *off* the μ_4 orbit) breaks $\mathbb{Z}_4$ -invariance and the chain fails. So the entire bedrock collapses to the *single* metric-realisation statement “the raw RP seam carries the uniformising flat pillowcase metric” — which is QGEO.SYM.01. A genuine reduction/sharpening (QGEO.ENERGY.03), [O] — it does *not* prove the raw seam must carry that metric; that converse is the irreducible postulate.

The minimal axiom: a rigidity theorem (`[verification/v267_qgeo_rigidity_minimal_axiom.py]`). One can sharpen the bedrock to its irreducible kernel. [E] the *single* bare statement “the seam admits the carrier’s order-4 clock as a conformal symmetry cyclically permuting its four marks” forces everything: an order-4 Möbius orbit $\{a, ia, -a, -ia\}$ has cross-ratio 2 *independent of a* ; cross-ratio 2 $\Leftrightarrow j=1728$ (only $\lambda \in \{-1, \frac{1}{2}, 2\}$) $\Leftrightarrow$ the order-4 CM automorphism $(x, y) \mapsto (-x, iy)$ on $y^2 = x^3 - x$; and from the square configuration the unique flat pillowcase metric (Troyanov), mark-locality, $[\rho, \Lambda] = 0$ and $\omega \circ \rho = \omega$ all follow (v214/v201/v264/v198). [E] neg controls: a generic four-mark set has $j \neq 1728$ and only a $\mathbb{Z}_2$ symmetry (no clock), the hexagonal point $j=0$ has a $\mathbb{Z}_6$ symmetry (order 6 $\neq 4$) — only the square (order-4) point realises the μ_4 clock. [C] a second, independent reason: $\{1, i, -1, -i\}$ is defined over $\mathbb{Q}$ with $j=1728$ (CM by $\mathbb{Z}[i]$), so by Belyi/dessins Galois-rigidity any deformation breaks the rational compiler readouts. [O] So QGEO.SYM.01 is now in its minimal sharpest form — the bare existence of that order-4 conformal symmetry, which (like $c=\text{const}$ in relativity) cannot be derived from nothing. The *reduction* is closed (axiom $\Rightarrow$ everything); the axiom itself stays the one foundational postulate (QGEO.SYM.02).

The flat geometry closes the commutator to all orders (`[verification/v276_qgeo_flat_closes_commutator.py]`). The state-level residual $[\rho, H] = 0$ — the one operator equation the bedrock rests on — is *not* a separate analytic gap once the flat $\tau=i$ pillowcase is granted. [E] if $H = f(\Delta_{\text{flat}})$ is a function of the intrinsic flat Laplacian (the DtN of a flat metric, no curvature corrections) and the order-4 deck ρ ($z \mapsto iz$) is a flat- $\tau=i$ isometry, then $[\rho, \Delta] = 0 \Rightarrow [\rho, H] = 0$ *exactly*, to all orders (an isometry commutes with every spectral function of its Laplacian; verified on the discretised square torus, with a generic non-isometry as a failing control). This *upgrades* the leading-order commutation v198 (principal symbol) and v201 (subprincipal) to the

full H — flatness removes the lower-order terms. [O] so QGEO.SYM.01 collapses to a *single geometric* statement, “the raw RP seam state is the flat $\tau=i$ pillowcase quasi-free state ($H=\sqrt{-\Delta_{\text{flat}}}$)”; everything downstream is [E], and the remaining human obligation is one constructive-QFT proof that the raw seam DtN state *is* that flat geometry (QGEO.SYM.03).

The obligation as one precise lemma, Lean-backed ([verification/v279_qgeo_obligation_lemma.py]). The bedrock is now written as a single constructive-QFT *lemma*: *given* the premise “the raw seam carries the flat $\tau=i$ pillowcase metric g_{flat} (Trojanov-unique, cone angles π), so $\Lambda_{\Sigma} = \sqrt{-\Delta_{g_{\text{flat}}}}$ ”, the conclusion $\omega_{\Sigma} \circ \rho = \omega_{\Sigma}$ (and hence mark-locality, the metric inclusion, E_8) follows. [E] the reduction DAG from “free RP seam” to this premise has *exactly one* un-discharged leaf — “the raw seam state *is* the flat state” — every other node being closed; and the geometric side is fixed (the flat DtN is Fourier-diagonal), so the obligation is a *state-identification*, not a geometric ambiguity, with two candidate routes ([C]): RP-state uniqueness on the flat $\tau=i$ orbifold, or Trojanov + OS reconstruction. [E] the all-orders closure step is now a *Lean theorem*: `TfptCarrier/SeamDeckClosure.lean` (`flat_all_orders_clock`) proves that any spectral function $g(\Delta_{\text{flat}})$ of the Fourier-diagonal flat Laplacian commutes with the carrier clock $\rho = \text{diag}(i^n)$ — upgrading v198/v201 to the full operator with only the standard axioms, no `sorry` (FORM.QGEO.03). [O] the one premise stays the honest axiom (QGEO.OBLIG.01).

Self-investigation: the experiment and the simplification ([verification/v280_pillowcase_steklov.py], [verification/v282_e8_tau_i_unification.py]). Two things can be done *ourselves*. (i) A direct numerical experiment on the actual flat $\tau=i$ pillowcase orbifold realises the whole chain: the order-4 deck permutes the four cone points (2 fixed + 1 swapped) and $[\rho, \Delta]=0 \Rightarrow [\rho, H]=0 \Rightarrow [\rho, C]=0 \Rightarrow \omega \circ \rho = \omega$ is numerically exact, with the Steklov DtN positive and metric-determined — route-(i) evidence (RP-state uniqueness), the premise still [O]. (ii) A candidate *simplification*: the $(E_8)_1$ character $\chi_{E_8} = \Theta_{E_8}/\eta^8 = j^{1/3}$ gives $\chi_{E_8}(i) = 1728^{1/3} = 12$, so $\tau=i$ is *both* the order-4 CM point (QGEO.SYM.01, flat geometry) *and* the $(E_8)_1$ partition-function modulus (QG.AMB.01, holomorphy) — the two open premises are two faces of one object, “the raw seam is $(E_8)_1$ at $\tau=i$ ”, dropping the obligation count $2 \rightarrow 1$ (QGAMB.UNIFY.01).

The two proof routes, decomposed ([verification/v284_route_i_rp_uniqueness.py], [verification/v285_route_ii_seam_condensation.py]). The single open lemma can be attacked from two sides, and both are now lemma chains with the dischargeable steps discharged. *Route (i)* (RP-state uniqueness) is a six-lemma chain — [E] five are already [E]/[O] (RP $\rightarrow$ DtN v194; marks $\rightarrow$ pillowcase v195/v216; the flat Steklov operator $= \sqrt{-\Delta_{\text{flat}}}$, principal symbol $|\xi|$ with no curvature term; covariance from the DtN v258; μ_4 -invariance v276), and [C] Trojanov ($\chi_{\text{orb}}=0$) reduces the residual to “the raw seam is the constant-curvature geometric state”. *Route (ii)* (seam-net condensation) is a four-lemma chain — [E] three are discharged (no abelian sector $\Leftrightarrow \det K=1 \Leftrightarrow$ holomorphic $\Leftrightarrow E_8$ v234/v277/v281; the tower $16 \rightarrow 4 \rightarrow 1$ v92), leaving “the free RP seam condenses the μ_4 -Lagrangian glue” = QGEO.SYM.01. [E] **the two open lemmas coincide** — both say “the raw seam is the $(E_8)_1$ at $\tau=i$ ” (v282), so the canonical equivalence between the raw seam KMS/DtN state and the holomorphic $(E_8)_1$ net at $\tau=i$ is the *single* [O] statement that closes both (QGEO.ROUTEI.01/QGAMB.ROUTEII.01).

The Seam Equivalence Theorem, named and firewalled ([verification/v286_seam_equivalence_contract.py]). That single statement is now the named target SEAM.EQUIV.01: *the raw RP seam state reconstructs canonically the holomorphic $(E_8)_1$ boundary net at the $\tau=i$ pillowcase* (respecting the DtN, the KMS flow, the μ_4 deck, the Q-system extension, $\det K=1$ and the $(E_8)_1$ character). [E] the contract types four objects and six arrows — two closed (the two routes), one conditional (v282), one open Gral — and enforces an *import firewall*: no Route-i module imports any Route-ii module or vice versa, so the two routes can converge only through SEAM.EQUIV.01 (this kills the “ E_8 smuggled into the geometry and pulled back out” attack). The two proof routes are then attacked directly: [E] Route A (AQFT, [verification/v287_free_rp_bulk_to_holomorphic_boundary.py]) reduces the Gral to *one* standard theorem, “invertible (SRE) Gaussian bulk $\Rightarrow$ single-sector boundary” (4/5 lemmas discharged); and [E] Route B (DtN, [verification/v288_full_l2_subprincipal_z4.py]) *proves* the full- L^2 lift of the sub-principal $\mathbb{Z}_4$ block-diagonality (a mark-sourced curvature has Fourier support only on $m \equiv 0 \pmod{4}$), so $[\rho, \Lambda]=0$ on all of L^2 , shrinking the residual to the single sharper question “why is the raw seam sub-principal term mark-local?” (SEAM.EQUIV.A01/SEAM.EQUIV.B01).

Route B sharpened to one analytic lemma; both routes precisely scoped. That last question is decomposed ([verification/v289_marklocal_raw.py], MARKLOCAL.RAW.01) into a five-lemma chain whose only open link is [O] *Flat-Away* — RP + gap + the four branch marks force the seam curvature

to vanish away from the marks — while the conic parametrix ([C], b-calculus/Melrose), the μ_4 -orbit source, the $4\mathbb{Z}$ Fourier support (v264/v288) and the full- L^2 commutation (v288) are discharged: 4/5 done, so Route B closes the moment Flat-Away does. The two claims are firewalled against over-reading: Route B does *not* assert “full- L^2 QGEO” — its exact-label split is Fourier-lemma (Formal/Exact), operator test (Numerical Exact), transfer to the raw conic DtN (Conditional, v264), mark-locality from the raw seam (Open Selection, v289); and Route A’s “one standard theorem” is itself a composition of citable results (Kitaev/Freed–Hopkins, Mürger/Kawahigashi–Longo–Mürger, Kawahigashi–Longo, [verification/v287_free_rp_bulk_to_holomorphic_boundary.py]) whose only open hypothesis — seam-bulk invertibility — coincides with Route B’s Flat-Away. Both routes therefore meet at the *same* geometric input, with no circularity (the import firewall of v286).

Red-team: $\mathbb{Z}_4$ -invariance is not mark-locality. A deliberate adversary ([verification/v290_z4_smooth_curvature_adversary.py]) shows the honest limit of Route B’s lift: a *smooth* $\mathbb{Z}_4$ -symmetric off-mark curvature $f = \varepsilon \cos(4\theta)$ passes the v288 commutator ($[[\rho, \Lambda]] = 0$, since $i^n - i^{n\pm 4} = 0$) yet is not mark-local — so $[\rho, \Lambda] = 0$ is necessary but not sufficient. The modulus is instead excluded by the spectral data: it shifts the DtN spectrum off the flat ladder and the heat trace ($\max|\Delta\lambda| = 0.155$, $\Delta\text{Tr} = 0.019$ at $\varepsilon = 0.3$), i.e. the KMS weight and $(E_8)_1$ character that SEAM.EQUIV.01 must match. Consequently the open lemma is named in its own right ([verification/v291_flataway_contract.py], FLATAWAY.RP.01): *the smooth curvature density in the DtN sub-principal symbol vanishes away from the four branch marks*, with three independent attack routes — spectral-rigidity and heat-kernel (the flat seam is the *unique* spectral/heat match of the smooth $\mathbb{Z}_4$ family) and RP-energy/Troyanov (the constant-curvature minimiser) — closing any one of which closes Route B.

All three routes attacked; Flat-Away reduced to one selection principle. Each route is now carried to its precise reduction: [E] the *heat* route ([verification/v292_flataway_heat_reduction.py]) shows the heat-trace deviation $\Delta\text{Tr}(e^{-t\Lambda}) = c(t)\varepsilon^2 + O(\varepsilon^4)$ is a positive-definite quadratic form in the off-mark curvature (leading invariant $\|f\|_{L^2}$), so Flat-Away $\Leftarrow$ “the seam fixes the a_2 heat coefficient”; [E] the *spectral* route ([verification/v293_flataway_spectral_hessian.py]) lifts this to the full smooth- $\mathbb{Z}_4$ space — mode $4k$ splits the degenerate Steklov level $|2k|$, the mismatch Hessian over $\{4, 8, 12\}$ is positive-definite ($\{0.497, 0.500, 0.503\}$), so the flat seam is a *strict* isolated minimum (answering “only one direction”); and [E]/[C] the *variational* route ([verification/v294_flataway_rp_energy.py]) gives, with the rigid mark divisor and Troyanov uniqueness, the flat cone metric as the unique strictly-convex curvature-energy minimiser. The three converge on *one* external selection principle (fix a_2 / pin the Steklov spectrum / realise the energy-minimiser); closing any one closes Route B, and with Route A’s holomorphy half, SEAM.EQUIV.01.

The heat route made exact and Lean-formalised. The positive-definiteness underpinning the heat route is now a theorem, not a numerical scan ([verification/v295_flataway_a2_exact.py]): the first variation of $\text{Tr } e^{-t\Lambda}$ vanishes (the smooth off-mark curvature has zero mean, Gauss–Bonnet), and since $\Lambda \mapsto \text{Tr } e^{-t\Lambda}$ is convex (Klein/Peierls), $\varepsilon=0$ is the *global* minimum — so $\Delta\text{Tr} \geq 0$ for every deformation, $= 0$ iff $f = 0$; the exact second-order coefficient is the divided-difference kernel of e^{-tx} (validated to 10^{-6}). The algebraic $\Delta=0 \Leftrightarrow$ flat step is machine-formalised in Lean (FORM.FLATAWAY.01, SeamDeckClosure.lean: a positive-weighted sum of squares is positive-definite; axiom-clean). So Flat-Away’s heat route is reduced to the single external invariant “the seam fixes a_2 ”, with its positivity both proved and formalised. The a_2 coefficient itself is now in *closed form* ([verification/v296_flataway_a2_closed.py]): the operator tails telescope geometrically to $W_k^{\text{tail}} = t(1 - e^{-4kt})/(8k(1 - e^{-t}))$, leaving a finite $(4k-1)$ -term middle, e.g. $W_1(t) = t(2te^t + e^{3t} + 3e^{2t} + e^t - 1)e^{-3t}/8$ (manifestly positive; small- t $W_k = t/2 + O(t^3)$). Symmetrically, Route A’s “one standard import” is now a precise, citable theorem stack ([verification/v297_route_a_literature_stack.py]): LIT-A (Kitaev; Freed–Hopkins) $\rightarrow$ LIT-B (Mürger; Kawahigashi–Longo–Mürger) $\rightarrow$ LIT-C (Conway–Sloane; Dong–Xu) compose into “invertible bulk $\Rightarrow (E_8)_1$ ”, all three legs established, the single open hypothesis (seam-bulk invertibility) coinciding with Route B’s Flat-Away — so both routes are reduced to citable/established machinery plus the *one* shared geometric input.

One surface for seam, carrier and lattice: the Kummer/K3 ([verification/v260_k3_kummer_unification.py]). The pillowcase is the elliptic-involution quotient $T_{\tau=i}^2/(z \mapsto -z)$, its four marks the four 2-torsion fixed points; squaring the construction to complex dimension two gives the *Kummer surface* $\text{Km}(T^2 \times T^2)$, a K3, whose three facets reproduce three TFPT objects at once. [E] the 16 nodes (the 2-torsion $|A[2]| = 2^4 = 16$ of the abelian surface) resolve to 16 exceptional $\mathbb{P}^1$ ’s $= \dim S^+ = \Lambda^{\text{even}}(\mathbb{C}^5) = 16$ (v197), the carrier-16 as the *square* of the seam-4; [E] the

cup-product form on $H^2(K3, \mathbb{Z})$ is the even unimodular lattice $U^3 \oplus E_8(-1)^2$ of signature $(3, 19)$, so the same E_8 the carrier glues to (v1) is the surface's intersection form. (Honestly: the 16 node classes span the rank-16 Kummer lattice, a sublattice *distinct* from the $E_8(-1)^2$ summand — both exact, not identified.) Thus the v1 “glue $D_5 \oplus A_3 + \mu_4 \Rightarrow E_8$ ” is the lattice shadow of a single smooth object that already carries the seam and the carrier; E_8 appears both *locally* as the du Val $\mathbb{C}^2/2I$ singularity (the Poincaré-sphere link, v232/v236) and *globally* as this H^2 summand. A unification (ARCH.K3.01), *not* a closure: the lattice arithmetic is [E], the carrier $\leftrightarrow$ node identification [C].

Synthesis — the Modular Spectral Closure ([verification/v261_modular_spectral_closure.py]). The ten steps above are not ten independent results: they are facets of one relative spectral datum

$$\text{TFPT}_{\text{QFT}} = (\mathcal{A}_\Sigma, \omega_\Sigma, \Delta_\Sigma, \rho, A_F, H_F, D_F, J, \gamma, S_{\text{rel}})$$

(seam algebra, KMS state, modular generator, μ_4 clock; the carrier finite triple; the relative spectral action), governed by a single *Modular Spectral Universality* statement: *the unique reflection-positive, holomorphic, index-4 KMS seam state ω_Σ minimises the modular clock defect, induces via the Kasparov/covariance map (v258) the 96-dim KO-6 finite triple D_F (v252), whose inner fluctuations (v254) generate the gauged Pati–Salam spectral action with the seam KMS cutoff (v259) — so Dirac, cutoff, gauging, glue and orientability are not separate choices but readouts of one seam state. The certificate v261 pins the cross-consistency: one number $4 = [B:A] = |\mu_4| = 2\chi = |(\mathbb{Z}/2)^2|$ (index = marks = fixed points = glue order), one carrier-16 ($\dim S^+ = \Lambda^{\text{even}}(\mathbb{C}^5) = \text{Kummer nodes}$, $H_F = 96$), and one gap $\Delta = 6 \log(3/2)$ (OS mass gap = recovery rate = Lindblad gap, static and dynamic). The honest account of completeness is therefore:*

- **[E]/[C] QFT-complete (modulo one premise).** The admissible OS sector, the boundary $(E_8)_1$ net, the relative spectral action, D_F (as covariance induction), the cutoff (as the KMS weight) and the gauging (as inner fluctuations) are one relative object — closed once QGEO.SYM.01 (the seam realisation: *the raw quasi-free seam state is μ_4 -invariant*) is granted. That single [O] premise is the only structural input the QFT layer still consumes.
- **[O] Ambient QG — separate by design.** The full covariant metric-sector / ambient quantum-gravity measure (QG.AMB.01) is gap-decoupled ($\Delta_{\text{eff}} = 1.648 > 0$) and is *not* a blocker for the QFT layer: it is a different (certification) object, not a missing piece of the boundary QFT.

So “solving the QFT” here means exactly this: the boundary QFT is a single relative object reduced to one foundational symmetry postulate, with ambient QG held apart — not a diffuse list of open interfaces. The MSC closes the boundary relative QFT object modulo QGEO.SYM.01; it does *not* assert SM-only 4D gauge unification, and the canonical 4D reading is boundary only.

4D-QFT fork policy — a frozen decision tree, not an open ambiguity ([verification/v265_qft4d_fork_freeze.py])

The 4D reading is a *branch policy*, machine-frozen in qft4d_fork_freeze.json:

- **A (canonical) — boundary only.** [C] *Boundary-only is the canonical 4D reading*: TFPT delivers the boundary QFT, the relative spectral action and the SM readout; a 4D-GUT is *not claimed by default* (E_8 is the audit hull, never an unbroken 4D gauge group).
- **SM-only 4D-GUT unification is killed.** [X] the plain SM couplings do not meet (at the $\alpha_1=\alpha_2$ scale α_3^{-1} is off by ~ 5.6 ; spread ~ 8.8 at 2×10^{16} GeV, v246). This is the canonical *expectation*, not a failure.
- **B (conditional) — carrier-native Pati–Salam UV branch.** [C]/[O] permitted *only* with the E_8 -allowed content $\{1, 10, 16, 45\}$ (no 126, v247), $\sin^2 \theta_W = 3/8$ (v248), $\kappa = M_{PS}/M_s$ in the frozen $O(1)$ band (v249/v253), and a threshold model with a proton-decay kill test; carrier gauging itself stays the contract CONTRACT.QFT4D.DIRAC.01.
- **Proton-decay rule, now applied** ([verification/v266_ps_threshold_proton.py]). [X] the explicit two-step thresholds give $d=6$ $\tau_p(p \rightarrow e^+ \pi^0)$: the *minimal* 16-Higgs content ($M_{GUT} \sim 2.4 \times 10^{15}$) gives $\tau_p \sim 4.4 \times 10^{33}$ yr < Super-K (2.4×10^{34}) and is *killed*; the E_8 -allowed $(15, 1, 1)$ [the single 45] raises $M_{GUT} \sim 5.9 \times 10^{15} \Rightarrow \tau_p \sim 1.6 \times 10^{35}$ yr, proton-safe *now* but sitting at the Hyper-K reach ($\sim 1.4 \times 10^{35}$ yr) — so branch B is *content-selected* (must be $+(15, 1, 1)$) and *Hyper-K-decisive within the decade*, not an open-ended hope (only one 45 in the hull, so it stays structurally marginal; τ_p carries an $O(3)$ hadronic band).

The fork is thus a frozen decision tree with explicit promotion/demotion rules, *not* an open ambiguity.

And the bedrock is now a falsifiable kill-test, not a proof-promise ([`verification/v215_seam_deck_killtest.py`]). A foundational symmetry cannot be derived from nothing (Bisognano–Wichmann would presuppose the very covariance it should produce); the honest move is to recast QGEO.SYM.01 as four independently-falsifiable predictions about the raw seam DtN. The non-circular reduction (v198/v199/v201) gives $\omega \circ \rho = \omega \Leftrightarrow [\rho, C] = 0$ for the quasi-free covariance, with the principal symbol $|k|$ commuting exactly, so the whole residual sits in the bounded sub-principal M_f (equivalently: the raw Gaussian bulk form A is μ_4 -deck-invariant). The four levers are **K1** exactly four marks ($b_1 = \# \text{marks} - 1 = 3 = N_{\text{fam}} \Rightarrow \# \text{marks} = 4 = |\mu_4|$); **K2** the *square* configuration (cross-ratio $2 \Rightarrow j = 1728$, $\text{Aut} = \mathbb{Z}/4$; the clock order $h(A_3) = 4$ selects $\mathbb{Z}/4$, not the $j = 0$ $\mathbb{Z}/6$, and a generic mark set has only $\mathbb{Z}/2$); **K3** the mod-4 sub-principal support ($[\rho, M_f] = 0 \Leftrightarrow f$ is $\mathbb{Z}_4$ -invariant, automatic for a μ_4 -mark sum); and **K4** the transfer gap $(2/3)^6 = 64/729$. These are frozen in `freeze_file.csv` (`seam_deck_symmetry`) and carrier-side green, so any future drift fails the suite. The *decisive* kill is deferred to QGEO.REALIZE.01: a first-principles raw-seam-collar DtN must *produce* four square marks without inputting them — an *emergence* test, the non-circular alternative to assuming the symmetry. [E] carrier-side (the four levers), [O] seam-side (the emergence): a kill-test, *not* a closure (QGEO.KILL.01).

And the mark count is now *derived*, not assumed: four marks from Gauss–Bonnet ([`verification/v216_marks_gauss_bonnet.py`], [`verification/v217_marks_emergence_scan.py`]). The K1 lever (“exactly four marks”) is the one that can be *executed* today. If the marks are $\mathbb{Z}_2$ branch points (cone angle π , deficit π) on the flat seam sphere ($\chi = 2$ — the same χ that supplies the 8 in c_3), Gauss–Bonnet *forces* their number: $n\pi = 2\pi\chi = 4\pi \Rightarrow n = 2\chi = 4 = |\mu_4| = N_{\text{fam}} + 1$. The closed Euclidean sphere 2-orbifolds are exactly $(2, 3, 6)$, $(2, 4, 4)$, $(3, 3, 3)$, $(2, 2, 2, 2)$, and *three independent* criteria — all cone orders = 2 (the $|\mathbb{Z}_2|$ sheet branch), rank $H^1 = \# \text{marks} - 1 = 3 = N_{\text{fam}}$ (the three generations pick the 4-mark square over the 3-mark hexagonal $(3, 3, 3)$), and a free order-4 deck — converge on the pillowcase $(2, 2, 2, 2)$. The numerical sibling v217 confirms it on the actual DtN/state: scanning the number n and the positions freely, the 4-equally-spaced (square) configuration is the unique joint minimiser of {clock-invariance, 3-family cohomology, transfer gap $(2/3)^6$ }. So the mark *count* and *equality* move from input to derived; only the square *modulus* (cross-ratio 2, the order-4 clock) stays the one carrier input, and the from-zero raw-seam emergence QGEO.REALIZE.01 stays [O]. [E] (count + equality), [O] (the square modulus and the raw-seam realisation) — QGEO.MARKS.03 / QGEO.EMERGE.01.

QGEO.SYM.01 is *not* a theorem — it is the final seam-identification premise

The single load-bearing bolt of the whole construction, on one page.

- What is identified.** That the conformal structure of the raw RP seam double *is* the μ_4 -deck structure of the carrier clock — i.e. the carrier (P2) and the seam’s conformal geometry are the same object. Equivalently, in Riemann–Roch *index* language: the seam’s degree-4 mark divisor on the genus-0 double has $h^0 = \deg + 1 = 5 = g_{\text{car}}$ and rank $H_1 = \deg - 1 = 3 = N_{\text{fam}}$ (with $\Lambda^{\text{even}}(\mathbb{C}^5) = 1 + 10 + 5 = 16 = \dim S^+$), so P2 is the genus-0 mode count of the four-mark seam divisor ([`verification/v228_rr_index_gate.py`]) — the same definitional identification, in index form, conditional on the seam producing that degree-4 divisor.
- Why it is weaker than QGEO.CONF.01/ISO.01.** It does *not* name $\mathbb{P}^1$ or μ_4 (uniformisation *produces* them) and asks only for a conformal *symmetry*, not a metric isometry or a pre-named conformal class — strictly the mildest of the three.
- What follows automatically (the conditional theorem).** *Given* QGEO.SYM.01, equivariant uniformisation puts the order-4 clock in the form $z \mapsto iz$ on $\mathbb{P}^1$, the four parabolic marks are its free orbit μ_4 , H^1 carries the $(1, 2, 3)$ grading, the seam Calderón contraction is the rank-8 K_Σ polarisation, and the index-4 simple-current extension is $(E_8)_1$ — the *entire* AQFT closure of v175–v181.
- What would falsify or unseat it.** A seam double of genus > 0 (uniformisation to $\mathbb{P}^1$ fails); a carrier clock that is *not* a conformal symmetry of the seam geometry (a non-deck transport); or a fourth chiral generation ($N_{\text{fam}} \neq 3$ collapses $b_1 = N_{\text{fam}}$ and the four-mark μ_4 structure). The first two are constructive-geometry questions about the raw seam; the third is the standard family-count kill test.

Honest framing. The correct external statement is *conditional*: “Given QGEO.SYM.01, the RP seam boundary realises the μ_4 deck structure, and the AQFT closure to $(E_8)_1$ follows through the verified chain v175–v181” — *not* “the theory *derives* the seam geometry”. The premise stays [O]; everything downstream of it is [E].




Figure 1: The emergent-QFT skeleton on the seam (v238–v246): time as the modular flow, the arrow as the recovery generator, particles as DHR sectors, the S-matrix as the braiding, and the 4d matter content as the carrier’s $SO(10)$ **16**. Each step is machine-checked; the two open items (QGEO.SYM.01 postulate and the unification tension) are in red. The whole skeleton is *conditional* on QGEO.SYM.01 and is neither solved nor experimentally confirmed.

This is not a gap to apologise for — it is TFPT’s one fundamental postulate. Every fundamental theory rests on an irreducible premise (the constancy of c in relativity, the equivalence principle, the canonical commutator). QGEO.SYM.01 is that premise for TFPT: a single geometric identification (seam conformal structure = carrier μ_4 deck) from which the entire algebraic and topological structure unfolds *by theorem*. The achievement is not that there is no postulate — it is that the whole theory has been compressed onto *exactly one*, sharply located and explicitly falsifiable.

The Seam-Holomorphy Theorem — the one named closing target [E]/[O]
([verification/v234_seam_holomorphy_selection.py])

The bedrock above can be stated as *one condition with three provably-equivalent faces*, so that “closing the theory” becomes a single, sharply-bounded analytic theorem. **The condition:** the seam carries *no nontrivial abelian sector*. Its three faces all force E_8 :

1. **AQFT** (v83/v154): the boundary chiral net is *holomorphic* (μ -index 1, single vacuum sector). Holomorphic + $c = 8 \Rightarrow$ the unique even unimodular rank-8 lattice net $(E_8)_1$, and $c = 8 = g_{\text{car}} + N_{\text{fam}}$ is itself structural.
2. **Geometry** (v232): the seam link is a *homology 3-sphere* $\Leftrightarrow$ its binary group Γ is *perfect* $\Leftrightarrow \Gamma = 2I$ (the Poincaré sphere $S^3/2I$).
3. **Representation theory** (v219): exactly *one* 1-dim irrep (the trivial one).




Figure 2: Honest data cross-check (v246): the spectral-action prediction $g_1=g_2=g_3$, $\sin^2 \theta_W=3/8$ versus the measured couplings run with the SM RGEs (one-loop solid, two-loop dashed). The three lines do *not* meet at any scale (1-loop residual $\Delta\alpha^{-1} \approx 9$ at 2×10^{16} GeV; 2-loop minimal spread ≈ 3.2 at 1.7×10^{14} GeV) — a falsifiable tension. With SM content and no new states there is no admissible threshold source: not a fit, and not rescued by a cascade.

These are the *same* number: $\#(1\text{-dim irreps}) = |\Gamma^{\text{ab}}| = |H_1(S^3/\Gamma)| = \#(\text{mark-1 affine-Dynkin nodes})$, which equals 1 *only* for E_8 ($A_n \rightarrow n+1$, $D_n \rightarrow 4$, $E_6 \rightarrow 3$, $E_7 \rightarrow 2$, $E_8 \rightarrow 1$). So $2I$ is the unique nontrivial perfect finite $SU(2)$ subgroup and $S^3/2I$ the unique homology-sphere space form. **Consequence:** Research Contract 2 (G_{net} /Target A), the carrier P2, and the Kleinian seam are *not* separate gates — they are this *one* condition.

The closing chain (what a complete closure would be), [O] at one step:

$$\begin{array}{c} \text{RP} + \text{gap} + \text{chirality} \xrightarrow{\text{v160}} \text{quasi-free bulk} \xrightarrow{?} \text{holomorphic boundary net} \\ \xrightarrow{[\text{E}]} (E_8)_1 \xrightarrow{\text{v219/v228}} 2I, (2,3,5), \text{P2} \xrightarrow{\text{v179-v181+Kleinian genus-0}} \text{QGEO.SYM.01.} \end{array}$$

The free-bulk premise is already a fixed-point theorem (v160–v165); the Kleinian model supplies the genus-0 of the uniformisation chain for free. **The single remaining analytic step is “free Gaussian RP bulk $\Rightarrow$ holomorphic (single-sector) boundary net”** — the bulk–boundary statement that an invertible (Gaussian) bulk has a holomorphic chiral boundary. That one theorem, if proven, closes the entire structural layer; v_{geo} stays the No-Unit metrology primitive and F_{transfer} stays external physics. This box certifies only that the three faces coincide and force E_8 ([E]); the analytic step itself is the open [O].

A physical realisation of the step ([verification/v235_seam_chern_simons.py]). The step lands in standard physics. A free gapped *bosonic* 2+1d bulk is an abelian Chern–Simons / K -matrix theory (even integer K), where $\#\text{anyons} = |\det K|$, the edge chiral central charge is $c = \text{signature}(K)$, and the edge net is *holomorphic* $\Leftrightarrow |\det K| = 1$ — so “no nontrivial abelian sector” is precisely $\det K = 1$. The extension tower of v92 is the anyon-condensation tower read by determinant: $D_5 \oplus A_3$ ($\det 16$, 16 anyons) $\rightarrow SO(16)_1 = D_8$ ($\det 4$) $\rightarrow (E_8)_1$ ($\det 1$, holomorphic), all rank 8 with $c = \text{signature} = 8 = g_{\text{car}} + N_{\text{fam}}$, so the holomorphic end is the *Kitaev E_8 quantum-Hall state*. Holomorphy $\Leftrightarrow$ condensing the order- $|\mu_4| = 4$ *Lagrangian glue* ($\det 16 \rightarrow 16/4^2 = 1$); a partial order-2 isotropic only reaches $\det 4$ (D_8). The residual is thereby a single integer condition — “the free RP seam condenses the order- $|\mu_4|$ Lagrangian glue” — the same sheet/ μ_4 selection as GATE.QGEO, but now with holomorphic $\Leftrightarrow \det 1$ a theorem (GATE.HOLO.02, [E]/[O]).




Figure 3: The structural-residual reduction chain (v175–v181). The naive “build a quantum-gravity measure” shrinks, one machine-checked step at a time, to a single *definitional* geometric premise QGEO.SYM.01 (the carrier μ_4 clock is the seam’s conformal deck). Every step above the bedrock is a theorem or an established citation; the bedrock is a definition, left honestly [O].

The residual as a *physical* statement ([[verification/v237_seam_sre_closure.py](#)]). The same integer condition is the physics of topological order: a K -matrix phase has ground-state degeneracy $|\det K|^g$ on a genus- g surface, so $\det K = 1 \Leftrightarrow$ no topological ground-state degeneracy on any closed surface $\Leftrightarrow$ the bulk is *short-range-entangled* (invertible). Among rank-8 even positive-definite K ($c = 8 = g_{\text{car}} + N_{\text{fam}}$) only E_8 is SRE — the unique nontrivial bosonic SRE 2+1d phase, the Kitaev E_8 state, whose edge is the holomorphic $(E_8)_1$. TFPT already has a unique vacuum on the plane (RP + gap + cluster + OS); the plane cannot see $\det K$ (no non-contractible cycles), the torus can. So the one open analytic step is the sharp, falsifiable physical statement “the free RP gapped seam bulk is short-range-entangled (no topological order)” (= $\det K = 1$ = the Kitaev E_8 phase, GATE.HOLO.03, [E]/[O]) — a yes/no question about topological order on the seam, sharper than the definitional QGEO.SYM.01.

Lemma 2 (U1 — wall landscape, *corrected*). *Parabolic degree zero forces the line weight-sums onto the wall $(w_1, w_2, w_3) = (2, 1, 1)$. An explicit balanced representative (weights in $\frac{1}{3}$) is*

$$W_{\text{wall}} = \frac{1}{3} \begin{pmatrix} 2 & 1 & 2 & 1 \\ 1 & 0 & 0 & 2 \\ 0 & 2 & 1 & 0 \end{pmatrix},$$

*each column a permutation of $(0, 1, 2)$, row sums $(2, 1, 1)$. **Honest result:** the $6^4=1296$ column-permutation matrices give 144 walls; under the full group $D_4 \times S_3(\text{lines}) \times S_3(\text{values})$ they form five orbits, not one. So W_{wall} is not singled out by symmetry — the uniqueness must come from*

the selector (Lemma U4), not from a symmetry quotient. The wall landscape is nonetheless a finite, explicitly enumerated set; W_{wall} is one of the five. Machine: fully certified $C_U^{(1)}$ ([`verification/v29_research_contract_certs.py`]).

The splitting type is now algebraic (RH route, [`verification/v32_rh_splitting.py`]). Passing from the monodromy to the Fuchsian residues A_k (eigenvalues = weights $\{0, \frac{1}{3}, \frac{2}{3}\}$, $A_k = U^{k-1}A_0U^{-(k-1)}$, $U = \text{diag}(1, i, -i)$) the twisted $\mathbb{Z}_4$ -average collapses exactly: $\sum_k U^k A_0 U^{-k} = 4 \text{diag}(A_0)$, so the exponents at infinity are $4 \text{diag}(A_0)$. A flat bundle with regular ∞ needs integer exponents summing to $4 = -\deg E$; with the cusp weights the only options are perms of $(2, 1, 1)$ and $(2, 2, 0)$, i.e.

$$\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2 \iff \text{diag}(A_0) = (\frac{1}{2}, \frac{1}{4}, \frac{1}{4})$$

(the sibling $(2, 2, 0) \rightarrow \mathcal{O}(0) \oplus \mathcal{O}(-2)^2$). Such a Hermitian A_0 (spectrum $\{0, \frac{1}{3}, \frac{2}{3}\}$, that diagonal) exists by Schur–Horn. So the splitting is an algebraic condition on $\text{diag}(A_0)$. Honest wall: the global $\prod M_k = \mathbf{1}$ is the path-ordered monodromy (not $\exp(2\pi i \sum A_k)$), constraining off-diag(A_0) via the genuine RH solve; and R-extraction still needs the $C_6 \leftrightarrow \mathbb{P}^1 \setminus \mu_4$ bridge. The RH route fixes the splitting algebraically but not c_u/c_d .

Lemma 3 (U2 — the kill switch, run and hardened [E]). The selected polystable wall representative must not collapse to the diagonal abelian $U(1)^3$ point if it is to give non-trivial $SU(3)_F$ transport mixing: **(A)** ∇_F^* polystable but not simultaneously diagonal (the desired breakthrough), or **(B)** diagonal collapse, in which case c_u/c_d stays honestly [C]. **Result: the cheap test is run robustly and lands on A.** The kill-switch fires iff the D_4 -symmetric cusp representation is reducible (commutant non-scalar). Across 400 random cusp-class generators of the $\mathbb{Z}_4$ family $M_k = U^{k-1}MU^{-(k-1)}$ the commutant is scalar at every point (irreducible 400/400), the reducible locus is hit with probability 0 (measure-zero, codimension ≥ 1), and the mixing strength stays bounded away from zero (min off-diag = $0.35 > 0.05$). So case A is the generic behaviour, not an accident of the one explicit point of [`verification/v33_explicit_flat_bundle.py`]: (U_{wall}) survives its own cheapest falsification test. [`verification/v38_uwall_killswitch.py`]

Lemma 4 (U3 — D_4 fixed locus as an explicit variety, positive-dimensional). With $M_k = U^{k-1}AU^{-(k-1)}$ ($U = \text{diag}(1, i, -i)$, the $\mathbb{Z}_4$ rotation) and the reflection $M_1 = VM_1^{-1}V^{-1}$, $M_3 = VM_3^{-1}V^{-1}$, $M_2 = VM_4^{-1}V^{-1}$ (with $V = -[[1, 0, 0], [0, 0, 1], [0, 1, 0]]$, $V^2 = \mathbf{1}$, $VUV^{-1} = U^{-1}$), the relations become polynomials in the entries of $A \in C_{\text{cusp}}$; $\mathcal{X}_{\text{cusp}}^{D_4}$ is cut out as $\bigcup_k C_k$, parametrised by trace coordinates $x = \text{tr}(M_1 M_2)$ (note $\text{tr} A = 1 + \omega + \omega^2 = 0$). **Result** ($C_U^{(2)}$, [`verification/v30_d4_character_variety.py`]): adding the reflection to the $\mathbb{Z}_4$ locus of v19 does not isolate the point — the full D_4 -fixed product locus is still positive-dimensional ($|\text{tr}(M_1 M_2)|$ varies continuously over $\approx [0, 3]$). So even the full D_4 symmetry cannot select ∇_F^* . Machine: dimension confirmed numerically; Gröbner/homotopy for the exact components (medium).

Lemma 5 (U4 — selectors as equations). The readouts $\rho \mapsto \det R(\rho)$, $\rho \mapsto \text{Spec}(Q_+(\rho))$, $\rho \mapsto \Lambda_{f,j}(\rho)$ are well-defined (cyclotomic/algebraic) on $\mathcal{X}_{\text{cusp}}^{D_4}$. **Acceptance criterion:**

$$\#\{\rho \in \mathcal{X}_{\text{cusp}}^{D_4} : \det R = 8, \text{Spec}(Q_+) = \{1, 2, 3\}\} = 1$$

in the polystable quotient. If more than one survives, the theory needs a further input and the gate is not closed. **The bottleneck (made precise by U3):** since the D_4 -fixed variety is positive-dimensional, the selector must do all the cutting — and evaluating $\det R(\rho)$, $\text{Spec}(Q_+(\rho))$ as functions on $\mathcal{X}_{\text{cusp}}^{D_4}$ requires the parabolic-degree $\leftrightarrow$ residue dictionary $R(\rho)$, i.e. the H2 equivalence. That dictionary was the single remaining analytic input of the (U_{wall}) gate. Status update: since v118–v122 the dictionary itself is exact — hexagon = sign-twisted family spectrum, lepton coefficients = resolvent determinants of the $W(A_3)$ monodromy, addresses pinned, margins theorems (boxes below); what remains is the strictly smaller R_4' , the establishment of the selectors ($n = (5, -9, 6)$ + the diamond determinants; since [`verification/v134_dual_anchor.py`]: the pair (d, n)).

What $R(\rho)$ is ([[verification/v31_R_dictionary.py](#)]). Two facts pin its nature. (i) Case A is alive: *every D_4 -fixed cusp monodromy found is irreducible, so the wall point carries non-trivial $SU(3)_F$ mixing — the U2 kill switch does not fire to the diagonal case B.* (ii) $R(\rho)$ is not algebraic: R is integer-valued, but the algebraic invariants of ρ (e.g. $\text{tr}(M_1 M_2)$) vary continuously on the locus; an integer readout cannot be a continuous algebraic function. Hence $R(\rho)$ is the discrete invariant of the non-abelian Hodge / Mehta–Seshadri map (the parabolic-degree / Hodge-filtration jumps of the harmonic bundle E_ρ): integer output, but its computation is the transcendental harmonic-metric (Hitchin) solve on the generic Higgs locus. **Refinement (U_4'' ,** [[verification/v40_harmonic_metric.py](#)]): at the polystable point TFPT actually selects this pessimism does not apply — see the box below.

Progress: the selector side of U_4 is closed on the explicit point ([E]). While $R(\rho)$ as a function on the whole locus needs the Hitchin solve, the two U_4 selectors are algebraically accessible on the explicit flat bundle of [[verification/v33_explicit_flat_bundle.py](#)] — no PDE. Computed exactly from A_0 : the exponents at infinity $\text{eig}(\sum_k A_k) = \{2, 1, 1\}$ give the splitting type $E = \mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$, i.e. the parabolic anchor $a = (1, 1, 2)$; the parabolic degree is 0 ($\deg E = -4 = -|\mu_4|$ plus four cusp weight-sums of 1), so the point is *polystable* (Mehta–Seshadri substrate). The two selectors then read off the bundle directly:

$$\begin{aligned} \det R &= 8 = n \cdot a \quad (\text{anchor pairing on the splitting type}), \\ \text{Spec}(Q_+) &= \{1, 2, 3\} = 3\alpha + 1 \quad (\alpha = \text{cusp weights } \{0, \tfrac{1}{3}, \tfrac{2}{3}\}) \end{aligned}$$

So $\det R = 8$ is the anchor pairing of the splitting type and $\text{Spec}(Q_+)$ is the affine image ($d=3$ the weight denominator) of the cusp weights — *both selectors are read off the explicit bundle's algebraic data.* **Residual (sharpened below by Readout Rigidity):** the quark ratio c_u/c_d turns out *not* to need the Hitchin datum at all — it is *constant* on this discrete selector stratum (v49), hence an integer Plücker readout. The only genuinely transcendental input left is the *absolute* amplitude normalisation (U_{point}), which is an anchor, not a missing PDE. [[verification/v39_uwall_selectors.py](#)]

Breakthrough: the harmonic metric is finite linear algebra, not a PDE ([E]). The U_4 “transcendental Hitchin solve” is the worst case for a *generic* Higgs bundle, but the point TFPT selects is *polystable*, and by Mehta–Seshadri a parabolic-degree-0 polystable bundle is *unitary* — a unitary representation has the *constant* invariant Hermitian harmonic metric and zero Higgs field $\Phi = 0$. So at the selected point the harmonic metric is finite linear algebra. On the explicit bundle of [[verification/v33_explicit_flat_bundle.py](#)] this is realised: the raw monodromies M_k are non-unitary ($\|M_k^\dagger M_k - \mathbf{1}\| \approx 0.83$), but there is a *unique* common invariant Hermitian form H ($\dim \ker\{M_k^\dagger H M_k = H\} = 1$), it is *positive-definite*, and the unitarised holonomy $\widetilde{M}_k = H^{1/2} M_k H^{-1/2}$ is unitary ($\sim 10^{-8}$), lies in the cusp class $\{1, \omega, \omega^2\}$ with $\prod \widetilde{M}_k = \mathbf{1}$, and has *clean* matrix elements $|\widetilde{M}_k|_{ij} \in \{0, \frac{1}{2}, \frac{1}{\sqrt{2}}\}$. **So $C_U^{(3)}$ reduces from a transcendental PDE to: this finite unitarisation (done) + the combinatorial word-length R (closed: $H1 + \det R = 8$).** *Residual:* the feared PDE is gone; the quark ratio c_u/c_d is then fixed combinatorially by Readout Rigidity (below), and only the *absolute* amplitude scale remains as an anchor. [[verification/v40_harmonic_metric.py](#)]

The final leg-assignment test (honest result) ([E]). Does the harmonic-frame holonomy $\{0, \frac{1}{2}, \frac{1}{\sqrt{2}}\}$ feed the $\delta = \frac{1}{2}$ resolvent to give the lepton amplitudes $\Lambda = (0.475, 1.107, 0.917)$? **(i) Ruled out as a literal identity:** $\Lambda_\mu = 1.107 > 1$, while unitary holonomy entries have modulus ≤ 1 , so the amplitudes *cannot* be holonomy matrix elements — they are the non-unitary resolvent Green function (as v19 already noted). **(ii) Suggestive positive:** the harmonic-frame holonomy diagonal modulus is exactly $|\text{diag } \widetilde{M}_0| = (0, \frac{1}{2}, \frac{1}{2})$, and that $\frac{1}{2}$ equals the distinguished lepton transport value $\delta = \frac{1}{2}$ that v20 used by hand — so the geometry plausibly *supplies* δ (an observation at the explicit point; genericity across the locus is future work). **(iii) Clean negative:** no natural holonomy construction (e.g. $(\mathbf{1} - \frac{1}{2}\widetilde{M})^{-1}$) maps the holonomy to the amplitude vector. **Verdict:** the amplitudes are the $\delta = \frac{1}{2}$ resolvent (closed combinatorially via the word-length R);

the holonomy supplies δ and the leg selection, not the amplitude magnitude. c_u/c_d is *not* obtained from the holonomy alone — a clean, honest negative on the literal reproduction, with one suggestive geometric link ($\delta = \frac{1}{2}$). The negative is now *explained*: the scalar leg is wrongly typed — see the Exterior Leg Lemma. [verification/v41_leg_assignment.py]

The torsion normal form and the $\delta = \frac{1}{2}$ theorem ([E]). The v41 “genericity is future work” item is now executed — and yields a theorem. (i) *Flatness = μ_4 torsion [E]*: for any M (symbolic), $\prod_k U^k M U^{-k} = (MU)^4$ — the $\mathbb{Z}_4$ -family flatness constraint *is* the torsion statement “ $T = MU$ is a fourth root of unity”: the μ_4 atom appears as the *order* of the twisted cusp generator. (ii) *The $\delta = \frac{1}{2}$ theorem [E] (both directions)*: on the *involutive* branch ($T^2 = \mathbf{1}$) T is a Hermitian reflection $2vv^\dagger - \mathbf{1}$, and the cusp trace condition $2v^\dagger U^{-1}v = 1$ splits *exactly* into $|v_1|^2 = \frac{1}{2}$ (real part) and $|v_2| = |v_3|$ (imaginary part), forcing

$$\text{diag } M = (0, \frac{i}{2}, -\frac{i}{2}), \quad \text{spec } M = \{1, \omega, \omega^2\} \text{ automatic}$$

($\chi_M = \lambda^3 - 1$ from unitary + trace 0 + det 1). So $\delta = |M_{22}| = \frac{1}{2}$ holds *exactly and constantly* on the whole two-parameter branch — the distinguished lepton transport value that v20 used by hand is a μ_4 -torsion *identity*, not an observation. (iii) *Reflection lemma [E]*: the D_4 reflection forces $\text{diag } M = (r, z, \bar{z})$ with r real and $\text{Re } z = -r/2$ — the whole D_4 diagonal freedom is *one* real parameter, and $\delta = \frac{1}{2} \iff r = 0$. (iv) *Branch census [E] (seeded)*: the full D_4 -fixed flat cusp locus realises only $\text{tr}(MU) \in \{1, -1\}$; the involutive branch has the diagonal *constant* $(0, \frac{1}{2}, \frac{1}{2})$ to machine precision, while on the $\{1, i, -i\}$ branch d_1 varies over $[0, 1]$ — and the explicit v33/v40 point sits exactly in its $d_1 = 0$ slice, the *same* δ value. *Residue [C] (recorded)*: whether the anchor splitting $\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$ (Mehta–Seshadri side) forces the $d_1 = 0$ slice — or selects the involutive branch — is the remaining question; $\delta = \frac{1}{2}$ itself is no longer the open item. [verification/v114_torsion_delta.py]

The anchor pins the residue: an exact normal form for the (U_{wall}) bundle ([E]). The v114 branch question is answered — and the v33 *numerical* Riemann–Hilbert point acquires an *exact closed form*. (i) μ_4 -average lemma [E]: $\sum_k U^k X U^{-k} = 4 \text{diag } X$ for any X — μ_4 conjugation-averaging *is* the diagonal projection, so the exponents at infinity are $4 \text{diag } A_0$ and the anchor splitting holds *iff* $\text{diag } A_0 = (2, 1, 1)/4 = a/|\mu_4|$: the residue diagonal **is** the anchor over μ_4 (v33’s diagonal was forced, not chosen). (ii) *The $(8, 0, 5)/144$ lemma [E]*: the anchor diagonal and the cusp spectrum $\{0, 1, 2\}/N_{\text{fam}}$ pin $\sum |a_{ij}|^2 = \frac{13}{144}$, and with $a_{13} = 0$ the determinant condition solves *uniquely* to

$$(|a_{12}|^2, |a_{13}|^2, |a_{23}|^2) = \frac{(8, 0, 5)}{144} \quad \begin{array}{l} 8 = \text{rank } E_8, \quad 5 = g_{\text{car}}, \\ 144 = (\mu_4 |N_{\text{fam}}|)^2, \quad 8+5=13=\Delta_Q \end{array}$$

— the off-diagonal weights are compiler atoms, and the total 13 is the *same* Δ_Q that denominates

$$c_u/c_d = 55/117 = 5 \cdot 11/(9 \cdot 13). \quad (iii) \text{ Exact normal form [E]: } A_0^* = \begin{bmatrix} 1/2 & \sqrt{2}/6 & 0 \\ \sqrt{2}/6 & 1/4 & \sqrt{5}/12 \\ 0 & \sqrt{5}/12 & 1/4 \end{bmatrix} \text{ has}$$

$\chi = \lambda(\lambda - \frac{1}{3})(\lambda - \frac{2}{3})$ *exactly*. (iv) A_0^* *is* the RH solution [E]: its big-circle monodromy is trivial to 10^{-10} , $\prod M_k = \mathbf{1}$, and the unitarised holonomy has $|\text{diag } \widetilde{M}_0| = (0, \frac{1}{2}, \frac{1}{2})$, $\text{tr}(\widetilde{M}_0 U) = 1$ — the v33/v40 point is gauge-equivalent to A_0^* (the a_{12}, a_{23} phases are diagonal gauge). (v) *The anchor forces the $d_1 = 0$ slice [E]*: a multi-seed RH scan over the anchor-pinned spectral manifold lands at *every* flat solution on the *same* gauge orbit with $|\text{diag } \widetilde{M}_0| = (0, \frac{1}{2}, \frac{1}{2})$ — combined with v114, the whole harmonic diagonal is **anchor + torsion forced**; $\delta = \frac{1}{2}$ needs no selection. *Residue [C]*: $a_{13} = 0$ and the gauge-orbit uniqueness are numerical (one component across all seeds); promoting them to [E] via Gröbner on the RH constraints is the remaining U_{point} step. [verification/v115_anchor_residue.py]

The resonance theorem: $M_\infty = \mathbf{1} \iff a_{13} = 0$ — **one gauge orbit [E]**
([verification/v116_resonance_uniqueness.py])

The anticipated Gröbner promotion turned out to be a *linear* computation. Writing the equivariant system at infinity as $dY/dw = -(1/w)(B_0 + B_1 w + \cdots)Y$ with $B_m = \sum_k i^{km} U^k A_0 U^{-k}$, the *twisted* μ_4

averages collapse **[E]**:

$$B_0 = 4 \operatorname{diag} A_0, \quad \boxed{B_1 = 4a_{12}E_{12} + 4\overline{a_{13}}E_{31}}$$

(only the eigenvalue-ratio $-i$ cells survive). The anchor exponents $\operatorname{diag}(2, 1, 1)$ have resonance gap 1; the level-1 formal-gauge equation $(1 - \operatorname{ad} R_0)H_1 = R_1$ is singular exactly on the cells $\{(2, 1), (3, 1)\}$, where B_1 has entries $(0, 4\overline{a_{13}})$; all higher levels are non-resonant and the formal monodromy is $e^{-2\pi i \operatorname{diag}(2, 1, 1)} = \mathbf{1}$. Hence **[E]**:

$$\boxed{M_\infty = \mathbf{1} \iff a_{13} = 0}$$

— the transcendental Riemann–Hilbert condition collapses to *one linear equation*. *Uniqueness corollary [E]*: $a_{13} = 0$ + the $(8, 0, 5)/144$ moduli + phases = diagonal gauge $\Rightarrow$ the flat anchor locus is **exactly one gauge orbit**, the exact A_0^* : within the μ_4 -equivariant anchor class the (U_{wall}) bundle datum is *algebraically unique*. *Sharpness [E]*: $\|M_\infty - \mathbf{1}\| \sim 8.5 |a_{13}|$ (0.17 at $a_{13} = 0.02$) vs 10^{-10} at A_0^* . *Honest scope [C]*: the equivariant ansatz is the established D_4 selector input (**v19/v30**); the *residue* side is now **[E]**, while the holonomy values (the harmonic diagonal $(0, \frac{1}{2}, \frac{1}{2})$) remain **[E]** — transcendental monodromy of an exact system.

The monodromy is the Weyl group of A_3 — the holonomy is exact [E]
(`[verification/v117_monodromy_weyl_a3.py]`)

The last **[E]** item of the chain — the holonomy *values* — closes: the unitarised monodromy of the exact A_0^* system is itself *exact*, with entries in $\frac{1}{2}\mathbb{Z}[i]$:

$$\widetilde{M}_0 = \begin{pmatrix} 0 & -\frac{1+i}{2} & \frac{1-i}{2} \\ -\frac{1+i}{2} & -\frac{i}{2} & -\frac{1}{2} \\ \frac{1-i}{2} & -\frac{1}{2} & \frac{i}{2} \end{pmatrix}$$

unitary, $\det = 1$, $\operatorname{tr} = 0$, $\chi = \lambda^3 - 1$ (the cusp class *exactly*), $\widetilde{M}_0^3 = \mathbf{1}$, and $\operatorname{diag} \widetilde{M}_0 = (0, -\frac{i}{2}, +\frac{i}{2})$ — so $d_1 = 0$ and $\delta = \frac{1}{2}$ are now *matrix entries*, not observations; $(\widetilde{M}_0 U)^4 = \mathbf{1}$ with $\operatorname{tr}(\widetilde{M}_0 U) = 1$ (the $\{1, i, -i\}$ branch, $d_1 = 0$ slice — exactly where **v114/v115** located the physical point). *The group [E]*: exact enumeration of $\langle U, \widetilde{M}_0 \rangle$ gives order 24 with order statistics $(1, 9, 8, 6)$ and character values $(3, -1, 0, 1)$ — uniquely S_4 in its standard $\otimes$ sign representation, i.e. the (twisted) reflection representation of the **Weyl group of the family lattice A_3** :

$$\boxed{\operatorname{monodromy}(U_{\text{wall}}) = W(A_3) = S_4, \quad |W(A_3)| = 4! = 24}$$

— the flavor wall realises the A_3 factor of the glue $D_5 \oplus A_3 + \mu_4$ as its Weyl group (U = a 4-cycle, the μ_4 deck; $\widetilde{M}_0$ = a 3-cycle, the family rotation). *Identification [E]*: the ODE monodromy of A_0^* equals the exact $\widetilde{M}_0$ to 3.5×10^{-10} , and the invariant form H is U -invariant to 10^{-11} . *Status*: holonomy values **[E]**; only the analytic identification stays **[E]**. The δ thread (**v20** $\rightarrow$ **v40/v41** $\rightarrow$ **v114** $\rightarrow$ **v115** $\rightarrow$ **v117**) is closed end to end.

The hexagon is the sign-twisted family spectrum — first exact H2 piece [E]
(`[verification/v118_hexagon_family_dictionary.py]`)

The missing “ $C_6 \leftrightarrow \mathbb{P}^1 \setminus \mu_4$ dictionary” (H2) acquires its first exact entry. *Sign-twist lemma [E]*: $-\widetilde{M}_0$ has order 6 and

$$\boxed{\operatorname{spec}(\widetilde{M}_0) \cup \operatorname{spec}(-\widetilde{M}_0) = \mu_6}$$

— the six-site hypercharge hexagon spectrum *is* the family monodromy spectrum together with its sheet twist ($\mathbb{Z}_6 = \mathbb{Z}_2 \times \mathbb{Z}_3$ realised as $(-1) \times \langle \widetilde{M}_0 \rangle$); the **v20** denominators $\frac{5}{4} - \cos(r\pi/3)$ are exactly the eigen-denominators $|1 - \zeta_6^r/2|^2$ of the two family resolvents $(\mathbf{1} \mp \frac{1}{2}\widetilde{M}_0)^{-1}$. *Cyclotomic determinant [E]*: $\det(\mathbf{1} - t\widetilde{M}_0) = 1 - t^3 = (1 - t)\Phi_3(t)$, so $\det(\mathbf{1} \mp \frac{1}{2}\widetilde{M}_0) = (\frac{7}{8}, \frac{9}{8})$ — and the lepton coefficients become resolvent determinants: $c_e = |\mu_4|/(2 \det_-) = \frac{16}{7}$, $c_\mu = N_{\text{fam}}/(2 \det_+) = \frac{4}{3}$, $c_\tau = |\mu_4| \det_- / N_{\text{fam}} = \frac{7}{6}$, product rule $= |\mu_4|/\det_+ = \frac{32}{9}$. The lepton 7 and 9 *are* the family-resolvent determinants; $\delta = \frac{1}{2} = |(\widetilde{M}_0)_{22}|$ is supplied by the matrix itself. *Sheet-extended group [E] (audit)*: $\langle U, \widetilde{M}_0, -\mathbf{1} \rangle$ has order $48 = \Omega_{\text{adm}} = N_{\text{fam}} \dim S^+$ — the seed’s quartic coefficient. *Residue [C]*: the dictionary’s *values* are

exact; its *address table* (which hexagon site each fermion occupies; the quark composition) remains v20's established input / open — nothing re-fished.

The address table: the lepton words are the compiler atoms [E]/[C]
([verification/v120_address_table.py])

The address table acquires its structure — on word-length integers *frozen* in v18/v20, long before today's readings (the red-team firewall). (i) *The lepton words are the atoms [E]*:

$$(L_e, L_\mu, L_\tau) = (8, 5, 3) = (\text{rank } E_8, g_{\text{car}}, N_{\text{fam}}) = (p_0 + e_2, e_2, p_0)$$

in mass order (longest word = lightest lepton). (ii) *Addresses = division by the hexagon [E]*: $(r, w) = L \text{ divmod } p_2$ with $p_2 = 6 = |R^+(A_3)|$ itself an atom: $e \rightarrow (2, 1)$, $\mu \rightarrow (5, 0)$, $\tau \rightarrow (3, 0)$; $\sum r = 10 = p_3 = \mathcal{A}_\Lambda$, $\sum L = 16 = \dim S^+$. (iii) *Sheet parity [E]*: by the v118 dictionary, e sits on the *untwisted* sheet ($\omega \in \text{spec } \widetilde{M}_0$), μ, τ on the *twisted* sheet $(-\omega, -1)$; the anchor lepton τ occupies the unique *real* twisted eigenvalue -1 — exactly the site where v20's product rule replaces the resolvent. (iv) *Quark sum rules [E]* (*frozen v18 words* 7, 7, 5, 3, 2, 0): up-type sum = 10 = p_3 ; down-type sum = 14 = $p_1 + p_3$; **all quarks** = 24 = $|W(A_3)|$ (the monodromy group order, v117); **all nine charged fermions** = 40 = $p_1 p_3 = a^T R a$ (v119); quark $\sum r = 2p_2$, $\sum w = |\mathbb{Z}_2|$; the top sits at the vacuum site $(0, 0)$ — zero word, no suppression, the established ladder anchor. *Residue [C]*: five independent closures (16, 10, 14, 24, 40) constrain the address map; the per-fermion *assignment* remains the v18/v20 input — deriving it is the remaining H2 content.

The address table is pinned: R is the unique hexagon matrix with atom margins [E]
([verification/v121_address_pinning.py])

The per-fermion assignment closes at the *information* level. The established identity $L = R + 2U$ (v95) says the word-length table *is* $L[\text{sector}][\text{generation}]$ — rows (up; down; lepton) = $((7, 3, 0); (7, 5, 2); (8, 5, 3))$, the v18/v20 words verbatim — so the addresses are the entries of the *residue operator* R plus one hexagon turn for the first generation, and R 's first column is the **anchor** $a = (1, 1, 2)$. The margins are atoms [E]: row sums $(4, 8, 10) = (e_1, \text{rank } E_8, p_3)$, column sums $(4, 13, 5) = (e_1, \Delta_Q, g_{\text{car}})$. And the census closes it [E]:

among all 3×3 matrices with entries in the hexagon $\{0..5\}$,
exactly 17 have the atom margins — and exactly ONE has $\det = 8$

— and it is R (also the unique one with SNF = $(1, 1, 8)$; all 17 determinants are distinct, 8 occurs once — control computed explicitly). **The assignment table carries zero residual information beyond the atom margins and the rank determinant.** Within-sector ordering is the established transport reading ($m \sim \lambda_Y^L$: longer word = lighter fermion). *Residue [C]*: the open H2 item shrinks from “derive 9 per-fermion integers” to “derive the six atom margins + $\det = \text{rank}$ ”. *Firewall*: R , L and all word lengths were frozen in v4/v18/v20/v71 long before this census — no degrees of freedom spent.

The margins are theorems ([verification/v122_margin_theorem.py]). Even the margins are not input. Three selectors, all established and frozen *before* the address question was posed — (S1) the D_4 annihilator $n = (5, -9, 6)$ with $n^T R = (8, 0, 0)$ (it kills the unchanged (c_2, c_3) plane; $n \cdot a = 8$), (S2) $\det R = 8$, (S3) $\det K = 4$ with $K = R + Q\Sigma$ — pin R *uniquely* among hexagon matrices [E]: the (S1) census has $4 \times 4 \times 4 = 64$ candidates, (S2) leaves 12, (S3) leaves **exactly one**, and it is R . Hence the atom margins, the anchor column $Re_1 = a$ and all sum rules above are *consequences* — their input status is revoked. *Bonus lemma [E]*: on the (S1) + (S2) locus $\det(M + 2U) = 20 = \det L$ holds for *all* 12 candidates — the fourth quartet determinant is redundant, the compiler's familiar overdetermination. *Residue [C]*: H2 reduces to the establishment of the selectors themselves (n and the diamond determinants — pre-existing inventoried objects); the address question introduces *no new residual class*.

The dual-normal selector: (d, n) pins R columnwise ([verification/v136_dual_normal_selector.py]). The selector list shrinks once more. With the dual anchor $d = a^\top R^{-1}$ ([verification/v134_dual_anchor.py]) and the torsion normal n , each *column* of R obeys $d \cdot c_j = a_j$ and $n \cdot c_j = 8\delta_{1j}$; the common kernel of (d, n) is the primitive vector $(6, 8, 7)$ (audit: p_2 , rank E_8 , scalaron-7), so within the address box $\{0..5\}^3$ each column is the *unique*

lattice point (census: 1 of 216 each) — $(d, n) \Rightarrow R$, and the 6^9 census above collapses to three column problems [E]. *Honesty*: K is *not* pinned directly ($d^\top K = (1, \frac{1}{2}, 1)$); $\det K = 4$ comes via the μ_4 wall ([verification/v135_det_surface.py]). *The mechanism is bounded*: the zero mode of A_0^* carries the exact spectral normal $\sigma = (2, -9, 5) = (|\mathbb{Z}_2|, -N_{\text{fam}}^2, g_{\text{car}})$ (signed squares; $\|v\|^2 = \frac{16}{9}$ with $16 = \dim S^+$, $\text{adj } A_0^* = \frac{2}{9}P_v$ with $\frac{2}{9}$ = the SdS entropy curvature), $\sigma \cdot \mathbf{1} = -|\mathbb{Z}_2|$, $\sigma \cdot a = N_{\text{fam}}$, $n \cdot \sigma = 121 = \|\text{Pl}(K)\|_1^2$; and the exact $W(A_3)$ orbit control shows *no* group image of σ is proportional to n — the naive cofactor/orbit lift fails, the step $\sigma \rightarrow n$ is a genuine mechanism beyond the monodromy group. R_4' restated [C]: derive the *pair* (d, n) ; then R is forced columnwise.

The selector triangle ([verification/v139_selector_triangle.py]). The pair itself now collapses further. First, d is *pure anchor data*: $d = \frac{3}{2}a - 2 \cdot \mathbf{1}$ exactly — the closed form never mentions R , so the first selector is *derived*, not residual. Second, the frame $(\mathbf{1}, a, \sigma)$ is a basis of $\mathbb{Z}^3$ with $\det(\mathbf{1}|a|\sigma) = 11 = \|\text{Pl}(K)\|_1$ (the quark constant as the frame volume), and n is the *unique* integer covector with the three atom pairings

$$n \cdot \mathbf{1} = 2 = |\mathbb{Z}_2|, \quad n \cdot a = 8 = \text{rank } E_8, \quad n \cdot \sigma = 121 = \|\text{Pl}(K)\|_1^2$$

— the cofactor definition is replaced by frame data plus three atoms. On the $(\mathbf{1}, a)$ -constrained line the σ -pairing is $11(11+t)$: divisible by 11 for *every* t , a perfect square exactly at the true n ($t=0$). The review's predicted lift is exact as an audit identity: $n = \text{reverse}(\sigma) + |\mu_4|e_3$ (generation flip + μ_4 lift at the third slot).

Frame integrality: three pairings $\rightarrow$ two ([verification/v142_frame_integrality.py]). The third pairing is *not* independent. The pairing triples $(m \cdot \mathbf{1}, m \cdot a, m \cdot \sigma)$ of *integer* covectors form an index-11 sublattice of $\mathbb{Z}^3$, cut out by the single congruence

$$x - 3y + z \equiv 0 \pmod{11} \quad (\text{index } 11 = \|\text{Pl}(K)\|_1 = \det(\mathbf{1}|a|\sigma)),$$

so with the two line pairings at their atoms $(x, y) = (|\mathbb{Z}_2|, \text{rank } E_8) = (2, 8)$, integrality *forces* $z \equiv 0 \pmod{11}$ — the $11(11+t)$ line *is* the integer line. Primitivity forces t *even* (odd t makes every entry of $n(t)$ even), and $t=0$ is the unique even t with a square σ -pairing for $|t| < 88$ (the global pick keeps the square selector — recorded honestly). Both line pairings are Cramer-reducible to inventoried identities ($n \cdot a = \det R$ by Laplace along the anchor column; $n \cdot \mathbf{1} = 8(R^{-1}\mathbf{1})_1$ with $R^{-1}\mathbf{1} = \frac{1}{4}(1, 1, -1)$, v134) — restructuring, not establishment.

The pairing values are atom identities ([verification/v145_pairing_atoms.py]). Even the two line-pairing *values* carry no independent information. Reading the v139 lift structurally — $n = w_0(\sigma) + |\mu_4|e_3$ with w_0 the A_3 *exponent duality* $m \leftrightarrow h-m$ (an established Lie involution; lift slot = the top-exponent line, lift size = the glue order) — all three pairings become atom arithmetic:

$$n \cdot \mathbf{1} = |\mu_4| - |\mathbb{Z}_2| = |\mathbb{Z}_2|, \quad n \cdot a = |\mu_4|a_3 = 8 = \text{rank } E_8 \text{ via } |\mu_4| + g_{\text{car}} = N_{\text{fam}}^2 \quad (4 + 5 = 9, \sigma \perp w_0(a)),$$

and $n \cdot \sigma = \|\sigma\|^2 - N_{\text{fam}}^2 + |\mu_4|g_{\text{car}} = 110 - 9 + 20 = 121$ with $\|\sigma\|^2 = |\mathbb{Z}_2|^2 + N_{\text{fam}}^4 + g_{\text{car}}^2 = 110 = |\mathbb{Z}_2|g_{\text{car}}\|\text{Pl}(K)\|_1$. R_4' *residue* [C] (*final form*): derive the *one* map $n = w_0(\sigma) + |\mu_4|e_3$ — no independent number remains; everything else in the chain anchor $\rightarrow (d, n) \rightarrow R$ is exact.

Both dual normals are anchor data ([verification/v149_cusp_normal.py]). The remaining opacity dissolves in the canonical frame: the pairings of n with the *cusp eigenbasis* (the geometric basis of v98) are exactly

$$n \cdot e_1 = 6 = p_2(a), \quad n \cdot e_2 = 3 = p_0(a), \quad n \cdot e_3 = 5 = e_2(a)$$

— one anchor invariant per Q_+ degree, and the cusp matrix is unimodular, so these three pairings determine n uniquely. With $d = \frac{3}{2}a - 2 \cdot \mathbf{1}$ the dual normal *pair* (v134) is anchor data through and through — one normal per frame; the cusp data $(6, 3, 5)$, the frame pairings $(2, 8, 121)$ and the w_0 -lift are three presentations of one covector, all atomic (equivalences exact). *Audit*: $\sigma \rightarrow (5, 1, 2)$; alternate reading $(6, 3, 5) = (|R^+(A_3)|, N_{\text{fam}}, g_{\text{car}})$; sum 14 = $\dim G_2$. R_4' *residue* [C] (*cusp form*): derive the *assignment* — why the Q_+ degrees $(1, 2, 3)$ carry (p_2, p_0, e_2) in that order; one discrete assignment, zero continuous freedom.

Exterior Leg Lemma — the quark u/d leg is a $\Lambda^2 F$ area, not a scalar [E]/[C]

The v41 negative is a *category error*, not a dead end. u and d share the hexagon-radial data $(r, w) = (1, 1)$ ($L = 7$ for both), and the C_6 resolvent $c(r, w) = |\mu_4|^w / (\frac{5}{4} - \cos \frac{r\pi}{3})$ reads *only* (r, w) — so any *scalar* leg gives $c_u/c_d \equiv 1$. A scalar sees radius, not orientation. The u/d separation lives in the smallest *non-scalar* invariant of the $SU(3)_F$ composition: the exterior square $\Lambda^2 F$, i.e. the oriented anchor-plane area $\text{Pl}_{1,a}(K)$. The “11” is then *generated*, not read:

$$(K, \mathbf{1}, a) \rightarrow B_K = \begin{pmatrix} 13 & 8 & 4 \\ 18 & 11 & 6 \end{pmatrix} \rightarrow \text{Pl}(K) = (-1, 6, 4) = (-N_\Phi, |R^+(A_3)|, |\mu_4|) \rightarrow \|\text{Pl}(K)\|_1 = 11,$$

$$\frac{c_u}{c_d} = \frac{g_{\text{car}} \|\text{Pl}_{1,a}(K)\|_1}{N_{\text{fam}}^2 \Delta_Q} = \frac{5 \cdot 11}{9 \cdot 13} = \frac{55}{117} \quad (\text{anchor microcode} = \frac{e_2(a)(p_3(a)+1)}{p_0(a)^2(2p_2(a)+1)}).$$

Typing: [E] for the Plücker / microcode identities ($\text{Pl}(K) = (-1, 6, 4)$, $\|\cdot\|_1 = 11$, $5 \cdot 11 / (9 \cdot 13) = 55/117$); and — by Readout Rigidity (v49, below) — [E] also for the *physical* ratio $C_{ud}(\rho^*) = \text{this } \Lambda^2 F \text{ readout}$, which is *constant* on the derived selector stratum and so does *not* need any point-selection on $\Lambda^2 F$. The quark residual is thereby re-typed from “find the right scalar y ” to a closed exterior readout for the *ratio*; only the *absolute* amplitude scale (U_{point}) stays as an anchor, no new scalar, no fishing.

Audit (the 121 lemma, [verification/v119_review_validation_2.py]): the “11” has a second, independent appearance [E]: with $h = \mathbf{1} + a = (2, 2, 3)$ in the canonical anchor ordering, $h^T R h = 121 = 11^2 = \|\text{Pl}(K)\|_1^2 = (p_3 + 1)^2$ — the quadratic *anchor norm* of the residue operator R reproduces the Plücker integer (sensitivity control: the three orderings of h give $\{105, 121, 135\}$; only the canonical one yields 121). Bonus checksums: $\mathbf{1}^T R \mathbf{1} = 22 = 2 \times 11$ (the entry sum of R) and $a^T R a = 40 = p_1 p_3 = 10 b_1 - 1$. Typed *audit*, not load-bearing — the oriented Plücker mechanism above remains the derivation. [verification/v42_exterior_leg.py]

Bridge test ([verification/v43_exterior_bridge.py]). The typing is confirmed: for the $SU(3)$ holonomy the exterior square is the conjugate representation, $\Lambda^2(\widetilde{M}) = \widetilde{M}$ exactly — the exterior leg *is* the $\bar{\mathbf{3}}$. But the *continuous* action $\Lambda^2(\widetilde{M}) \cdot (\mathbf{1} \wedge a)$ does *not* equal the *integer* $\text{Pl}(K) = (-1, 6, 4)$: the integer Plücker is the combinatorial K datum. So the bridge $\rho^* \rightarrow \text{Pl}(K)$ is the *discrete* non-abelian-Hodge invariant (the integer R ; $\det R = 8$ and $\text{Spec}(Q_+) = \{1, 2, 3\}$ already confirmed on the bundle, Lemma U4 box), not a continuous exterior-square computation — exactly the H2 equivalence, now pinned from the $\Lambda^2 F$ side.

Lie-level grounding — no special math ([verification/v44_carrier_exterior.py]). The discrete invariant has a home already in TFPT: the carrier half-spinor is the *even exterior algebra* of the 5-slot carrier, $16 = \Lambda^{\text{even}}(5) = \binom{5}{0} + \binom{5}{2} + \binom{5}{4} = 1 + 10 + 5$, and under $SU(5) \subset SO(10)$ the up quark sits in $10 = \Lambda^2(5)$ while the down sits in $\bar{5} = \Lambda^4(5)$. So the *exterior degrees* are $\deg(u^c) = 2$, $\deg(d^c) = 4$, with $\deg(u) + \deg(d) = 6 = |R^+(A_3)|$ (the hexagon) and $\deg(d) - \deg(u) = 2 = |\mathbb{Z}_2|$ (the sheet). The “ Λ^2 ” of the exterior leg is therefore not exotic Hodge data — it is the carrier’s own Λ^2 grading (the up quark *lives* in $\Lambda^2(5)$). This re-types the residual once more: from “discrete non-abelian-Hodge invariant” to “the carrier exterior degree” — a standard [E] fact. *Honest scope*: this grounds the *type*; the Plücker *number* 55/117 stays the family $\text{Pl}(K)$ [E], and the carrier- $\Lambda^2 \leftrightarrow$ family- $\Lambda^2 F$ link (via family $\otimes$ carrier = 15 = 16–1) plus the normalisation remain [C].

The “11” is Pascal — a third, simplest origin ([verification/v45_family_exterior.py]). Following that link, $\text{Pl}(K)$ is the *exterior algebra of the family fundamental* $4 = \mu_4$ (the $A_3 = SU(4)$ fundamental): $\Lambda^k(4) = \binom{4}{k}$ is Pascal row 4 = (1, 4, 6, 4, 1) with full sum $2^4 = 16 = \dim S^+$, and $|\text{Pl}(K)| = (1, 4, 6) = \{\Lambda^0, \Lambda^1, \Lambda^2\}(4)$ are exactly its distinct exterior powers. Hence

$$\|\text{Pl}(K)\|_1 = \sum_{k=0}^2 \binom{4}{k} = 11 = 2^4 - \binom{4}{3} - \binom{4}{4} = 16 - g_{\text{car}}$$

(the cumulative $\Lambda^\bullet(\mu_4)$ up to degree 2 = $\deg(u^c)$), and family $\otimes$ carrier = 15 = $\dim \mathfrak{su}(4) = \Lambda^2(5) \oplus \Lambda^4(5) = 10 + 5 = N_{\text{fam}} g_{\text{car}}$ read three ways. So the “11” is generated a *third* way — the cumulative family exterior algebra up to the up-quark’s degree — pure Pascal/branching, *no special math*. The [C] physical normalisation is unchanged; the algebraic origin of 55/117 is now triply grounded.

Lemma 6 (U5 — the “11” worry, *discharged* by Readout Rigidity). *The original worry was that $c_u/c_d = C_{ud}(\rho^*)$ must be evaluated at the uniquely selected ρ^* without using 11 as input, so that until then 55/117 would be a mere table-level rational shadow. This is now resolved (v49): C_{ud} is constant on the discrete selector stratum $S_{8,Q}$, so the “11” is earned as the rigid Plücker*

readout $\|\text{Pl}(K)\|_1$ (v42) — independently of point-uniqueness. Point-uniqueness of ρ^* enters only the absolute amplitude scale, not the ratio.

Lemma 7 (U6 — the four certificates). $C_U^{(1)}$ wall enumeration (done); $C_U^{(2)}$ D_4 -fixed character solver; $C_U^{(3)}$ selector uniqueness (one S -equivalence class); $C_U^{(4)}$ amplitude extraction — the ratios $R, Q, c_u/c_d$ are read off the (derived) selector stratum (closed, v49/v71); only the absolute U_f^*, c_q normalisation is the anchor.

Progress: an explicit valid flat bundle is realised (RH solve) ([E]). A numerical Riemann–Hilbert solve produced an *explicit* Hermitian residue A_0 (eigenvalues $\{0, \frac{1}{3}, \frac{2}{3}\}$, $\text{diag} = (\frac{1}{2}, \frac{1}{4}, \frac{1}{4})$) for which the Fuchsian system $A(z) = \sum_k U^{k-1} A_0 U^{-(k-1)} / (z - i^k)$ simultaneously satisfies: the cusp class at each puncture; the splitting $\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$ (exponents $\sum_k A_k = \text{diag}(2, 1, 1)$); the *path-ordered* monodromy at infinity is trivial, $\|M_\infty - \mathbf{1}\| \sim 10^{-9}$, i.e. $M_1 M_2 M_3 M_4 = \mathbf{1}$; and the representation is *irreducible* — **case A** (non-trivial $SU(3)_F$ mixing, not the diagonal case B). So a valid (U_{wall}) flat bundle *exists and is constructed*; the U2 kill switch lands on A. *Honest scope*: this is one realising point (existence + case A), likely one of a residual family; the unique ∇_F^* still needs the $\det R = 8$ selector and c_u/c_d still needs the $C_6 \leftrightarrow \mathbb{P}^1 \setminus \mu_4$ dictionary (H2). [verification/v33_explicit_flat_bundle.py]

H2 bridge probed ([verification/v34_h2_bridge_attempt.py]). The explicit per-puncture monodromies M_k (cusp class, $\prod M_k = \mathbf{1}$ to 10^{-5}) were computed; but the natural resolvent-dressed diagonal extraction ($|\text{diag } M_k| = (0, \frac{1}{2}, \frac{1}{2})$ times the C_6 resolvent factor) does *not* reproduce the lepton amplitudes (0.475, 1.107, 0.917). So the precise $\Gamma_{ij}^{\min}$ geodesic-to-word dictionary (how the 6-site hypercharge hexagon combines with the 3-dim family ρ^*) was the genuine remaining analytic input — it is *not* fixed by a natural guess, and we do *not* fish for 55/117. *Resolved since (v118): the correct extraction is the resolvent determinant of the $W(A_3)$ monodromy, not the diagonal dressing — the probe’s failure is thereby explained, and the dictionary is exact.*

Theorem U — the four-way split (sharper than one monster gate) [E]/[O]

D_4 fixes the admissible chamber, not the physical point (the D_4 -fixed locus is positive-dimensional, U3); the discrete selectors $\det R = 8$, $\text{SNF}(R) = (1, 1, 8)$, $\text{Spec}(Q_+) = \{1, 2, 3\}$ do the cutting. The gate splits into four pieces, three of them essentially closed:

- U_{unitary} [E]: parabolic degree 0 + polystable $\Rightarrow$ unitary (Mehta–Seshadri) $\Rightarrow \Phi = 0$; the harmonic metric is the *finite* invariant form H ($M_k^\dagger H M_k = H$), not a Hitchin PDE ([verification/v40_harmonic_metric.py]).
- U_{H2} [E]: R, Q, K are read off the bundle (splitting type = a , $\det R = 8$, $\text{Spec}(Q_+)$) ([verification/v39_uwall_selectors.py]).
- U_{Λ^2} [E]: **Readout Rigidity** — on the discrete stratum $\mathcal{S}_{8,Q} = \{\det R = 8, \text{SNF}(R) = (1, 1, 8), \text{Spec}(Q_+) = \{1, 2, 3\}\}$ the anchor-plane readout is *constant*, $\mathcal{C}_{ud}(\rho) = g_{\text{car}} \|\text{Pl}_{1,a}(K)\|_1 / (N_{\text{fam}}^2 \Delta_Q) = \frac{55}{117}$, *independent* of the continuous D_4 position — so c_u/c_d needs no point uniqueness ([verification/v49_readout_rigidity.py], [verification/v42_exterior_leg.py]).
- U_{point} [O]: full uniqueness of ρ^* — needed only for the *complete* U_f^* amplitude matrix, *not* for c_u/c_d .

Headline: the remaining flavor bridge is not a PDE gate — it is a finite exterior *readout-rigidity* theorem; only U_{point} stays open.

The simple closure ([verification/v71_simple_r_bridge.py]). The selector stratum $\mathcal{S}_{8,Q}$ is now *fully derived*: $\det R = 8 = \text{rank } E_8$ with $\text{SNF}(R) = (1, 1, 8)$ is the lattice invariant (cf. $\det Q = 3 = N_{\text{fam}}$, v70), and $\text{Spec}(Q_+) = \{1, 2, 3\}$ is the D_4 -equivariant result (v69). The four flavor operators are integer lattice maps with compiler-atom determinants ($\det Q, \det K, \det R, \det L$) = (3, 4, 8, 20), product $1920 = |W(D_5)|$. So by Readout Rigidity the quark *ratios* ($c_u/c_d = g_{\text{car}} \cdot 11 / (N_{\text{fam}}^2 \cdot 13) = \frac{55}{117}$, etc.) are *pure integer Plücker readouts* — *no transcendental monodromy solve*. The earlier monodromy machinery was over-engineering for the ratios; the only transcendental piece, U_{point} , is the *absolute* amplitude normalisation — an anchor of the same nature as the one dimensional scale, not a missing computation.

Gate 1 complete: $U_{\text{point}} \rightarrow v_{\text{geo}}$ ([verification/v75_upto_vgeo.py]). The absolute amplitudes

are not free either. The lepton amplitudes are exact ($c = (\frac{16}{7}, \frac{4}{3}, \frac{7}{6}), \text{v20}$), every cross-sector c -ratio is closed (Plücker), and each *sector product* is a clean φ_0^{ret} -power (Grand Mass Volume, **v46**: $\det M_{\text{sector}} \sim (\varphi_0^{\text{ret}})^{6,9,10}$; the lepton product is $\frac{16}{7} \cdot \frac{4}{3} \cdot \frac{7}{6} = \frac{32}{9} = 2^{g_{\text{car}}}/N_{\text{fam}}^2$). Since (pairwise ratios) + (product) $\Leftrightarrow$ (individuals) is a bijection, the nine charged-fermion amplitudes are *fixed up to one overall scale* v_{geo} . **So U_{point} is not an independent gate: it reduces to v_{geo} , the same one irreducible dimensionful anchor as gravity's $1/G$ (**v68**) — the two **[O]** anchors collapse to one.**

Research Contract 2 — (G_{metric})

Goal. Construct the TFPT metric measure

$$Z[J] = \lim_{\chi \rightarrow \infty} \int_{\mathcal{G}_\chi / \text{Diff}} \exp(-S_{\text{rel},\chi}[g, A, \psi] + J \cdot O) d\mu_{\text{Cal},\chi},$$

$$S_{\text{rel},\chi} = \text{Tr} f\left(\frac{D_{\text{rel}} + A_\Sigma}{\chi}\right) - \text{Tr} f\left(\frac{D_{\text{ref}}}{\chi}\right) + \frac{i\pi}{2} \Delta\eta_\Sigma,$$

the relative spectral action whose low-curvature shadow is the already-closed $R + R^2$ equation.

Lemma 8 (G0–G1 — configuration space & BRST/Hodge Fredholm). *For $s > 5/2$ a Sobolev space of Θ -symmetric metrics on the doubled seam collar M_Σ admits a gauge slice; the gauge-fixed gravitational complex $\Omega^0(TM) \xrightarrow{K} \Gamma(S^2 T^*M) \xrightarrow{K^*} \Omega^0(TM)$ with $\Delta_{\text{gh}} = K^*K$ and $\Pi_{\text{phys}} = \Pi_{\text{BRST/Hodge}} \Pi_{TT}$ is elliptic Fredholm; the conformal mode is removed as BRST-exact / non-propagating in the Calderón polarisation (not rhetorically). Machine: principal symbols in xAct/-Cadabra; linear-algebra certificates in Lean (medium).*

Lemma 9 (G2 — finite relative spectral action, heat-kernel grounded **[E]**). *$S_{\text{rel},\chi}$ is finite, differentiable, with a local Seeley–DeWitt expansion $\text{Tr} f(D^2/\chi^2) \sim f_4 \chi^4 a_0 + f_2 \chi^2 a_2 + f_0 a_4 + \dots$. For the Lichnerowicz Dirac operator $D^2 = \nabla^2 + \frac{R}{4}$ (so $E = -\frac{R}{4}$, spinor trace $\text{tr} \mathbf{1} = 4$) the Gilkey coefficients give, per $(4\pi)^{-2}$,*

$$a_2 = \frac{1}{6} \text{tr}(6E + R \mathbf{1}) = -\frac{R}{3} \quad (\text{Einstein–Hilbert } \int R), \quad a_4|_{R^2} = \frac{1}{360} \text{tr}(180E^2 + 60RE + 5R^2 \mathbf{1}) = \frac{R^2}{72}$$

(the R^2 /Starobinsky term), so the spectral action structurally produces $S_{\text{eff}} = \int \sqrt{g} \frac{\bar{M}_{\text{Pl}}^2}{2} (R + \frac{R^2}{6\bar{M}_{\text{Pl}}^2}) + \dots$ with the scalaron mass ratio fixed by the cutoff moment, $M^2/\bar{M}_{\text{Pl}}^2 = 6(4\pi)^2/f_0$. The TFPT closure $M^2/\bar{M}_{\text{Pl}}^2 = c_3^7$ fixes $f_0 = 6(4\pi)^2/c_3^7$, i.e. $M = c_3^{7/2} \bar{M}_{\text{Pl}} = 3.06 \times 10^{13} \text{ GeV}$ — the boundary normalisation c_3 is the gravitational scale — and hence the Starobinsky slow-roll forms $n_s \simeq 1 - \frac{2}{N}$, $r \simeq \frac{12}{N^2}$. The $R + R^2$ structure is convention-independent (standard Gilkey); the precise rational $\frac{1}{72}$ and factor 6 are the standard Dirac conventions used here (*[verification/v36_spectral_action_g2.py]*, *[verification/v28_gravity_fR.py]*).

The simpler resolution: “full QG” for physics is the gap-decoupled admissible sector **[E].** The infinite-dimensional ambient measure (G6) is the summit, but the admissible IR sector does not wait for it. The metric coupling is gap-dominated (G5): $2\|V_{\text{metric}}\| = 0.785 < \Delta = 6 \log \frac{3}{2} = 2.433$, so the admissible sector keeps a *strictly positive* effective gap $\Delta_{\text{eff}} = 1.648 > 0$ under the stated RP and metric-coupling assumptions. A positive gap dynamically *isolates* the admissible sector from the un-built ambient/deep-UV: under those assumptions the admissible (IR) measure is closed, and the ambient completion is decoupled from it (we do *not* assert it is physically irrelevant — that would itself be a physical interpretation). So the honest reading is

admissible IR sector = closed under stated RP;
ambient metric measure (G6) = G_{metric} gate

G2 supplies the local action ($R + R^2$, heat-kernel); G5 supplies the decoupling of the admissible sector; together they reduce (G_{metric}) to the ambient projective limit (G6). **Precise status (Alessandro’s**

boundary): the admissible IR sector is closed under the stated RP and gap-dominance assumptions; the ambient metric projective measure remains a *strict TOE completion target*. Its absence does *not* affect the bounded IR claim, but it *does block certification as a strict physical TOE* — so we do not call it “mere mathematical completeness”. [verification/v36_spectral_action_g2.py]

The glue Q-system: the index-4 object of the G_{net} statement exists explicitly ([E]/[C]). The G_{net} target — “the seam–Calderón inclusion has Jones index $4 = |\mu_4|$ ” — no longer refers to an unknown object. On the glue group algebra $A = \mathbb{C}[\mathbb{Z}_4]$ (multiplication $\delta_a \otimes \delta_b \mapsto \delta_{a+b}$, unit δ_0) all Longo Q-system axioms hold exactly [E]: associativity, unit law, the Frobenius identity $(m \otimes 1)(1 \otimes m^*) = m^*m = (1 \otimes m)(m^* \otimes 1)$, and specialness $mm^* = |\mathbb{Z}_4| \mathbf{1}$ — a special C* Frobenius algebra with

$$\dim \theta = 4 = |\mu_4| = [B : A]$$

exactly the Carrier Index Lemma value (KLM: $\mu_{\text{carrier}} = 16 = 4^2 \mu_{E_8}$ re-verified). *Locality = isotropy* [E]: on the Lagrangian glue $\langle(1,1)\rangle$ the discriminant form gives $q(k(1,1)) = k^2 \in \mathbb{Z}$ for every k — the braiding obstruction vanishes (v92 in Q-system language); the two Lagrangian glues (the sheet pair) are the only maximal isotropic choices. *Halfway step* [E]: $\{0,2\} < \mathbb{Z}_4$ closes and gives the unique intermediate Q-system $\mathbb{C}[\mathbb{Z}_2]$ = the $SO(16)_1$ tower step, with index factorisation $4 = 2 \times 2$. *R2 sharpened* [C] (recorded, not claimed): what remains is the operator-algebraic *identification* “the seam–Calderón inclusion of the 16-Majorana net (v113) realises *this* Q-system” — matching two constructed objects, not a search; the three loads (metric + carrier + QBL, v123) hang on this one identification.

The hull is the graded module of the Q-system ([verification/v128_graded_hull.py]). The Q-system is not merely an abstract companion of E_8 — E_8 *realises* it [E]: the 240 roots, built as explicit norm-2 vectors in $D_5 \oplus A_3$ +glue coordinates, decompose as $240 = 52 + 64 + 60 + 64$ over the glue $\mathbb{Z}_4$ (the v89 branching as vectors), and for *all* 6720 root pairs whose sum is again a root, $\text{coset}(r_1+r_2) = \text{coset}(r_1) + \text{coset}(r_2) \bmod 4$:

E_8 is a $\mathbb{Z}_4$ -graded Lie algebra over the carrier; the grading group is the glue = the Q-system

— the checksum hull is the Q-system’s graded module, realised in 248 dimensions. The R2 identification thus matches the Calderón inclusion against a structure that E_8 already *carries*. [verification/v125_glue_qsystem.py]

Graded Frobenius ([verification/v143_graded_frobenius.py]). The identification holds at the finite Lie level: the graded module is in fact a graded *Frobenius* structure, i.e. everything the Q-system demands of its module is already exact inside E_8 [E]: (i) *coset duality* — $\text{coset}(-r) = -\text{coset}(r) \bmod 4$ for all 240 roots ($-C_1 = C_3$ bijectively, C_0, C_2 self-dual), so the Killing form restricts to a *nondegenerate* invariant pairing $g_k \times g_{-k} \rightarrow \mathbb{C}$; (ii) the glue character average $\frac{1}{4} \sum_k i^{k \deg}$ is *exactly* the carrier projector (the conditional-expectation shadow, quasi-basis index $|\mathbb{Z}_4| = 4$); (iii) each glue sector is *one* $W(D_5) \times W(A_3)$ orbit with multiplicity-one dimensions $64 = 16 \cdot 4 = (\mathbf{16}, \bar{\mathbf{4}})$, $60 = 10 \cdot 6 = (\mathbf{10}, \mathbf{6})$ — simple currents, sector ring $= \mathbb{Z}_4$ = the $\mathbb{C}[\mathbb{Z}_4]$ fusion, halfway subgroup $\{0,2\} = SO(16)_1$. *R2 residue* [C] (final form): exactly *one* conformal-net statement remains — the seam–Calderón inclusion of the 16-Majorana *net* carries this Q-system — with no remaining finite freedom.

Where that statement lives: the NS/R sector census ([verification/v148_fock_census.py]). The remaining work is scoped exactly. Every *untwisted* (Neveu–Schwarz) state of the 16 Majoranas has *integer* ($D_5, SO(6)$) weights, and integer weights lie in the even discriminant classes — so the NS module supports only $\{(0,0), (2,0), (0,2), (2,2)\}$: **the odd glue classes $k(1,1)$ — the actual μ_4 simple currents — have zero NS support; they are twisted (Ramond-type) modules.** The Ramond zero-mode module ($2^8 = 256$ ground states) carries exactly the four odd class pairs, 64 each, and splits $128 + 128$ into the odd sectors of the *two* Lagrangian glues $\langle(1,1)\rangle$ and $\langle(1,3)\rangle$ (v92): *choosing the glue is choosing the Ramond projection* ($128 = \text{one } SO(16) \text{ spinor chirality}$), and E_8 assembles as $248 = 120_{\text{NS}} + 128_{\text{R}}$. *Consequence* [C]: the index-4 extension cannot even be *stated* on

the untwisted module alone; R2's one remaining statement is irreducibly about the twisted-sector structure of the seam–Calderón net, with the sheet choice realised as the R-projection; the finite shadow is exhausted (v113/v125/v143 + this census).

Lemma 10 (G3–G4 — reflection positivity, fixed and integrated). *For each Θ -invariant g in the Calderón ball the physical boundary kernel $K_g = P_{\text{adm}} \Pi_{\text{phys}} \mathcal{N}_g^{TT} \Pi_{\text{phys}} P_{\text{adm}}$ is reflection positive, $\langle \Theta F, F \rangle_{K_g} \geq 0$ (G3); and $d\mu_{\text{Cal},\chi}$ is Θ -invariant, supported where G3 holds uniformly, so $\int \langle \Theta F, F \rangle_{K_g} d\mu_{\text{Cal},\chi} \geq 0$ (G4). **G4 is the decisive step:** RP after integrating over dynamical metrics, not merely at fixed background.*

Lemma 11 (G5 — gap dominance [E]). $2\|V_{\text{metric}}\|_{\text{rel}} < \Delta = 6 \log \frac{3}{2}$, with $\|V_{\text{metric}}\|_{\text{rel}} \leq 248c_3^2 = \frac{31}{8\pi^2} = 0.39262$, hence $2\|V\| = 0.7852 < 2.4328$ and $\Delta_{\text{eff}} \geq \Delta - 2\|V\| > 1.647 > 0$. The numerical inequality is trivial ([verification/v29_research_contract_certs.py]); the work is the operator-norm bound (hard).

Sugawara reading + atomic OS moment ([verification/v171_os_moment_cluster.py]). The G5 budget is the E_8 Sugawara constant: $248c_3^2 = \frac{31}{8\pi^2}$ with $31 = 1 + h^\vee(E_8)$ and $c((E_8)_1) = 248/31 = 8$, so gap dominance reads $|R^+(A_3)| \log(N_{\text{fam}}/|\mathbb{Z}_2|) > \frac{1+h^\vee(E_8)}{4\pi^2}$ — the A_3 transfer gap beats the E_8 Sugawara metric budget. Independently, the G3 reflection positivity is a *positive moment problem*: $\text{spec}(T) = \{1, (2/3)^6, (1/3)^6\}$ makes every two-point correlator $C(n) = A + B(64/729)^n + C(1/729)^n$ the moment sequence of a positive atomic measure, so each OS Hankel matrix factorises as a Vandermonde Gram and is positive semidefinite (a clean RP certificate; cluster constant 729/665, correlation length $\xi = 1/(6 \log \frac{3}{2}) = 0.4111$).

Theorem 1 (Decoupling theorem — physical low-energy claims do *not* need the full G_{metric}). *If the admissible sector is reflection-positive (G4) and the metric coupling is gap-dominated, $2\|V_{\text{metric}}\| < \Delta = 6 \log \frac{3}{2}$ (G5), then the admissible IR measure is stable under the ambient completion: a strictly positive effective gap $\Delta_{\text{eff}} = \Delta - 2\|V\| > 0$ protects the admissible sector from the (un-built) ambient/deep-UV, so every physical low-energy read-out (masses, mixings, α^{-1} , the $R+R^2$ regime) is independent of the open ambient measure. Symbolically*

$$\boxed{\text{RP} + 2\|V\| < \Delta \implies \text{admissible IR measure stable under ambient completion.}}$$

Thus (G_{metric}) is not a diffuse “full quantum gravity missing” block: it is the sharply delimited ambient projective limit (G6), a strict-TOE certification target whose closure the falsifiable IR physics does not depend on. **Typing:** the inequality is machine-checked [E]; the operator-norm bound on $\|V_{\text{metric}}\|$ is the conditional input.

Lemma 12 (G6–G7 — projective limit & Ward identities). *A projective system μ_{χ_n} with $\pi_{m,n*}\mu_{\chi_m} = \mu_{\chi_n}$, uniform moment bounds $\sup_n \int \|g\|_{H^s}^p d\mu_{\chi_n} < \infty$, tightness and regulator-independence yields $\mu_{\text{QG}} = \lim \mu_{\chi_n}$ on $\mathcal{G}/\text{Diff}$ (G6, the hardest step); the limit obeys the BRST Ward identities $\langle s\mathcal{O} \rangle = 0$, whence $\nabla^\mu T_{\mu\nu} = 0$ and the η boundary phase cancels the APS anomalies (G7).*

Gate 2 reduction: G6 is a boundary projective limit, not a bulk one ([verification/v76_gmetric_reduction.py])

The exact decoupling margin is $\Delta_{\text{eff}} = \Delta - 2\|V\| = 6 \log \frac{3}{2} - \frac{31}{4\pi^2} = 1.648 > 0$, with $\|V_{\text{metric}}\|_{\text{rel}} \leq 248c_3^2 = \dim(E_8)c_3^2 = \frac{31}{8\pi^2}$ ($31 = 2^{g_{\text{car}}} - 1$). **Holographic reduction:** because the seam is a *finite causal boundary*, the ambient *bulk* metric measure is reconstructed from a finite-codimension seam (Calderón) boundary measure — so G6 is a *boundary* projective limit. The conditional theorem is then

$$\boxed{\text{RP}(\text{seam-Calderón boundary kernel}) + \text{tightness} \implies G_{\text{metric}} \text{ closes}}$$

i.e. the dimension of the open problem drops from bulk QG to a finite seam-boundary measure. **Residual:** the constructive boundary projective limit (constructive-QFT grade) — reduced, not closed. So Gate 2 is now *two clean tiers*: Tier A (IR physics) closed under RP+gap; Tier B (strict TOE) reduced to a boundary measure, no longer a diffuse “full QG missing”. [C]/[O]

And that boundary measure is a *rigorously constructed* net ([verification/v77_e8_conformal_net.py]). The level-1 central charges $c = \dim \mathfrak{g}/(1 + h^\vee)$ give $c(E_8)_1 = \frac{248}{31} = 8 = \text{rank } E_8$, $c(D_5)_1 = \frac{45}{9} = 5$, $c(A_3)_1 = \frac{15}{5} = 3$, and $c(D_5) + c(A_3) = 5 + 3 = 8 = c(E_8)$ is a *conformal embedding* $(D_5)_1 \times (A_3)_1 \subset (E_8)_1$ (coset $c = 0$). Now $(E_8)_1$ is the holomorphic $c = 8$ E_8 -lattice VOA / conformal net — one of the most rigorously constructed objects in mathematical QFT (Frenkel–Kac–Segal; Kawahigashi–Longo / Carpi). **So the conditional theorem becomes concrete:** *identify the seam-Calderón boundary measure with the $(E_8)_1$ lattice net* (carrier = $(D_5)_1 \times (A_3)_1$ subnet), and the gap $\Delta_{\text{eff}} > 0$ supplies the clustering/tightness — so G_6 is imported into existing RCFT/conformal-net rigor rather than built from scratch. Residual = the net identification + bulk reconstruction. [E] (charges, embedding) + [C] (identification)

The Simple-Current Extension Theorem — the central G_{net} closing statement [E]/[C] ([verification/v154_simple_current_theorem.py])

The conditional theorem can now be stated as one clean *simple-current extension*. Let $A = (D_5)_1 \otimes (A_3)_1$ be the carrier net and $L = \langle(1, 1)\rangle \subset \text{Disc}(D_5) \oplus \text{Disc}(A_3) = \mathbb{Z}_4 \oplus \mathbb{Z}_4$ the isotropic μ_4 glue line ($q(k(1, 1)) = k^2 \in \mathbb{Z}$). Then the local extension $B = A \rtimes L$ has

$$[B : A] = |L| = 4 = |\mu_4|, \quad c(B) = 5 + 3 = 8, \quad \mu(B) = \frac{\mu(A)}{|L|^2} = \frac{16}{16} = 1$$

so B is *holomorphic* (KLM μ -additivity), and a holomorphic $c=8$ chiral net is the lattice net of the *unique* even-unimodular rank-8 lattice — hence $B \cong (E_8)_1$ (the same- c rival $SO(16)_1$, $\mu=4$, is excluded). Every algebraic ingredient is machine-checked: the $\mathbb{C}[\mathbb{Z}_4]$ Longo Q-system ([verification/v125_glue_qsystem.py]), the $\mathbb{Z}_4$ -graded Frobenius E_8 hull ([verification/v143_graded_frobenius.py]), the Ramond glue sectors ([verification/v148_fock_census.py]). **Consequence:** G_{net} is internally *closed* if the seam-Calderón boundary net is *defined* to be A ; it stays the *one* external identification “the RP seam-Calderón net is A ” if the boundary net is to be *derived* from the seam kernel. Either way this is the single strict-TOE certification theorem still carrying weight — a clean extension statement, not “find a quantum gravity”.

And that one identification reduces to a single shared premise ([verification/v155_quasifree_boundary.py]). The identification splits (v110) into (a) “the seam Calderón involution is sheet-odd” — the double-cover deck, the same $|\mathbb{Z}_2|$ that halves $c_3 = \frac{1}{2.4\pi}$ (structural) — and (b) “the seam 2-point kernel is quasi-free”. Part (b) is *not independent*: the boundary marginal of a reflection-positive *Gaussian* bulk is the compression PTP of the bulk covariance, and the compression of a contraction ($i\Gamma$ with spectrum in $[-1, 1]$) is a contraction — so the boundary/seam state is automatically quasi-free, with Wick inherited (every boundary correlator a Pfaffian of Γ_d). Hence (b) *follows* from “the seam bulk is a reflection-positive free field” — the *same* premise already load-bearing in the area law ([verification/v59_area_law_evidence.py]) and the EH mechanism ([verification/v150_replica_eh_model.py]/[verification/v151_bfk_split.py]). **So the entire seam-side residual is one physical premise** — the seam bulk is a free RP field — shared by G_{net} , the area law and R3, and given it the chain Gaussian bulk $\Rightarrow$ quasi-free boundary $\Rightarrow$ one kernel = net (v113) $\Rightarrow$ carrier $A \Rightarrow (E_8)_1$ is exact. It is not reducible by a finite computation — it is the one place TFPT meets functional analysis.

The construction, exhibited and verified ([verification/v156_seam_net_construction.py]). The “ $B \cong (E_8)_1$ ” step is no longer an assertion — the chiral net is built explicitly and the identification *checked*. The Dirichlet-to-Neumann (Calderón) operator of the 2d Laplacian acts on a boundary mode e^{ikx} by the harmonic extension $e^{ikx - |k|y}$: $-\partial_n u|_0 = |k|u$, so for a free bulk the seam operator is the massless chiral dispersion $\omega_k = |k|$ that defines the free-fermion net on S^1 . Sixteen Majoranas give $c = 8 = 5 + 3$; the weight-1 currents are $248 = 120_{\text{NS}} + 128_{\text{R}}$ (the $SO(16)$ bilinears plus the Ramond spinor, Frenkel–Kac); and the holomorphic character is

$$\chi = \frac{E_4}{\eta^8} = q^{-1/3} (1 + 248q + 4124q^2 + 34752q^3 + 213126q^4 + \dots) = j^{1/3}$$

(verified by q -series), the 248 at level 1 being $\dim E_8$ — so the net *is* $(E_8)_1$. *What remains* is exactly (i) the free-bulk premise above and (ii) the operator-algebraic existence / uniqueness of the net generated by

the seam Calderón kernel (a Kawahigashi–Longo-type net-construction theorem). Route 3 thus *exhibits and verifies* the target; the irreducible residual is the one premise + the standard net-existence step — where TFPT meets constructive QFT, now sharply located.

And the free-bulk premise itself reframes: freeness is a rigid fixed point, not a postulate ([`verification/v157_rigid_fixed_point.py`]). Three facts click together. (i) The DtN/Calderón symbol is *universally* $|k|$: the boundary operator of *any* second-order elliptic bulk is an order-1 Ψ DO with principal symbol $|\xi|_g$ (Lee–Uhlmann), so the free chiral dispersion is bulk-detail-independent and any interaction lives in the lower-order (irrelevant) symbol. (ii) A holomorphic $c=8$ theory is *rigid*: every field has weight $(h, \bar{h}=0)$, an exactly-marginal deformation needs $(1, 1)$, and with $\bar{h}=0$ *no* $(1, 1)$ field exists — the fixed point is isolated, there is no continuous knob along which a non-Gaussian interaction could turn on. (iii) The *same* $|\mathbb{Z}_2|$ that halves the one-sided S^2 Gauss–Bonnet ($c_3 = \frac{1}{|\mathbb{Z}_2| 2\pi \chi(S^2)} = \frac{1}{8\pi}$) labels the sheet *and* is the Ramond/GSO projection giving $\mu=1$ ($248=120_{\text{NS}}+128_{\text{R}}$) — one object, three roles, so the geometry fixing c_3 is the projection making the net holomorphic. **Consequence:** premise (A) shrinks from “prove the seam is Gaussian” to “the gapped ($\Delta=6 \log \frac{3}{2} > 0$) boundary flow reaches the holomorphic fixed point” — given which, freeness is *forced* by rigidity + uniqueness ($(E_8)_1$, v83), not fine-tuned. A strictly smaller and simpler residual.

And the destination is a proven stable fixed point ([`verification/v158_fixed_point_stable.py`]). An exact operator-dimension count settles it: the chiral Majorana has $h=\frac{1}{2}$, the bilinear current $h=1$, the quartic $h=2$. The relevant window $0 < h < 1$ for bosonic deformations is *empty* (bosonic chiral operators have integer h , lowest 1) — nothing relevant flows the free theory away; the $h=1$ currents are chiral (v157), not $(1, 1)$ -marginal; and the lowest genuinely-interacting operator (the quartic, $h=2 > 1$) is *irrelevant*, as are all higher $2k$ -fermi operators ($h=k$). So the free chiral $c=8$ fixed point is *isolated and stable* against all interactions. Premise (A) thereby reduces to a *basin-of-attraction* statement with a proven stable target — the gapped ($\Delta=6 \log \frac{3}{2}$) flow lands here; the convergence is conformal-perturbation analysis, the destination is settled.

And the closure can be stated as one fixed-point theorem — which relocates (A) to a single internal premise ([`verification/v160_seam_gaussianity_from_pf.py`]). Phrase the argument internally: (i) a quasi-free state has *all* higher connected cumulants $\kappa_{2n}=0$ (the signed set-partition/Pfaffian inversion of $\langle \psi_1 \cdots \psi_{2n} \rangle = \text{Pf}(\Gamma)$ leaves only the degree-2 kernel — the v113 “one kernel = the net” property at the cumulant level); (ii) a kernel-preserving transport fixes the whole quasi-free state by Wick functoriality; (iii) a gapped primitive Perron–Frobenius transport has a *unique* fixed ray, so a non-Gaussian piece is impossible — if κ_{2n} were fixed it would be a *second* fixed ray (killing uniqueness), and if not fixed it sits in the gapped complement and decays to 0 (any gap > 0 suffices; the value $(2/3)^6$ only sets the rate). *The honest residual:* v56’s gapped PF uniqueness is established on the *three-dimensional* scalar transfer spectrum $\{1, (2/3)^6, (1/3)^6\}$ (the flavour gap / horizon Page-recovery rates), whereas the 16-Majorana seam state already carries $\binom{16}{4}=1820$ and $\binom{16}{6}=8008$ higher-cumulant directions — the Schwinger cone the transport must contract has dimension $\gg 3$, and uniqueness must hold there (not merely on the quasi-free sub-cone) for the argument to be non-circular. **So (A) is equivalent to one strictly smaller, internal, exact-testable statement:** the admissible TFPT seam transport is a *primitive, spectrally gapped, RP-cone-preserving* map on the *full* Schwinger cone, not only on the scalar readout spectrum of v56 — which the suite does *not* yet establish [E]/[O].

And that full-cone premise is now discharged to a single finite one-particle statement ([`verification/v161_one_particle_reduction.py`]). The lever is Bogoliubov second quantisation: a quasi-free transport is $T_\Sigma = \Gamma(t)$, the second quantisation of a one-particle contraction t on the 16-dimensional Majorana space H_Σ , and the Fock space grades by correlator degree, so $\Gamma(t)$ acts on the degree- m sector as the exterior power $\Lambda^m(t)$. Three exact consequences follow (all machine-checked): (i) $\text{spec}(\Gamma(t)|_{\deg m})$ is the set of products of m one-particle eigenvalues — the whole correlation spectrum is generated by the finite one-particle spectrum; (ii) **the cone gap equals the one-particle gap:** the sub-leading eigenvalue of $\Gamma(t)$ over *every* degree is $(2/3)^6$ (pad the largest contracted mode with fixed factors), so premise 5 on the cumulant space *is* premise 5 on the 16-dim space, the same number; (iii) the eigenvalue-1 sector is Λ^* of the fixed subspace — the disconnected Wick powers of K_Σ — so with a strict one-particle gap no independent connected cumulant is fixed, and an independent fixed cumulant *on* the fixed subspace is excluded by v113 (one polarisation $\Leftrightarrow$ one vacuum). Premise 1 (quasi-freeness) is itself *forced*, not assumed: the $120 = \dim \mathfrak{so}(16)$ weight-1 chiral currents *are* the Bogoliubov generators on the 16 Majoranas, and v158 makes every non-Gaussian deformation irrelevant. **So the three full-cone premises collapse to one finite, 16-dimensional statement:** the seam RG transport’s one-particle symbol is the gapped Calderón Bogoliubov map — fixed eigenspace = the rank-8 K_Σ polarisation (v113),

sub-leading gap $= (2/3)^6$ (v56). Every *number* in it is already supplied by the suite; the one irreducible step is the physical identification that the actual seam RG generator equals this Bogoliubov map — the same operator-algebraic (Kawahigashi–Longo-class) statement as the net-existence residual of (A_2) , now with the infinite-dimensional cone eliminated [E]/[O].

And that finite statement carries no new open content: it factors into the two already-open items ([verification/v162_seam_transport_identification.py]). Take the one-particle symbol apart into its two data, the gap value (β) and the fixed space (α) . (β) is forced: a μ_4 -equivariant operator is diagonal in the cusp basis — the deck average is the diagonal projection, $\sum_{k=0}^3 U^k X U^{-k} = |\mu_4| \text{diag}(X)$ for $U = \text{diag}(1, i, -i)$, and the commutant of U is exactly the diagonal algebra — so its spectrum is the cusp weights $\text{spec}(A_0^*) = \{0, \frac{1}{3}, \frac{2}{3}\}$ (the parabolic residue, v115), and the transfer eigenvalue $(1-\alpha)^{p_2}$ with $p_2=6=|R_+(A_3)|$ gives $\{1, (2/3)^6, (1/3)^6\}$, gap $6 \log \frac{3}{2}$ — every entry carrier data, no free knob; the lift to the 16 Majoranas is the established A_3 subnet grading (v137). (α) is the rank-8 Calderón polarisation of the $\omega_k=|k|$ free dispersion (v113/v156), the IR conformal free-fermion vacuum. **What is left** is only that the seam RG generator is μ_4 -equivariant and admissible — which is the R1 “seam clock” identification (the transfer operator as the seam’s gapped generator) — together with the A_2 net-existence theorem. Both are pre-existing open items, so premise (A) introduces *no independent open content*: it factors exactly into $A_2 + R_1$ with every number forced. The “free-bulk cloud” is gone; what remains is the same two identifications the programme already names — (A) is closed *conditionally* on them (a unification, honestly not an unconditional proof) [E]/[O].

And both residual identifications are pinned to their floor, so (A) has no independent open gate left ([verification/v165_premise_a_closure.py]). A_2 (net existence) is not open research but *assembly* of established mathematics: the 16 free Majoranas define a free-fermion conformal net (Buchholz–Mack–Todorov), the holomorphic $c=8$ lattice net is $(E_8)_1$ (Frenkel–Kac–Segal / Dong–Xu / Kawahigashi–Longo), the μ_4 extension is a Longo Q-system of index 4 (v125); the one TFPT-specific input — that the seam Calderón kernel is the free-fermion two-point function — is done because the DtN symbol is universally $|k|$ (v156/v157). So A_2 re-types [O] $\rightarrow$ [C] (closed by assembly). R_1 (the seam clock) reduces to GATE.QGEO: a μ_4 -equivariant flat gapped generator on the carrier is *unique* ($= A_0^*$, v115/v116), so its gap $(2/3)^6$ is forced once the seam generator lies in that class, and the residual is exactly one geometric identification — the seam collar boundary is the μ_4 -punctured sphere $\mathbb{P}^1 \setminus \mu_4$ — whose cohomological half is already established (v137, $b_1=3=N_{\text{fam}}$). That is GATE.QGEO, an already-open realisation premise, *not* a new gate. By the No-Unit Theorem (v153) the irreducibles stay exactly $\{\pi, v_{\text{geo}}\}$ — closing (A) adds none. **Final accounting:** premise (A) = (A_2 assembly, [C]) + (R_1 =GATE.QGEO, already open) + the $\{\pi, v_{\text{geo}}\}$ irreducible core (a theorem) — it carries *zero* independent open gates and ceases to be its own gate [E]/[O].

And the two structural residuals — net existence and full-cone reflection positivity — are now discharged to a theorem, leaving the seam realisation as the one irreducible open premise ([verification/v175_net_existence_full_cone.py]). The CAR second-quantisation functor $\Gamma(t) = \bigoplus_m \Lambda^m(t)$ lifts the one-particle seam contraction t to the *complete* Fock space $\Lambda^\bullet(\mathbb{R}^{16})$ ($\dim 2^{16}=65536$); by the exterior-power spectrum theorem every eigenvalue of $\Gamma(t)$ is a subset product of $\text{spec}(t)$, so checking all 2^{16} products shows $\Gamma(t)$ is a *contraction* (full-cone RP), with eigenvalue-1 multiplicity $2^8=256$ (the wedges of the rank-8 K_Σ polarisation) and sub-leading eigenvalue exactly the one-particle gap $(2/3)^6$ — so full-cone RP/gap reduce *for every* m (not only $m \leq 6$, v161) to the one-particle contraction (Shale–Stinespring), *discharging the v160 scope premise by a theorem*. And A_2 is now an *assembled and verified* certificate: the $(E_8)_1$ character $E_4/\eta^8=1+248q+4124q^2+34752q^3+213126q^4+1057504q^5$ (level-1=248), the embedding $248=120+128$, and the E_8 Cartan matrix even with $\det 1$ (the unique rank-8 even unimodular lattice, v83) — the net *exists* by cited theorems. **Both residuals then collapse to the same input:** that the actual seam Calderón kernel is this gapped one-particle contraction on $\mathbb{P}^1 \setminus \mu_4$, i.e. QGEO.REALIZE.01, whose finite half is proven (v168). That realisation half is a constructive-geometry/AQFT statement, *not* closeable by a finite computation, and is left honestly [O] — the single irreducible structural premise; net existence and full-cone RP are [E].

The No-Unit Theorem — v_{geo} is an irreducible metrology primitive [E]/[O] ([verification/v153_no_unit_theorem.py])

v_{geo} is not an open gap. Every load-bearing compiler datum — $g_{\text{car}}, |\mu_4|, N_{\text{fam}}, \text{rank } E_8, \alpha^{-1}$, the determinant ladder $(3, 4, 8, 20)$, $c_3 = \frac{1}{8\pi}$, φ_0^{ret} — is mass-dimension 0 and *invariant* under a unit rescaling $L \mapsto \lambda L$, whereas a mass / energy / $1/G$ carries nonzero dimension. A map from dimension-0 inputs

to a dimension- $d \neq 0$ output is simultaneously invariant and covariant: impossible unless the unit is introduced. **So a dimensionless boundary compiler provably cannot select an absolute scale** — v_{geo} is a *forced input*. The three apparent residuals then collapse onto the one unit, differing only in mass-dimension:

$$U_{\text{point}} \sim v_{\text{geo}}, \quad \frac{1}{G} \sim v_{\text{geo}}^2, \quad m/\mu = e^{3/4} \text{ (a pure ratio),}$$

so the irreducibles stay exactly {one dimensionful unit, π }. *Re-typing*: v_{geo} is an *irreducible metrology primitive* (a declared unit, like choosing a metre), not a missing derivation.

Theorem G — the closure statement [O] (target)

The relative spectral boundary kernel over the diffeomorphism-quotiented metric sector admits a regulator-independent, reflection-positive projective-limit measure μ_{QG} . Its Ward identities imply covariant conservation, and its low-curvature expansion is the TFPT $R + R^2$ seam action

$$f_R R_{\mu\nu} - \frac{1}{2} f(R) g_{\mu\nu} + (g_{\mu\nu} \square - \nabla_\mu \nabla_\nu) f_R = \bar{M}_{\text{Pl}}^{-2} T_{\mu\nu} + O(R^3/M^4),$$

so the closed covariant equation becomes the controlled IR shadow of the full measure. *Status: open; the deepest item. G5 certified; the regime equation closed [verification/v28_gravity_fR.py].*

Certifiability and recommended order

Building block	Machine	Hardness
W_{wall} enumeration ($C_U^{(1)}$)	Lean / Python / Wolfram	easy (<i>done</i>)
D_4 fixed-locus polynomial ($C_U^{(2)}$)	Sage / Mathematica	medium
selector uniqueness ($C_U^{(3)}$)	interval arithmetic	medium
SNF, determinants, spectra	Lean	easy-medium
principal-symbol ellipticity (G1)	xAct / Cadabra / Lean	medium
gap inequality (G5)	Lean	easy (<i>done</i>)
operator-norm bound (G5)	analysis + numerics	hard
projective limit (G6)	not primarily machine	very hard
Ward identities (G7)	BV/BRST formal + checks	hard

Recommended order.

Selector-triangle pairings $\rightarrow v_{\text{geo}}$ scale anchor $\rightarrow G_{\text{net}}$ inclusion theorem

The flavor interface (historically U_{wall}) is now reduced to the *selector-triangle pairings* (the dual normal pair (d, n) forcing R columnwise, v136/v139); it is finite, falsifiable and the cheapest visible upgrade. The v_{geo} scale anchor is the one dimensionful input (same nature as $1/G$). G_{net} — the index-4 seam-net inclusion certifying the metric sector — is more fundamental but a *programme*, not a single proof: heavy machinery, with the projective limit the genuine summit. F_{transfer} (source $\rightarrow$ pole / relic / cosmology) is the remaining downstream interface.

F_{transfer} **is a typed functor, not a bag of open topics.** It is standard physics fed TFPT *source* data, $F_{\text{transfer}} = F_{\text{observable}} \circ F_{\text{threshold}} \circ F_{\text{RG}}$, with exactly four interfaces:

Interface	Map	Status	Source data → observable
F_{pole}	source→pole masses	[E] / [C]	Koide $53/54 = a^T(R+Q)\mathbf{1}/(2\mathbf{1}^T Ra)$ is an exact operator factor [E] (the missing Sheet-Diamond corner); the pole-transfer interpretation is [C] ([verification/v183_koide_f_corner_transfer.py])
$F_{\text{Boltzmann}}$	CP source→ η_B	[C]	$\tilde{m}_1=m_3/A_\Lambda$ anchors the washout, but $M_1=M_R\varphi_0^4$ only relocates the free input (M_R is the seesaw scale, not a compiler power), so η_B stays a <i>sharper scenario</i> , not a closure ([verification/v184_etaB_anchored_boltzmann.py])
F_{relic}	$(f_a, m_a, \theta_i, N_{\text{DW}}) \rightarrow \Omega_a$	[C]	the determinant-line axion can be all of DM, but only on the $\theta_i \approx 170^\circ$ hilltop where the relic is exponentially sensitive — a viable scenario ([verification/v185_axion_relic_solver.py])
F_{QCD}	$(\alpha_s, b_3, \Lambda_{\text{QCD}}, \text{lattice}) \rightarrow m_p/m_e$	[O]	a QCD matching contract ($b_3 = -7$ a carrier output), <i>not</i> a compiler number — the rejected 1836 near-formula is the discipline working

A machine guard enforces the typing: the four families must always carry a [C]/[O] marker and must never be promoted to a primitive [E] compiler prediction ([verification/v187_ftransfer_laws.py]). Exact *algebraic* sub-parts (the Koide factor, $b_3 = -7$) may be [E]; the *physical* prediction never is.

The functor contract CONTRACT.F.01, four axioms ([verification/v213_ftransfer_functor.py]).

Beyond the typing guard, the functor itself satisfies four checkable structural axioms, each instance a *discrete* compiler kernel [E] composed with an *external* continuous solver [C]: (1) **μ_4 -deck equivariance** — the discrete kernels are carrier/deck invariants $\{|\mu_4|, N_{\text{fam}}, g_{\text{car}}, A_\Lambda\}$, and the Koide multiplier $\lambda_2=(2/3)^6=64/729$ is the μ_4 -deck transfer eigenvalue (v82/v54); (2) **Plücker preservation** — the dimensionless readouts are Plücker readings ($53=a^T(R+Q)\mathbf{1}$, $\text{Pl}(F)=(1, 22, 30)$; $\|\text{Pl}(K)\|_1=11$); (3) **positivity/stochasticity** — $\text{spec } T=\{1, (2/3)^6, (1/3)^6\}$ is positive (RP/OS), $\kappa_f \in (0, 1)$; (4) **external modules explicit** — the continuous content (pole transfer, the flavored Boltzmann network, the finite- T $\chi(T)$ ODE, the lattice C_N) is named and external, never a compiler number. So CONTRACT.F.01 is the third research contract alongside (U_{wall}) and (G_{metric}): a typed functor, *not* a bag of open topics — but a consolidation, *not* a closure. So the visible residual is one geometric postulate (QGEO.SYM.01), one unit gauge (v_{geo}), and this one typed transfer functor — everything else is out-of-sample data.