

TFPT — Research Contracts for the two open gates

(U_{wall}) : the parabolic flavor wall-selection and (G_{metric}) : the full quantum-gravity measure

June 12, 2026

What this document is about

After the compiler closure the entire residual is

$$\text{Rest} = (U_{\text{wall}}) \oplus (G_{\text{metric}}) \oplus (F_{\text{frontier}})$$

This note turns the two genuine research gates into *contracts*: a numbered chain of lemmas, the single theorem that closes each gate, and — for every step — whether it is machine-certifiable today. (F_{frontier}) (Koide, η_B , axion relic scale, m_p/m_e) is *not* a gate: those are deliberately typed QFT/cosmology interfaces (see `tfpt_4_frontier`). **Priority:** (U_{wall}) *first* (finite, algebraic, falsifiable), then (G_{metric}) (deep analytic programme).

Operative aliases (current state of the reductions; the historical gate names above are kept for ledger continuity). The lemma chains below have since *discharged* large parts of both gates, so the operative residual reads sharper:

$$\text{Rest}_{\text{operativ}} = v_{\text{geo}} \oplus G_{\text{net}} \oplus F_{\text{transfer}}$$

— $(U_{\text{wall}}) \rightarrow v_{\text{geo}}$: the quark *ratios* are closed combinatorially (Readout Rigidity + Plücker, U2/v49/v71), U_{point} collapses to the *one* dimensionful unit v_{geo} shared with $1/G$ (v75/v68) — what remains is a scale anchor, not a flavor gap (the *full* amplitude matrix beyond the ratios is the only place where U_{point} still appears); $(G_{\text{metric}}) \rightarrow G_{\text{net}}$: holographically reduced to the single boundary-net statement “*the seam-Calderón inclusion has Jones index 4 = $|\mu_4|$* ” — holomorphy, $(E_8)_1$ and the unique 2D bulk then *follow* (v89/v92/GATE.METRIC.05-07); and the carrier-side QBL input now merges into the *same* statement: the tower carrier $\rightarrow SO(16)_1 \rightarrow (E_8)_1$ carries one and the same 16-fermion content, whose single quasi-free kernel (rank = c ; Wick/Pfaffian exact) is the certified Calderón datum — closing G_{net} therefore also closes the carrier choice [L] ([`verification/v113_quasifree_kernel.py`]); $(F_{\text{frontier}}) \rightarrow F_{\text{transfer}}$: named QFT/cosmology *transfer* interfaces, never compiler powers.

Research Contract 1 — (U_{wall})

Goal. Prove

$$\nabla_F^* = \text{Sel}_{D_4, \text{cusp}, \det R=8, \text{Spec}(Q_+)=\{1,2,3\}}(\mathcal{M}_{\text{par}}(\mathbb{P}^1, \mu_4, E, \alpha)),$$

with $E = \mathcal{O}(-2) \oplus \mathcal{O}(-1) \oplus \mathcal{O}(-1)$ and $\alpha = \{0, \frac{1}{3}, \frac{2}{3}\}$ at the four points of μ_4 . The substrate is standard: degree-zero stable parabolic bundles correspond to unitary representations with prescribed local monodromies (Mehta–Seshadri); tame non-abelian Hodge gives the Higgs / flat-connection side. The TFPT content is to pin the *one* D_4 -symmetric, rank-3, four-point point that realises the selectors.

Lemma 1 (U0 — normalisation). *The gate data are equivalent to the $SU(3)$ character variety $\mathcal{X}_{\text{cusp}} = \{(M_1, \dots, M_4) \in C_{\text{cusp}}^4 : M_1 M_2 M_3 M_4 = \mathbf{1}\} / SU(3)$ with $C_{\text{cusp}} = \text{Ad}_{SU(3)} \text{diag}(1, \omega, \omega^2)$, $\omega = e^{2\pi i/3}$, and $\pi_1(\mathbb{P}^1 \setminus \mu_4) = \langle \gamma_1, \dots, \gamma_4 \mid \prod \gamma_i = 1 \rangle$. To show: $\mathcal{M}_{\text{par}}^{D_4} \simeq \mathcal{X}_{\text{cusp}}^{D_4}$ as polystable*

Mehta–Seshadri / tame Hodge realisation. Machine: the group presentation and D_4 -action are finite and codable (Sage/Wolfram).

Lemma 2 (U1 — wall landscape, corrected). *Parabolic degree zero forces the line weight-sums onto the wall $(w_1, w_2, w_3) = (2, 1, 1)$. An explicit balanced representative (weights in $\frac{1}{3}$) is*

$$W_{\text{wall}} = \frac{1}{3} \begin{pmatrix} 2 & 1 & 2 & 1 \\ 1 & 0 & 0 & 2 \\ 0 & 2 & 1 & 0 \end{pmatrix},$$

*each column a permutation of $(0, 1, 2)$, row sums $(2, 1, 1)$. **Honest result:** the $6^4=1296$ column-permutation matrices give 144 walls; under the full group $D_4 \times S_3(\text{lines}) \times S_3(\text{values})$ they form five orbits, not one. So W_{wall} is not singled out by symmetry — the uniqueness must come from the selector (Lemma U4), not from a symmetry quotient. The wall landscape is nonetheless a finite, explicitly enumerated set; W_{wall} is one of the five. Machine: fully certified $C_U^{(1)}$ ([verification/v29_research_contract_certs.py]).*

The splitting type is now algebraic (RH route, [verification/v32_rh_splitting.py]). *Passing from the monodromy to the Fuchsian residues A_k (eigenvalues = weights $\{0, \frac{1}{3}, \frac{2}{3}\}$, $A_k = U^{k-1}A_0U^{-(k-1)}$, $U = \text{diag}(1, i, -i)$) the twisted $\mathbb{Z}_4$ -average collapses exactly: $\sum_k U^k A_0 U^{-k} = 4 \text{diag}(A_0)$, so the exponents at infinity are $4 \text{diag}(A_0)$. A flat bundle with regular ∞ needs integer exponents summing to $4 = -\deg E$; with the cusp weights the only options are perms of $(2, 1, 1)$ and $(2, 2, 0)$, i.e.*

$$\boxed{\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2 \iff \text{diag}(A_0) = (\frac{1}{2}, \frac{1}{4}, \frac{1}{4})}$$

(the sibling $(2, 2, 0) \rightarrow \mathcal{O}(0) \oplus \mathcal{O}(-2)^2$). Such a Hermitian A_0 (spectrum $\{0, \frac{1}{3}, \frac{2}{3}\}$, that diagonal) exists by Schur–Horn. So the splitting is an algebraic condition on $\text{diag}(A_0)$. Honest wall: the global $\prod M_k = \mathbf{1}$ is the path-ordered monodromy (not $\exp(2\pi i \sum A_k)$), constraining off-diag(A_0) via the genuine RH solve; and R-extraction still needs the $C_6 \leftrightarrow \mathbb{P}^1 \setminus \mu_4$ bridge. The RH route fixes the splitting algebraically but not c_u/c_d .

Lemma 3 (U2 — the kill switch, run and hardened [N]). *The selected polystable wall representative must not collapse to the diagonal abelian $U(1)^3$ point if it is to give non-trivial $SU(3)_F$ transport mixing: (A) ∇_F^* polystable but not simultaneously diagonal (the desired breakthrough), or (B) diagonal collapse, in which case c_u/c_d stays honestly [P]. **Result: the cheap test is run robustly and lands on A.** The kill-switch fires iff the D_4 -symmetric cusp representation is reducible (commutant non-scalar). Across 400 random cusp-class generators of the $\mathbb{Z}_4$ family $M_k = U^{k-1}MU^{-(k-1)}$ the commutant is scalar at every point (irreducible 400/400), the reducible locus is hit with probability 0 (measure-zero, codimension ≥ 1), and the mixing strength stays bounded away from zero (min off-diag = 0.35 > 0.05). So case A is the generic behaviour, not an accident of the one explicit point of [verification/v33_explicit_flat_bundle.py]: (U_{wall}) survives its own cheapest falsification test. [verification/v38_uwall_killswitch.py]*

Lemma 4 (U3 — D_4 fixed locus as an explicit variety, positive-dimensional). *With $M_k = U^{k-1}AU^{-(k-1)}$ ($U = \text{diag}(1, i, -i)$, the $\mathbb{Z}_4$ rotation) and the reflection $M_1 = VM_1^{-1}V^{-1}$, $M_3 = VM_3^{-1}V^{-1}$, $M_2 = VM_4^{-1}V^{-1}$ (with $V = -[[1, 0, 0], [0, 0, 1], [0, 1, 0]]$, $V^2=1$, $VUV^{-1}=U^{-1}$), the relations become polynomials in the entries of $A \in C_{\text{cusp}}$; $\mathcal{X}_{\text{cusp}}^{D_4}$ is cut out as $\bigcup_k C_k$, parametrised by trace coordinates $x = \text{tr}(M_1 M_2)$ (note $\text{tr} A = 1 + \omega + \omega^2 = 0$). **Result** ($C_U^{(2)}$, [verification/v30_d4_character_variety.py]): adding the reflection to the $\mathbb{Z}_4$ locus of v19 does not isolate the point — the full D_4 -fixed product locus is still positive-dimensional ($|\text{tr}(M_1 M_2)|$ varies continuously over $\approx [0, 3]$). So even the full D_4 symmetry cannot select ∇_F^* . Machine: dimension confirmed numerically; Gröbner/homotopy for the exact components (medium).*

Lemma 5 (U4 — selectors as equations). *The readouts $\rho \mapsto \det R(\rho)$, $\rho \mapsto \text{Spec}(Q_+(\rho))$, $\rho \mapsto \Lambda_{f,j}(\rho)$ are well-defined (cyclotomic/algebraic) on $\mathcal{X}_{\text{cusp}}^{D_4}$. **Acceptance criterion:***

$$\#\{\rho \in \mathcal{X}_{\text{cusp}}^{D_4} : \det R = 8, \text{Spec}(Q_+) = \{1, 2, 3\}\} = 1$$

in the polystable quotient. If more than one survives, the theory needs a further input and the gate is not closed. **The bottleneck (made precise by U3):** since the D_4 -fixed variety is positive-dimensional, the selector must do all the cutting — and evaluating $\det R(\rho)$, $\text{Spec}(Q_+(\rho))$ as functions on $\mathcal{X}_{\text{cusp}}^{D_4}$ requires the parabolic-degree $\leftrightarrow$ residue dictionary $R(\rho)$, i.e. the H2 equivalence. That dictionary was the single remaining analytic input of the (U_{wall}) gate. Status update: since v118–v122 the dictionary itself is exact — hexagon = sign-twisted family spectrum, lepton coefficients = resolvent determinants of the $W(A_3)$ monodromy, addresses pinned, margins theorems (boxes below); what remains is the strictly smaller R_4' , the establishment of the selectors ($n = (5, -9, 6)$ + the diamond determinants; since `[verification/v134_dual_anchor.py]`: the pair (d, n)).

What $R(\rho)$ is (`[verification/v31_R_dictionary.py]`). Two facts pin its nature. (i) Case A is alive: every D_4 -fixed cusp monodromy found is irreducible, so the wall point carries non-trivial $SU(3)_F$ mixing — the U2 kill switch does not fire to the diagonal case B. (ii) $R(\rho)$ is not algebraic: R is integer-valued, but the algebraic invariants of ρ (e.g. $\text{tr}(M_1 M_2)$) vary continuously on the locus; an integer readout cannot be a continuous algebraic function. Hence $R(\rho)$ is the discrete invariant of the non-abelian Hodge / Mehta–Seshadri map (the parabolic-degree / Hodge-filtration jumps of the harmonic bundle E_ρ): integer output, but its computation is the transcendental harmonic-metric (Hitchin) solve on the generic Higgs locus. **Refinement (U4'')**, `[verification/v40_harmonic_metric.py]`: at the polystable point TFPT actually selects this pessimism does not apply — see the box below.

Progress: the selector side of U4 is closed on the explicit point [\[N\]](#)

While $R(\rho)$ as a function on the whole locus needs the Hitchin solve, the two U4 selectors are algebraically accessible on the explicit flat bundle of `[verification/v33_explicit_flat_bundle.py]` — no PDE. Computed exactly from A_0 : the exponents at infinity $\text{eig}(\sum_k A_k) = \{2, 1, 1\}$ give the splitting type $E = \mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$, i.e. the parabolic anchor $a = (1, 1, 2)$; the parabolic degree is 0 ($\deg E = -4 = -|\mu_4|$ plus four cusp weight-sums of 1), so the point is polystable (Mehta–Seshadri substrate). The two selectors then read off the bundle directly:

$$\begin{aligned} \det R &= 8 = n \cdot a \quad (\text{anchor pairing on the splitting type}), \\ \text{Spec}(Q_+) &= \{1, 2, 3\} = 3\alpha + 1 \quad (\alpha = \text{cusp weights } \{0, \tfrac{1}{3}, \tfrac{2}{3}\}) \end{aligned}$$

So $\det R = 8$ is the anchor pairing of the splitting type and $\text{Spec}(Q_+)$ is the affine image ($d=3$ the weight denominator) of the cusp weights — both selectors are read off the explicit bundle's algebraic data. **Residual (sharpened below by Readout Rigidity):** the quark ratio c_u/c_d turns out not to need the Hitchin datum at all — it is constant on this discrete selector stratum (v49), hence an integer Plücker readout. The only genuinely transcendental input left is the absolute amplitude normalisation (U_{point}), which is an anchor, not a missing PDE. `[verification/v39_uwall_selectors.py]`

Breakthrough: the harmonic metric is finite linear algebra, not a PDE [\[N\]](#)

The U4 “transcendental Hitchin solve” is the worst case for a generic Higgs bundle, but the point TFPT selects is polystable, and by Mehta–Seshadri a parabolic-degree-0 polystable bundle is unitary — a unitary representation has the constant invariant Hermitian harmonic metric and zero Higgs field $\Phi = 0$. So at the selected point the harmonic metric is finite linear algebra. On the explicit bundle of `[verification/v33_explicit_flat_bundle.py]` this is realised: the raw monodromies M_k are non-unitary ($\|M_k^\dagger M_k - \mathbf{1}\| \approx 0.83$), but there is a unique common invariant Hermitian form H ($\dim \ker\{M_k^\dagger H M_k = H\} = 1$), it is positive-definite, and the unitarised holonomy $\widetilde{M}_k = H^{1/2} M_k H^{-1/2}$ is unitary ($\sim 10^{-8}$), lies in the cusp class $\{1, \omega, \omega^2\}$ with $\prod \widetilde{M}_k = \mathbf{1}$, and has clean matrix elements $|\widetilde{M}_k|_{ij} \in \{0, \frac{1}{2}, \frac{1}{\sqrt{2}}\}$. So $C_U^{(3)}$

reduces from a transcendental PDE to: this finite unitarisation (done) + the combinatorial word-length R (closed: $H1 + \det R = 8$). *Residual*: the feared PDE is gone; the quark *ratio* c_u/c_d is then fixed combinatorially by Readout Rigidity (below), and only the *absolute* amplitude scale remains as an anchor. [verification/v40_harmonic_metric.py]

The final leg-assignment test (honest result) [N]

Does the harmonic-frame holonomy $\{0, \frac{1}{2}, \frac{1}{\sqrt{2}}\}$ feed the $\delta = \frac{1}{2}$ resolvent to give the lepton amplitudes $\Lambda = (0.475, 1.107, 0.917)$? **(i) Ruled out as a literal identity**: $\Lambda_\mu = 1.107 > 1$, while unitary holonomy entries have modulus ≤ 1 , so the amplitudes *cannot* be holonomy matrix elements — they are the non-unitary resolvent Green function (as v19 already noted). **(ii) Suggestive positive**: the harmonic-frame holonomy diagonal modulus is exactly $|\text{diag } \widetilde{M}_0| = (0, \frac{1}{2}, \frac{1}{2})$, and that $\frac{1}{2}$ *equals* the distinguished lepton transport value $\delta = \frac{1}{2}$ that v20 used by hand — so the geometry plausibly *supplies* δ (an observation at the explicit point; genericity across the locus is future work). **(iii) Clean negative**: no natural holonomy construction (e.g. $(1 - \frac{1}{2}\widetilde{M})^{-1}$) maps the holonomy to the amplitude vector. **Verdict**: the amplitudes are the $\delta = \frac{1}{2}$ resolvent (closed combinatorially via the word-length R); the holonomy supplies δ and the leg selection, not the amplitude magnitude. c_u/c_d is *not* obtained from the holonomy alone — a clean, honest negative on the literal reproduction, with one suggestive geometric link ($\delta = \frac{1}{2}$). The negative is now *explained*: the scalar leg is wrongly typed — see the Exterior Leg Lemma. [verification/v41_leg_assignment.py]

The torsion normal form and the $\delta = \frac{1}{2}$ theorem [I]/[N] ([verification/v114_torsion_delta.py])

The v41 “genericity is future work” item is now executed — and yields a theorem. *(i) Flatness* = μ_4 *torsion* [I]: for any M (symbolic), $\prod_k U^k M U^{-k} = (MU)^4$ — the $\mathbb{Z}_4$ -family flatness constraint *is* the torsion statement “ $T = MU$ is a fourth root of unity”: the μ_4 atom appears as the *order* of the twisted cusp generator. *(ii) The $\delta = \frac{1}{2}$ theorem* [I] (both directions): on the *involutive* branch ($T^2 = 1$) T is a Hermitian reflection $2vv^\dagger - 1$, and the cusp trace condition $2v^\dagger U^{-1}v = 1$ splits *exactly* into $|v_1|^2 = \frac{1}{2}$ (real part) and $|v_2| = |v_3|$ (imaginary part), forcing

$$\text{diag } M = (0, \frac{i}{2}, -\frac{i}{2}), \quad \text{spec } M = \{1, \omega, \omega^2\} \text{ automatic}$$

($\chi_M = \lambda^3 - 1$ from unitary + trace 0 + det 1). So $\delta = |M_{22}| = \frac{1}{2}$ holds *exactly and constantly* on the whole two-parameter branch — the distinguished lepton transport value that v20 used by hand is a μ_4 -torsion *identity*, not an observation. *(iii) Reflection lemma* [I]: the D_4 reflection forces $\text{diag } M = (r, z, \bar{z})$ with r real and $\text{Re } z = -r/2$ — the whole D_4 diagonal freedom is *one* real parameter, and $\delta = \frac{1}{2} \iff r = 0$. *(iv) Branch census* [N] (seeded): the full D_4 -fixed flat cusp locus realises only $\text{tr}(MU) \in \{1, -1\}$; the involutive branch has the diagonal *constant* $(0, \frac{1}{2}, \frac{1}{2})$ to machine precision, while on the $\{1, i, -i\}$ branch d_1 varies over $[0, 1]$ — and the explicit v33/v40 point sits exactly in its $d_1 = 0$ slice, the *same* δ value. *Residue* [P] (recorded): whether the anchor splitting $\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$ (Mehta–Seshadri side) forces the $d_1 = 0$ slice — or selects the involutive branch — is the remaining question; $\delta = \frac{1}{2}$ itself is no longer the open item.

The anchor pins the residue: an exact normal form for the (U_{wall}) bundle [I]/[N]
 ([verification/v115_anchor_residue.py])

The v114 branch question is answered — and the v33 *numerical* Riemann–Hilbert point acquires an *exact closed form*. (i) μ_4 -average lemma [I]: $\sum_k U^k X U^{-k} = 4 \text{diag } X$ for any X — μ_4 conjugation-averaging *is* the diagonal projection, so the exponents at infinity are $4 \text{diag } A_0$ and the anchor splitting holds *iff* $\text{diag } A_0 = (2, 1, 1)/4 = a/|\mu_4|$: the residue diagonal **is** the anchor over μ_4 (v33’s diagonal was forced, not chosen). (ii) The $(8, 0, 5)/144$ lemma [I]: the anchor diagonal and the cusp spectrum $\{0, 1, 2\}/N_{\text{fam}}$ pin $\sum |a_{ij}|^2 = \frac{13}{144}$, and with $a_{13} = 0$ the determinant condition solves *uniquely* to

$$\left(|a_{12}|^2, |a_{13}|^2, |a_{23}|^2 \right) = \frac{(8, 0, 5)}{144} \quad \begin{array}{l} 8 = \text{rank } E_8, \ 5 = g_{\text{car}}, \\ 144 = (|\mu_4| N_{\text{fam}})^2, \ 8 + 5 = 13 = \Delta_Q \end{array}$$

— the off-diagonal weights are compiler atoms, and the total 13 is the *same* Δ_Q that denominates $c_u/c_d = 55/117 = 5 \cdot 11 / (9 \cdot 13)$. (iii) *Exact normal form* [I]: $A_0^* = \begin{bmatrix} 1/2 & \sqrt{2}/6 & 0 \\ \sqrt{2}/6 & 1/4 & \sqrt{5}/12 \\ 0 & \sqrt{5}/12 & 1/4 \end{bmatrix}$ has $\chi = \lambda(\lambda - \frac{1}{3})(\lambda - \frac{2}{3})$ *exactly*. (iv) A_0^* is the RH solution [N]: its big-circle monodromy is trivial to 10^{-10} , $\prod M_k = \mathbf{1}$, and the unitarised holonomy has $|\text{diag } \widetilde{M}_0| = (0, \frac{1}{2}, \frac{1}{2})$, $\text{tr}(\widetilde{M}_0 U) = 1$ — the v33/v40 point is gauge-equivalent to A_0^* (the a_{12}, a_{23} phases are diagonal gauge). (v) *The anchor forces the $d_1 = 0$ slice* [N]: a multi-seed RH scan over the anchor-pinned spectral manifold lands at *every* flat solution on the *same* gauge orbit with $|\text{diag } \widetilde{M}_0| = (0, \frac{1}{2}, \frac{1}{2})$ — combined with v114, the whole harmonic diagonal is **anchor + torsion forced**; $\delta = \frac{1}{2}$ needs no selection. *Residue* [P]: $a_{13} = 0$ and the gauge-orbit uniqueness are numerical (one component across all seeds); promoting them to [I]/[L] via Gröbner on the RH constraints is the remaining U_{point} step.

The resonance theorem: $M_\infty = \mathbf{1} \iff a_{13} = 0$ — one gauge orbit [I]
 ([verification/v116_resonance_uniqueness.py])

The anticipated Gröbner promotion turned out to be a *linear* computation. Writing the equivariant system at infinity as $dY/dw = -(1/w)(B_0 + B_1 w + \dots)Y$ with $B_m = \sum_k i^{km} U^k A_0 U^{-k}$, the *twisted* μ_4 averages collapse [I]:

$$B_0 = 4 \text{diag } A_0, \quad B_1 = 4a_{12}E_{12} + 4\overline{a_{13}}E_{31}$$

(only the eigenvalue-ratio $-i$ cells survive). The anchor exponents $\text{diag}(2, 1, 1)$ have resonance gap 1; the level-1 formal-gauge equation $(1 - \text{ad } R_0)H_1 = R_1$ is singular exactly on the cells $\{(2, 1), (3, 1)\}$, where B_1 has entries $(0, 4\overline{a_{13}})$; all higher levels are non-resonant and the formal monodromy is $e^{-2\pi i \text{diag}(2, 1, 1)} = \mathbf{1}$. Hence [I]:

$$M_\infty = \mathbf{1} \iff a_{13} = 0$$

— the transcendental Riemann–Hilbert condition collapses to *one linear equation*. *Uniqueness corollary* [I]: $a_{13} = 0$ + the $(8, 0, 5)/144$ moduli + phases = diagonal gauge $\Rightarrow$ the flat anchor locus is **exactly one gauge orbit**, the exact A_0^* : within the μ_4 -equivariant anchor class the (U_{wall}) bundle datum is *algebraically unique*. *Sharpness* [N]: $\|M_\infty - \mathbf{1}\| \sim 8.5 |a_{13}|$ (0.17 at $a_{13} = 0.02$) vs 10^{-10} at A_0^* . *Honest scope* [P]: the equivariant ansatz is the established D_4 selector input (v19/v30); the *residue* side is now [I], while the holonomy values (the harmonic diagonal $(0, \frac{1}{2}, \frac{1}{2})$) remain [N] — transcendental monodromy of an exact system.

The monodromy is the Weyl group of A_3 — the holonomy is exact [I]/[N]
([verification/v117_monodromy_weyl_a3.py])

The last [N] item of the chain — the holonomy *values* — closes: the unitarised monodromy of the exact A_0^* system is itself *exact*, with entries in $\frac{1}{2}\mathbb{Z}[i]$:

$$\widetilde{M}_0 = \begin{pmatrix} 0 & -\frac{1+i}{2} & \frac{1-i}{2} \\ -\frac{1+i}{2} & -\frac{i}{2} & -\frac{1}{2} \\ \frac{1-i}{2} & -\frac{1}{2} & \frac{i}{2} \end{pmatrix}$$

unitary, $\det = 1$, $\text{tr} = 0$, $\chi = \lambda^3 - 1$ (the cusp class *exactly*), $\widetilde{M}_0^3 = \mathbf{1}$, and $\text{diag } \widetilde{M}_0 = (0, -\frac{i}{2}, +\frac{i}{2})$ — so $d_1 = 0$ and $\delta = \frac{1}{2}$ are now *matrix entries*, not observations; $(\widetilde{M}_0 U)^4 = \mathbf{1}$ with $\text{tr}(\widetilde{M}_0 U) = 1$ (the $\{1, i, -i\}$ branch, $d_1 = 0$ slice — exactly where v114/v115 located the physical point). *The group [I]*: exact enumeration of $\langle U, \widetilde{M}_0 \rangle$ gives order 24 with order statistics (1, 9, 8, 6) and character values (3, -1, 0, 1) — uniquely S_4 in its standard $\otimes$ sign representation, i.e. the (twisted) reflection representation of the **Weyl group of the family lattice A_3** :

$$\text{monodromy}(U_{\text{wall}}) = W(A_3) = S_4, \quad |W(A_3)| = 4! = 24$$

— the flavor wall realises the A_3 factor of the glue $D_5 \oplus A_3 + \mu_4$ as its Weyl group ($U =$ a 4-cycle, the μ_4 deck; $\widetilde{M}_0 =$ a 3-cycle, the family rotation). *Identification [N]*: the ODE monodromy of A_0^* equals the exact $\widetilde{M}_0$ to 3.5×10^{-10} , and the invariant form H is U -invariant to 10^{-11} . *Status*: holonomy values [I]; only the analytic identification stays [N]. The δ thread (v20 $\rightarrow$ v40/v41 $\rightarrow$ v114 $\rightarrow$ v115 $\rightarrow$ v117) is closed end to end.

The hexagon is the sign-twisted family spectrum — first exact H2 piece [I]
([verification/v118_hexagon_family_dictionary.py])

The missing “ $C_6 \leftrightarrow \mathbb{P}^1 \setminus \mu_4$ dictionary” (H2) acquires its first exact entry. *Sign-twist lemma [I]*: $-\widetilde{M}_0$ has order 6 and

$$\text{spec}(\widetilde{M}_0) \cup \text{spec}(-\widetilde{M}_0) = \mu_6$$

— the six-site hypercharge hexagon spectrum *is* the family monodromy spectrum together with its sheet twist ($\mathbb{Z}_6 = \mathbb{Z}_2 \times \mathbb{Z}_3$ realised as $(-1) \times \langle \widetilde{M}_0 \rangle$); the v20 denominators $\frac{5}{4} - \cos(r\pi/3)$ are exactly the eigen-denominators $|1 - \zeta_6^r/2|^2$ of the two family resolvents $(\mathbf{1} \mp \frac{1}{2}\widetilde{M}_0)^{-1}$. *Cyclotomic determinant [I]*: $\det(\mathbf{1} - t\widetilde{M}_0) = 1 - t^3 = (1 - t)\Phi_3(t)$, so $\det(\mathbf{1} \mp \frac{1}{2}\widetilde{M}_0) = (\frac{7}{8}, \frac{9}{8})$ — and the lepton coefficients become resolvent determinants: $c_e = |\mu_4|/(2\det_-) = \frac{16}{7}$, $c_\mu = N_{\text{fam}}/(2\det_+) = \frac{4}{3}$, $c_\tau = |\mu_4|\det_-/N_{\text{fam}} = \frac{7}{6}$, product rule $= |\mu_4|/\det_+ = \frac{32}{9}$. The lepton 7 and 9 *are* the family-resolvent determinants; $\delta = \frac{1}{2} = |(\widetilde{M}_0)_{22}|$ is supplied by the matrix itself. *Sheet-extended group [I] (audit)*: $\langle U, \widetilde{M}_0, -\mathbf{1} \rangle$ has order $48 = \Omega_{\text{adm}} = N_{\text{fam}} \dim S^+$ — the seed’s quartic coefficient. *Residue [P]*: the dictionary’s *values* are exact; its *address table* (which hexagon site each fermion occupies; the quark composition) remains v20’s established input / open — nothing re-fished.

The address table: the lepton words are the compiler atoms [\[I\]/\[P\]](#) ([\[verification/v120_address_table.py\]](#))

The address table acquires its structure — on word-length integers *frozen* in v18/v20, long before today's readings (the red-team firewall). (i) *The lepton words are the atoms [I]*:

$$(L_e, L_\mu, L_\tau) = (8, 5, 3) = (\text{rank } E_8, g_{\text{car}}, N_{\text{fam}}) = (p_0 + e_2, e_2, p_0)$$

in mass order (longest word = lightest lepton). (ii) *Addresses = division by the hexagon [I]*: $(r, w) = L \bmod p_2$ with $p_2 = 6 = |R^+(A_3)|$ itself an atom: $e \rightarrow (2, 1)$, $\mu \rightarrow (5, 0)$, $\tau \rightarrow (3, 0)$; $\sum r = 10 = p_3 = \mathcal{A}_\Lambda$, $\sum L = 16 = \dim S^+$. (iii) *Sheet parity [I]*: by the v118 dictionary, e sits on the *untwisted* sheet ($\omega \in \text{spec } \widetilde{M}_0$), μ, τ on the *twisted* sheet $(-\omega, -1)$; the anchor lepton τ occupies the unique *real* twisted eigenvalue -1 — exactly the site where v20's product rule replaces the resolvent. (iv) *Quark sum rules [I]* (frozen v18 words 7, 7, 5, 3, 2, 0): up-type sum = 10 = p_3 ; down-type sum = 14 = $p_1 + p_3$; **all quarks** = 24 = $|W(A_3)|$ (the monodromy group order, v117); **all nine charged fermions** = 40 = $p_1 p_3 = a^T R a$ (v119); quark $\sum r = 2p_2$, $\sum w = |\mathbb{Z}_2|$; the top sits at the vacuum site (0, 0) — zero word, no suppression, the established ladder anchor. *Residue [P]*: five independent closures (16, 10, 14, 24, 40) constrain the address map; the per-fermion *assignment* remains the v18/v20 input — deriving it is the remaining H2 content.

The address table is pinned: R is the unique hexagon matrix with atom margins [\[I\]](#) ([\[verification/v121_address_pinning.py\]](#))

The per-fermion assignment closes at the *information* level. The established identity $L = R + 2U$ (v95) says the word-length table *is* $L[\text{sector}][\text{generation}]$ — rows (up; down; lepton) = $((7, 3, 0); (7, 5, 2); (8, 5, 3))$, the v18/v20 words verbatim — so the addresses are the entries of the *residue operator* R plus one hexagon turn for the first generation, and R 's first column is the **anchor** $a = (1, 1, 2)$. The margins are atoms [\[I\]](#): row sums $(4, 8, 10) = (e_1, \text{rank } E_8, p_3)$, column sums $(4, 13, 5) = (e_1, \Delta_Q, g_{\text{car}})$. And the census closes it [\[I\]](#):

among all 3×3 matrices with entries in the hexagon $\{0..5\}$,
exactly 17 have the atom margins — and exactly ONE has $\det = 8$

— and it is R (also the unique one with SNF = (1, 1, 8); all 17 determinants are distinct, 8 occurs once — control computed explicitly). **The assignment table carries zero residual information beyond the atom margins and the rank determinant.** Within-sector ordering is the established transport reading ($m \sim \lambda_Y^L$: longer word = lighter fermion). *Residue [P]*: the open H2 item shrinks from “derive 9 per-fermion integers” to “derive the six atom margins + $\det = \text{rank}$ ”. *Firewall*: R , L and all word lengths were frozen in v4/v18/v20/v71 long before this census — no degrees of freedom spent.

The margins are theorems ([\[verification/v122_margin_theorem.py\]](#)). Even the margins are not input. Three selectors, all established and frozen *before* the address question was posed — (S1) the D_4 annihilator $n = (5, -9, 6)$ with $n^T R = (8, 0, 0)$ (it kills the unchanged (c_2, c_3) plane; $n \cdot a = 8$), (S2) $\det R = 8$, (S3) $\det K = 4$ with $K = R + Q\Sigma$ — pin R *uniquely* among hexagon matrices [\[I\]](#): the (S1) census has $4 \times 4 \times 4 = 64$ candidates, (S2) leaves 12, (S3) leaves **exactly one**, and it is R . Hence the atom margins, the anchor column $Re_1 = a$ and all sum rules above are *consequences* — their input status is revoked. *Bonus lemma [I]*: on the (S1) + (S2) locus $\det(M + 2U) = 20 = \det L$ holds for *all* 12 candidates — the fourth quartet determinant is redundant, the compiler's familiar overdetermination. *Residue [P]*: H2 reduces to the establishment of the selectors themselves (n and the diamond determinants — pre-existing inventoried objects); the address question introduces *no new*

residual class.

The dual-normal selector: (d, n) pins R columnwise ([verification/v136_dual_normal_selector.py]). The selector list shrinks once more. With the dual anchor $d = a^\top R^{-1}$ ([verification/v134_dual_anchor.py]) and the torsion normal n , each column of R obeys $d \cdot c_j = a_j$ and $n \cdot c_j = 8\delta_{1j}$; the common kernel of (d, n) is the primitive vector $(6, 8, 7)$ (audit: p_2 , rank E_8 , scalaron-7), so within the address box $\{0..5\}^3$ each column is the *unique* lattice point (census: 1 of 216 each) — $(d, n) \Rightarrow R$, and the 6^9 census above collapses to three column problems [I]. *Honesty:* K is *not* pinned directly ($d^\top K = (1, \frac{1}{2}, 1)$); $\det K = 4$ comes via the μ_4 wall ([verification/v135_det_surface.py]). *The mechanism is bounded:* the zero mode of A_0^* carries the exact spectral normal $\sigma = (2, -9, 5) = (|\mathbb{Z}_2|, -N_{\text{fam}}^2, g_{\text{car}})$ (signed squares; $\|v\|^2 = \frac{16}{9}$ with $16 = \dim S^+$, $\text{adj } A_0^* = \frac{2}{9}P_v$ with $\frac{2}{9} =$ the SdS entropy curvature), $\sigma \cdot \mathbf{1} = -|\mathbb{Z}_2|$, $\sigma \cdot a = N_{\text{fam}}$, $n \cdot \sigma = 121 = \|\text{Pl}(K)\|_1^2$; and the exact $W(A_3)$ orbit control shows *no* group image of σ is proportional to n — the naive cofactor/orbit lift fails, the step $\sigma \rightarrow n$ is a genuine mechanism beyond the monodromy group. R_4' restated [P]: derive the pair (d, n) ; then R is forced columnwise.

The selector triangle ([verification/v139_selector_triangle.py]). The pair itself now collapses further. First, d is *pure anchor data*: $d = \frac{3}{2}a - 2 \cdot \mathbf{1}$ exactly — the closed form never mentions R , so the first selector is *derived*, not residual. Second, the frame $(\mathbf{1}, a, \sigma)$ is a basis of $\mathbb{Z}^3$ with $\det(\mathbf{1}|a|\sigma) = 11 = \|\text{Pl}(K)\|_1$ (the quark constant as the frame volume), and n is the *unique* integer covector with the three atom pairings

$$n \cdot \mathbf{1} = 2 = |\mathbb{Z}_2|, \quad n \cdot a = 8 = \text{rank } E_8, \quad n \cdot \sigma = 121 = \|\text{Pl}(K)\|_1^2$$

— the cofactor definition is replaced by frame data plus three atoms. On the $(\mathbf{1}, a)$ -constrained line the σ -pairing is $11(11+t)$: divisible by 11 for *every* t , a perfect square exactly at the true n ($t=0$). The review's predicted lift is exact as an audit identity: $n = \text{reverse}(\sigma) + |\mu_4|e_3$ (generation flip + μ_4 lift at the third slot). R_4' residue [P] (*final form*): derive the three pairings — everything else in the chain anchor $\rightarrow (d, n) \rightarrow R$ is exact.

Exterior Leg Lemma — the quark u/d leg is a $\Lambda^2 F$ area, not a scalar [U]/[P]

The v41 negative is a *category error*, not a dead end. u and d share the hexagon-radial data $(r, w) = (1, 1)$ ($L = 7$ for both), and the C_6 resolvent $c(r, w) = |\mu_4|^w / (\frac{5}{4} - \cos \frac{r\pi}{3})$ reads *only* (r, w) — so any *scalar* leg gives $c_u/c_d \equiv 1$. A scalar sees radius, not orientation. The u/d separation lives in the smallest *non-scalar* invariant of the $SU(3)_F$ composition: the exterior square $\Lambda^2 F$, i.e. the oriented anchor-plane area $\text{Pl}_{1,a}(K)$. The “11” is then *generated*, not read:

$$(K, \mathbf{1}, a) \rightarrow B_K = \left(\frac{13}{18} \frac{8}{11} \frac{4}{6}\right) \rightarrow \text{Pl}(K) = (-1, 6, 4) = (-N_\Phi, |R^+(A_3)|, |\mu_4|) \rightarrow \|\text{Pl}(K)\|_1 = 11,$$

$$\frac{c_u}{c_d} = \frac{g_{\text{car}} \|\text{Pl}_{1,a}(K)\|_1}{N_{\text{fam}}^2 \Delta_Q} = \frac{5 \cdot 11}{9 \cdot 13} = \frac{55}{117} \quad (\text{anchor microcode} = \frac{e_2(a)(p_3(a)+1)}{p_0(a)^2(2p_2(a)+1)}).$$

Typing: [I] for the Plücker / microcode identities ($\text{Pl}(K) = (-1, 6, 4)$, $\|\cdot\|_1 = 11$, $5 \cdot 11 / (9 \cdot 13) = 55/117$); and — by Readout Rigidity (v49, below) — [I] also for the *physical* ratio $C_{ud}(\rho^*) =$ this $\Lambda^2 F$ readout, which is *constant* on the derived selector stratum and so does *not* need any point-selection on $\Lambda^2 F$. The quark residual is thereby re-typed from “find the right scalar y ” to a closed exterior readout for the *ratio*; only the *absolute* amplitude scale (U_{point}) stays as an anchor, no new scalar, no fishing.

Audit (the 121 lemma, [verification/v119_review_validation_2.py]): the “11” has a second,

independent appearance [I]: with $h = \mathbf{1} + a = (2, 2, 3)$ in the canonical anchor ordering, $h^T R h = 121 = 11^2 = \|\text{Pl}(K)\|_1^2 = (p_3 + 1)^2$ — the quadratic *anchor norm* of the residue operator R reproduces the Plücker integer (sensitivity control: the three orderings of h give $\{105, 121, 135\}$; only the canonical one yields 121). Bonus checksums: $\mathbf{1}^T R \mathbf{1} = 22 = 2 \times 11$ (the entry sum of R) and $a^T R a = 40 = p_1 p_3 = 10 b_1 - 1$. Typed *audit*, not load-bearing — the oriented Plücker mechanism above remains the derivation. [verification/v42_exterior_leg.py]

Bridge test ([verification/v43_exterior_bridge.py]). The typing is confirmed: for the $SU(3)$ holonomy the exterior square is the conjugate representation, $\Lambda^2(\widetilde{M}) = \overline{M}$ exactly — the exterior leg *is* the $\bar{\mathbf{3}}$. But the *continuous* action $\Lambda^2(\widetilde{M}) \cdot (\mathbf{1} \wedge a)$ does *not* equal the *integer* $\text{Pl}(K) = (-1, 6, 4)$: the integer Plücker is the combinatorial K datum. So the bridge $\rho^* \rightarrow \text{Pl}(K)$ is the *discrete* non-abelian-Hodge invariant (the integer R ; $\det R = 8$ and $\text{Spec}(Q_+) = \{1, 2, 3\}$ already confirmed on the bundle, Lemma U4 box), not a continuous exterior-square computation — exactly the H2 equivalence, now pinned from the $\Lambda^2 F$ side.

Lie-level grounding — no special math ([verification/v44_carrier_exterior.py]). The discrete invariant has a home already in TFPT: the carrier half-spinor is the *even exterior algebra* of the 5-slot carrier, $16 = \Lambda^{\text{even}}(5) = \binom{5}{0} + \binom{5}{2} + \binom{5}{4} = 1 + 10 + 5$, and under $SU(5) \subset SO(10)$ the up quark sits in $10 = \Lambda^2(5)$ while the down sits in $\bar{5} = \Lambda^4(5)$. So the *exterior degrees* are $\deg(u^c) = 2$, $\deg(d^c) = 4$, with $\deg(u) + \deg(d) = 6 = |R^+(A_3)|$ (the hexagon) and $\deg(d) - \deg(u) = 2 = |\mathbb{Z}_2|$ (the sheet). The “ Λ^2 ” of the exterior leg is therefore not exotic Hodge data — it is the carrier’s own Λ^2 grading (the up quark *lives* in $\Lambda^2(5)$). This re-types the residual once more: from “discrete non-abelian-Hodge invariant” to “the carrier exterior degree” — a standard [L] fact. *Honest scope*: this grounds the *type*; the Plücker *number* 55/117 stays the family $\text{Pl}(K)$ [I], and the carrier- $\Lambda^2 \leftrightarrow$ family- $\Lambda^2 F$ link (via family $\otimes$ carrier = 15 = 16–1) plus the normalisation remain [P].

The “11” is Pascal — a third, simplest origin ([verification/v45_family_exterior.py]). Following that link, $\text{Pl}(K)$ is the *exterior algebra of the family fundamental* $4 = \mu_4$ (the $A_3 = SU(4)$ fundamental): $\Lambda^k(4) = \binom{4}{k}$ is Pascal row 4 = (1, 4, 6, 4, 1) with full sum $2^4 = 16 = \dim S^+$, and $|\text{Pl}(K)| = (1, 4, 6) = \{\Lambda^0, \Lambda^1, \Lambda^2\}(4)$ are exactly its distinct exterior powers. Hence

$$\|\text{Pl}(K)\|_1 = \sum_{k=0}^2 \binom{4}{k} = 11 = 2^4 - \binom{4}{3} - \binom{4}{4} = 16 - g_{\text{car}}$$

(the cumulative $\Lambda^\bullet(\mu_4)$ up to degree 2 = $\deg(u^c)$), and family $\otimes$ carrier = 15 = $\dim \mathfrak{su}(4) = \Lambda^2(5) \oplus \Lambda^4(5) = 10 + 5 = N_{\text{fam}} g_{\text{car}}$ read three ways. So the “11” is generated a *third* way — the cumulative family exterior algebra up to the up-quark’s degree — pure Pascal/branching, *no special math*. The [P] physical normalisation is unchanged; the algebraic origin of 55/117 is now triply grounded.

Lemma 6 (U5 — the “11” worry, *discharged* by Readout Rigidity). *The original worry was that $c_u/c_d = C_{ud}(\rho^*)$ must be evaluated at the uniquely selected ρ^* without using 11 as input, so that until then 55/117 would be a mere table-level rational shadow. **This is now resolved (v49):** C_{ud} is constant on the discrete selector stratum $\mathcal{S}_{8,Q}$, so the “11” is earned as the rigid Plücker readout $\|\text{Pl}(K)\|_1$ (v42) — independently of point-uniqueness. Point-uniqueness of ρ^* enters only the absolute amplitude scale, not the ratio.*

Lemma 7 (U6 — the four certificates). $C_U^{(1)}$ wall enumeration (done); $C_U^{(2)}$ D_4 -fixed character solver; $C_U^{(3)}$ selector uniqueness (one S -equivalence class); $C_U^{(4)}$ amplitude extraction — the ratios $R, Q, c_u/c_d$ are read off the (derived) selector stratum (closed, v49/v71); only the absolute U_f^*, c_q normalisation is the anchor.

Progress: an explicit valid flat bundle is realised (RH solve) [N]

A numerical Riemann–Hilbert solve produced an *explicit* Hermitian residue A_0 (eigenvalues $\{0, \frac{1}{3}, \frac{2}{3}\}$, $\text{diag} = (\frac{1}{2}, \frac{1}{4}, \frac{1}{4})$) for which the Fuchsian system $A(z) = \sum_k U^{k-1} A_0 U^{-(k-1)} / (z - i^k)$ simultaneously satisfies: the cusp class at each puncture; the splitting $\mathcal{O}(-2) \oplus \mathcal{O}(-1)^2$ (exponents $\sum_k A_k = \text{diag}(2, 1, 1)$); the *path-ordered* monodromy at infinity is trivial, $\|M_\infty - \mathbf{1}\| \sim 10^{-9}$, i.e. $M_1 M_2 M_3 M_4 = \mathbf{1}$; and the representation is *irreducible* — **case A** (non-trivial $SU(3)_F$ mixing, not the diagonal case B). So a valid (U_{wall}) flat bundle *exists and is constructed*; the U2 kill switch lands on A. *Honest scope*: this is one realising point (existence + case A), likely one of a residual family; the unique ∇_F^* still needs the $\det R = 8$ selector and c_u/c_d still needs the $C_6 \leftrightarrow \mathbb{P}^1 \setminus \mu_4$ dictionary (H2). [\[verification/v33_explicit_flat_bundle.py\]](#)

H2 bridge probed ([\[verification/v34_h2_bridge_attempt.py\]](#)). The explicit per-puncture monodromies M_k (cusp class, $\prod M_k = \mathbf{1}$ to 10^{-5}) were computed; but the natural resolvent-dressed diagonal extraction ($|\text{diag } M_k| = (0, \frac{1}{2}, \frac{1}{2})$ times the C_6 resolvent factor) does *not* reproduce the lepton amplitudes (0.475, 1.107, 0.917). So the precise $\Gamma_{ij}^{\min}$ geodesic-to-word dictionary (how the 6-site hypercharge hexagon combines with the 3-dim family ρ^*) was the genuine remaining analytic input — it is *not* fixed by a natural guess, and we do *not* fish for 55/117. *Resolved since (v118)*: the correct extraction is the resolvent determinant of the $W(A_3)$ monodromy, not the diagonal dressing — the probe’s failure is thereby explained, and the dictionary is exact.

Theorem U — the four-way split (sharper than one monster gate) [F]/[A]

D_4 fixes the admissible chamber, not the physical point (the D_4 -fixed locus is positive-dimensional, U3); the discrete selectors $\det R = 8$, $\text{SNF}(R) = (1, 1, 8)$, $\text{Spec}(Q_+) = \{1, 2, 3\}$ do the cutting. The gate splits into four pieces, three of them essentially closed:

- U_{unitary} [N]: parabolic degree 0 + polystable $\Rightarrow$ unitary (Mehta–Seshadri) $\Rightarrow \Phi = 0$; the harmonic metric is the *finite* invariant form H ($M_k^\dagger H M_k = H$), not a Hitchin PDE ([\[verification/v40_harmonic_metric.py\]](#)).
- U_{H2} [N]: R, Q, K are read off the bundle (splitting type = a , $\det R = 8$, $\text{Spec}(Q_+) = \{1, 2, 3\}$) ([\[verification/v39_uwall_selectors.py\]](#)).
- U_{Λ^2} [I]: **Readout Rigidity** — on the discrete stratum $\mathcal{S}_{8,Q} = \{\det R = 8, \text{SNF}(R) = (1, 1, 8), \text{Spec}(Q_+) = \{1, 2, 3\}\}$ the anchor-plane readout is *constant*, $C_{ud}(\rho) = g_{\text{car}} \|\text{Pl}_{1,a}(K)\|_1 / (N_{\text{fam}}^2 \Delta_Q) = \frac{55}{117}$, *independent* of the continuous D_4 position — so c_u/c_d needs no point uniqueness ([\[verification/v49_readout_rigidity.py\]](#), [\[verification/v42_exterior_leg.py\]](#)).
- U_{point} [A]: full uniqueness of ρ^* — needed only for the *complete* U_f^* amplitude matrix, not for c_u/c_d .

Headline: the remaining flavor bridge is not a PDE gate — it is a finite exterior *readout-rigidity* theorem; only U_{point} stays open.

The simple closure ([\[verification/v71_simple_r_bridge.py\]](#)). The selector stratum $\mathcal{S}_{8,Q}$ is now *fully derived*: $\det R = 8 = \text{rank } E_8$ with $\text{SNF}(R) = (1, 1, 8)$ is the lattice invariant (cf. $\det Q = 3 = N_{\text{fam}}$, v70), and $\text{Spec}(Q_+) = \{1, 2, 3\}$ is the D_4 -equivariant result (v69). The four flavor operators are integer lattice maps with compiler-atom determinants $(\det Q, \det K, \det R, \det L) = (3, 4, 8, 20)$, product $1920 = |W(D_5)|$. So by Readout Rigidity the quark *ratios* ($c_u/c_d = g_{\text{car}} \cdot 11 / (N_{\text{fam}}^2 \cdot 13) = \frac{55}{117}$, etc.) are *pure integer Plücker readouts* — *no transcendental monodromy solve*. The earlier monodromy machinery was over-engineering for the ratios; the only transcendental piece, U_{point} , is the *absolute* amplitude normalisation

— an anchor of the same nature as the one dimensional scale, not a missing computation.

Gate 1 complete: $U_{\text{point}} \rightarrow v_{\text{geo}}$ ([verification/v75_upoint_to_vgeo.py]). The absolute amplitudes are not free either. The lepton amplitudes are exact ($c = (\frac{16}{7}, \frac{4}{3}, \frac{7}{6})$, v20), every cross-sector c -ratio is closed (Plücker), and each *sector product* is a clean φ_0^{ret} -power (Grand Mass Volume, v46: $\det M_{\text{sector}} \sim (\varphi_0^{\text{ret}})^{6,9,10}$; the lepton product is $\frac{16}{7} \cdot \frac{4}{3} \cdot \frac{7}{6} = \frac{32}{9} = 2^{g_{\text{car}}} / N_{\text{fam}}^2$). Since (pairwise ratios) + (product) $\Leftrightarrow$ (individuals) is a bijection, the nine charged-fermion amplitudes are *fixed up to one overall scale* v_{geo} . **So U_{point} is not an independent gate: it reduces to v_{geo} , the *same* one irreducible dimensionful anchor as gravity's $1/G$ (v68) — the two $[A]$ anchors collapse to one.**

Research Contract 2 — (G_{metric})

Goal. Construct the TFPT metric measure

$$Z[J] = \lim_{\chi \rightarrow \infty} \int_{\mathcal{G}_{\chi}/\text{Diff}} \exp(-S_{\text{rel},\chi}[g, A, \psi] + J \cdot O) d\mu_{\text{Cal},\chi},$$

$$S_{\text{rel},\chi} = \text{Tr} f\left(\frac{D_{\text{rel}} + A_{\Sigma}}{\chi}\right) - \text{Tr} f\left(\frac{D_{\text{ref}}}{\chi}\right) + \frac{i\pi}{2} \Delta\eta_{\Sigma},$$

the relative spectral action whose low-curvature shadow is the already-closed $R + R^2$ equation.

Lemma 8 (G0–G1 — configuration space & BRST/Hodge Fredholm). *For $s > 5/2$ a Sobolev space of Θ -symmetric metrics on the doubled seam collar M_{Σ} admits a gauge slice; the gauge-fixed gravitational complex $\Omega^0(TM) \xrightarrow{K} \Gamma(S^2 T^*M) \xrightarrow{K^*} \Omega^0(TM)$ with $\Delta_{\text{gh}} = K^*K$ and $\Pi_{\text{phys}} = \Pi_{\text{BRST/Hodge}} \Pi_{TT}$ is elliptic Fredholm; the conformal mode is removed as BRST-exact / non-propagating in the Calderón polarisation (not rhetorically). Machine: principal symbols in xAct-/Cadabra; linear-algebra certificates in Lean (medium).*

Lemma 9 (G2 — finite relative spectral action, *heat-kernel grounded* [N]). $S_{\text{rel},\chi}$ is finite, differentiable, with a local Seeley–DeWitt expansion $\text{Tr} f(D^2/\chi^2) \sim f_4 \chi^4 a_0 + f_2 \chi^2 a_2 + f_0 a_4 + \dots$. For the Lichnerowicz Dirac operator $D^2 = \nabla^2 + \frac{R}{4}$ (so $E = -\frac{R}{4}$, spinor trace $\text{tr} \mathbf{1} = 4$) the Gilkey coefficients give, per $(4\pi)^{-2}$,

$$a_2 = \frac{1}{6} \text{tr}(6E + R \mathbf{1}) = -\frac{R}{3} \quad (\text{Einstein–Hilbert } \int R), \quad a_4|_{R^2} = \frac{1}{360} \text{tr}(180E^2 + 60RE + 5R^2 \mathbf{1}) = \frac{R^2}{72}$$

(the R^2 /Starobinsky term), so the spectral action structurally produces $S_{\text{eff}} = \int \sqrt{g} \frac{\bar{M}_{\text{Pl}}^2}{2} (R + \frac{R^2}{6\bar{M}^2}) + \dots$ with the scalaron mass ratio fixed by the cutoff moment, $M^2/\bar{M}_{\text{Pl}}^2 = 6(4\pi)^2/f_0$. The TFPT closure $M^2/\bar{M}_{\text{Pl}}^2 = c_3^7$ fixes $f_0 = 6(4\pi)^2/c_3^7$, i.e. $M = c_3^{7/2} \bar{M}_{\text{Pl}} = 3.06 \times 10^{13} \text{ GeV}$ — the boundary normalisation c_3 is the gravitational scale — and hence the Starobinsky slow-roll forms $n_s \simeq 1 - \frac{2}{N}$, $r \simeq \frac{12}{N^2}$. The $R + R^2$ structure is convention-independent (standard Gilkey); the precise rational $\frac{1}{72}$ and factor 6 are the standard Dirac conventions used here ([verification/v36_spectral_action_g2.py], [verification/v28_gravity_fR.py]).

The simpler resolution: “full QG” for physics is the gap-decoupled admissible sector [F]

The infinite-dimensional ambient measure (G6) is the summit, but the admissible IR sector does not wait for it. The metric coupling is gap-dominated (G5): $2\|V_{\text{metric}}\| = 0.785 < \Delta = 6 \log \frac{3}{2} = 2.433$, so the admissible sector keeps a *strictly positive* effective gap $\Delta_{\text{eff}} = 1.648 > 0$ under the stated RP and metric-coupling assumptions. A positive gap dynamically *isolates* the admissible sector from the un-built ambient/deep-UV: under those assumptions the admissible (IR) measure is closed, and the ambient completion is decoupled from it (we do

not assert it is physically irrelevant — that would itself be a physical interpretation). So the honest reading is

$$\begin{aligned} \text{admissible IR sector} &= \text{closed under stated RP}; \\ \text{ambient metric measure (G6)} &= G_{\text{metric gate}} \end{aligned}$$

G2 supplies the local action ($R + R^2$, heat-kernel); G5 supplies the decoupling of the admissible sector; together they reduce (G_{metric}) to the ambient projective limit (G6).

Precise status (Alessandro’s boundary): the admissible IR sector is closed under the stated RP and gap-dominance assumptions; the ambient metric projective measure remains a *strict TOE completion target*. Its absence does *not* affect the bounded IR claim, but it *does block certification as a strict physical TOE* — so we do not call it “mere mathematical completeness”. [verification/v36_spectral_action_g2.py]

The glue Q-system: the index-4 object of the G_{net} statement exists explicitly [I]/[P] (verification/v125_glue_qsystem.py)

The G_{net} target — “the seam–Calderón inclusion has Jones index $4 = |\mu_4|$ ” — no longer refers to an unknown object. On the glue group algebra $A = \mathbb{C}[\mathbb{Z}_4]$ (multiplication $\delta_a \otimes \delta_b \mapsto \delta_{a+b}$, unit δ_0) all Longo Q-system axioms hold exactly [I]: associativity, unit law, the Frobenius identity $(m \otimes 1)(1 \otimes m^*) = m^*m = (1 \otimes m)(m^* \otimes 1)$, and specialness $mm^* = |\mathbb{Z}_4| \mathbf{1}$ — a special C^* Frobenius algebra with

$$\dim \theta = 4 = |\mu_4| = [B : A]$$

exactly the Carrier Index Lemma value (KLM: $\mu_{\text{carrier}} = 16 = 4^2 \mu_{E_8}$ re-verified). *Locality = isotropy* [I]: on the Lagrangian glue $\langle(1, 1)\rangle$ the discriminant form gives $q(k(1, 1)) = k^2 \in \mathbb{Z}$ for every k — the braiding obstruction vanishes (v92 in Q-system language); the two Lagrangian glues (the sheet pair) are the only maximal isotropic choices. *Halfway step* [I]: $\{0, 2\} < \mathbb{Z}_4$ closes and gives the unique intermediate Q-system $\mathbb{C}[\mathbb{Z}_2]$ = the $SO(16)_1$ tower step, with index factorisation $4 = 2 \times 2$. *R2 sharpened* [P] (recorded, not claimed): what remains is the operator-algebraic *identification* “the seam–Calderón inclusion of the 16-Majorana net (v113) realises *this* Q-system” — matching two constructed objects, not a search; the three loads (metric + carrier + QBL, v123) hang on this one identification.

The hull is the graded module of the Q-system ([verification/v128_graded_hull.py]).

The Q-system is not merely an abstract companion of E_8 — E_8 realises it [I]: the 240 roots, built as explicit norm-2 vectors in $D_5 \oplus A_3$ +glue coordinates, decompose as $240 = 52 + 64 + 60 + 64$ over the glue $\mathbb{Z}_4$ (the v89 branching as vectors), and for all 6720 root pairs whose sum is again a root, $\text{coset}(r_1 + r_2) = \text{coset}(r_1) + \text{coset}(r_2) \bmod 4$:

$$E_8 \text{ is a } \mathbb{Z}_4\text{-graded Lie algebra over the carrier; the grading group is the glue = the Q-system}$$

— the checksum hull is the Q-system’s graded module, realised in 248 dimensions. The R2 identification thus matches the Calderón inclusion against a structure that E_8 already carries.

Lemma 10 (G3–G4 — reflection positivity, fixed and integrated). *For each Θ -invariant g in the Calderón ball the physical boundary kernel $\mathcal{K}_g = P_{\text{adm}} \Pi_{\text{phys}} \mathcal{N}_g^{TT} \Pi_{\text{phys}} P_{\text{adm}}$ is reflection positive, $\langle \Theta F, F \rangle_{\mathcal{K}_g} \geq 0$ (G3); and $d\mu_{\text{Cal}, \chi}$ is Θ -invariant, supported where G3 holds uniformly, so $\int \langle \Theta F, F \rangle_{\mathcal{K}_g} d\mu_{\text{Cal}, \chi} \geq 0$ (G4). **G4 is the decisive step:** RP after integrating over dynamical metrics, not merely at fixed background.*

Lemma 11 (G5 — gap dominance [F]). $2\|V_{\text{metric}}\|_{\text{rel}} < \Delta = 6 \log \frac{3}{2}$, with $\|V_{\text{metric}}\|_{\text{rel}} \leq 248c_3^2 = \frac{31}{8\pi^2} = 0.39262$, hence $2\|V\| = 0.7852 < 2.4328$ and $\Delta_{\text{eff}} \geq \Delta - 2\|V\| > 1.647 > 0$. The numerical inequality is trivial ([verification/v29_research_contract_certs.py]); the work is the operator-norm bound (hard).

Theorem 1 (Decoupling theorem — physical low-energy claims do *not* need the full G_{metric}). If the admissible sector is reflection-positive (G4) and the metric coupling is gap-dominated, $2\|V_{\text{metric}}\| < \Delta = 6 \log \frac{3}{2}$ (G5), then the admissible IR measure is stable under the ambient completion: a strictly positive effective gap $\Delta_{\text{eff}} = \Delta - 2\|V\| > 0$ protects the admissible sector from the (un-built) ambient/deep-UV, so every physical low-energy read-out (masses, mixings, α^{-1} , the $R+R^2$ regime) is independent of the open ambient measure. Symbolically

$$\boxed{\text{RP} + 2\|V\| < \Delta \implies \text{admissible IR measure stable under ambient completion .}}$$

Thus (G_{metric}) is not a diffuse “full quantum gravity missing” block: it is the sharply delimited ambient projective limit (G6), a strict-TOE certification target whose closure the falsifiable IR physics does not depend on. **Typing:** the inequality is machine-checked [F]; the operator-norm bound on $\|V_{\text{metric}}\|$ is the conditional input.

Lemma 12 (G6–G7 — projective limit & Ward identities). A projective system μ_{χ_n} with $\pi_{m,n*}\mu_{\chi_m} = \mu_{\chi_n}$, uniform moment bounds $\sup_n \int \|g\|_{H^s}^p d\mu_{\chi_n} < \infty$, tightness and regulator-independence yields $\mu_{\text{QG}} = \lim \mu_{\chi_n}$ on $\mathcal{G}/\text{Diff}$ (G6, the hardest step); the limit obeys the BRST Ward identities $\langle s\mathcal{O} \rangle = 0$, whence $\nabla^\mu T_{\mu\nu} = 0$ and the η boundary phase cancels the APS anomalies (G7).

Gate 2 reduction: G6 is a boundary projective limit, not a bulk one
 ([verification/v76_gmetric_reduction.py])

The exact decoupling margin is $\Delta_{\text{eff}} = \Delta - 2\|V\| = 6 \log \frac{3}{2} - \frac{31}{4\pi^2} = 1.648 > 0$, with $\|V_{\text{metric}}\|_{\text{rel}} \leq 248c_3^2 = \dim(E_8)c_3^2 = \frac{31}{8\pi^2}$ ($31 = 2^{g_{\text{car}}} - 1$). **Holographic reduction:** because the seam is a *finite causal boundary*, the ambient *bulk* metric measure is reconstructed from a finite-codimension seam (Calderón) *boundary* measure — so G6 is a *boundary* projective limit. The conditional theorem is then

$$\boxed{\text{RP}(\text{seam-Calderón boundary kernel}) + \text{tightness} \implies G_{\text{metric}} \text{ closes}}$$

i.e. the dimension of the open problem drops from bulk QG to a finite seam-boundary measure. **Residual:** the constructive boundary projective limit (constructive-QFT grade) — reduced, not closed. So Gate 2 is now *two clean tiers*: Tier A (IR physics) closed under RP+gap; Tier B (strict TOE) reduced to a boundary measure, no longer a diffuse “full QG missing”. [P]/[A]

And that boundary measure is a rigorously constructed net ([verification/v77_e8_conformal_net.py]). The level-1 central charges $c = \dim \mathfrak{g}/(1 + h^\vee)$ give $c(E_8)_1 = \frac{248}{31} = 8 = \text{rank } E_8$, $c(D_5)_1 = \frac{45}{9} = 5$, $c(A_3)_1 = \frac{15}{5} = 3$, and $c(D_5) + c(A_3) = 5 + 3 = 8 = c(E_8)$ is a *conformal embedding* $(D_5)_1 \times (A_3)_1 \subset (E_8)_1$ (coset $c = 0$). Now $(E_8)_1$ is the holomorphic $c = 8$ E_8 -lattice VOA / conformal net — one of the most rigorously constructed objects in mathematical QFT (Frenkel–Kac–Segal; Kawahigashi–Longo / Carpi). **So the conditional theorem becomes concrete:** identify the seam-Calderón boundary measure with the $(E_8)_1$ lattice net (carrier = $(D_5)_1 \times (A_3)_1$ subnet), and the gap $\Delta_{\text{eff}} > 0$ supplies the clustering/tightness — so G6 is imported into existing RCFT/conformal-net rigor rather than built from scratch. Residual = the net identification + bulk reconstruction. [I] (charges, embedding) + [P] (identification)

Theorem G — the closure statement [A] (target)

The relative spectral boundary kernel over the diffeomorphism-quotiented metric sector admits a regulator-independent, reflection-positive projective-limit measure μ_{QG} . Its Ward identities imply covariant conservation, and its low-curvature expansion is the TFPT $R + R^2$ seam action

$$f_R R_{\mu\nu} - \frac{1}{2} f(R) g_{\mu\nu} + (g_{\mu\nu} \square - \nabla_\mu \nabla_\nu) f_R = \bar{M}_{\text{Pl}}^{-2} T_{\mu\nu} + O(R^3/M^4),$$

so the closed covariant equation becomes the controlled IR shadow of the full measure. *Status: open; the deepest item. G5 certified; the regime equation closed [verification/v28_gravity_fR.py].*

Certifiability and recommended order

Building block	Machine	Hardness
W_{wall} enumeration ($C_U^{(1)}$)	Lean / Python / Wolfram	easy (<i>done</i>)
D_4 fixed-locus polynomial ($C_U^{(2)}$)	Sage / Mathematica	medium
selector uniqueness ($C_U^{(3)}$)	interval arithmetic	medium
SNF, determinants, spectra	Lean	easy-medium
principal-symbol ellipticity (G1)	xAct / Cadabra / Lean	medium
gap inequality (G5)	Lean	easy (<i>done</i>)
operator-norm bound (G5)	analysis + numerics	hard
projective limit (G6)	not primarily machine	very hard
Ward identities (G7)	BV/BRST formal + checks	hard

Recommended order

$$(U_{\text{wall}}) \rightarrow \text{freeze the frontier status} \rightarrow (G_{\text{metric}})$$

(U_{wall}) can visibly upgrade the theory — it hits the last concrete flavor ambiguity, including the “11” — and it is finite and falsifiable (the U2 kill switch can destroy it cheaply). (G_{metric}) is more fundamental but a *programme*, not a single proof: heavy machinery, with the projective limit (G6) the genuine summit.