

Assignment 1 Question 1

Sheen Thusoo

c. Investigate if $\alpha(\mathcal{P})$ is location-scale invariant, location-scale equivariant, or neither.

For $m > 0$ and $b \in \mathbb{R}$, attribute $\alpha(\mathcal{P})$ is location-scale invariant if

$$\alpha(my_1 + b, \dots, my_N + b) = \alpha(y_1, \dots, y_N)$$

and is location-scale equivariant if

$$\alpha(my_1 + b, \dots, my_N + b) = m\alpha(y_1, \dots, y_N) + b$$

To begin,

$$y'_u = my_u + b$$

and

$$\bar{y}' = m\bar{y} + b$$

Thus,

$$\alpha(my_1 + b, \dots, my_N + b) = \frac{\frac{1}{N} \sum_{u \in \mathcal{P}} (my_u + b - m\bar{y} + b)^4}{[\frac{1}{N} \sum_{u \in \mathcal{P}} (my_u + b - m\bar{y} + b)^2]^2} - 3$$

$$\alpha(my_1 + b, \dots, my_N + b) = \frac{\frac{m^4}{N} \sum_{u \in \mathcal{P}} (y_u - \bar{y})^4}{[\frac{m^2}{N} \sum_{u \in \mathcal{P}} (y_u - \bar{y})^2]^2} - 3$$

$$\alpha(my_1 + b, \dots, my_N + b) = \frac{m^4 \frac{1}{N} \sum_{u \in \mathcal{P}} (y_u - \bar{y})^4}{m^4 [\frac{1}{N} \sum_{u \in \mathcal{P}} (y_u - \bar{y})^2]^2} - 3$$

$$\alpha(my_1 + b, \dots, my_N + b) = \frac{\frac{1}{N} \sum_{u \in \mathcal{P}} (y_u - \bar{y})^4}{[\frac{1}{N} \sum_{u \in \mathcal{P}} (y_u - \bar{y})^2]^2} - 3$$

$$\alpha(my_1 + b, \dots, my_N + b) = \frac{\frac{1}{N} \sum_{u \in \mathcal{P}} (y_u - \bar{y})^4}{\sigma^4} - 3$$

$$\alpha(my_1 + b, \dots, my_N + b) = \alpha(y_1, \dots, y_N)$$

Therefore, the excess kurtosis attribute is location-scale invariant.