

Elevated seawater temperatures have a limited impact on the coral immune response following physical damage

Jeroen A. J. M. van de Water · William Leggat ·
David G. Bourne · Madeleine J. H. van Oppen ·
Bette L. Willis · Tracy D. Ainsworth

Received: 15 April 2014 / Revised: 15 February 2015 / Accepted: 2 March 2015 / Published online: 17 March 2015
© Springer International Publishing Switzerland 2015

Abstract Recurrent disturbances on coral reefs that cause injuries, like predation and storm damage, and elevated seawater temperatures reduce coral fitness and immunocompetence. An effective immune response is essential to prevent disease and enhance colony survival. To evaluate how elevated seawater temperatures affect the coral immune response following injury, fragments of *Acropora aspera* were exposed to ambient (27–29°C) or elevated (32–33.5°C) seawater temperatures for 8 days and subsequently experimentally injured. Expression patterns for 15 immune genes 24 h post-injury revealed that most genes involved in the Toll-like receptor pathway were unaffected by elevated seawater

temperatures. Exceptions to this pattern were *cFos* and *cJun*, which were upregulated and likely played a role in repair processes, and *TRAF-6* and *NFκB*, which were downregulated suggesting reduced immune function. Components of the complement system were upregulated (*millelectin*, *C3*) or downregulated (*Bf*, *Tx60*, *apexrin*) in corals at high temperatures. However, corals that also sustained injury, showed normal *Tx60* and *apexrin* expression, suggesting roles in the wounding response. Overall, basal expression levels of immune genes are sufficient to mount a response to injury in the short term, and the immune response of *A. aspera* following injury is not significantly affected by minor elevations in seawater temperatures.

Guest editors: Yehuda Benayahu, Oren Levy & Tamar Lotan /
Coelenterate Biology: Advanced Studies on Cnidaria and
Ctenophora

J. A. J. M. van de Water (✉) · W. Leggat ·
M. J. H. van Oppen · B. L. Willis · T. D. Ainsworth
ARC Centre of Excellence for Coral Reef Studies, James
Cook University, Townsville, QLD, Australia
e-mail: jeroen.vandewater@my.jcu.edu.au

J. A. J. M. van de Water · B. L. Willis
College of Marine and Environmental Sciences, James
Cook University, Townsville, QLD, Australia

J. A. J. M. van de Water · D. G. Bourne ·
M. J. H. van Oppen · B. L. Willis
AIMS@JCU, James Cook University, Townsville, QLD,
Australia

Keywords Coral · Immunity · Heat stress · Injury ·
Gene expression

J. A. J. M. van de Water · D. G. Bourne ·
M. J. H. van Oppen
Australian Institute of Marine Science, Townsville, QLD,
Australia

W. Leggat
College of Public Health, Medical & Veterinary Sciences,
James Cook University, Townsville, QLD, Australia

Introduction

Physical injury presents a major challenge to an organism's immune system, providing a point-of-entry for pathogens and requiring energy for tissue repair. On coral reefs, as many as 21–56% of coral colonies are commonly damaged (Lindsay, 2010). Predictions of increasing severity and/or frequency of impacts that cause loss of tissue integrity, including physical damage following severe tropical storms and dredging associated with escalating coastal developments, as well as more localised impacts (e.g. destructive fishing techniques, anchoring, reef-based tourism activities and entangled fishing line) (Lamb & Willis, 2011; Lamb et al., 2014; Pollock et al., 2014), highlight the importance of understanding the capacity of the coral's immune system to respond to injury. In addition, demands on a coral's energy budget associated with climate change-related stressors, particularly ocean warming and acidification, are likely to increase in coming decades. Given the important role of the immune system in wound healing, understanding how the coral's immune response to injury is likely to be affected by additional stressors is becoming increasingly important.

Maintaining tissue integrity is of vital importance for the maintenance of fitness in any animal, including corals. To re-establish tissue integrity following injury, corals have a wound healing process similar to those in higher organisms (Palmer et al., 2011b). In this process, the immune system aids in wound sealing, elimination of microbes and tissue regeneration by activating transglutaminase and phenoloxidase and recruiting immune cells to the lesion site (Palmer et al., 2011a, b, 2012). In addition, molecular studies have revealed that corals possess components of a range of innate immune mechanisms that are involved in eliminating potential harmful microbes and maintaining healthy microbial communities in other invertebrates (Fraune & Bosch, 2007; Franzenburg et al., 2012, 2013). Various genes encoding the microbe-associated molecular patterns (MAMP)-detecting Toll-like receptors (TLR) and nucleotide-binding oligomerisation domain (NOD)-like receptors (NLR) have been reported in corals, as well as components of their respective downstream signalling pathways, such as the JNK, MAPK p38, ERK-2 and NF- κ B pathways (Miller et al., 2007; Shinzato et al., 2011; Hamada et al., 2013) (Fig. 1I).

Signalling via these pathways results in the transcription of genes involved in the regulation of an immune response, as well as cell proliferation (Fig. 1II). The existence of genes involved in these pathways in corals suggests that they may play a significant role in their immune responses following wounding. Nevertheless, few studies have investigated the functions of various homologues and their roles in innate immunity, defence and disease in corals, and therefore little is known about the molecular mechanisms that regulate the injury-induced immune response.

Discovery of various components of the complement system in corals, including complement C3, Factor B (Bf), lectins and mannose-binding lectin (MBL)-associated serine protease (MASP) (Miller et al. 2007, Kvennefors et al., 2010, Shinzato et al., 2011), suggests that corals also possess a functional complement system (Fig. 1III). Binding of an MBL to the microbial cell wall results in the activation of MASPs, which subsequently cleave complement C3 and Bf (Cerenius et al., 2010) (Fig. 1III). Together, C3 and Bf form a C3-convertase that hydrolyses C3 into C3a and C3b. Whereas C3a functions as a chemotactic cytokine, C3b is deposited on the microbial surface (Fig. 1III), promoting phagocytosis of the microbes by C3a-recruited immune cells (Cerenius et al., 2010) (Fig. 1IV). Moreover, it initiates the formation of a microbicidal membrane attack complex (MAC) (Cerenius et al., 2010) (Fig. 1V), for which several candidate MAC/PF (membrane attack complex/perforin) domain-containing genes have been identified in *Acropora millepora* (Miller et al. 2007). Studies of two complement system proteins in corals have shown that the MBL millelectin agglutinates bacteria and is upregulated upon challenge with immunostimulants (Kvennefors et al., 2008). In contrast, complement C3 was upregulated following injury (Kvennefors et al., 2010). The presence of multiple components of the TLR pathway and complement system in corals suggests that they possess complex wound healing and immune systems, similar to those seen in higher organisms, although few studies have focused on functional responses of these pathways.

Various environmental and biological factors are known to significantly affect the process of wound healing in corals, including elevated nutrient levels (Renegar et al., 2008; Denis et al., 2013), sedimentation (Mascarelli & Bunkley-Williams, 1999; Cróquer




Fig. 1 Toll-like receptor signalling pathways and the complement system. (I) Activation of Toll-like receptors result in the activation of various signal transduction pathways, including: (1) signalling via TRAM leads indirectly to the activation of TBK1, which relays the signal through to IRFs and NF-κB; (2) signalling via the MyD88-dependent pathway results in the activation of TRAF6 and ECSIT which activate the ERK-2, JNK and MAPK p38 pathways. Through activation of Tak1 and MEKK1, TRAF6 also indirectly activates the NF-κB pathway via the IKK complex. In addition, TRAF6 may induce apoptosis via a caspase-8-mediated pathway. NLRs indirectly activate MEKK1 thereby activating the ERK-2, JNK, MAPK p38 and NF-κB pathways. (II) Signalling results in the transcription of immune response regulatory and effector genes as well as genes involved in cell survival and apoptosis. (III) The complement system is initiated following the binding of lectins to microbes, resulting in the conversion of C3 into C3b and C3a. In addition, Bf is activated and forms the C3b:Bf complex, an efficient C3 convertase. (IV) Lectins and C3b promote phagocytosis of the targeted microbe leading to its destruction. (V) C3b also induces the formation of the membrane attack complex (MAC) that is

inserted into the membrane and cause lysis of the microbe. *TLR* Toll-like receptor, *MyD88* myeloid differentiation primary response, *IRAK* interleukin-1 receptor-associated kinase, *TRAF* TNF receptor-associated factor, *TNF* tumor necrosis factor, *ECSIT* evolutionarily conserved signalling intermediate in Toll pathways, *TAK1* transforming growth factor β-activated kinase-1, *MEKK* MAPK/ERK kinase kinase, *MKK* MAPK kinase, *MAPK* mitogen-activated protein kinase, *ERK* extracellular signal-regulated kinase, *JNK* c-Jun N-terminal kinase, *CREB* cAMP response element-binding protein, *AP-1* activator protein-1, *NLR* nucleotide-binding oligomerisation domain (NOD-like) receptor, *RIP2* receptor-interacting serine/threonine-protein kinase-2, *IKK* I-kappa-B kinase, *NF-κB* nuclear factor kappa B, *TRAM* TRIF-related adaptor molecule, *TRIF* TIR-domain-containing adapter-inducing interferon-β, *TIR* Toll/interleukin-1 receptor, *TBK1* TANK-binding kinase, *TANK* TRAF-family member-associated NFκB activator, *IRF* interferon regulatory factor. *Bf* factor B, *C3* complement C3. Asterisk (*) indicates genes that are included in the gene expression analysis in this study

et al., 2002), light intensity (Titlyanov & Titlyanova, 2009), cyanobacteria (Titlyanov et al., 2005) and algae (Mascarelli & Bunkley-Williams, 1999; Titlyanov et al., 2005), and the climate change-related variables pCO₂ and temperature. Increases in both pCO₂ levels and seawater temperatures have been shown to reduce wound healing rates (Hess et al., 2004; Henry & Hart, 2005; Renegar et al. 2008; Edmunds & Lenihan, 2010;

Lenihan & Edmunds, 2010; Denis et al., 2013). For example, corals that partially bleached following exposure to high temperatures healed at significantly lower rates than controls, and corals that fully bleached were unable to regenerate following injury (Meesters et al., 1997; Mascarelli & Bunkley-Williams, 1999; Fine et al. 2002). In contrast, regeneration capacity may also increase due to higher metabolic rates

associated with a Q_{10} effect (Kramarsky-Winter & Loya, 2000). Increasing seawater temperatures predicted for the coming decades could have dire consequences for corals living near their thermal thresholds (Berkelmans & Willis, 1999), with large scale losses of coral reefs predicted (Hoegh-Guldberg et al., 2007). In addition, as seawater temperatures continue to rise, elevated seawater temperatures have also been linked to the recent rise in coral disease (Willis et al., 2004; Sokolow, 2009; Ruiz-Moreno et al., 2012). Shifts in coral-associated bacterial communities (Ritchie, 2006; Bourne et al., 2008; Mouchka et al., 2010; Littman et al., 2011; Witt et al., 2011), increases in coral pathogen virulence and a change in coral immune function (Mydlarz et al., 2008; Palmer et al., 2011a; Vidal-Dupiol et al. 2011a, b) have all been demonstrated to occur following exposure to elevated seawater temperatures. However, the combined effects of heat stress and injury on the various biochemical components and pathways involved in the coral's immune response are still largely unknown.

In this study, we used a manipulative aquarium-based experiment to assess the impacts of elevated seawater temperatures on the coral immune response exhibited in the first 24 h following injury. In particular, we analysed the expression of several genes involved in the TLR pathway and complement system, to determine how the short-term wounding-induced immune response of corals is compromised by heat stress.

Material and methods

Experimental design and sampling

Branches (approximately 10 cm in length, $n = 56$) of the scleractinian coral *Acropora aspera* were collected from 3 large patches of *A. aspera* (approximately 100 m²) across a 50 m length of the reef flat of Heron Island, Great Barrier Reef, Australia (23°26'31.20"S, 151°54'50.40"E) and transported in seawater to holding tanks at the Heron Island Research Station. Branches were held upright in 4 racks ($n = 14$ fragments per rack) and acclimated at ambient reef flat temperatures [$27.1 \pm 1.3^{\circ}\text{C}$ (mean $\pm$ SE)] for a period of 7 days in flow-through seawater. Racks were randomly assigned to 50 litre aquaria ($n = 4$) (1 rack per aquarium) supplied with seawater at ambient

temperatures. Daytime seawater temperature in two aquaria was gradually increased by 1°C per day over 4 days using thermostat-controlled heaters until the desired sub-lethal heat stress temperature of 32°C was reached. To simulate day-night temperature fluctuations on the reef flat, heaters were turned on at 07:00 and off at 19:00 h each day. Therefore, temperatures in the elevated treatment tanks fluctuated between 27 and 33.5°C (mid-day maximum because of solar heating) each day once the target temperature was achieved. This temperature regimen has previously been described to induce heat stress responses in *Acropora aspera* (Leggat et al., 2011; Ainsworth et al., 2011). Similarly, because of natural fluctuations in daily ambient temperatures of seawater supplied from the reef, corals in the control temperature treatment were exposed to temperature fluctuations of $26\text{--}28.5^{\circ}\text{C}$ (the maximum daily water temperature on the reef flat at time of collection). After 8 days in each temperature treatment, half of the coral fragments in each tank were injured ($n = 7$ uninjured; $n = 7$ injured per aquaria) by removing tissue using a high-pressure airgun at 15.00 h. Lesions comprised a

Fig. 2 Representative image of a branch of *Acropora aspera*. Injuries of approximately 4 cm in length were artificially inflicted using an airgun



strip of exposed skeleton approximately 1 cm wide along a third of the length of each branch (~4 cm) (Fig. 2). Whole branches were collected 24 h post-injury and snap-frozen in liquid nitrogen.

Messenger RNA isolation

Frozen samples of each experimental branch of *A. aspera* were crushed in a stainless steel mortar and pestle, pre-chilled with liquid nitrogen, using a hydraulic press. Messenger RNA (mRNA) was isolated from the crushed coral using the Dynabeads mRNA DIRECT kit (Invitrogen Dynal AS, Oslo, Norway) according to a modified protocol based on the manufacturer's recommendations. In summary, approximately 100 mg of crushed coral was added to 400 µl lysis buffer, incubated on a vortex at low speed for 7 min and centrifuged for 2 min at 12,000×g. The supernatant was added to oligo(dT)-Dynabeads (pre-washed in lysis buffer) and incubated on the vortex at medium speed for 8 min to allow mRNA annealing. Tubes were placed on a DynaMag-2 magnetic particle concentrator for 5 min and supernatant was removed. Using the DynaMag-2, oligo(dT) Dynabead/mRNA complexes were washed twice with 300 µl of Buffer A and subsequently twice with 400 µl of Buffer B. Complexes were resuspended in 27 µl ice-cold 10 mM Tris–HCl, incubated at 80°C for 2 min and rapidly cooled down on ice. Oligo(dT)-Dynabeads were concentrated on the DynaMag-2 and mRNA-containing supernatant was collected and stored at –80°C until use.

Quantification of gene expression

Expression levels of 20 immune system-related genes (Siboni et al., 2012) and 4 reference genes (Csaszar et al., 2009; Seneca et al., 2010) (Table 1) were analysed using the GenomeLab GeXP Start Kit and the CEQ-8800 Genetic Analysis System (Beckman-Coulter, Brea, CA, United States of America) following the protocols described in Siboni et al. (2012) with minor modifications. In summary, cDNA was generated from 9 ng of mRNA. Reverse primer concentrations were optimised for the multiplex to ensure signals in the electropherogram were within the CEQ-8800 detection range: 500 nM for *TRAM*, *TIR-1*, *MEKK-1*, *TRAF6*, *Tx60*, *ERK2*, *Bf*, *MAPK p38*, *Millectin*, *ctg_1913*, *Apextrin*, *cJun*, *HL-1*, *HL-2*,

HL-3, *CTL-1* and *CTL-2*; 250 nM for *GAPDH* and *RPL9*; 62.5 nM for *NF-κB* and *C3/A2M-2*; 12.5 nM for *cFos* and *ATF4/5*; 3.75 nM for *RPS7*. Forward primer concentrations were 200 nM. All details for genes of interest and primer sequences can be found in Table 1. Prior to loading on the CEQ-8800 Genetic Analysis System, PCR products were pre-diluted 1:20. Data were filtered and analysed using the GeXP Fragment Analysis module and the eXpress Profiler software (Beckman-Coulter, Brea, CA, United States of America) (Souter et al., 2011). Gene expression levels were normalised to the internal control (Kan^R) and to the geometric mean of the expression levels of three reference genes (*RPL9*, *RPS7* and *ctg_1913*) selected using geNorm (Vandesompele et al., 2002). Data were obtained for three technical replicates per sample.

Statistical analysis

For each gene, expression data were analysed for differences in expression using a two-factor analysis of variance (ANOVA), where health status (healthy versus injured) and temperature (ambient versus elevated) were both treated as fixed factors. Differences between treatments were analysed using Tukey's HSD post hoc tests. Differences were considered significant when $p < 0.05$ (ANOVA) or when the value 0 did not fall within the 95% confidence interval (Tukey's HSD). All analyses were performed in the statistical software package S-PLUS 8.0.

Results

Genes involved in the Toll-like receptor pathway were largely unaffected by injury or heat stress in the first 24 h, with no significant differences in expression patterns found between health states or temperature treatments for *TIR-1*, *MAPK p38*, *MEKK-1* and *ERK-2* (Fig. 3A, C–E). In contrast, injured corals had significantly lower *TRAF-6* expression compared to uninjured corals at ambient temperatures, although expression levels did not differ between health states under mild heat stress (Fig. 3B). The transcriptomic response of both *ATF4/5* to injury was dependent on seawater temperature, being downregulated at ambient temperatures, but upregulated in the elevated temperature treatment as evidenced by a significant

Table 1 List of immune genes

Gene name	Acronym	Accession number	PCR product length	Sequence of primer coding region	References
<i>Reference gene</i>					
Glyceraldehyde 3 phosphate dehydrogenase	<i>GAPDH</i>	EZ026309	187	F: AGGTGGAGCCAAGAAAGTCA R: TTAGCTAGAGGAGCCAGGCA	Csaszar et al. (2009) and Seneca et al. (2010)
Ribosomal protein S7	<i>RPS7</i>	EZ031290	197	F: CAGGCATGCTTACAACCAAA R: TCAACCTCCTTTGCTCCAGT	
Ribosomal protein L9	<i>RPL9</i>	EZ026324	292	F: CGTGTAACGTTGTGGTTTGC R: TTTGACACCTGAATTCGCAC	
Unknown transcript	<i>Ctg_1913</i>	EZ040581	280	F: GATTTAACCACCGGCAGTGT R: ATGGTAGGGAGGAGGCTGTT	
<i>Toll-like receptor pathway</i>					
Toll interleukin receptor	<i>TIR-1</i>	EF090256	137	F: AAAGCCGCAGTCATCAGTTT R: GAAATTGGCGTTGAATTCGT	Miller et al. (2007)
TNF receptor-associated factor 6	<i>TRAF-6</i>	DY583189	127	F: TGATGAATGTCCTTTCGCAG R: ACATGCTTTGCAAGCTGATG	
Mitogen-activated protein kinase/ERK kinase kinase-1	<i>MEKK-1</i>	DY581208 DY581138 DY582675	117	F: CTGCGGATATTTGGTCTCTGT R: TTTCTTTGTGCGTTGATCCC	
Extracellular signal-regulated kinase	<i>ERK-2</i>	EZ025389	217	F: CCAAAGGTTACAGCAAGGCT R: TGCGTGCCTTTTCATTCATA	
Mitogen-activated protein kinase p38	<i>MAPK p38</i>	EZ031759	237	F: AAAATCAGCAGTGAATCCGC R: TCGGGGTCTGAATACGTAGC	
Component of AP-1 transcription factor	<i>cFos</i>	EZ016042	177	F: CTGGAAAGAGAATTGCTGGC R: GACGATTGCACTTCGGACTT	
Component of AP-1 transcription factor	<i>cJun</i>	EZ020860	366	F: TCGATCGAAGGGACAGTTCT R: GTGCTAGTTGCGGTGTTCAA	
Activating transcription factor 4	<i>ATF4/5</i>	DY577805	247	F: GGCCAGAACGTATACCAAT R: TCTTCGAAATCAAACCCCTG	
Nuclear factor kappa B	<i>NF-kB</i>	G0000491 G0002043 DY582971 DY580118	152	R: CTCATATGCAGGTTGGTGGA F: GATGTTGCAGGCTCAGTTCA	
TRIF-related adaptor molecule	<i>TRAM</i>	EZ047194	157	F: AAGCTAACGGCTCACCAAGA R: TGTGCCATGCACAAGAAAAAT	
<i>Alternative complement pathway</i>					
Complement factor B	<i>Bf</i>	GO001635	227	F: TTATCCATCCCGACGCTAAC R: AGGATCATCTTTTCCTGCGA	Kimura et al. (2009)
Complement C3	<i>C3</i>	EF090257	167	F: CCGCTACACGCTAGACAACA R: CCGCAGAGTCGATGTACAAA	
<i>Lectins</i>					
Mannose-binding lectin	<i>millectin</i>	EU717895	257	F: AGCGAGTATCCACAACACCC R: GGCTTTTTTCGATGTTTCCA	Kvennefors et al. (2008)

Table 1 continued

Gene name	Acronym	Accession number	PCR product length	Sequence of primer coding region	References
C type lectin-1	<i>CTL-1</i>	GO001638	312	F: GGGTTGTGTACAACGGCTTT R: CTTTCCATTCGGTTCTCCTG	Grasso et al. (2008) and Grasso et al. (2011)
C type lectin-2	<i>CTL-2</i>	GS01UH10	267	F: CAGGTCTGGATCGGACTCAT R: CATGTCCAGTGGTTGTACGC	
Hemolytic lectin-1	<i>HL-1</i>	EU863776	335	F: TTCGCTCCAGAGGGAACTA R: GCAGAAATGCCTTTGGTTGT	Grasso et al. (2008, 2011)
Hemolytic lectin-2	<i>HL-2</i>	EU863776	302	F: AACAGTTGAGATAACCGCCG R: TTGATTCCTGGTGCATTGA	
Hemolytic lectin-3	<i>HL-3</i>	EU863777	379	F: TTCTGGAGATTGGGTAACGC R: TCGTTCTCAGCGTGTGTTC	
<i>Membrane attack complex/perforin</i>					
Apical extracellular protein	<i>Apextrin</i>	EF091848	352	F: GGATTCGTACCAAAAAGGCA R: GAGGGGTCTGATATGGGGTT	Miller et al. (2007)
60 kDa proteinaceous toxin	<i>Tx60-A</i>	DY579588	207	F: TACTGCCCTTGAGGTTTGCT R: CTGAAAATCCCGCTGACTGT	

At the 5'-end, all primers contained universal fluorescent tags. Forward primer: AGGTGACACTATAGAATA. Reverse primer: GTACGACTCACTATAGGGA

interaction effect between temperature and injury (Fig. 3B, F). The transcription factors *cFos* and *cJun* were significantly upregulated in response to injury, by up to 1.6-fold for *cJun* (Fig. 3G, H). Moreover, exposure to high temperatures caused further increases in expression of *cJun* in injured corals in comparison to controls (Fig. 3H). *NF-κB* was downregulated 24 h post-injury, but this downregulation was only significant when corals were exposed to high seawater temperatures (Fig. 3I).

Expression levels of genes involved in the complement system were affected by both elevated seawater temperatures and injury. Corals downregulated Factor B (*Bf*) by 1.2-fold, as well as the MAC/PF domain-containing genes *apextrin* and *Tx60* by 2.2- and 2.1-fold, respectively, when exposed to elevated temperatures (Fig. 4C, E–F). Interestingly, no difference in *apextrin* and *Tx60* expression was observed between corals injured under heat stress compared to injury at ambient temperatures (Fig. 4E–F). The lectins *millelectin* and *HL-2* were both significantly upregulated under heat stress, but expression was not altered following injury (Fig. 4A, B). Although, complement C3 was upregulated in response to heat stress for both healthy and injured corals (Fig. 4D),

expression levels were significantly lower in injured corals relative to those that were uninjured. Expression of *HL-1*, *HL-3*, *CTL-1*, *CTL-2* and *TRAM* was undetectable in all coral samples, potentially due to low expression levels. All results from statistical analyses can be found in Table 2.

Discussion

Studies of the wounding response in invertebrates have addressed the cellular and molecular processes in tissue regeneration, but the effects of confounding environmental stressors on the immune response are largely unknown. Here, we demonstrate that components of the Toll-like receptor pathway and complement system involved in the wounding response are robust to mild increases in seawater temperatures in *A. aspera* 24 h post-injury.

Toll-like receptor pathway

In the current study, we found that exposure of corals to mild heat stress for 8 days did not significantly affect the regulation of the TLR pathway, except for an




Fig. 3 Relative expression of genes involved in the Toll-like receptor pathway in injured and uninjured corals under ambient or elevated seawater temperatures. **A** *TIR-1*, **B** *TRAF-6*, **C** *MEKK-1*, **D** *MAPK p38*, **E** *ERK2*, **F** *ATF4/5*, **G** *cFos*, **H** *cJun* and **I** *NF-κB*. Capital letters (A, B, C) indicate

experimental treatment groups that are statistically similar. For figures without letters, no significant differences were detected among the four treatment groups. The asterisk (*) identifies the genes for which expression levels differed between the two health states (but not between the two temperature treatments)

increase in *cJun* expression. Injury, on the other hand, resulted in upregulation of both *cJun* and *cFos*, whereas *TRAF-6* was downregulated. Interestingly, the combination of injury and heat stress increased

cJun expression in corals further, suggesting that it is involved in both the thermal stress response and wound repair. The transcription factors *cJun* and *cFos* together form the AP-1 transcription factor complex




Fig. 4 Relative expression of genes involved in the complement system in injured and uninjured corals under ambient or elevated seawater temperatures. **A** *HL-2*, **B** *Millectin*, **C** *Bf*, **D** complement *C3*, **E** *Apextrin* and **F** *Tx60*. Capital letters (A,

B) indicate experimental treatment groups that are statistically similar. The asterisk (*) identifies the gene for which expression levels differed between the two temperature treatments (but not between the two health states)

(Hess et al., 2004). AP-1 is involved not only in the TLR pathway, but also plays a role in the regulation of cell proliferation, differentiation and apoptosis. These processes are essential for sealing lesions and development of tissue layers and polyp structures, and are regulated via different signal transduction pathways (Hess et al., 2004). Since no other genes in the TLR pathway were upregulated following injury, our interpretation is that *cJun* and *cFos* are primarily involved in lesion regeneration rather than an immune response in the initial stages following injury.

Although no major changes in the expression of genes involved in the TLR pathway were detected under mild heat stress, injured heat-stressed corals did have reduced levels of NF- κ B. Because the TLR signalling pathway regulates the expression of anti-microbial peptides (AMP) primarily via NF- κ B, these

corals may have reduced AMP production and may thus be more susceptible to disease. Although no AMP has been identified in *Acropora* yet, AMP-encoding genes are present in scleractinian corals and other cnidarians (Fraune & Bosch, 2007; Fraune et al., 2011; Vidal-Dupiol et al. 2011a; Burge et al., 2013; Vidal-Dupiol et al., 2014). Studies on the (septic) wounding response in a range of invertebrates (Fehlbaum et al., 1994; Rodriguez de la Vega et al., 2004; Pujol et al., 2008; Xu et al., 2010) have also shown upregulation of AMPs. The limited transcriptomic response of genes involved in the TLR pathway under heat stress observed in this study could indicate that these corals were capable of maintaining stable bacterial communities under these conditions. While no evidence of active regulation of bacterial community composition by corals has been presented to date, AMPs have been

Table 2 Overview of statistical analysis. Statistically significant differences are indicated in italics for the main ANOVA results

Gene	ANOVA	<i>p</i> value	Tukey's HSD	95 % CI	
Apextrin	Temperature	<i>0.0074</i>	Ambient healthy—32°C healthy	0.0147	0.9240
	Injury	<i>0.0234</i>	Ambient injured—32°C healthy	0.1790	1.0400
	Temperature: Injury	0.2128			
ATF4/5	Temperature	0.3668			
	Injury	0.6372			
	Temperature: Injury	<i>0.0133</i>			
Bf	Temperature	<i>0.0323</i>			
	Injury	0.1765			
	Temperature: Injury	0.3414			
C3	Temperature	<i>0.0002</i>	Ambient healthy—ambient injured	0.0971	1.7200
	Injury	<i>0.0007</i>	Ambient healthy—32°C healthy	−1.7700	−0.0593
	Temperature: Injury	0.8464	Ambient injured—32°C healthy	−2.6400	−1.0100
			Ambient injured—32°C injured	−1.8100	−0.1830
cFos	Temperature	0.7165			
	Injury	<i>0.0196</i>			
	Temperature: Injury	0.3523			
cJun	Temperature	<i>0.0003</i>	Ambient healthy—ambient injured	−0.4250	−0.0520
	Injury	<i>0.0001</i>	Ambient healthy—32°C healthy	−0.4280	−0.0351
	Temperature: Injury	0.8737	Ambient healthy—32°C injured	−0.6810	−0.2880
			Ambient injured—32°C injured	−0.4330	−0.0601
			32°C healthy—32°C injured	−0.4500	−0.0568
ERK-2	Temperature	0.5531			
	Injury	0.4315			
	Temperature: Injury	0.4086			
HL-2	Temperature	<i>0.0001</i>	Ambient healthy—32°C healthy	−1.5700	−0.0754
	Injury	0.1499	Ambient injured—32°C healthy	−1.9200	−0.5010
	Temperature: Injury	0.4899	Ambient injured—32°C injured	−1.7900	−0.3640
MAPK p38	Temperature	0.8432			
	Injury	0.6897			
	Temperature: Injury	0.8290			
MEKK-1	Temperature	0.2473			
	Injury	0.1549			
	Temperature: Injury	0.7194			
Millectin	Temperature	<i><0.0001</i>	Ambient healthy—32°C healthy	−2.8000	−0.7920
	Injury	0.1680	Ambient healthy—32°C injured	−2.2500	−0.2410
	Temperature: Injury	0.4201	Ambient injured—32°C healthy	−2.9100	−1.0000
			Ambient injured—32°C injured	−2.3600	−0.4510
NFκB	Temperature	0.3367	Ambient healthy—32°C injured	0.0266	1.2200
	Injury	<i>0.0060</i>			
	Temperature: injury	0.7836			
TIR-1	Temperature	<i>0.0556</i>			
	injury	0.3996			
	Temperature: injury	0.5339			

Table 2 continued

Gene	ANOVA	<i>p</i> value	Tukey's HSD	95 % CI	
TRAF6	Temperature	0.4919	Ambient healthy—ambient injured	0.3880	1.2000
	Injury	0.0003	Ambient healthy—32°C injured	0.1700	1.0300
	Temperature: Injury	0.0086			
Tx60	Temperature	0.0003	Ambient healthy—32°C healthy	0.0673	0.5040
	Injury	0.5243	Ambient injured—32°C healthy	0.0742	0.4890
	Temperature: Injury	0.4509			

The 95 % CI is given for pair-wise comparisons where the difference was considered statistically different (i.e. it did not contain the value 0)

shown to be crucial in maintaining healthy tissue-associated bacterial communities in the cnidarian *Hydra* (Fraune & Bosch, 2007; Franzenburg et al., 2012, 2013).

In summary, we were unable to establish a link between the TLR pathway and the response to injury in *A. aspera*. As our study is preliminary, consisting of a single post-injury time point, potential earlier or later changes in gene expression levels may have been overlooked. Based on this study, the TLR pathway in *A. aspera* was uncompromised, suggesting that the coral has good potential to respond to injury under mild heat stress. However, understanding the impact of heat stress on the full dynamics of the wounding-induced immune response in corals requires study over longer time-frames and with greater sampling resolution.

Complement system

The expression of genes involved in the complement system was affected by both elevated seawater temperatures and injury. Whereas heat stress caused downregulation of the MAC/PF domain-containing *Tx60* and *apextrin*; at ambient temperatures, expression of *Tx60* and *apextrin* was similar between healthy and injured corals. This suggests that not only are *Tx60* and *apextrin* involved in the wounding response, but also that heat-stressed corals are capable of inducing an immune response similar to those at ambient temperatures. Our results are in line with a previous study on the bumble bee *Bombus terrestris*, where upregulation of genes encoding MAC proteins were observed following injury (Erler et al., 2011). The function of these genes in the recovery process in corals remains to be determined; however, the

presence of a MAC/PF domain suggests a role in the anti-microbial response and could indicate infection of the lesion by microbes. The involvement of *Tx60* and *apextrin* in the wounding response, but the absence of upregulation in expression at ambient temperatures in response to injury, indicates that the initial gene expression levels under normal conditions may be sufficient to prevent infection. This suggests that normal expression levels for other genes may also be sufficient, and that potentially, the immune protein expression at 24 h post-injury is sufficient for an appropriate response. Alternatively, protein expression may be primarily regulated through increased mRNA stability (Wu & Brewer, 2012) rather than an increased transcriptomic response. This would enable the coral to direct its limited resources towards wound sealing and regeneration in the initial stages of the wounding response, although further studies are required to address this question.

The upregulation in expression of complement C3 and millectin observed in response to 8 days of exposure to elevated seawater temperatures may indicate that these heat-stressed corals were directing resources to preserve the coral-*Symbiodinium* endosymbiosis. This is in accordance with previous studies that have implicated these proteins in the maintenance of the coral-*Symbiodinium* endosymbiosis in *Acropora millepora* (Kvennefors et al., 2008, 2010). The downregulation of *Bf* under heat stress suggests that this gene, which is a key player in the alternative complement pathway, may not be involved in the activation of C3 for the maintenance of symbiosis. This confirms a role for the lectin-complement pathway, with millectin as the MBL. In addition, our results for *millectin* are consistent with reports that injury does not change expression of this gene

(Kvennefors et al., 2010). However, we show reduced C3 expression 24 h post-injury, which is in contrast to results from a recent study showing upregulation of C3 within 12 h after corals were injured (Kvennefors et al., 2010). Although this may appear inconsistent, very little is known about expression dynamics of immune genes in corals, and the 12-h time difference between sampling in the two studies could explain differences in results. Investigations into expression levels over time will be required to gain more insight into these dynamics.

Conclusion

In this study, we demonstrate that the immune response of *A. aspera* exhibited 24 h post-injury is not significantly affected by mild temperature stress. While heat stress did influence the coral's immune effector capacity, responses to injury under heat stress were as strong as those of non-heat-stressed corals. Gene expression levels of both the TLR pathway components and other immune effectors were largely unaffected at 24 h post-injury, while genes potentially involved in wound repair and tissue regeneration were upregulated. In addition, upregulation of components of the lectin-complement system, which is involved in the coral-*Symbiodinium* symbiosis, under heat stress suggests that corals were attempting to maintain the endosymbiosis. Normal expression levels of immune genes in the initial response following injury may be sufficient to ward off invading microbes, enabling the coral to invest more heavily in tissue repair. This study provides evidence that some corals are capable of withstanding the adverse effects of physical damage under elevated temperatures.

Acknowledgments The authors would like to thank the Australian Research Council for the funding provided.

References

- Ainsworth, T. D., K. Wasmund, K. Wasmund, L. Ukani, F. Seneca, D. Yellowlees, D. Miller & W. Leggat, 2011. Defining the tipping point. A complex cellular life/death balance in corals in response to stress. *Scientific Reports* 1: 160.
- Berkelmans, R. & B. L. Willis, 1999. Seasonal and local spatial variation in the upper thermal limits of corals in the central Great Barrier Reef. *Coral Reefs* 18: 219–228.
- Bourne, D., Y. Iida, S. Uthicke & C. Smith-Keune, 2008. Changes in coral-associated microbial communities during a bleaching event. *The ISME Journal* 2: 350–363.
- Burge, C. A., M. E. Mouchka, C. D. Harvell & S. Roberts, 2013. Immune response of the Caribbean sea fan, *Gorgonia ventalina*, exposed to an *Aplanochytrium* parasite as revealed by transcriptome sequencing. *Frontiers in Physiology* 4: 180.
- Cerenius, L., S. Kawabata, B. L. Lee, M. Nonaka & K. Söderhäll, 2010. Proteolytic cascades and their involvement in invertebrate immunity. *Trends in Biochemical Sciences* 35: 575–583.
- Cróquer, A., E. Villamizar & N. Noriega, 2002. Environmental factors affecting tissue regeneration of the reef—building coral *Montastraea annularis* (Faviidae) at Los Roques National Park, Venezuela. *Revista de Biología Tropical* 50: 1055–1065.
- Csaszar, N. B. M., F. O. Seneca & M. van Oppen, 2009. Variation in antioxidant gene expression in the scleractinian coral *Acropora millepora* under laboratory thermal stress. *Marine Ecology Progress Series* 392: 93–102.
- Denis, V., M. M. M. Guillaumme, M. Goutx, S. de Palmas, J. Debreuil, A. C. Baker, R. K. Boonstra & J. H. Bruggemann, 2013. Fast growth may impair regeneration capacity in the branching coral *Acropora muricata*. *PLoS One* 8: e72618.
- Edmunds, P. & H. Lenihan, 2010. Effect of sub-lethal damage to juvenile colonies of massive *Porites* spp. under contrasting regimes of temperature and water flow. *Marine Biology* 157: 887–897.
- Erlor, S., M. Popp & H. M. Lattorff, 2011. Dynamics of immune system gene expression upon bacterial challenge and wounding in a social insect (*Bombus terrestris*). *Plos One* 6: e18126.
- Fehlbaum, P., P. Bulet, L. Michaut, M. Lagueux, W. F. Broekaert, C. Hetru & J. A. Hoffmann, 1994. Insect immunity. Septic injury of *Drosophila* induces the synthesis of a potent antifungal peptide with sequence homology to plant antifungal peptides. *The Journal of Biological Chemistry* 269: 33159–33163.
- Fine, M., U. Oren & Y. Loya, 2002. Bleaching effect on regeneration and resource translocation in the coral *Oculina patagonica*. *Marine Ecology Progress Series* 234: 119–125.
- Franzenburg, S., S. Fraune, S. Kunzel, J. F. Baines, T. Domazet-Loso & T. C. G. Bosch, 2012. MyD88-deficient Hydra reveal an ancient function of TLR signaling in sensing bacterial colonizers. *Proceedings of the National Academy of Sciences of the United States of America* 109: 19374–19379.
- Franzenburg, S., J. Walter, S. Kunzel, J. Wang, J. F. Baines, T. C. G. Bosch & S. Fraune, 2013. Distinct antimicrobial peptide expression determines host species-specific bacterial associations. *Proceedings of the National Academy of Sciences of the United States of America* 110: E3730–E3738.
- Fraune, S., R. Augustin & T. C. G. Bosch, 2011. Embryo protection in contemporary immunology: Why bacteria matter. *Communicative & Integrative Biology* 4: 369–372.
- Fraune, S. & T. C. G. Bosch, 2007. Long-term maintenance of species-specific bacterial microbiota in the basal metazoan Hydra. *Proceedings of the National Academy of Sciences of the United States of America* 104: 13146–13151.

- Grasso, L., J. Maindonald, S. Rudd, D. C. Hayward, R. Saint, D. J. Miller & E. E. Ball, 2008. Microarray analysis identifies candidate genes for key roles in coral development. *BMC Genomics* 9: 540.
- Grasso, L. C., A. P. Negri, S. Foret, R. Saint, D. C. Hayward, E. E. Ball & D. J. Miller, 2011. The biology of coral metamorphosis: molecular responses of larvae to inducers of settlement and metamorphosis. *Developmental Biology* 353: 411–419.
- Hamada, M., E. Shoguchi, C. Shinzato, T. Kawashima, D. J. Miller & N. Satoh, 2013. The complex NOD-like receptor repertoire of the coral *Acropora digitifera* includes novel domain combinations. *Molecular Biology and Evolution* 30: 167–176.
- Henry, L.-A. & M. Hart, 2005. Regeneration from injury and resource allocation in sponges and corals—a review. *International Review of Hydrobiology* 90: 125–158.
- Hess, J., P. Angel & M. Schorpp-Kistner, 2004. AP-1 subunits: quarrel and harmony among siblings. *Journal of Cell Science* 117: 5965–5973.
- Hoegh-Guldberg, O., P. Mumby, A. Hooten, R. Steneck, P. Greenfield, E. Gomez, C. Harvell, P. Sale, A. Edwards & K. Cadeira, 2007. Coral reefs under rapid climate change and ocean acidification. *Science* 318: 1737–1742.
- Lenihan, H. S. & P. J. Edmunds, 2010. Response of *Pocillopora verrucosa* to corallivory varies with environmental conditions. *Marine Ecology Progress Series* 409: 51–63.
- Kimura, A., E. Sakaguchi & M. Nonaka, 2009. Multi-component complement system of Cnidaria: C3, Bf, and MASP genes expressed in the endodermal tissues of a sea anemone, *Nematostella vectensis*. *Immunobiology* 214: 165–178.
- Kramarsky-Winter, E. & Y. Loya, 2000. Tissue regeneration in the coral *Fungia granulosa*: the effect of extrinsic and intrinsic factors. *Marine Biology* 137: 867–873.
- Kvennefors, E. C. E., W. Leggat, O. Hoegh-Guldberg, B. M. Degnan & A. C. Barnes, 2008. An ancient and variable mannose-binding lectin from the coral *Acropora millepora* binds both pathogens and symbionts. *Developmental and Comparative Immunology* 32: 1582–1592.
- Kvennefors, E. C. E., W. Leggat, C. C. Kerr, T. D. Ainsworth, O. Hoegh-Guldberg & A. C. Barnes, 2010. Analysis of evolutionarily conserved innate immune components in coral links immunity and symbiosis. *Developmental and Comparative Immunology* 34: 1219–1229.
- Lamb, J. B. & B. L. Willis, 2011. Using coral disease prevalence to assess the effects of concentrating tourism activities on offshore reefs in a tropical marine park. *Conservation Biology* 25: 1044–1052.
- Lamb, J. B., J. D. True, S. Piromvaragorn & B. L. Willis, 2014. Scuba diving damage and intensity of tourist activities increases coral disease prevalence. *Biological Conservation* 178: 88–96.
- Leggat, W., F. Seneca, K. Wasmund, L. Ukani, D. Yellowlees & T. D. Ainsworth, 2011. Differential responses of the coral host and their algal symbiont to thermal stress. *PLoS One* 6: e26687.
- Lindsay, S. M., 2010. Frequency of injury and the ecology of regeneration in marine benthic invertebrates. *Integrative and Comparative Biology* 50: 479–493.
- Littman, R., B. L. Willis & D. G. Bourne, 2011. Metagenomic analysis of the coral holobiont during a natural bleaching event on the Great Barrier Reef. *Environmental Microbiology Reports* 3: 651–660.
- Mascarelli, P. E. & L. Bunkley-Williams, 1999. An experimental field evaluation of healing in damaged, unbleached and artificially bleached star coral, *Montastraea annularis*. *Bulletin of Marine Science* 65: 577–586.
- Meesters, E. H., M. Pauchli & R. P. M. Bak, 1997. Predicting regeneration of physical damage on a reef-building coral by regeneration capacity and lesion shape. *Marine Ecology Progress Series* 146: 91–99.
- Miller, D. J., G. Hemmrich, E. E. Ball, D. C. Hayward, K. Khalturin, N. Funayama, K. Agata & T. C. G. Bosch, 2007. The innate immune repertoire in Cnidaria—ancestral complexity and stochastic gene loss. *Genome Biology* 8: R59.
- Mouchka, M. E., I. Hewson & C. D. Harvell, 2010. Coral-associated bacterial assemblages: current knowledge and the potential for climate-driven impacts. *Integrative and Comparative Biology* 50: 662–674.
- Mydlarz, L. D., S. F. Holthouse, E. C. Peters & C. D. Harvell, 2008. Cellular responses in sea fan corals: granular amoebocytes react to pathogen and climate stressors. *Plos One* 3: e1811.
- Palmer, C. V., J. C. Bythell & B. L. Willis, 2011a. A comparative study of phenoloxidase activity in diseased and bleached colonies of the coral *Acropora millepora*. *Developmental and Comparative Immunology* 35: 1098–1101.
- Palmer, C. V., N. G. Traylor-Knowles, B. L. Willis & J. C. Bythell, 2011b. Corals use similar immune cells and wound-healing processes as those of higher organisms. *Plos One* 6(8): e23992.
- Palmer, C. V., J. C. Bythell & B. L. Willis, 2012. Enzyme activity demonstrates multiple pathways of innate immunity in Indo-Pacific anthozoans. *Proceedings of the Royal Society B* 279: 3879–3887.
- Pollock, F. J., J. B. Lamb, S. N. Field, S. F. Heron, B. Schaffelke, G. Shedrawi, D. G. Bourne & B. L. Willis, 2014. Sediment and turbidity associated with offshore dredging increase coral disease prevalence on nearby reefs. *PLoS ONE* 9: e102498.
- Pujol, N., S. Cypowyj, K. Ziegler, A. Millet, A. Astrain, A. Goncharov, Y. Jin, A. D. Chisholm & J. J. Ewbank, 2008. Distinct innate immune responses to infection and wounding in the *C. elegans* epidermis. *Current Biology* 18: 481–489.
- Renegar, D. A., P. L. Blackwelder & A. L. Moulding, (2008). Coral ultrastructural response to elevated pCO₂ and nutrients during tissue repair and regeneration. In *Proceedings of the 11th International Coral Reef Symposium*, Ft. Lauderdale, Florida, 7–11 July 2008. 2: 1320–1324.
- Ritchie, K. B., 2006. Regulation of microbial populations by coral surface mucus and mucus-associated bacteria. *Marine Ecology Progress Series* 322: 1–14.
- Rodriguez de la Vega, R. C., B. I. Garcia, C. D'Ambrosio, E. Diego-Garcia, A. Scaloni & L. D. Possani, 2004. Antimicrobial peptide induction in the haemolymph of the Mexican scorpion *Centruroides limpidus limpidus* in response

- to septic injury. *Cellular and Molecular Life Sciences* 61: 1507–1519.
- Ruiz-Moreno, D., B. L. Willis, A. C. Page, E. Weill, A. Croquer, B. Vargas-Angel, A. G. Jordan-Garza, E. Jordan-Dahlgren, L. Raymundo & C. D. Harvell, 2012. Global coral disease prevalence associated with sea temperature anomalies and local factors. *Diseases of Aquatic Organisms* 100: 249–261.
- Seneca, F. O., S. Foret, E. E. Ball, C. Smith-Keune, D. J. Miller & M. J. H. van Oppen, 2010. Patterns of gene expression in a scleractinian coral undergoing natural bleaching. *Marine Biotechnology* 12: 594–604.
- Shinzato, C., E. Shoguchi, T. Kawashima, M. Hamada, K. Hisata, M. Tanaka, M. Fujie, M. Fujiwara, R. Koyanagi, T. Ikuta, A. Fujiyama, D. J. Miller & N. Satoh, 2011. Using the *Acropora digitifera* genome to understand coral responses to environmental change. *Nature* 476: U320–U382.
- Siboni, N., D. Abrego, F. O. Seneca, C. A. Motti, N. Andreakis, J. Tebben, L. L. Blackall & T. Harder, 2012. Using bacterial extract along with differential gene expression in *Acropora millepora* larvae to decouple the processes of attachment and metamorphosis. *Plos One* 7(5): e37774.
- Sokolow, S., 2009. Effects of a changing climate on the dynamics of coral infectious disease: a review of the evidence. *Diseases of Aquatic Organisms* 87(1–2): 5–18.
- Souter, P., L. K. Bay, N. Andreakis, N. Csaszar, F. O. Seneca & M. J. H. van Oppen, 2011. A multilocus, temperature stress-related gene expression profile assay in *Acropora millepora*, a dominant reef-building coral. *Molecular Ecology Resources* 11: 328–334.
- Titlyanov, E. A. & T. V. Titlyanova, 2009. The dynamics of the restoration of mechanical damage to colonies of the scleractinian coral *Porites lutea* under conditions of competition with algal settlers for substratum. *Russian Journal of Marine Biology* 35: 230–235.
- Titlyanov, E. A., T. V. Titlyanova, I. M. Yakovleva, Y. Nakano & R. Bhagooli, 2005. Regeneration of artificial injuries on scleractinian corals and coral/algal competition for newly formed substrate. *Journal of Experimental Marine Biology and Ecology* 323: 27–42.
- Vandesompele, J., K. De Preter, F. Pattyn, B. Poppe, N. van Roy, A. de Paepe & F. Speleman, 2002. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biology* 3: Research0034.
- Vidal-Dupiol, J., O. Ladriere, D. Destoumieux-Garzon, P. E. Sautiere, A. L. Meistertzheim, E. Tambutte, S. Tambutte, D. Duval, L. Foure, M. Adjeroud & G. Mitta, 2011a. Innate immune responses of a scleractinian coral to vibriosis. *Journal of Biological Chemistry* 286: 22688–22698.
- Vidal-Dupiol, J., O. Ladriere, A. L. Meistertzheim, L. Foure, M. Adjeroud & G. Mitta, 2011b. Physiological responses of the scleractinian coral *Pocillopora damicornis* to bacterial stress from *Vibrio coralliilyticus*. *Journal of Experimental Biology* 214: 1533–1545.
- Vidal-Dupiol, J., N. M. Dheilly, R. Rondon, C. Grunau, C. Cosseau, K. M. Smith, M. Freitag, M. Adjeroud & G. Mitta, 2014. Thermal stress triggers broad *Pocillopora damicornis* transcriptomic remodeling, while *Vibrio coralliilyticus* infection induces a more targeted immunosuppression response. *PLoS One* 9(9): e107672.
- Willis, B. L., C. A. Page & E. A. Dinsdale, 2004. Coral disease on the Great Barrier Reef. In Rosenberg, E. & Y. Loya (eds), *Coral Health and Disease*. Springer, Berlin: 69–104.
- Witt, V., C. Wild, K. R. Anthony, G. Diaz-Pulido & S. Uthicke, 2011. Effects of ocean acidification on microbial community composition of, and oxygen fluxes through, biofilms from the Great Barrier Reef. *Environmental Microbiology* 13: 2976–2989.
- Wu, X. & G. Brewer, 2012. The regulation of mRNA stability in mammalian cells: 2.0. *Gene* 500: 10–21.
- Xu, Q., G. Wang, H. Yuan, Y. Chai & Z. Xiao, 2010. cDNA sequence and expression analysis of an antimicrobial peptide, theromacin, in the triangle-shell pearl mussel *Hyriopsis cumingii*. *Comparative Biochemistry and Physiology Part B* 157: 119–126.