

Impacts of fragment genotype, habitat, and size on outplanted elkhorn coral success under thermal stress

Rachel E. Pausch*, Dana E. Williams, Margaret W. Miller

*Corresponding author: rpausch@ucsc.edu

Marine Ecology Progress Series 592: 109–117 (2018)

Supplement. Additional data, including detailed bleaching scores of the genotype experiment, survivorship curves by site, and site coordinates




Fig. S1. Bleaching score distribution over time of genotype experiment fragments during September, October, and November 2015.

Table S1. The attached [.kml](#) file gives the location of *Acropora palmata* donor genotype sites (denoted with a g), as well as outplant sites.

Site name	Latitude	Longitude
Conch_Reef_1_g	24° 57'36.60"N	80° 27'25.02"W
Conch_Reef_2_g	24° 57'35.16"N	80° 27'21.96"W
Horseshoe_Reef_g	25° 8'23.76"N	80° 17'40.32"W
Snapper_Ledge_g	24° 58'53.46"N	80° 25'19.80"W
AAA_Reef	25° 10'6.32"N	80° 16'0.32"W
French_Reef	25° 2'3.06"N	80° 20'56.36"W
Horseshoe_Reef	25° 8'23.76"N	80° 17'40.32"W
Molasses_Reef	25° 0'40.85"N	80° 22'25.45"W
North_Dry_Rocks_Reef	25° 7'47.27"N	80° 17'37.68"W
Pickles_Reef	24° 59'10.21"N	80° 24'54.96"W




Fig. S2. Survival curves of the size and genotype experiments by site. Survivorship of the small and large fragments was not significantly different at any of the sites of the size experiment. Wilcoxon post-hoc pairwise comparisons are indicated by letters for the genotype experiment. Sites are labeled using two letter abbreviations given in methods.