



Suppl. Figure 1: Nursery structure at Goulding Cay where PVC plates with coral plugs with settled recruits were outplanted after having received different aquaculture regimes and culturing periods. The structure consisted of 4 vertically placed star pickets (1.5 m long) connected by 2 horizontally placed stainless-steel rods (2m long, diameter 1 cm).

Culture period 1:								
<i>1 September - 28 September</i>								
	temperature			salinity			pH	
ambient	27.9	±	0.7	33.9	±	0.2	8.14	± 0.03
ambient x feeding	27.9	±	0.8	33.9	±	0.2	8.13	± 0.04
high TA	27.9	±	0.8	34.1	±	0.2	8.03	± 0.03
high TA x feeding	27.9	±	0.9	34.1	±	0.2	8.03	± 0.04
Culture period 2:								
<i>29 September - 4 October (before hurricane)</i>								
	temperature			salinity			pH	
ambient	28.5	±	0.4	33.9	±	0.8	8.10	± 0.04
ambient x feeding	28.4	±	0.3	34.0	±	0.7	8.10	± 0.04
high TA	28.6	±	0.4	34.0	±	0.9	8.01	± 0.05
high TA x feeding	28.7	±	0.4	34.0	±	0.8	8.01	± 0.05
<i>4 - 14 October (during hurricane, in aquarium container)</i>								
	temperature			salinity			pH	
ambient	28.5	±	0.3	34.0	±	0.1	8.06	± 0.06
high TA	28.4	±	0.3	34.0	±	0.3	8.04	± 0.07
<i>15 October - 4 November (after hurricane)</i>								
	temperature			salinity			pH	
ambient	27.6	±	0.6	33.0	±	0.4	8.08	± 0.03
ambient x feeding	27.5	±	0.6	33.0	±	0.4	8.07	± 0.04
high TA	27.3	±	0.7	33.0	±	0.5	7.99	± 0.03
high TA x feeding	27.5	±	0.6	33.0	±	0.5	7.98	± 0.03
Culture period 3:								
<i>5 November - 9 December</i>								
	temperature			salinity			pH	
ambient	27.5	±	0.7	34.2	±	0.2	8.08	± 0.03
ambient x feeding	27.4	±	0.8	34.2	±	0.3	8.07	± 0.05
high TA	27.4	±	0.7	34.4	±	0.3	7.98	± 0.02
high TA x feeding	27.5	±	0.7	34.3	±	0.3	7.98	± 0.03

Suppl. Table 1: Temperature (°C), salinity (ppt) and pH during the experiment. Culture period 1 is the period from settlement till the first outplacement, culture period 2 is the period between the first and second outplacement, and culture period 3 is the period between the second and last outplacement.

Survival during aquaculture period

Fit: glm (cbind(p, q) ~ treatment * period, family=binomial, data=data14wks)

Analysis of Deviance Table (Type III tests)

Response: cbind(p,q)

	LR Chisq	Df	Pr(>Chisq)	
treatment	9.4142	3	0.024262	*
period	16.6004	2	0.000249	***
treatment:period	8.7499	6	0.18814	NS

dispersion parameter for binomial family taken as 1 (0.4909627, Chisquare test: p=0.48)

Multiple Comparisons of Means: Tukey Contrasts

Fit: glm(formula = cbind(p, q) ~ treatment, family = binomial, data = data14wks)

Linear Hypotheses:

	Estimate	Std. Error	z value	Pr(> z)	
ambientxfeeding - ambient == 0	0.0461	0.04464	1.033	1	NS
highTA - ambient == 0	0.13987	0.04536	3.084	0.0123	*
highTAxfeeding - ambient == 0	0.08447	0.04495	1.879	0.3613	NS
highTA - ambientxfeeding == 0	0.09377	0.04796	1.955	0.3033	NS
highTAxfeeding - ambientxfeeding == 0	0.03837	0.04757	0.807	1	NS
highTAxfeeding - highTA == 0	-0.0554	0.04825	-1.148	1	NS

Fit: glm(formula = cbind(p, q) ~ period, family = binomial, data = data14wks)

Linear Hypotheses:

	Estimate	Std. Error	z value	Pr(> z)	
0-4wks - 0-14wks == 0	-0.49978	0.04348	-11.494	< 2e-16	***
0-9wks - 0-14wks == 0	-0.39524	0.0504	-7.843	1.33E-14	***
0-9wks - 0-4wks == 0	0.10454	0.03986	2.623	0.0262	*

Suppl. Table 1: Results from a factorial logistic regression analysis testing the effect of aquaculture conditions ("treatment") and aquaculture duration("period") on the survivorship of coral plugs with minimum one recruit throughout the aquaculture period. The test is followed by Tukey contrasts testing for significance between treatment levels. Level of significance is indicated using asterisks: *** $p < 0.0001$, ** $p < 0.001$, * $p < 0.05$, NS not significant.

Average number of polyps per recruit during aquaculture

Kruskal-Wallis rank sum test

polyps~time

Kruskal-Wallis chi-squared = 97.502, df = 5, p-value < 2.2e-16

polyps~TA

Kruskal-Wallis chi-squared = 0.53462, df = 1, p-value = 0.4647

polyps~feeding

Kruskal-Wallis chi-squared = 14.15, df = 1, p-value = 0.0001688

Dunn (1964) Kruskal-Wallis multiple comparison

Comparison	Z	P.unadj	P.adj	
time1 - time2	-6.8646055	6.7E-12	2.0E-11	***
time1 - time3	-7.3690572	1.7E-13	6.4E-13	***
time2 - time3	-0.5390453	0.59	0.88	NS
time1 - time4	-7.5641209	3.9E-14	2.0E-13	***
time2 - time4	-0.7713569	0.44	0.83	NS
time3 - time4	-0.2340721	0.81	0.87	NS
time1 - time5	-7.7122083	1.2E-14	9.3E-14	***
time2 - time5	-0.958013	0.34	0.72	NS
time3 - time5	-0.4228379	0.67	0.84	NS
time4 - time5	-0.1890746	0.85	0.85	NS
time1 - time6	-7.9719499	1.6E-15	2.3E-14	***
time2 - time6	-1.298498	0.19	0.49	NS
time3 - time6	-0.7680452	0.44	0.74	NS
time4 - time6	-0.5352569	0.59	0.81	NS
time5 - time6	-0.3465223	0.73	0.84	NS

p-values adjusted with the Benjamini-Hochberg method

Suppl. Table 2: Results from Kruskal-Wallis rank sum tests, testing the effect of aquaculture duration ("time"), bicarbonate addition ("TA") and feeding ("feeding") on recruit size as number of polyps per recruit. Level of significance is indicated using asterisks: *** $p < 0.0001$, ** $p < 0.001$, * $p < 0.05$, NS not significant.

Average living tissue area during aquaculture

Kruskal-Wallis rank sum test

area~time

Kruskal-Wallis chi-squared = 16.144, df = 5, p-value = 0.006445

area~TA

Kruskal-Wallis chi-squared = 2.0746, df = 1, p-value = 0.1498

area~feeding

Kruskal-Wallis chi-squared = 0.10159, df = 1, p-value = 0.7499

Dunn (1964) Kruskal-Wallis multiple comparison

Comparison	Z	P.unadj	P.adj	
time1 - time2	-2.09925394	0.04	0.11	NS
time1 - time3	-2.56783524	0.01	0.04	*
time2 - time3	-0.47753667	0.63	0.79	NS
time1 - time4	-3.30501845	0.001	0.014	*
time2 - time4	-1.23474933	0.22	0.54	NS
time3 - time4	-0.75862820	0.45	0.75	NS
time1 - time5	-3.10980532	0.002	0.009	**
time2 - time5	-1.04052191	0.30	0.56	NS
time3 - time5	-0.56541359	0.57	0.78	NS
time4 - time5	0.19110292	0.85	0.98	NS
time1 - time6	-3.21759028	0.001	0.010	*
time2 - time6	-1.17295045	0.24	0.52	NS
time3 - time6	-0.70291073	0.48	0.72	NS
time4 - time6	0.04615273	0.96	0.96	NS
time5 - time6	-0.14271499	0.89	0.95	NS

p-values adjusted with the Benjamini-Hochberg method

Suppl. Table 3: Results from Kruskal-Wallis rank sum tests, testing the effect of aquaculture duration ("time"), bicarbonate addition ("TA") and feeding ("feeding") on recruit size as living tissue area per recruit. Level of significance is indicated using asterisks: *** $p < 0.0001$, ** $p < 0.001$, * $p < 0.05$, NS not significant.

Survival after outplacement after 44 weeks

Fit: glm (cbind(p, q) ~ periodxtreatment, family=binomial, data=data44wks)

Analysis of Deviance Table (Type III tests)

Analysis of Deviance Table (Type III tests)

Response: cbind(p, q)

	IR Chisq	Df	Pr(>Chisq)	
periodxtreatment	128.33	8	<2.2e-16	***

dispersion parameter for binomial family taken as 1 (0.8681502, Chisquare test: p=0.72)

Multiple Comparisons of Means: User-defined Contrasts

Fit: glm(formula = cbind(p, q) ~ periodxtreatment, family = binomial, data = data44wks)

Linear Hypotheses:

	Estimate	Std. Error	z value	Pr(> z)	
t=0 vs. t=1 ambient == 0	-0.5306	0.2386	-2.224	0.47	NS
t=0 vs. t=2 ambient == 0	-0.9854	0.2761	-3.569	0.01	**
t=1 vs t=2 ambient == 0	-0.4548	0.2999	-1.516	1	NS
t=1 vs. t=2 high TAxfeeding == 0	-0.5572	0.175	-3.184	0.03	*
t=1 vs. t=2 ambientxfeeding == 0	-1.0736	0.2401	-4.471	0.0001	***
t=1 vs. t=2 high TA == 0	-1.4214	0.3044	-4.67	5.4E-05	***
t=1 ambient vs high TAxfeeding == 0	1.2335	0.2168	5.689	2.3E-07	***
t=1 ambient vs ambientxfeeding == 0	0.8873	0.2258	3.929	0.002	**
t=1 ambient vs high TA == 0	0.6592	0.2335	2.823	0.09	NS
t=1 high TAxfeeding vs ambientxfeeding ==	-0.3463	0.1654	-2.093	0.65	NS
t=1 high TAxfeeding vs highhigh TA == 0	-0.5743	0.1758	-3.266	0.02	*
t=1 ambientxfeeding vs high TA == 0	-0.228	0.1868	-1.221	1	NS
t=2 ambient vs high TAxfeeding == 0	1.1311	0.2712	4.171	0.001	***
t=2 ambient vs ambientxfeeding == 0	0.2685	0.3108	0.864	1	NS
t=2 ambient vs high TA == 0	-0.3074	0.3578	-0.859	1	NS
t=2 high TAxfeeding vs ambientxfeeding ==	-0.8626	0.2468	-3.495	0.009	**
t=2 high TAxfeeding vs high TA == 0	-1.4385	0.3039	-4.734	4.0E-05	***
t=2 ambientxfeeding vs high TA == 0	-0.5759	0.3397	-1.695	1	NS

Suppl. Table 5: Results from a factorial logistic regression analysis testing the effect of aquaculture condition x aquaculture duration ("periodxtreatment") on the survivorship of coral plugs with minimum one recruit measured at 44 weeks after settlement. The test is followed by multiple comparisons using user-defined contrasts. Level of significance is indicated using asterisks: *** p<0.0001, ** p<0.001, * p<0.05, NS not significant.

Survival after outplacement at 27 weeks

Fit: glm (cbind(p, q) ~ periodxtreatment, family=quasibinomial,data=data27wks)

Analysis of Deviance Table (Type III tests)

Response: cbind(p,q)

Error estimate based on deviance

	SS	Df	F	Pr(>F)	
periodxtreatment	101.11	8	5.6776	5.35E-05	***
Residuals	100.17	45			

dispersion parameter for binomial family was 2.192082, Chisquare test: p<0.0001)

Multiple Comparisons of Means: User-defined Contrasts

Fit: glm(formula = rv27wks ~ periodxtreatment, family = quasibinomial, data = data27wks)

Linear Hypotheses:

	Estimate	Std. Error	z value	Pr(> z)	
t=0 vs. t=1 ambient == 0	-0.01316	0.19376	-0.068	1	NS
t=0 vs. t=2 ambient == 0	-0.30106	0.20599	-1.462	1	NS
t=1 vs t=2 ambient == 0	-0.28791	0.20645	-1.395	1	NS
t=1 vs. t=2 high TAxfedding == 0	-0.5974	0.19557	-3.055	0.04	*
t=1 vs. t=2 ambientxfedding == 0	-0.74059	0.21527	-3.44	0.01	*
t=1 vs. t=2 high TA == 0	-0.85605	0.23822	-3.593	0.006	**
t=1 ambient vs high TAxfedding == 0	0.32505	0.18359	1.771	1	NS
t=1 ambient vs ambientxfedding == 0	0.17369	0.18811	0.923	1	NS
t=1 ambient vs high TA == 0	0.01316	0.19376	0.068	1	NS
t=1 high TAxfedding vs ambientxfedding ==	-0.15136	0.17708	-0.855	1	NS
t=1 high TAxfedding vs highhigh TA == 0	-0.3119	0.18307	-1.704	1	NS
t=1 ambientxfedding vs high TA == 0	-0.16054	0.18761	-0.856	1	NS
t=2 ambient vs high TAxfedding == 0	0.01556	0.21718	0.072	1	NS
t=2 ambient vs ambientxfedding == 0	-0.27899	0.23147	-1.205	1	NS
t=2 ambient vs high TA == 0	-0.55499	0.24866	-2.232	0.46	NS
t=2 high TAxfedding vs ambientxfedding ==	-0.29455	0.23072	-1.277	1	NS
t=2 high TAxfedding vs high TA == 0	-0.57054	0.24796	-2.301	0.39	NS
t=2 ambientxfedding vs high TA == 0	-0.276	0.26057	-1.059	1	NS

Suppl. Table 4: Results from a factorial logistic regression analysis testing the effect of aquaculture condition x aquaculture duration ("periodxtreatment") on the survivorship of coral plugs with minimum one recruit measured at 27 weeks after settlement. The test is followed by multiple comparisons using user-defined constraints. Level of significance is indicated using asterisks: *** p<0.0001, ** p<0.001, * p<0.05, NS not significant.

Recruit size (number of polyps per recruit) at 44 weeks after settlement

Anova Table (Type III tests)

Response: sqrt(polypperrecruit)					
	Sum Sq	Df	F value	Pr(>F)	
(Intercept)	16.3082	1	222.088	3.00E-14	***
periodxtreat	7.1006	8	12.087	5.05E-07	***
Residuals	1.9092	26			

Multiple Comparisons of Means: User-defined Contrasts

Fit: aov(formula = sqrt(data\$polypperrecruit) ~ periodxtreat)

Linear Hypotheses:

	Estimate	Std. Error	z value	Pr(> z)	
t=0 vs. t=1 ambient == 0	-0.008326	0.191613	-0.043	1	NS
t=0 vs. t=2 ambient == 0	0.299555	0.206966	1.45	1	NS
t=1 vs t=2 ambient == 0	0.307882	0.206966	1.49	1	NS
t=1 vs. t=2 high TAxfedding == 0	-0.201922	0.191613	-1.054	1	NS
t=1 vs. t=2 ambientxfedding == 0	0.280923	0.191613	1.466	1	NS
t=1 vs. t=2 high TA == 0	0.696635	0.191613	3.64	0.02	*
t=1 ambient vs high TAxfedding == 0	-0.760073	0.191613	-3.97	0.009	**
t=1 ambient vs ambientxfedding == 0	-0.511917	0.191613	-2.7E+00	0.23	NS
t=1 ambient vs high TA == 0	-0.183937	0.191613	-0.9600	1	NS
t=1 high TAxfedding vs ambientxfedding ==	0.248155	0.191613	1.295	1	NS
t=1 high TAxfedding vs highhigh TA == 0	0.576135	0.191613	3.007	0.10	NS
t=1 ambientxfedding vs high TA == 0	0.32798	0.191613	1.7E+00	1	NS
t=2 ambient vs high TAxfedding == 0	-1.269877	0.206966	-6.136	3.1E-05	***
t=2 ambient vs ambientxfedding == 0	-0.538876	0.206966	-2.604	0.27	NS
t=2 ambient vs high TA == 0	0.204816	0.206966	9.9E-01	1	NS
t=2 high TAxfedding vs ambientxfedding ==	0.731	0.191613	3.8150	0.01	*
t=2 high TAxfedding vs highhigh TA == 0	1.474693	0.191613	7.696	6.5E-07	***
t=2 ambientxfedding vs high TA == 0	0.743692	0.191613	3.9E+00	0.01	*

Suppl. Table 7: Results from ANOVA and multiple comparison tests testing the effect aquaculture duration x aquaculture condition ("periodxtreat") on the number of polyps per recruit at 44 weeks after settlement. Aquaculture periods are labeled as t=0 (4 weeks aquaculture), t=1 (9 weeks aquaculture) and t=2 (14 weeks aquaculture). Level of significance is indicated using asterisks: *** $p < 0.0001$, ** $p < 0.001$, * $p < 0.05$, NS not significant.

Recruit size (number of polyps per recruit) at 27 weeks after settlement

Anova Table (Type III tests)

Response: sqrt(polyp_per_recruit)

	Sum Sq	Df	F value	Pr(>F)	
(Intercept)	17.7536	1	736.75	< 2.2e-16	***
periodxtreat	5.277	8	27.373	7.38E-15	***
Residuals	1.0844	45			

Multiple Comparisons of Means: User-defined Contrasts

Fit: aov(formula = sqrt(data\$polyp_per_recruit) ~ periodxtreat)

Linear Hypotheses:

	Estimate	Std. Error	z value	Pr(> z)	
t=0 vs. t=1 ambient == 0	0.054325	0.089624	0.606	1	NS
t=0 vs. t=2 ambient == 0	0.254481	0.089624	2.839	0.12	NS
t=1 vs t=2 ambient == 0	0.200156	0.089624	2.233	0.55	NS
t=1 vs. t=2 high TAxfed == 0	-0.163189	0.089624	-1.821	1	NS
t=1 vs. t=2 ambientxfed == 0	0.173227	0.089624	1.933	1	NS
t=1 vs. t=2 high TA == 0	0.263911	0.089624	2.945	0.09	NS
t=1 ambient vs high TAxfed == 0	-0.580104	0.089624	-6.473	1.1E-06	***
t=1 ambient vs ambientxfed == 0	-0.413822	0.089624	-4.617	0.0006	***
t=1 ambient vs high TA == 0	-0.069988	0.089624	-0.781	1	NS
t=1 high TAxfed vs ambientxfed == 0	0.166282	0.089624	1.855	1	NS
t=1 high TAxfed vs highhigh TA == 0	0.510116	0.089624	5.692	1.6E-05	***
t=1 ambientxfed vs high TA == 0	0.343834	0.089624	3.836	0.007	**
t=2 ambient vs high TAxfed == 0	-0.943449	0.089624	-10.527	1.8E-12	***
t=2 ambient vs ambientxfed == 0	-0.44075	0.089624	-4.918	0.0002	***
t=2 ambient vs high TA == 0	-0.006232	0.089624	-0.07	1	NS
t=2 high TAxfed vs ambientxfed == 0	0.502699	0.089624	5.609	2.1E-05	***
t=2 high TAxfed vs highhigh TA == 0	0.937216	0.089624	10.457	2.3E-12	***
t=2 ambientxfed vs high TA == 0	0.434518	0.089624	4.848	0.0003	***

Suppl. Table 6: Results from ANOVA and multiple comparison tests testing the effect aquaculture duration x aquaculture condition ("periodxtreat") on the number of polyps per recruit at 27 weeks after settlement. Aquaculture periods are labeled as t=0 (4 weeks aquaculture), t=1 (9 weeks aquaculture) and t=2 (14 weeks aquaculture). Level of significance is indicated using asterisks: ***p<0.0001, **p<0.001, *p<0.05, NS not significant.

Recruit size (living tissue area) at 44 weeks after settlement

Anova Table (Type III tests)					
Response: area ^{-0.1}					
	Sum Sq	Df	F value	Pr(>F)	
(Intercept)	5.1162	1	837.4337	< 2.2e-16	***
periodxtreat	0.1733	8	3.5461	0.004	**
Residuals	0.2199	36			

Multiple Comparisons of Means: User-defined Contrasts

Fit: aov(formula = area^{-0.1} ~ factors)

Linear Hypotheses:

	Estimate	Std. Error	t value	Pr(> t)	
t=0 vs. t=1 ambient == 0	0.006018	0.045127	0.133	1	NS
t=0 vs. t=2 ambient == 0	-0.07164	0.04733	-1.514	1	NS
t=1 vs t=2 ambient == 0	-0.07766	0.04733	-1.641	1	NS
t=1 vs. t=2 high TAxfedding == 0	-0.05828	0.052433	-1.112	1	NS
t=1 vs. t=2 ambientxfedding == 0	-0.03422	0.055269	-0.619	1	NS
t=1 vs. t=2 high TA == 0	0.074625	0.053937	1.384	1	NS
t=1 ambient vs high TAxfedding == 0	0.07248	0.04733	1.531	1	NS
t=1 ambient vs ambientxfedding == 0	0.077427	0.045127	1.716	1	NS
t=1 ambient vs high TA == 0	-0.13197	0.055269	-2.388	0.40	NS
t=1 high TAxfedding vs ambientxfedding ==	0.004946	0.04733	0.105	1	NS
t=1 high TAxfedding vs high TA == 0	-0.20445	0.057082	-3.582	0.02	*
t=1 ambientxfedding vs high TA == 0	-0.20939	0.055269	-3.789	0.01	*
t=2 ambient vs high TAxfedding == 0	0.091856	0.052433	1.752	1	NS
t=2 ambient vs ambientxfedding == 0	0.120867	0.057082	2.117	0.74	NS
t=2 ambient vs high TA == 0	0.020315	0.045767	0.444	1	NS
t=2 high TAxfedding vs ambientxfedding ==	0.029011	0.059698	0.486	1	NS
t=2 high TAxfedding vs high TA == 0	-0.07154	0.048991	-1.46	1	NS
t=2 ambientxfedding vs high TA == 0	-0.10055	0.053937	-1.864	1	NS

Suppl. Table 9: Results from ANOVA and multiple comparison tests testing the effect aquaculture duration x aquaculture condition ("periodxtreat") on recruit size as living tissue area at 44 weeks after settlement. Aquaculture periods are labeled as t=0 (4 weeks aquaculture), t=1 (9 weeks aquaculture) and t=2 (14 weeks aquaculture). Level of significance is indicated using asterisks: ***p<0.0001, **p<0.001, *p<0.05, NS not significant.

Recruit size (living tissue area) at 27 weeks after settlement

Anova Table (Type III tests)

Response: area^{-0.1}

	Sum Sq	Df	F value	Pr(>F)	
(Intercept)	6.4045	1	1908.661	< 2.2e-16	***
periodxtreat	0.1586	8	5.9094	4.77E-06	***
Residuals	0.2953	88			

Multiple Comparisons of Means: User-defined Contrasts

Fit: aov(formula = area^{-0.1} ~ periodxtreatment)

Linear Hypotheses:

	Estimate	Std. Error	t value	Pr(> t)	
t=0 vs. t=1 ambient == 0	-0.05653	0.028007	-2.019	0.84	NS
t=0 vs. t=2 ambient == 0	-0.04555	0.02625	-1.735	1	NS
t=1 vs t=2 ambient == 0	0.010982	0.022688	0.484	1	NS
t=1 vs. t=2 high TAxfedding == 0	-0.04139	0.025673	-1.612	1	NS
t=1 vs. t=2 ambientxfedding == 0	0.006869	0.027477	0.25	1	NS
t=1 vs. t=2 high TA == 0	-0.03079	0.024365	-1.264	1	NS
t=1 ambient vs high TAxfedding == 0	0.125493	0.026916	4.662	0.0002	***
t=1 ambient vs ambientxfedding == 0	0.076581	0.026916	2.845	0.10	NS
t=1 ambient vs high TA == 0	0.035833	0.02531	1.416	1	NS
t=1 high TAxfedding vs ambientxfedding == 0	-0.04891	0.028963	-1.689	1	NS
t=1 high TAxfedding vs high TA == 0	-0.08966	0.027477	-3.263	0.03	*
t=1 ambientxfedding vs high TA == 0	-0.04075	0.027477	-1.483	1	NS
t=2 ambient vs high TAxfedding == 0	0.073121	0.021199	3.449	0.02	*
t=2 ambient vs ambientxfedding == 0	0.072468	0.023351	3.103	0.046	*
t=2 ambient vs high TA == 0	-0.00594	0.021629	-0.275	1	NS
t=2 high TAxfedding vs ambientxfedding == 0	-0.00065	0.023984	-0.027	1	NS
t=2 high TAxfedding vs high TA == 0	-0.07906	0.022311	-3.544	0.01	*
t=2 ambientxfedding vs high TA == 0	-0.07841	0.024365	-3.218	0.03	*

Suppl. Table 8: Results from ANOVA and multiple comparison tests testing the effect aquaculture duration x aquaculture condition ("periodxtreat") on recruit size as living tissue area at 27 weeks after settlement. Aquaculture periods are labeled as t=0 (4 weeks aquaculture), t=1 (9 weeks aquaculture) and t=2 (14 weeks aquaculture). Level of significance is indicated using asterisks: *** $p < 0.0001$, ** $p < 0.001$, * $p < 0.05$, NS not significant.

Survival after outplacement between 27 and 44 weeks

Fit: glm (cbind(p, q) ~ periodxtreatment, family=binomial,data=data2744wks)

Analysis of Deviance Table (Type III tests)

Analysis of Deviance Table (Type III tests)

Response: cbind(p, q)

	LR Chisq	Df	Pr(>Chisq)	
periodxtreatment	35.656	8	2.03E-05	***

dispersion parameter for binomial family taken as 1 (0.861526, Chisquare test: p=0.67)

Multiple Comparisons of Means: User-defined Contrasts

Fit: glm(formula = rv2744wks ~ periodxtreatment, family = binomial, data = data2744wks)

Linear Hypotheses:

	Estimate	Std. Error	z value	Pr(> z)	
t=0 vs. t=1 ambient == 0	-0.49062	0.240549	-2.04	0.75	NS
t=0 vs. t=2 ambient == 0	-0.59262	0.272527	-2.175	0.53	NS
t=1 vs t=2 ambient == 0	-0.102	0.292244	-0.349	1	NS
t=1 vs. t=2 high TAxfedding == 0	0.009662	0.187558	0.052	1	NS
t=1 vs. t=2 ambientxfedding == 0	-0.29375	0.24442	-1.202	1	NS
t=1 vs. t=2 high TA == 0	-0.67084	0.325905	-2.058	0.71	NS
t=1 ambient vs high TAxfedding == 0	0.89994	0.216808	4.151	0.0006	***
t=1 ambient vs ambientxfedding == 0	0.700252	0.226026	3.098	0.04	*
t=1 ambient vs high TA == 0	0.615362	0.235042	2.618	0.16	NS
t=1 high TAxfedding vs ambientxfedding ==	-0.19969	0.17057	-1.171	1	NS
t=1 high TAxfedding vs highhigh TA == 0	-0.28458	0.18235	-1.561	1	NS
t=1 ambientxfedding vs high TA == 0	-0.08489	0.193218	-0.439	1	NS
t=2 ambient vs high TAxfedding == 0	1.011601	0.271254	3.729	0.003	**
t=2 ambient vs ambientxfedding == 0	0.508497	0.306692	1.658	1	NS
t=2 ambient vs high TA == 0	0.04652	0.369291	0.126	1	NS
t=2 high TAxfedding vs ambientxfedding ==	-0.5031	0.256564	-1.961	0.90	NS
t=2 high TAxfedding vs high TA == 0	-0.96508	0.328847	-2.935	0.06	NS
t=2 ambientxfedding vs high TA == 0	-0.46198	0.358639	-1.288	1	NS

Suppl. Table 10: Results from a factorial logistic regression analysis testing the effect of aquaculture condition x aquaculture duration ("periodxtreatment") on the survivorship of coral plugs with minimum one recruit measured at 44 weeks after settlement. The test is followed by multiple comparisons using user-defined contrasts. Aquaculture periods are labeled as t=0 (4 weeks aquaculture), t=1 (9 weeks aquaculture) and t=2 (14 weeks aquaculture). Level of significance is indicated using asterisks: *** p<0.0001, ** p<0.001, * p<0.05, NS not significant.

The effect of recruit size at 27 weeks on being dead or alive at 44 weeks

Anova Table (Type III tests)

Response: sqrt(1/polypperrecruit)					
	Sum Sq	Df	F value	Pr(>F)	
(Intercept)	1.17784	1	850.3422	< 2.2e-16	***
periodxtreatment	0.30882	8	27.8692	< 2.2e-16	***
deadALIVE	0.01316	1	9.4977	0.003261	**
periodxtreatment:deadALIVE	0.01869	8	1.6867	0.123406	NS
Residuals	0.07341	53			

Suppl. Table 11: Results from an ANOVA testing the effect of aquaculture duration x aquaculture condition ("periodxtreatment"), being dead or alive at 44 weeks ("deadALIVE") and their interaction on recruit size (as number of polyps per recruit) at 27 weeks. Level of significance is indicated using asterisks: *** $p < 0.0001$, ** $p < 0.001$, * $p < 0.05$, NS not significant.