



Local Polynomial Regression

Statistical Machine Learning - individual project


Leonardo Stincone

Università degli Studi di Trieste

18th July 2019



Problem statement: LIDAR dataset



LIDAR = Light Detection And Ranging


- it is a surveying method that measures distance to a target by illuminating the target with laser light and measuring the reflected light with a sensor
- x : distance travelled before the light is reflected back to its source
- y : logarithm of the ratio of received light from two laser sources

The objective is to estimate:

$$f(x) = E[Y | X = x]$$



Problem statement: LIDAR dataset



LIDAR = Light Detection And Ranging

- it is a surveying method that measures distance to a target by illuminating the target with laser light and measuring the reflected light with a sensor
- x : distance travelled before the light is reflected back to its source
- y : logarithm of the ratio of received light from two laser sources


The objective is to estimate:

$$f(x) = E[Y \mid X = x]$$




What does local means?

If we had enough point with $x = x_0$




What does local means?

If we had enough point with $x = x_0$




We can consider points "close" to x_0




k nearest neighbors

$$\hat{f}(x) = \frac{1}{k} \sum_{i=1}^n y_i I_{N_k(x)}(x_i)$$

Neighborhood of radius h

$$\hat{f}(x) = \frac{\sum_{i=1}^n y_i I_{[0,h]}(|x - x_i|)}{\sum_{i=1}^n I_{[0,h]}(|x - x_i|)}$$



$$\hat{f}(x) = \sum_{i=1}^n \ell_i(x) y_i$$

with:

$$\ell_i(x) = \frac{K\left(\frac{x-x_i}{h}\right)}{\sum_{j=1}^n K\left(\frac{x-x_j}{h}\right)}$$


where $K(\cdot)$ is a kernel function that satisfies:

- $K(x) \geq 0$
- $\int K(x) dx = 1$
- $\int x K(x) dx = 0$
- $\int x^2 K(x) dx > 0$




Some proposed kernels


Uniform
 $\frac{1}{2}I_{[-1,1]}(u)$




Triangle
 $(1 - |u|)I_{[-1,1]}(u)$




Triweight
 $\frac{35}{32}(1 - u^2)^3I_{[-1,1]}(u)$




Quartic
 $\frac{15}{16}(1 - u^2)^2I_{[-1,1]}(u)$




Cosine
 $\frac{\pi}{4} \cos\left(\frac{\pi}{2}u\right)I_{[-1,1]}(u)$




Epanechnikov
 $\frac{3}{4}(1 - u^2)I_{[-1,1]}(u)$




Gaussian
 $\frac{1}{\sqrt{2\pi}}e^{-u^2/2}$




Tricube
 $\frac{70}{81}(1 - |u|^3)^3I_{[-1,1]}(u)$



Design bias, boundary bias and concavity bias



Design bias: what's happening?



Locally the Nadaraya-Watson estimator is a Weighted Least Square Estimator:

$$\hat{f}_{NW}(x_0) = \underset{a}{\operatorname{argmin}} \sum_{i=1}^n K\left(\frac{x_i - x_0}{h}\right) (y_i - a)^2$$

Idea: instead of approximating $f(x_0)$ with a constant value a , we could approximate it with a polynomial $p_{x_0}(u, \mathbf{a})$.

Taylor polynomial approximation:

$$p_{x_0}(u; \mathbf{a}) = a_0 + a_1(u - x_0) + \frac{a_2}{2!}(u - x_0)^2 + \dots + \frac{a_d}{d!}(u - x_0)^d$$



Locally the Nadaraya-Watson estimator is a Weighted Least Square Estimator:

$$\hat{f}_{NW}(x_0) = \underset{a}{\operatorname{argmin}} \sum_{i=1}^n K\left(\frac{x_i - x_0}{h}\right) (y_i - a)^2$$

Idea: instead of approximating $f(x_0)$ with a constant value a , we could approximate it with a polynomial $p_{x_0}(u, \mathbf{a})$.

Taylor polynomial approximation:

$$p_{x_0}(u; \mathbf{a}) = a_0 + a_1(u - x_0) + \frac{a_2}{2!}(u - x_0)^2 + \dots + \frac{a_d}{d!}(u - x_0)^d$$



Local polynomial regression: estimation

We can estimate the coefficients of $p_{x_0}(u; \mathbf{a})$ as:

$$\hat{\mathbf{a}}(x_0) = \underset{\mathbf{a}}{\operatorname{argmin}} \sum_{i=1}^n K\left(\frac{x_i - x_0}{h}\right) (y_i - p_{x_0}(x_i; \mathbf{a}))^2$$


Thus, we can define the estimator for $f(x)$ in x_0 just computing $p_{x_0}(u; \hat{\mathbf{a}})$ in x_0 :

$$\hat{f}(x_0) = p_{x_0}(x_0; \hat{\mathbf{a}})$$


Then we can repeat the process for each value of x in a grid and obtain $\hat{f}(x)$.




Local polynomial regression: example




Local polynomial regression: example




Local polynomial regression: example




Local polynomial regression: example



Design bias, boundary bias and concavity bias




Local Polynomial Regression on LIDAR dataset




Local polynomial regression: linear vs quadratic

Linear loess




Quadratic loess



Fan, Gijbels

Local Polynomial Modelling and its Applications

Springer (1996)



Ruppert, Wand, Carroll

Semiparametric Regression

Cambridge University Press (2003)

